
Angioinvasive fungal infections impacting the skin



Background, epidemiology, and clinical presentation

Bridget E. Shields, MD,^a Misha Rosenbach, MD,^b Zoe Brown-Joel, BS,^c Anthony P. Berger, MD,^d
Bradley A. Ford, MD, PhD,^e and Karolyn A. Wanat, MD^f
Madison, Wisconsin; Philadelphia, Pennsylvania; and Iowa City, Iowa

Learning objectives

After completing this learning activity, participants should be able to identify and describe the epidemiology, initial presentation and subsequent clinical manifestations of angioinvasive fungal infections; characterize at-risk populations for angioinvasive fungal infection; and classify frequently encountered opportunistic organisms in dermatology and compare their distinguishing cutaneous features.

Disclosures

Editors

The editors involved with this CME activity and all content validation/peer reviewers of the journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

Authors

The authors involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

Planners

The planners involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s). The editorial and education staff involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

Angioinvasive fungal infections cause significant morbidity and mortality because of their propensity to invade blood vessel walls, resulting in catastrophic tissue ischemia, infarct, and necrosis. While occasionally seen in immunocompetent hosts, opportunistic fungi are emerging in immunosuppressed hosts, including patients with hematologic malignancy, AIDS, organ transplant, and poorly controlled diabetes mellitus. The widespread use of antifungal prophylaxis has led to an “arms race” of emerging fungal resistance patterns. As the at-risk population expands and new antifungal resistance patterns develop, it is critical for dermatologists to understand and recognize angioinvasive fungal pathogens, because they are often the first to encounter the cutaneous manifestations of these diseases. Rapid clinical recognition, histopathologic, and culture confirmation can help render a timely, accurate diagnosis to ensure immediate medical and surgical intervention. Superficial dermatophyte infections and deep fungal infections, such as blastomycosis and histoplasmosis, have been well characterized within the dermatologic literature, and therefore this article will focus on the severe infections acquired by angioinvasive fungal species, including an update on new and emerging pathogens. In the first article in this continuing medical education series, we review the epidemiology and cutaneous manifestations. The second article in the series focuses on diagnosis, treatment, and complications of these infections. (J Am Acad Dermatol 2019;80:869-80.)

Key words: angioinvasive fungus; aspergillosis; mucormycetes; Candidiasis; Fusarium; Scedosporium.

From the Department of Dermatology,^a University of Wisconsin, Madison; Department of Dermatology,^b University of Pennsylvania, Philadelphia; University of Iowa Carver College of Medicine^c and the Departments of Dermatology^d and Pathology and Clinical Microbiology,^e University of Iowa Hospitals and Clinics, and the Department of Dermatology and Pathology,^f University of Iowa, Iowa City.

Dr Wanat is currently affiliated with the Medical College of Wisconsin, Department of Dermatology, Milwaukee, WI.

Dr Wanat received a Dermatology Foundation Career Development Award.

Conflicts of interest: None disclosed.

Accepted for publication April 19, 2018.

Correspondence to: Karolyn A. Wanat, MD, Department of Dermatology and Pathology, University of Iowa, 200 Hawkins Dr, Iowa City, IA 52246. E-mail: kwanat@gmail.com.

0190-9622/\$36.00

© 2018 by the American Academy of Dermatology, Inc.

<https://doi.org/10.1016/j.jaad.2018.04.059>

Date of release: April 2019

Expiration date: April 2022

EPIDEMIOLOGY AND CLINICAL PRESENTATION

Many of the causative organisms of angioinvasive fungal infections are ubiquitous in the environment and rarely cause clinical infection in patients with an intact immune system, but have the potential to result in serious, often fatal disease in immunocompromised patients.¹⁻⁴ Fungi previously thought to be nonpathogenic are becoming increasingly medically significant as the incidence of invasive fungal infections rises along with extended survival in critically ill patients.⁵ The widespread use of antifungal prophylaxis in immunosuppressed individuals is further altering the resistance patterns of these organisms.⁶ The medical community is called upon to recognize individuals who are at risk for infection, identify variations in clinical presentation, and understand factors involved in prevention and treatment.⁷ Invasive fungal organisms have a propensity to invade blood vessels, and can either be inoculated from local trauma or reach the skin via dissemination in the blood stream.^{8,9} Cutaneous manifestations include pink to purple macules, papules, patches, and plaques, regardless of the causative agent or underlying etiology (Fig 1, A-C). Hemorrhagic, ecthyma gangrenosum-like lesions can occur in infection with systemic candidiasis,¹⁰ aspergillosis,⁹ mucormycetes,¹¹ phaeohyphomycosis, and *Fusarium* infection.¹²⁻¹⁴ Necrotic papulonodules and deep subcutaneous nodules are common to candidiasis,^{15,16} aspergillosis,¹⁷ mucormycosis,¹⁸ and hyalohyphomycosis.^{13,19,20} Patients with mucormycosis^{18,21} and hyalohyphomycosis^{19,22} can both present with cellulitic plaques as well. Unique clinical manifestations of each angioinvasive fungus will be detailed in their individual sections.

CANDIDIASIS

Key points

- **Invasive candidiasis is the leading cause of mycosis-associated mortality in the United States**
- **There is a growing number of medically relevant *Candida* species, including the recently recognized *Candida auris***
- **Disseminated candidiasis is more severe in immunocompromised hosts and neonates**

Invasive candidiasis is the leading cause of mycosis-associated mortality in the United States and is globally recognized as a cause of significant morbidity.²³ The majority of invasive disease is caused by *Candida albicans*, *Candida glabrata*, *Candida tropicalis*, *Candida parapsilosis*, and *Candida krusei*. While *C albicans* remains the most common cause of invasive candidiasis, *C glabrata*

has now emerged as the second most common pathogen in the United States, and *C parapsilosis* and *C tropicalis* remain the second most common causative agents of blood stream infections in other countries.²³ As microbial detection methods become more sensitive, the number of clinically relevant candida species continues to grow.²³ Changing fungal resistance patterns, specifically the development of multidrug resistance in *C glabrata*, makes its rise especially problematic.²⁴ A new angioinvasive species, *Candida auris*, has been recently recognized and exhibits multidrug resistance and reduced susceptibility to fluconazole and amphotericin B.^{25,26}

Humans are known carriers of commensal candidal species within the gastrointestinal tract and vagina, while a subset of the population also carries oral candida asymptomatically.²⁷ Colonizing candida from the oral and gastrointestinal mucosa in the setting of immunosuppression related to chemotherapy can result in invasive infection. Disseminated candidiasis has been cited as the fourth leading cause of hospital-acquired bloodstream infection in the United States²⁸ and has been touted as the causative agent of the majority of cases of invasive fungal infections worldwide.²⁹ The incidence of candidemia in the United States varies by region at 14 per 100,000 people in the Baltimore area and 10 per 100,000 in the Atlanta area.³⁰ The high incidence of infection has been attributed to the increased use of aggressive immunosuppressive therapy, blood transfusion, total parenteral nutrition,³¹ central venous catheters, and immune modifying drugs in well-developed nations,²⁹ with the risk in developing countries attributed to limited medical care, difficulties in implementing infection control in hospitals, and less aggressive antifungal treatments in at-risk patients.³²

Candida spp are not a known commensal organism of the skin and when present are able to trigger an inflammatory response via Toll-like receptor 2, altering keratinocyte expression of key defense cytokines.^{33,34} Interleukins, such as interleukin-17 (IL-17), are recognized for their ability to fight *Candida spp* infection.³⁵⁻³⁷ In addition, certain genetic mutations, including those seen in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy, can predispose to infection with *Candida spp*. There has been an association between the development of autoantibodies against IL-17A with severity of candidal infection in patients with autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy.³⁸ In addition, the use of IL-17 inhibitors (secukinumab and ixekizumab) also can increase the risk of *Candida spp* infections.³⁹




Fig 1. Variable cutaneous manifestations of angioinvasive fungal infections. Presentations include pink to red and violaceous papules and plaques. **A**, *Fusarium* infection involving the left index finger. **B**, *Lichtheimia* on the right thigh. **C**, *Aspergillus* spp infection on the left thigh.

Common cutaneous manifestations caused by *Candida* spp are highlighted in Table I.⁴⁰⁻¹⁰²

In individuals with underlying immunosuppression, candidiasis can be much more severe. Disseminated candidiasis generally presents as a triad of fever, rash, and diffuse muscle tenderness¹⁰³; however, it can involve any organ, including the kidneys, liver, spleen, myocardium, eyes, and brain.²⁹ Cutaneous findings in disseminated disease are broad, but the most commonly reported are asymptomatic, erythematous to violaceous papules with pale, vesicular centers (Supplemental Fig 1; available at <http://www.jaad.org>) that can progress to necrotic, purpuric plaques and tense hemorrhagic bullae.¹⁵ Less common clinical presentations of adult and pediatric candidiasis are shown in Table I. Clinicians should always consider candidemia in the differential when areas of central necrosis and hemorrhage are present.

ASPERGILLOSIS

Key points

- **Invasive aspergillosis infections have increased since 1980**
- **Infection often occurs at sites of skin breakdown, trauma, adhesive dressings, venipuncture or catheter insertion sites, and burn scars**
- **Presentation in disseminated disease can be atypical in immunocompromised hosts**

Aspergillus spp are omnipresent in the environment and inhaled as conidial spores daily.² Aspergillosis infection is most commonly caused by *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus terreus*, and *Aspergillus niger*, all of which generate the clinically distinct entities of allergic bronchopulmonary aspergillosis, allergic fungal

rhinosinusitis, chronic pulmonary aspergillosis, chronic necrotizing pulmonary aspergillosis, and primary cutaneous aspergillosis. Slow-sporulating, difficult-to-culture species *Aspergillus lentulus*, *Aspergillus calidoustus*, and *Neosartorya (Aspergillus) udagawae* are emerging pathogens that have also been implicated in causing invasive infection.¹⁰⁴ Angioinvasion by hyphae is usually rapid and causes necrosis and tissue hemorrhage. Invasive disease was estimated at 1 to 2 cases per 100,000 people in the San Francisco Bay area in one of the first population-based estimates.¹⁰⁵

Cutaneous manifestations can vary depending on whether they are the result of primary inoculation or secondary infection caused by dissemination. Primary cutaneous aspergillosis is rare and is most frequently encountered in immunocompromised patients as described in Table I. Primary cutaneous aspergillosis is also seen in those with chronic granulomatous disease who have a predisposition to acquiring refractory infections caused by *Aspergillus nidulans*, an otherwise rare species.^{2,49} Infection is often nosocomial, occurring at sites of trauma, such as adhesive tape dressings, venipuncture or catheter insertion sites, or burn scars.^{9,50,53,106,107} Burn victims most frequently experience infection between 10 and 35 days postinjury, and environmental triggers have been questioned, with almost 60% of cases of aspergillosis attributed to hospital construction or air conditioner ducts and filters being serviced, according to some reports.^{108,109}

Premature neonates often exhibit some level of mechanical cutaneous barrier dysfunction via the use of adhesives and pulse oximetry. Neonates have diminished phagocytic ability, greatly increasing their risk of aspergillosis. Primary cutaneous infections typically occur between 5 and 30 days after

Table I. Common cutaneous manifestations caused by *Candida spp*

Fungus	Host risk factors	Cutaneous features
<i>Candida spp</i>	Immunosuppression, blood transfusion, total parenteral nutrition ³¹ central venous catheter, immune-modifying medications, ²⁹ limited medical care and infection control in developing countries, ³¹ genetic mutations, ³⁸ interleukin-17 inhibitor use, ³⁹ and autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy ³⁸	Adult: ecthyma gangrenosum—like lesions, ¹⁰ diaper and perianal dermatitis, oral and vaginal thrush, intertrigo, balanitis, angular cheilitis, erosio interdigitalis blastomycetica, ^{18,40-43} pink erythematous and violaceous papules and vesicles, necrotic purpuric plaques, tense hemorrhagic bullae, ¹⁵ folliculitis-like eruption, ⁴⁴ and subcutaneous nodules ¹⁶ Pediatric: erythematous dermatitis with vesicles and desquamation with collarettes of scale ^{45,46} and ecthyma gangrenosum—like lesions ⁴⁷
<i>Aspergillus spp</i>	Immunosuppression, hematologic malignancy, HIV/AIDS, solid organ transplantation, burns, corticosteroid use, diabetes mellitus, neonates, ⁴⁸ chronic granulomatous disease, ^{2,49} and trauma ⁹	Ecthyma gangrenosum—like lesions, ⁹ necrotic papulonodules, ¹⁷ deep subcutaneous nodules with central eschar, ^{50,51} violaceous macules, papules, or plaques, ^{9,52} pinpoint pustules, ⁵² molluscum contagiosum—like lesions, ⁵³ dermatophyte-like lesions, ⁵³ panniculitis, ⁵⁴ subcutaneous granulomas and abscesses, ⁵⁵ erythematous macules with hemorrhagic, and necrotizing ulcerations ¹⁷
Mucormycetes (<i>Rhizopus spp</i> , <i>Mucor spp</i> , <i>Rhizomucor spp</i> , <i>Lichtheimia spp</i> , <i>Saksenaia spp</i> , <i>Cunninghamella spp</i> , and <i>Apophysomyces spp</i>)	Hematologic malignancy, hematopoietic stem cell transplant, renal disease, malnutrition, neutropenia, immunosuppressive medication use (chemotherapy, tumor necrosis factor inhibitors, and corticosteroids), ⁵⁶ iron overload, ^{57,58} diabetes mellitus, burns, traumatic injury, infected tape, wooden splints, ^{56,59-61} toxic epidermal necrolysis, ⁶² and autoimmune disease ²¹	Ecthyma gangrenosum—like lesions, ¹¹ mucocutaneous ulceration and eschar, ²¹ necrotic papulonodules, ¹⁸ gangrenous lesions, ⁶³ cellulitic plaques, ²¹ pink erythematous papules, pustules, vesicles, and subcutaneous nodules, "bull's-eye infarcts," ⁶⁴ targetoid plaques with central necrosis, ⁶⁴ "cotton-like" appearance within wounds, ⁶⁵ pyoderma gangrenosum—like lesions, ⁶⁶ dermatophyte-like lesions, ⁶⁷ and cutaneous vasculitis—like lesions ⁶⁸
<i>Fusarium spp</i>	Immunosuppression, neutropenia, lymphopenia, graft versus host disease, hematologic abnormalities, immunosuppressive medications, ⁷ trauma, burns, and foreign body ⁶⁹	Ecthyma gangrenosum—like lesions, ^{12,70,71} necrotic papulonodules, deep subcutaneous nodules, tinea pedis, intertrigo, onychomycosis, ^{12,13} granuloma annulare—like plaques, ¹⁴ facial granulomas, ⁷² cellulitis, abscesses, and umbilicated or necrotic papules and pustules ^{18,19}

Continued

Table I. Cont'd

Fungus	Host risk factors	Cutaneous features
<i>Scedosporium spp</i>	Immunosuppression, neutropenia, malignancy, hematopoietic or solid organ transplantation, ⁷³ lymphopenia, corticosteroid use, serum albumin <3 mg/dL, cytomegalovirus reactivation, and diabetes ⁷⁴	Eumycetomas with erythematous and necrotic papules, ²⁰ ulcerations, subcutaneous nodules, ^{2,75} cellulitis, ²² sporotrichoid spread, thrombophlebitis, ^{76,77} diffuse papules and pustules, ⁷⁵ cutaneous abscesses, ⁷⁵ pyoderma gangrenosum–like lesions, ⁷⁸ and malignant otitis externa ⁷⁹
<i>Paecilomyces spp</i>	Immunosuppression, ⁸⁰ postoperative, ^{81–83} and HIV/AIDS ^{84,85}	Subcutaneous swellings and nodules, ^{84,86} septic bursitis, ⁸⁷ red-brown papules, plaques, papules, abscesses, ⁸⁸ and eumycetoma ⁸⁹
Other hyalohyphomycoses (<i>Acremonium spp</i> and <i>Trichoderma spp</i>)	Immunosuppression, intravenous catheters, and trauma ⁹⁰	Cellulitic plaques, tender red papules, ⁹⁰ mycetoma, ^{90,91} and ulceronecrotic plaques at catheter sites ⁹²
Phaeohyphomycoses (<i>Alternaria spp</i> , <i>Bipolaris spp</i> , <i>Wangiella spp</i> , <i>Fonsecaea spp</i> , <i>Cladophialophora spp</i> , <i>Curvalaria spp</i> , <i>Exophiala spp</i> , and <i>Phialophora spp</i>)	Immunosuppression, malignancy, diabetes mellitus, corticosteroid use, HIV/AIDS, ^{73,93,94} solid organ transplantation, ⁹⁵ and hypercortisolism ⁹⁶	Ecthyma gangrenosum–like lesions, ⁹⁷ verrucous papulonodules, ^{98,99} eumycetoma, ² granulomatous nodules, ^{98,100,101} sporotrichoid spread, ⁹⁸ multiple pyogenic granuloma–like nodules, dermatophytosis-like plaques, ¹⁰² subcutaneous cysts, ¹⁰² cutaneous erythema, desquamation, eczematous patches that ulcerate and erode, ⁹⁷ and onychomycosis ²

birth and may present with papules, nodules, pustules, and ulcers.⁹

As the population of immunocompromised individuals expands, it is likely that the prevalence of primary cutaneous aspergillosis will increase.¹¹⁰ In primary lesions, the skin exhibits red to violaceous, broad, flat, hard, macules, papules, or plaques that may be painful or entirely asymptomatic.⁹ Aspergillosis disease progression leads to the development of vesicles with central hemorrhage, which subsequently ulcerate (Fig 2, A). Primary infection is usually limited to a few lesions but may rapidly expand and develop disseminated lesions with variable morphologies as outlined in Table I. In patients with AIDS or who are otherwise immunocompromised, atypical presentations can be seen as outlined in Table I. In immunocompetent patients, disease can arise from surgical wounds, traumatic inoculation, or via exposure to high spore counts, which are frequently encountered in farming.⁹

Dissemination of primary cutaneous lesions remains a risk in the immunocompromised host. Accompanying systemic symptoms, including



Fig 2. Cutaneous manifestations of *Aspergillus spp* infection. Primary cutaneous *Aspergillus spp* infections on the right forearm presenting as an impending ulcerated violaceous plaque.

respiratory disease, fever, stroke, and seizure, are observed in disseminated disease, which usually occurs from hematogenous spread from the lungs.² The Centers for Disease Control and Prevention report an increase in invasive aspergillosis infection of 357% since 1980.^{48,111} In systemic infections that

secondarily involve the skin, widespread lesions with deep, multiple, subcutaneous nodules, with or without central eschar, can be seen (Fig 2, B; Supplemental Fig 2, available at <http://www.jaad.org>).^{50,51} Importantly, cutaneous findings can be the first indicator of underlying systemic infection, and a high clinical index of suspicion is key to making this diagnosis in a timely fashion.⁵⁰

These patients should be treated rapidly because disseminated disease can be fatal. Prognosis can improve with discontinuation of immunosuppressant medications or reversal of neutropenia.¹¹² While aspergillosis in patients undergoing allogeneic hematopoietic stem cell transplants used to be rare, the incidence of invasive disseminated aspergillosis amongst this patient population has increased along with a shift from early to late postengraftment aspergillosis.¹¹³ Treatment with fludarabine or rituximab during the pretransplant period are extremely high risk factors for invasive aspergillosis, and these patients should be regularly assessed for this complication.¹¹⁴ Similarly, there are reported cases of transition from aspergilloma to invasive aspergillosis after therapy with sunitinib.¹¹⁵

MUCORMYCOSIS

Key points

- **Mucormycetes, previously referred to as zygomycetes, includes several genera: *Rhizopus*, *Mucor*, *Rhizomucor*, *Lichtheimia*, *Saksena*, *Cunninghamella*, and *Apophysomyces***
- **Diabetes is a major risk factor for disseminated mucormycosis**
- ***Mucor spp* sequester iron from their hosts, indicating iron overload syndromes as an additional risk factor in disseminated disease**

Mucormycetes include the genera *Rhizopus*, *Mucor*, and *Rhizomucor*,²¹ as well as *Lichtheimia spp*, *Saksena*, *Cunninghamella spp*, and *Apophysomyces spp*, and are so named for their ability to cause systemic mucormycosis. Previously referred to as zygomycetes after the phylum Zygomycota (which is no longer accepted), a genetic reconsideration of taxonomy suggests that mucormycosis (after the order *Mucorales*) is more appropriate than zygomycosis when referring to these fungi and constellation of symptoms.¹¹⁶

Mucormycetes are abundant in the environment and are routinely encountered by humans via direct contact with soil and decaying vegetation or through the inhalation of conidial spores.¹¹⁷ Mucormycetes are frequently implicated pathogens in patients with underlying immunocompromise as outlined in

Table I. A well-known virulence factor of the *Mucorales* is the ability to sequester iron from host organisms.¹¹⁸ Iron overload syndromes, including those in patients requiring frequent red blood cell transfusions in the setting of transplant or hematopoietic disorders^{57,58} or medications that chelate iron, such as deferoxamine, represent potential risk factors for infection; however, the novel iron chelators deferasirox and deferiprone do not predispose to mucormycosis.²¹ Although most often seen as a disease of the immunosuppressed host, reports of these species causing infection in healthy hosts without identifiable risk factors have been published.¹¹⁹⁻¹²¹ While epidemiologic data are difficult to obtain, small case reviews have cited the incidence of mucormycosis between 10.8%¹¹⁹ and 22.2%¹²⁰ in hosts without known risk factors for disease.

Recent epidemiologic research suggests that an environmental component also contributes to the infectivity of these organisms, citing natural disasters, contaminated adhesive dressings, water supplies, tongue depressors, hospital linens, and building construction in outbreaks of mucormycosis.¹²²⁻¹²⁵ Lower levels of economic and industrial development in the community serve as an appreciable risk for infection in individual hosts.²¹ This may be related to sanitation practices, temperature differences, stagnant water supplies, or another unidentified factor.

Mucormycosis can exhibit 5 main clinical presentations, including pulmonary, gastrointestinal, rhinocerebral, cutaneous, and disseminated infections.¹²⁶ Pulmonary disease arises after the inhalation of conidial spores, and pulmonary mucormycosis can develop into chest wall cellulitis via direct extension.^{21,127} Gastrointestinal mucor localizes to the stomach, ileum, and colon and typically affects malnourished patients, premature neonates, and neutropenic adults. The rapid fatality of gastrointestinal disease makes it frequently diagnosed postmortem¹²⁸ (Supplemental Fig 3, available at <http://www.jaad.org>). Rhinocerebral infection is most frequently encountered in patients with poorly controlled diabetes when infection of the paranasal sinuses occurs after the inhalation of sporangiospores.²¹ As rapid spread to contiguous structures occurs, patients exhibit facial pain, nasal discharge, periorbital cellulitis, blurred vision, proptosis, headache, mucocutaneous ulceration, and eschar, with a risk for cerebral extension.²¹

Disease is considered localized when it is limited to the skin and subcutaneous tissue and is considered disseminated when involvement includes muscle, tendon, bone, and noncontiguous organs.⁶⁷




Fig 3. *Mucormycosis spp* can present in a variety of ways. Purpuric plaques in the perinasal and periocular locations, which are concerning for extension from sinonasal locations, and in occluded areas such as (A) on the left forearm and (B) hemorrhagic bullae on the toes.

When cutaneous mucormycosis occurs secondarily, as a consequence of hematogenous dissemination, skin findings are the result of hyphae invading blood vessels, with rapid necrosis and tissue hemorrhage. The skin can exhibit pink, erythematous papules, pustules, and vesicles with suppurative inflammation and rapid ulceration that evolves to central crusting and eschar (Fig 3, A and B; Supplemental Fig 4, available at <http://www.jaad.org>). Coloration can be variable, ranging from pink-yellow nodules to black necrosis in immunosuppressed patients.^{129,130} A predilection for areas of trauma and distal extremities frequently occurs.¹¹⁷

OTHER HYALINE MOLDS

Key points

- Cutaneous involvement can be seen in approximately 75% of cases of disseminated *Fusarium* infection
- Hyalohyphomycoses often present with eumycetoma

Hyaline molds, or hyalohyphomycoses, include lightly pigmented, septate, branching filamentous fungi. *Fusarium spp*, *Scedosporium spp*, *Paecilomyces spp*, *Acremonium spp*, and *Trichoderma spp* are the most frequently encountered infectious agents. Infection can result from inhalation or traumatic implantation from the environment, including soil, wood, or contaminated water sources. It is often difficult to distinguish between true infection and colonization in immunocompromised patients.² Cutaneous hyalohyphomycosis can be difficult to detect clinically, presenting with numerous clinical morphologies and frequently evolving with time and host immune status as described in Table I.

Tissue breakdown is the primary risk factor in immunocompetent individuals, and *Fusarium* can

exhibit virulence via attachment to prosthetic material and suppression of the immune system by production of mycotoxins, such as trichothecenes.¹³¹ Table I outlines the risk factors for fusariosis. Inhalation of airborne conidial spores, as well as skin and mucosal membrane bypass, serve as the primary portals of entry for *Fusarium*; however, these species have been uncovered in hospital water systems,¹³² and it has been suggested that water-related activities could serve in acquisition of infection.

Cutaneous involvement can be seen in approximately 75% of cases of disseminated *Fusarium* infection, and manifestations include diffuse, erythematous to violaceous papules and nodules, often with central necrosis and gray eschar, which can resemble ecthyma gangrenosum (Fig 4; Supplemental Fig 5, available at <http://www.jaad.org>).^{12,70,71} Lesions can be exquisitely tender and evolve quickly over the course of a few days, likely because *Fusarium spp* have the ability to generate collagenases and proteases that result in tissue breakdown.¹³³ Hematogenous and lymphangitic spread of deep lesions may progress to disseminated fusariosis.¹³⁴ The prognosis for immunocompromised patients with widespread disease is poor, but an intact immune system serves to increase survival rates drastically.¹³

There are two medically significant *Scedosporium* species, including, *Scedosporium apiospermum* (*Pseudallescheria boydii*) and *Scedosporium prolificans*.⁷ Scedosporiosis is most frequently found in immunocompromised patients as reviewed in Table I. Targeted organs include the lungs, central nervous system, skin, and eyes. The lungs are the most frequently infected with *S apiospermum*, and patients with cystic fibrosis and chronic pulmonary disease are at an elevated risk of pulmonary infection.¹³⁵ In patients with pulmonary scedosporiosis, development of secondary brain abscesses has been



Fig 4. *Fusarium* infection presenting in the skin as a purpuric plaque on the right thigh.

reported.¹³⁶ Endophthalmitis, presenting as ocular floaters and decreased visual acuity, can be the first clinical manifestation of scedosporiosis, and clinicians should be wary of this symptom in immunocompromised hosts.¹³⁷ Cutaneous findings can be primary or caused by dissemination. In primary lesions, traumatic implantation can result in white-grain mycetomas leading to the variable clinical presentations highlighted in Table I.² If nodules develop they can ulcerate and crust, mimicking syphilitic gummas.

Paecilomyces spp are encountered in foods, soil, and wood; however, their ascospores are strongly heat- and treatment-resistant, making them a contaminant of pasteurized and treated food products and laboratory cultures.^{138,139} Infection most commonly occurs in immunocompromised individuals and after surgical procedures or implantation of cardiac valves,^{81,140} intraocular lenses,^{82,141} and dialysis catheters.^{80,83} Disseminated *Paecilomycosis* infection often presents with tender, subcutaneous, ulcerated nodules.⁸⁴ Cases of *Paecilomyces* infections are also reported in immunocompetent hosts, with features outlined in Table I. Coinfection with *Alternaria* also has been reported.^{142,143} Given their frequency as contaminants, *Paecilomyces* infections have been initially misdiagnosed as lupus on histopathology, so clinician awareness of this organism's potential for infection is important.¹³⁹

Trichoderma spp are most frequently responsible for peritonitis in patients undergoing peritoneal dialysis; however, reports of disseminated infection have surfaced more recently. Cutaneous presentations have included perianal ulceration and ulceronecrotic lesions at the sites of intravenous catheters.⁹²

Infections caused by *Acremonium spp* occur most frequently in the settings of immunosuppression, penetrating trauma, and intravascular catheters.⁹⁰ There are reports of cutaneous *Acremonium*

presenting with mycetoma in both immunocompromised and immunocompetent hosts after trauma.^{90,91} Less common presentations of *Acremonium* infection are outlined in Table I.

PHAEOHYPHOMYCOSIS

Key points

- **Phaeohyphomycoses are rare but can cause infection in both immunocompetent and immunocompromised hosts**
- **Melanin production in these organisms plays a key role in fungal infectivity and virulence**

Dematiaceous molds, also known as phaeohyphomycoses, encompass the genera *Alternaria spp*, *Bipolaris spp*, *Wangiella spp*, *Madurella spp*, *Fonsecaea spp*, *Cladophialophora spp*, *Curvularia spp*, *Exophiala spp*, and *Phialophora spp*. They are termed for their natural melanin pigment and subsequent colored cell walls. Their ability to produce melanin serves as an important virulence factor in fungal infectivity, presumably by reducing susceptibility to host defense mechanisms and antifungals.^{2,144} Much like the mucormycoses, they are ubiquitous in nature, present in soil and vegetation worldwide, with traumatic implantation frequently related in these infections. Risk factors for infection are described in Table I.

While most notable for their role in allergic fungal sinusitis, phaeohyphomycoses can cause the skin and soft tissue infections outlined in Table I. These fungal species also have a predilection for invasion of vessels with risk for hematogenous and lymphatic dissemination as well as invasion of bone, causing osteomyelitis.¹⁴⁵ In a review of 56 stem cell and solid organ transplant patients with identified phaeohyphomycosis, 55.4% of them exhibited disseminated disease, with *Alternaria* being the more frequent causative agent,⁹⁵ highlighting the importance of immune compromise as a risk factor for generalized disease. One of the main predisposing factors to alternariosis is hypercortisolism or Cushing syndrome.⁹⁷ Less frequent cutaneous manifestations of phaeohyphomycoses are outlined in Table I.

Disseminated *Cladophialophora* infection is rare but is notable for the formation of skin lesions and brain abscesses, and cases have been reported in both healthy and immunocompromised hosts.^{146,147} *Curvularia spp* have reportedly caused ecthyma gangrenosum-like lesions,⁹⁷ disseminated erythematous papules and nodules,¹⁴⁸ and deep, soft tissue abscesses with ulceration.¹⁴⁹ There are few reports of disseminated *Bipolaris* infection with cutaneous

involvement; however, this organism is known to cause localized cutaneous disease. Unfortunately, recent reports of brain abscesses and biliary tree invasion in an immunocompetent patient with *Bipolaris*¹⁵⁰ and brain abscesses in an immunocompetent patient with *Cladophialophora*¹⁵¹ make it clear that healthy hosts can be affected with phaeohyphomycoses.

In conclusion, the clinical presentations of the angioinvasive fungi can vary depending on a patient's immune status, but early recognition for prompt diagnosis and the fast initiation of therapy are imperative to prevent morbidity and high mortality in these conditions. Diagnostic work-up, therapies, and potential complications will be discussed in the second article in this continuing medical education series.

REFERENCES

- Low CY, Rotstein C. Emerging fungal infections in immunocompromised patients. *F1000 Med Rep*. 2011;3:14.
- Guarner J, Brandt ME. Histopathologic diagnosis of fungal infections in the 21st century. *Clin Microbiol Rev*. 2011;24:247-280.
- McNeil MM, Nash SL, Hajjeh RA, et al. Trends in mortality due to invasive mycotic diseases in the United States, 1980-1997. *Clin Infect Dis*. 2001;33:641-647.
- Mora C, Tittensor DP, Adl S, Simpson AG, Worm B. How many species are there on Earth and in the ocean? *PLoS Biol*. 2011;9:e1001127.
- Naggie S, Perfect JR. Molds: hyalohyphomycosis, phaeohyphomycosis, and zygomycosis. *Clin Chest Med*. 2009;30:337-353.
- Enoch DA, Ludlam HA, Brown NM. Invasive fungal infections: a review of epidemiology and management options. *J Med Microbiol*. 2006;55:809-818.
- Walsh TJ, Groll A, Hiemenz J, Fleming R, Roilides E, Anaissie E. Infections due to emerging and uncommon medically important fungal pathogens. *Clin Microbiol Infect*. 2004;10(suppl 1):48-66.
- Ben-Ami R, Lewis RE, Leventakos K, Latgé JP, Kontoyiannis DP. Cutaneous model of invasive aspergillosis. *Antimicrob Agents Chemother*. 2010;54:1848-1854.
- van Burik JA, Colven R, Spach DH. Cutaneous aspergillosis. *J Clin Microbiol*. 1998;36:3115-3121.
- Fine JD, Miller JA, Harrist TJ, Haynes HA. Cutaneous lesions in disseminated candidiasis mimicking ecthyma gangrenosum. *Am J Med*. 1981;70:1133-1135.
- Kramer BS, Hernandez AD, Reddick RL, Levine AS. Cutaneous infarction. Manifestation of disseminated mucormycosis. *Arch Dermatol*. 1977;113:1075-1076.
- Gupta AK, Baran R, Summerbell RC. Fusarium infections of the skin. *Curr Opin Infect Dis*. 2000;13:121-128.
- Nucci M, Anaissie E. Fusarium infections in immunocompromised patients. *Clin Microbiol Rev*. 2007;20:695-704.
- Ooi SP, Chen TT, Huang TH, Chang HS, Hsieh HY. Granuloma annulare-like skin lesion due to *Fusarium roseum*: therapy with ketoconazole. *Arch Dermatol*. 1987;123:167-168.
- Pedraz J, Delgado-Jiménez Y, Pérez-Gala S, Nam-Cha S, Fernández-Herrera J, García-Díez A. Cutaneous expression of systemic candidiasis. *Clin Exp Dermatol*. 2009;34:106-110.
- Arribi A, Martínez I, Alvarez P, Amondarain I, López S, Del Palacio A. Candidemia and cutaneous lesions in two patients without intravenous drug addiction [in Spanish]. *Rev Iberoam Micol*. 1999;16:235-237.
- Khadori N, Hayat S, Rolston K, Bodey GP. Cutaneous Rhizopus and Aspergillus infections in five patients with cancer. *Arch Dermatol*. 1989;125:952-956.
- Bologna JL, Jorizzo JL, Rapini RP, et al., eds. *Dermatology*. 2nd ed. Philadelphia, PA: Mosby Elsevier; 2008.
- Nucci M, Anaissie E. Cutaneous infection by *Fusarium* species in healthy and immunocompromised hosts: implications for diagnosis and management. *Clin Infect Dis*. 2002;35:909-920.
- Severo LC, Oliveira FeM, Londero AT. Subcutaneous scedosporiosis. Report of two cases and review of the literature. *Rev Inst Med Trop Sao Paulo*. 1997;39:227-230.
- Petrikos G, Skiada A, Lortholary O, Roilides E, Walsh TJ, Kontoyiannis DP. Epidemiology and clinical manifestations of mucormycosis. *Clin Infect Dis*. 2012;54(suppl 1):S23-S34.
- Misselbrook GP, Lillie P, Thomas CP. An unusual pathogen in ambulatory care: two cases of *Scedosporium* soft tissue infections presenting as "unresponsive cellulitis". *Acute Med*. 2016;15:88-91.
- Pfaller MA, Diekema DJ. Epidemiology of invasive candidiasis: a persistent public health problem. *Clin Microbiol Rev*. 2007;20:133-163.
- Sanglard D. Emerging threats in antifungal-resistant fungal pathogens. *Front Med (Lausanne)*. 2016;3:11.
- Chowdhary A, Sharma C, Meis JF. *Candida auris*: a rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog*. 2017;13:e1006290.
- Larkin E, Hager C, Chandra J, et al. The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob Agents Chemother*. 2017:61.
- Cannon RD, Chaffin WL. Oral colonization by *Candida albicans*. *Crit Rev Oral Biol Med*. 1999;10:359-383.
- Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, Edmond MB. Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. *Clin Infect Dis*. 2004;39:309-317.
- Enoch DA, Yang H, Aliyu SH, Micallef C. The changing epidemiology of invasive fungal infections. *Methods Mol Biol*. 2017;1508:17-65.
- Cleveland AA, Harrison LH, Farley MM, et al. Declining incidence of candidemia and the shifting epidemiology of *Candida* resistance in two US metropolitan areas, 2008-2013: results from population-based surveillance. *PLoS One*. 2015;10:e0120452.
- Mermutluoglu C, Deveci O, Dayan S, Aslan E, Bozkurt F, Tekin R. Antifungal susceptibility and risk factors in patients with Candidemia. *Eurasian J Med*. 2016;48:199-203.
- Warnock DW. Trends in the epidemiology of invasive fungal infections. *Nihon Ishinkin Gakkai Zasshi*. 2007;48:1-12.
- Li M, Chen Q, Shen Y, Liu W. *Candida albicans* phospholipomannan triggers inflammatory responses of human keratinocytes through Toll-like receptor 2. *Exp Dermatol*. 2009;18:603-610.
- Shiraki Y, Ishibashi Y, Hiruma M, Nishikawa A, Ikeda S. *Candida albicans* abrogates the expression of interferon-gamma-inducible protein-10 in human keratinocytes. *FEMS Immunol Med Microbiol*. 2008;54:122-128.
- Nielsen J, Kofod-Olsen E, Spaun E, Larsen CS, Christiansen M, Mogensen TH. A STAT1-gain-of-function mutation causing Th17 deficiency with chronic mucocutaneous candidiasis, psoriasiform hyperkeratosis and dermatophytosis. *BMJ Case Rep*. 2015;2015.

36. Campois TG, Zucoloto AZ, de Almeida Araujo EJ, et al. Immunological and histopathological characterization of cutaneous candidiasis. *J Med Microbiol*. 2015;64:810-817.
37. Conti HR, Whibley N, Coleman BM, Garg AV, Jaycox JR, Gaffen SL. Signaling through IL-17C/IL-17RE is dispensable for immunity to systemic, oral and cutaneous candidiasis. *PLoS One*. 2015;10:e0122807.
38. Sarkadi AK, Taskó S, Csorba G, Tóth B, Erdős M, Maródi L. Autoantibodies to IL-17A may be correlated with the severity of mucocutaneous candidiasis in APECED patients. *J Clin Immunol*. 2014;34:181-193.
39. Saunte DM, Mrowietz U, Puig L, Zachariae C. Candida infections in psoriasis and psoriatic arthritis patients treated with IL-17 inhibitors and their practical management. *Br J Dermatol*. 2017;177:47-62.
40. Bonifaz A, Rojas R, Tirado-Sánchez A, et al. Superficial mycoses associated with diaper dermatitis. *Mycopathologia*. 2016;181:671-679.
41. Bradford LL, Chibucos M, Ma B, Bruno V, Ravel J. Vaginal *Candida* spp. genomes from women with vulvovaginal candidiasis. *Pathog Dis*. 2017;75. <https://doi.org/10.1093/femspd/ftx061>.
42. Hedderwick S, Kauffman CA. Opportunistic fungal infections: superficial and systemic candidiasis. *Geriatrics*. 1997;52:50-54, 9.
43. Elder DE, Lever WF. *Lever's Histopathology of the Skin*. Philadelphia, PA: Lippincott-Raven; 1997.
44. Recio C, Piqué E, Lluch J, Vidal JM. Candida folliculitis in intravenous drug users. [in Spanish] *Enferm Infecc Microbiol Clin*. 2003;21:386-387.
45. Brenuchon C, Lebas D, Rakza T, Piette F, Storme L, Catteau B. Invasive fungal dermatitis in extremely premature newborns: a specific clinical form of systemic candidiasis. [in French] *Ann Dermatol Venerol*. 2006;133:341-346.
46. Rudolph N, Tariq AA, Reale MR, Goldberg PK, Kozinn PJ. Congenital cutaneous candidiasis. *Arch Dermatol*. 1977;113:1101-1103.
47. Agarwal S, Sharma M, Mehndirata V. Solitary ecthyma gangrenosum (EG)-like lesion consequent to *Candida albicans* in a neonate. *Indian J Pediatr*. 2007;74:582-584.
48. Steinbach WJ, Marr KA, Anaissie EJ, et al. Clinical epidemiology of 960 patients with invasive aspergillosis from the PATH Alliance registry. *J Infect*. 2012;65:453-464.
49. Segal BH, Romani LR. Invasive aspergillosis in chronic granulomatous disease. *Med Mycol*. 2009;47(suppl 1):S282-S290.
50. Schimmelpfennig C, Naumann R, Zuberbier T, et al. Skin involvement as the first manifestation of systemic aspergillosis in patients after allogeneic hematopoietic cell transplantation. *Bone Marrow Transplant*. 2001;27:753-755.
51. Ozcan D, Güleç AT, Haberal M. Multiple subcutaneous nodules leading to the diagnosis of pulmonary aspergillosis in a renal transplant recipient. *Clin Transplant*. 2008;22:120-123.
52. Vedder JS, Schorr WF. Primary disseminated pulmonary aspergillosis with metastatic skin nodules. Successful treatment with inhalation nystatin therapy. *JAMA*. 1969;209:1191-1195.
53. Hunt SJ, Nagi C, Gross KG, Wong DS, Mathews WC. Primary cutaneous aspergillosis near central venous catheters in patients with the acquired immunodeficiency syndrome. *Arch Dermatol*. 1992;128:1229-1232.
54. Colmenero I, Alonso-Sanz M, Casco F, Hernández-Martín A, Torrel A. Cutaneous aspergillosis mimicking pancreatic and gouty panniculitis. *J Am Acad Dermatol*. 2012;67:789-791.
55. Findlay GH, Roux HF, Simson IW. Skin manifestations in disseminated aspergillosis. *Br J Dermatol*. 1971;85:94-97.
56. Rammaert B, Lanternier F, Poirée S, Kania R, Lortholary O. Diabetes and mucormycosis: a complex interplay. *Diabetes Metab*. 2012;38:193-204.
57. Delea TE, Hagiwara M, Phatak PD. Retrospective study of the association between transfusion frequency and potential complications of iron overload in patients with myelodysplastic syndrome and other acquired hematopoietic disorders. *Curr Med Res Opin*. 2009;25:139-147.
58. Shander A, Cappellini MD, Goodnough LT. Iron overload and toxicity: the hidden risk of multiple blood transfusions. *Vox Sang*. 2009;97:185-197.
59. Hammond DE, Winkelmann RK. Cutaneous phycomycosis. Report of three cases with identification of *Rhizopus*. *Arch Dermatol*. 1979;115:990-992.
60. Gartenberg G, Bottone EJ, Keusch GT, Weitzman I. Hospital-acquired mucormycosis (*Rhizopus rhizopodiformis*) of skin and subcutaneous tissue: epidemiology, mycology and treatment. *N Engl J Med*. 1978;299:1115-1118.
61. Lalayanni C, Baliakas P, Xochelli A, et al. Outbreak of cutaneous zygomycosis associated with the use of adhesive tape in haematology patients. *J Hosp Infect*. 2012;81:213-215.
62. Razmi TM, Shivaprakash MR, Saikia UN, De D, Handa S. All that necroses is not toxic epidermal necrolysis. *J Cutan Med Surg*. 2017;21:172-173.
63. Boyd AS, Wiser B, Sams HH, King LE. Gangrenous cutaneous mucormycosis in a child with a solid organ transplant: a case report and review of the literature. *Pediatr Dermatol*. 2003;20:411-415.
64. Rubin AI, Grossman ME. Bull's-eye cutaneous infarct of zygomycosis: a bedside diagnosis confirmed by touch preparation. *J Am Acad Dermatol*. 2004;51:996-1001.
65. Chawla R, Sehgal S, Kumar SR, Mishra B. A rare case of mucormycosis of median sternotomy wound caused by *Rhizopus arrhizus*. *Indian J Med Microbiol*. 2007;25:419-421.
66. Kerr OA, Bong C, Wallis C, Tidman MJ. Primary cutaneous mucormycosis masquerading as pyoderma gangrenosum. *Br J Dermatol*. 2004;150:1212-1213.
67. de Oliveira-Neto MP, Da Silva M, Fialho Monteiro PC, et al. Cutaneous mucormycosis in a young, immunocompetent girl. *Med Mycol*. 2006;44:567-570.
68. Meyer RD, Kaplan MH, Ong M, Armstrong D. Cutaneous lesions in disseminated mucormycosis. *JAMA*. 1973;225:737-738.
69. Dignani MC, Anaissie E. Human fusariosis. *Clin Microbiol Infect*. 2004;10(suppl 1):67-75.
70. Martino P, Gastaldi R, Raccach R, Girmenia C. Clinical patterns of *Fusarium* infections in immunocompromised patients. *J Infect*. 1994;28(suppl 1):7-15.
71. Uludokumacı S, Balkan I, Mete B, Ozaras R, Saltoğlu N, Soysal T. Ecthyma gangrenosum-like lesions in a febrile neutropenic patient with simultaneous pseudomonas sepsis and disseminated fusariosis. *Turk J Haematol*. 2013;30:321-324.
72. Benjamin RP, Callaway L, Conant NF. Facial granuloma associated with *Fusarium* infection. *Arch Dermatol*. 1970;101:598-600.
73. Revankar SG, Patterson JE, Sutton DA, Pullen R, Rinaldi MG. Disseminated phaeohyphomycosis: review of an emerging mycosis. *Clin Infect Dis*. 2002;34:467-476.
74. Lamarinis GA, Chamilos G, Lewis RE, Safdar A, Raad II, Kontoyiannis DP. *Scedosporium* infection in a tertiary care cancer center: a review of 25 cases from 1989-2006. *Clin Infect Dis*. 2006;43:1580-1584.
75. Valenzuela Salas I, Martínez Peinado C, Fernández Miralbell A, et al. Skin infection caused by *Scedosporium apiospermum* in immunocompromised patients. Report of two cases. *Dermatol Online J*. 2013;19:20022.

76. Boyce Z, Collins N. *Scedosporium apiospermum*: an unreported cause of fungal sporotrichoid-like lymphocutaneous infection in Australia and review of the literature. *Australas J Dermatol*. 2015;56:e39-42.
77. Scott M, Wanat K, Pappas-Taffer L. Sporotrichoid eruption in a patient after lung transplantation. *Scedosporium apiospermum* fungal thrombophlebitis and sporotrichoid nodules. *JAMA Dermatol*. 2014;150:83-84.
78. Duretz C, Buchlin P, Huguenin A, et al. *Scedosporium apiospermum* skin infection mimicking a pyoderma gangrenosum. [in French] *Med Mal Infect*. 2018;48:212-214.
79. McLaren O, Potter C. *Scedosporium apiospermum*: a rare cause of malignant otitis externa. *BMJ Case Rep*. 2016;2016.
80. Tan TQ, Ogden AK, Tillman J, Demmler GJ, Rinaldi MG. *Paecilomyces lilacinus* catheter-related fungemia in an immunocompromised pediatric patient. *J Clin Microbiol*. 1992;30:2479-2483.
81. Allevalo PA, Ohorodnik JM, Mezger E, Eisses JF. *Paecilomyces javanicus* endocarditis of native and prosthetic aortic valve. *Am J Clin Pathol*. 1984;82:247-252.
82. Mosier MA, Lusk B, Pettit TH, Howard DH, Rhodes J. Fungal endophthalmitis following intraocular lens implantation. *Am J Ophthalmol*. 1977;83:1-8.
83. Nankivell BJ, Pacey D, Gordon DL. Peritoneal eosinophilia associated with *Paecilomyces variotii* infection in continuous ambulatory peritoneal dialysis. *Am J Kidney Dis*. 1991;18:603-605.
84. Martin CA, Roberts S, Greenberg RN. Voriconazole treatment of disseminated *paecilomyces* infection in a patient with acquired immunodeficiency syndrome. *Clin Infect Dis*. 2002;35:e78-e81.
85. Lovell RD, Moll M, Allen J, Cicci LG. Disseminated *Paecilomyces lilacinus* infection in a patient with AIDS. *AIDS Read*. 2002;12:212-213, 8, 21.
86. Permi HS, Sunil KY, Karnaker VK, Kishan PH, Teerthanath S, Bhandary SK. A rare case of fungal maxillary sinusitis due to *Paecilomyces lilacinus* in an immunocompetent host, presenting as a subcutaneous swelling. *J Lab Physicians*. 2011;3:46-48.
87. Schweitzer KM, Richard MJ, Leversedge FJ, Ruch DS. *Paecilomyces lilacinus* septic olecranon bursitis in an immunocompetent host. *Am J Orthop (Belle Mead NJ)*. 2012;41:E74-E75.
88. Kitami Y, Kagawa S, Iijima M. A case of cutaneous *Paecilomyces lilacinus* infection on the face [in Japanese]. *Nihon Ishinkin Gakkai Zasshi*. 2005;46:267-272.
89. Motswaledi HM, Matheka K, Sein PP, Nemutavhanani DL. *Paecilomyces lilacinus* eumycetoma. *Int J Dermatol*. 2009;48:858-861.
90. Hilmioglu S, Metin DY, Tasbakan M, Pullukcu H, Akalin T, Tumbay E. Skin infection on both legs caused by *Acremonium strictum* (case report). *Ann Saudi Med*. 2015;35:406-408.
91. Guevara D, Wongkittiroch K, Goodman M, Singh A, Weiss E, Glick B. *Acremonium* mycetoma: a case report and discussion. *Cutis*. 2011;88:293-295, 9.
92. Chouaki T, Lavarde V, Lachaud L, Raccurt CP, Hennequin C. Invasive infections due to *Trichoderma* species: report of 2 cases, findings of in vitro susceptibility testing, and review of the literature. *Clin Infect Dis*. 2002;35:1360-1367.
93. Brandt ME, Warnock DW. Epidemiology, clinical manifestations, and therapy of infections caused by dematiaceous fungi. *J Chemother*. 2003;15(suppl 2):36-47.
94. Wong EH, Revankar SG. Dematiaceous molds. *Infect Dis Clin North Am*. 2016;30:165-178.
95. McCarty TP, Baddley JW, Walsh TJ, et al. Phaeohyphomycosis in transplant recipients: results from the Transplant Associated Infection Surveillance Network (TRANSNET). *Med Mycol*. 2015;53:440-446.
96. Vennwald I, Wollina U. Cutaneous infections due to opportunistic molds: uncommon presentations. *Clin Dermatol*. 2005;23:565-571.
97. Bonduel M, Santos P, Turienzo CF, Chantada G, Paganini H. Atypical skin lesions caused by *Curvularia* spp. and *Pseudallescheria boydii* in two patients after allogeneic bone marrow transplantation. *Bone Marrow Transplant*. 2001;27:1311-1313.
98. Hsu CC, Chang SS, Lee PC, Chao SC. Cutaneous alternariosis in a renal transplant recipient: a case report and literature review. *Asian J Surg*. 2015;38:47-57.
99. Lyskova P, Kubanek M, Hubka V, et al. Successful posaconazole therapy of disseminated alternariosis due to *Alternaria* infectoria in a heart transplant recipient. *Mycopathologia*. 2017;182:297-303.
100. Iwatsu T. Cutaneous alternariosis. *Arch Dermatol*. 1988;124:1822-1825.
101. Mitchell AJ, Solomon AR, Beneke ES, Anderson TF. Subcutaneous alternariosis. *J Am Acad Dermatol*. 1983;8:673-676.
102. Khader A, Ambooken B, Binitha MP, Francis S, Kuttiyil AK, Sureshan DN. Disseminated cutaneous phaeohyphomycosis due to *Cladophialophora bantiana*. *Indian J Dermatol Venereol Leprol*. 2015;81:491-494.
103. Jarowski CI, Fialk MA, Murray HW, et al. Fever, rash, and muscle tenderness. A distinctive clinical presentation of disseminated candidiasis. *Arch Intern Med*. 1978;138:544-546.
104. Kauffman CA, Freifeld AG, Andes DR, et al. Endemic fungal infections in solid organ and hematopoietic cell transplant recipients enrolled in the Transplant-Associated Infection Surveillance Network (TRANSNET). *Transpl Infect Dis*. 2014;16:213-224.
105. Rees JR, Pinner RW, Hajjeh RA, Brandt ME, Reingold AL. The epidemiological features of invasive mycotic infections in the San Francisco Bay area, 1992-1993: results of population-based laboratory active surveillance. *Clin Infect Dis*. 1998;27:1138-1147.
106. Allo MD, Miller J, Townsend T, Tan C. Primary cutaneous aspergillosis associated with Hickman intravenous catheters. *N Engl J Med*. 1987;317:1105-1108.
107. Romero LS, Hunt SJ. Hickman catheter-associated primary cutaneous aspergillosis in a patient with the acquired immunodeficiency syndrome. *Int J Dermatol*. 1995;34:551-553.
108. Bryce EA, Walker M, Scharf S, et al. An outbreak of cutaneous aspergillosis in a tertiary-care hospital. *Infect Control Hosp Epidemiol*. 1996;17:170-172.
109. Stone HH, Cuzzell JZ, Kolb LD, Moskowitz MS, McGowan JE. Aspergillus infection of the burn wound. *J Trauma*. 1979;19:765-767.
110. Walsh TJ. Primary cutaneous aspergillosis—an emerging infection among immunocompromised patients. *Clin Infect Dis*. 1998;27:453-457.
111. Steinbach WJ. Pediatric invasive aspergillosis. *Pediatr Infect Dis J*. 2010;29:964-965.
112. Walsh TJ, Anaissie EJ, Denning DW, et al. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis*. 2008;46:327-360.
113. Marr KA, Carter RA, Boeckh M, Martin P, Corey L. Invasive aspergillosis in allogeneic stem cell transplant recipients: changes in epidemiology and risk factors. *Blood*. 2002;100:4358-4366.
114. Gil L, Kozłowska-Skrzypczak M, Mol A, Poplawski D, Styczynski J, Komarnicki M. Increased risk for invasive aspergillosis in patients with lymphoproliferative diseases


- after autologous hematopoietic SCT. *Bone Marrow Transplant*. 2009;43:121-126.
115. Kim YW, Lee HW, Cho J, et al. Conversion of aspergilloma to chronic necrotizing pulmonary aspergillosis following treatment with sunitinib: a case report. *Oncol Lett*. 2016;12:3472-3474.
 116. Kwon-Chung KJ. Taxonomy of fungi causing mucormycosis and entomophthoromycosis (zygomycosis) and nomenclature of the disease: molecular mycologic perspectives. *Clin Infect Dis*. 2012;54(suppl 1):S8-S15.
 117. Skiada A, Rigopoulos D, Larios G, Petrikos G, Katsambas A. Global epidemiology of cutaneous zygomycosis. *Clin Dermatol*. 2012;30:628-632.
 118. Ibrahim AS, Spellberg B, Edwards J. Iron acquisition: a novel perspective on mucormycosis pathogenesis and treatment. *Curr Opin Infect Dis*. 2008;21:620-625.
 119. Kennedy KJ, Daveson K, Slavin MA, et al. Mucormycosis in Australia: contemporary epidemiology and outcomes. *Clin Microbiol Infect*. 2016;22:775-781.
 120. Mohindra S, Gupta R, Bakshi J, Gupta SK. Rhinocerebral mucormycosis: the disease spectrum in 27 patients. *Mycoses*. 2007;50:290-296.
 121. Rahman A, Akter K, Hossain S, Rashid HU. Rhino-orbital mucormycosis in a non-immunocompromised patient. *BMJ Case Rep*. 2013;2013.
 122. Austin CL, Finley PJ, Mikkelsen DR, Tibbs B. Mucormycosis: a rare fungal infection in tornado victims. *J Burn Care Res*. 2014;35:e164-e171.
 123. Tribble DR, Warkentien T, Rodriguez C, Trauma Infectious Diseases Outcomes Study Group of the Infectious Disease Clinical Research Program. Mucormycosis after a tornado in Joplin, Missouri. *N Engl J Med*. 2013;368:1067.
 124. Duffy J, Harris J, Gade L, et al. Mucormycosis outbreak associated with hospital linens. *Pediatr Infect Dis J*. 2014;33:472-476.
 125. Rammaert B, Lanternier F, Zahar JR, et al. Healthcare-associated mucormycosis. *Clin Infect Dis*. 2012;54(suppl 1):S44-S4.
 126. Kontoyiannis DP, Marr KA, Park BJ, et al. Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001-2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis*. 2010;50:1091-1100.
 127. Connor BA, Anderson RJ, Smith JW. Mucor mediastinitis. *Chest*. 1979;75:525-526.
 128. Spellberg B. Gastrointestinal mucormycosis: an evolving disease. *Gastroenterol Hepatol (N Y)*. 2012;8:140-142.
 129. Kontoyiannis DP, Lewis RE. Invasive zygomycosis: update on pathogenesis, clinical manifestations, and management. *Infect Dis Clin North Am*. 2006;20:581-607.
 130. Kontoyiannis DP. Invasive mycoses: strategies for effective management. *Am J Med*. 2012;125(1 suppl):S25-38.
 131. Nelson PE, Dignani MC, Anaissie EJ. Taxonomy, biology, and clinical aspects of Fusarium species. *Clin Microbiol Rev*. 1994;7:479-504.
 132. Anaissie EJ, Kuchar RT, Rex JH, et al. Fusariosis associated with pathogenic fusarium species colonization of a hospital water system: a new paradigm for the epidemiology of opportunistic mold infections. *Clin Infect Dis*. 2001;33:1871-1878.
 133. Krátká J, Kováčiková E. The effect of temperature and age of strains of Fusarium oxysporum on its enzymatic activity. *Zentralbl Bakteriol Naturwiss*. 1979;134:154-158.
 134. Assi MA, Sandid MS, Baddour LM, Roberts GD, Walker RC. Systemic histoplasmosis: a 15-year retrospective institutional review of 111 patients. *Medicine (Baltimore)*. 2007;86:162-169.
 135. Padoan R, Poli P, Colombrita D, Borghi E, Timpano S, Berlucchi M. Acute scedosporium apiospermum endobronchial infection in cystic fibrosis. *Pediatr Infect Dis J*. 2016;35:701-702.
 136. Castiglioni B, Sutton DA, Rinaldi MG, Fung J, Kusne S. *Pseudallescheria boydii* (anamorph *Scedosporium apiospermum*). Infection in solid organ transplant recipients in a tertiary medical center and review of the literature. *Medicine (Baltimore)*. 2002;81:333-348.
 137. Caira M, Girmenia C, Valentini CG, et al. Scedosporiosis in patients with acute leukemia: a retrospective multicenter report. *Haematologica*. 2008;93:104-110.
 138. Houbraken J, Varga J, Rico-Munoz E, Johnson S, Samson RA. Sexual reproduction as the cause of heat resistance in the food spoilage fungus *Byssoschlamys spectabilis* (anamorph *Paecilomyces variotii*). *Appl Environ Microbiol*. 2008;74:1613-1619.
 139. Mehta V. Cutaneous hyalohyphomycosis in an immunocompetent host. *Indian J Dermatol*. 2013;58:411.
 140. McClellan JR, Hamilton JD, Alexander JA, Wolfe WG, Reed JB. *Paecilomyces varioti* endocarditis on a prosthetic aortic valve. *J Thorac Cardiovasc Surg*. 1976;71:472-475.
 141. Pettit TH, Olson RJ, Foos RY, Martin WJ. Fungal endophthalmitis following intraocular lens implantation. A surgical epidemic. *Arch Ophthalmol*. 1980;98:1025-1039.
 142. Lavergne RA, Cassaing S, Nocera T, et al. Simultaneous cutaneous infection due to *Paecilomyces lilacinus* and *Alternaria* in a heart transplant patient. *Transpl Infect Dis*. 2012;14:E156-E160.
 143. Saegeman VS, Dupont LJ, Verleden GM, Lagrou K. *Paecilomyces lilacinus* and *alternaria* infectoria cutaneous infections in a sarcoidosis patient after double-lung transplantation. *Acta Clin Belg*. 2012;67:219-221.
 144. Taborda CP, da Silva MB, Nosanchuk JD, Travassos LR. Melanin as a virulence factor of *Paracoccidioides brasiliensis* and other dimorphic pathogenic fungi: a minireview. *Mycopathologia*. 2008;165:331-339.
 145. Sharma NL, Sharma VC, Mahajan V, Shanker V, Sarin S. Chromoblastomycosis with underlying osteolytic lesion. *Mycoses*. 2007;50:517-519.
 146. Keyser A, Schmid FX, Linde HJ, Merk J, Birnbaum DE. Disseminated *Cladophialophora bantiana* infection in a heart transplant recipient. *J Heart Lung Transplant*. 2002;21:503-505.
 147. Petrini B, Farnebo F, Hedblad MA, Appelgren P. Concomitant late soft tissue infections by *Cladophialophora bantiana* and *Mycobacterium abscessus* following tsunami injuries. *Med Mycol*. 2006;44:189-192.
 148. Balla A, Pierson J, Hugh J, Wojewoda C, Gibson P, Greene L. Disseminated cutaneous *Curvularia* infection in an immunocompromised host; diagnostic challenges and experience with voriconazole. *J Cutan Pathol*. 2016;43:383-387.
 149. Rohwedder JJ, Simmons JL, Colfer H, Gatmaitan B. Disseminated *Curvularia lunata* infection in a football player. *Arch Intern Med*. 1979;139:940-941.
 150. Frank T, Esquenazi Y, Nigo M, Wanger A, Portnoy B, Shepard S. Disseminated phaeohyphomycosis with brain abscess and biliary invasion due to *Bipolaris* spp. in an immunocompetent patient. *Ann Clin Lab Sci*. 2016;46:439-442.
 151. Atalay MA, Koç AN, Koyuncu S, Ulu Kiliç A, Kurtsoy A, Alp Meşe E. *Cladophialophora bantiana* brain abscess treated with voriconazole in an immunocompetent patient [in Turkish]. *Mikrobiyol Bul*. 2014;48:501-506.




Supplemental Fig 1. Disseminated *Candida spp* infection, clinically characterized by several red papules and pustules in an immunosuppressed patient.




Supplemental Fig 2. Disseminated *Aspergillus spp* presenting as a firm nodule on the right thigh.



Supplemental Fig 3. Postmortem diagnosis of disseminated *Mucormycosis*. Involvement of the gastrointestinal tract resulted in hemorrhagic necrosis of the (A) stomach and (B) spleen. Photograph courtesy of Amanda Barrett, MD, and Stuti Shroff, MD.



Supplemental Fig 4. *Mucormycosis spp* can present in a variety of ways, including red papules on the foot.



Supplemental Fig 5. *Fusarium* infection presenting in the skin. **A**, On the left wrist, infection presented as an erythematous nodule with central eschar. **B**, An erythematous nodule with overlying bullae on the right great toe.