

Concentrations of melatonin, thyroxine, 17 β -estradiol and 11-ketotestosterone in round goby (*Neogobius melanostomus*) in different phases of the reproductive cycle

Tatiana Guellard, Hanna Kalamarz-Kubiak*, Ewa Kulczykowska

Institute of Oceanology, Polish Academy of Sciences, Sopot, Poland

ARTICLE INFO

Keywords:

Melatonin
Thyroxine
Gonadal steroids
Round goby
Reproductive cycle

ABSTRACT

The aim of this study was to determine changes in concentrations of melatonin (Mel) and thyroxine (T₄) in plasma, and 17 β -estradiol (E₂) and 11-ketotestosterone (11-KT) in plasma and gonads of female and male round gobies (*Neogobius melanostomus*) from the Southern Baltic Sea in four phases of the reproductive cycle classified as pre-spawning, spawning, late spawning and non-spawning periods. The concentrations of Mel, T₄ and E₂ were determined by radioimmunoassay (RIA) whereas 11-KT was quantified using an enzyme immunoassay (EIA). The maturity stage of gonads was determined using histological analyses. The pattern of changes in Mel concentrations of females and males was similar with the greatest concentrations in the spawning and non-spawning phases. In both sexes, there was a similar tendency of change in concentrations of T₄ and E₂ with the increase being in the pre-spawning and non-spawning phases. The greatest concentrations of 11-KT were observed in the plasma and gonads of males during the spawning phase. In females, there were no changes in 11-KT concentrations either in plasma or gonads during all phases where quantifications occurred. This is the first study for determination of the pattern of changes in Mel and T₄ concentrations as well as gonadal steroids E₂ and 11-KT, supported by histological analysis of gonads, in batch spawning fish during the reproductive cycle.

1. Introduction

Since the 1990s, the round goby (*Neogobius melanostomus*) became a significant invader in several regions of Europe (including the Baltic Sea) and in the Laurentian Great Lakes of North America (Kornis et al., 2012). In the Baltic Sea, this batch spawning species is most abundant in the Gulf of Gdańsk, where the spawning season lasts from the beginning of May to the end of September and females may lay eggs from two to four times during the season (Śapota, 2012).

Although considerable research on the biology of round goby has been conducted (Kornis et al., 2012), the hormonal regulation of the reproductive cycle of this species has received little attention. It is well recognized that sex steroids have important functions in fish reproduction, and there are also other hormones such as melatonin (Mel) and thyroxine (T₄) that are essential for regulation of reproductive cycle (Borg, 1994; Falcón et al., 2010; Leatherland, 1994; Lubzens et al., 2010). To the authors' knowledge, there is no study on Mel and T₄ in round goby. In fish reproduction, Mel has been reported to be an important hormone in regulation of occurrence of spawning, gametogenic and steroidogenic functions of gonads (Popek et al., 1991; Sokołowska et al., 2004; Maitra

* Corresponding author at: Institute of Oceanology, Polish Academy of Sciences, Powstańców Warszawy 55, 81-712, Sopot, Poland.
E-mail address: hkalamarz@iopan.gda.pl (H. Kalamarz-Kubiak).

et al., 2006). These actions of Mel occur as a result of effects at the brain – pituitary – gonad axis (Falcón et al., 2007; Popek et al., 2010; Maitra et al., 2013). This hormone transduces seasonal temperature and photoperiod signals that are perceived via sensory neurons that have an important role in synchronizing the stage of different biological rhythms also involved in reproduction (Ekström and Meissl, 1989; Falcón et al., 2010; Maitra et al., 2013). Melatonin can also protect cells from oxidative stress in the processes of the maturation and regression of gonads in an annual cycle (Bromage et al., 2001; Bayarri et al., 2004; Falcón et al., 2010; Maitra et al., 2013, 2015; Alvarado et al., 2015; Maitra and Hasan, 2016). In fish, T_4 has wide-ranging effects on gonadal development and maturation, stimulating and maintaining the functions during the reproductive cycle (Biswas et al., 2006; Swapna and Senthilkumaran, 2007). In females, T_4 stimulates the development of previtellogenic oocytes and early oocyte vitellogenesis (Wiegand, 1982; Cyr and Eales, 1988), up-regulates hepatic synthesis of vitellogenin - the protein sequestered by growing oocytes (Monteverdi and Giulio, 2000) that induces final oocyte maturation and ovulation (Hurlburt, 1977; Detlaff and Davydova, 1979). In males, T_4 modulates the actions of many factors that are important for maintaining spermatogenesis and spermiation (Sower and Schreck, 1982; Leatherland, 1994; Swapna and Senthilkumaran, 2007; Flood et al., 2013).

Furthermore, there are only a few studies focused on the sex steroids in round goby. There have been results reported describing the 17β -estradiol (E_2) and 11-ketotestosterone (11-KT) concentrations in non-reproductive and reproductive males and females and two different male reproductive morphs, namely: parental and sneaker males (Marentette et al., 2009; Bowley et al., 2010; Zeyl et al., 2014). In female fish, E_2 , which is produced mainly by the ovarian follicles, has an important role in supporting the hepatic synthesis and secretion of vitellogenin (Lubzens et al., 2010). There is also an essential function of E_2 in male reproduction (Borg, 1994; Schulz et al., 2010). There is synthesis of E_2 by Leydig and Sertoli cells in fish testes (Janz and Weber, 2000) and this hormone affects the regulation of spermatogenesis, especially spermatogonial proliferation (Schulz et al., 2010). In most teleosts males, 11-KT is the primary androgen, stimulating spermatogenesis and regulating growth and development of testis as well as secondary sexual characteristics and reproductive behaviour (Borg, 1994; Schulz et al., 2010). The function of 11-KT in females is not fully understood but *in vivo* and *in vitro* studies of short-finned eel (*Anguilla australis*; Rohr et al., 2001; Lokman et al., 2007) have reported that 11-KT contributes in oocyte growth.

The objective of the present study was to investigate the changes in Mel, T_4 , E_2 and 11-KT in four phases of the reproductive cycle of round goby characterized as the pre-spawning, spawning, late spawning and non-spawning phases. In addition to analyses of hormones, there were histological studies of gonads as a part of this study.

2. Materials and methods

2.1. Fish

Adult round gobies of both sexes ($n = 84$) were caught using fyke nets, near Hel Harbour ($54^\circ 36' 04.17''N$, $18^\circ 47' 56.06''E$) (Gulf of Gdańsk, southern Baltic Sea) in March, April, July and October. After capture, the fish were transferred to the Marine Station (Institute of Oceanography, University of Gdańsk, Poland) in Hel and were sexed by examination of the urogenital papilla. Each group of 21 individuals of similar size consisted of 14 males and seven females and were acclimated to the tank conditions for 6 weeks. During this time, fish were maintained in natural photoperiodic and temperature conditions (Table 1) in a 2640 L outdoor tank with an open circuit water system providing constant flow through of fresh brackish water (~7% salt) from the Gulf of Gdańsk. The temperature in the outdoor tank was measured using a portable meter CC-401 (Elmetron, Poland) every day at noon. The measurements were taken in the middle of the tank water column. In each tank, devices were inserted that facilitated fish seclusion that were made from PVC pipes so as to mimic natural living conditions of round goby. Fish were fed once daily with fresh fish meat (herring), frozen mussels or shrimps at 3:00 p.m. After the acclimatisation, fish were anaesthetized in MS-222 (tricaine methanesulfonate) water buffered solution (0.1 g/L; Sigma-Aldrich, USA). Sampling occurred between 10:00 a.m. and 2:00 p.m. The length of each individual was measured and weighing occurred. Blood samples were collected by cardiac puncture, using a heparinized syringe. The plasma samples were centrifuged at 3000 g for 10 min at $4^\circ C$ and frozen at $-70^\circ C$ until analyses. After blood sampling, fish were euthanized by transection of the spinal cord. Gonads were collected and one gonad of each individual was frozen at $-70^\circ C$ and stored until steroid analysis while the second gonad was preserved in 4% buffered formalin for histology. There were histological analyses of ovaries and testes for characterization of sexual maturity stage because the gonadosomatic index (GSI) is apparently a less reliable assessment of stage of sexual maturation in batch spawners (Guerriero, 2007; Zeyl et al., 2014).

Table 1

Average day length and range of water temperature during reproductive phases.

Phase	Day length [h]	Water temperature [$^\circ C$]
Pre-spawning	13	4.8 ± 0.5
Spawning	15	11.5 ± 2.5
Late spawning	14	19.6 ± 1.2
Non-spawning	10	11.0 ± 1.3

2.2. Analytical methods

2.2.1. Plasma analysis

Plasma Mel was assayed using a total melatonin radioimmunoassay (RIA) kit (IBL International, Germany), after conducting an extraction procedure using the previously reported method of [Kulczykowska et al. \(2007\)](#). The solid phase extraction was conducted using a 10 μ m Octadecyl C18 Speedisk Column, (J.T. Baker, USA). Eluates obtained after extraction were dried and then stored at -70°C until analysis. Before the RIA procedure, dried samples were re-suspended in Dulbecco's Phosphate Buffered Saline containing 0.01% Thimerosal (Sigma-Aldrich, USA). The labelled Mel with iodine-125 (^{125}I) was used as a tracer for this analysis. The assay was performed using polyclonal rabbit anti-Mel first antibodies and goat anti-rabbit second antibodies (double antibody method). The detection limit was 3.0 pg/mL of plasma. The intra-assay coefficients of variation were 6.5%.

There was quantification of plasma T_4 using a total thyroxine RIA-gnost kit (Cisbio Bioassays, France) without extraction procedures being conducted the method previously reported by [Kulczykowska et al. \(2007\)](#). The assay was performed using tubes coated with murine anti- T_4 monoclonal antibodies and 8-aniline-1-sulfonic acid as a displacement reagent. The T_4 labelled with ^{125}I was used as a tracer for RIA. The detection limit was 1.1 ng/mL of plasma. The intra-assay coefficients of variation were 5.6%.

Plasma E_2 was measured using the ESTR-CTRIA RIA kit (Cisbio Bioassays, France) without conducting extraction procedures according to the manufacturer's protocol validated in our laboratory ([Kalamarz-Kubiak et al., 2017](#)). Iodinated E_2 with ^{125}I was used as a tracer. The assay was conducted in RIA tubes pre-coated with polyclonal anti-rabbit antiserum according to the kit manufacturer's instructions. The detection limit of the assay was 4.25 pg/mL. The intra-assay coefficient of variation was 2.1%.

Plasma 11-KT was quantified using the competitive enzyme immunoassay (EIA) Cayman kit (Ann Arbor, USA) with extraction procedures occurring before conducting the assay following the methods previously reported by [Sokołowska et al. \(2013\)](#). Before analysis, extraction of plasma samples (100 μ L) was conducted using ethyl acetate/hexane (50:50 v/v) at $4 \times$ the sample volume utilizing the method recommended in the EIA kit protocol with slight modifications. Dried extracts were stored at -20°C until further analysis. The recovery of extraction was between 98% and 115%. Extracts were dissolved in 2 mL of EIA buffer and 50 μ L of the diluted samples were used for the EIA analysis. The 11-KT-acetylcholinesterase conjugate was used as a tracer. The assay was performed in the pre-coated microplate with mouse anti-rabbit IgG using rabbit anti-11KT antibodies. The microplate was read at 412 nm using a Sunrise Absorbance Reader (Tecan, Austria). All samples were assayed in duplicate. The detection limit of the assay was 0.7 pg/mL. The intra-assay coefficient of variation was 0.6%. For the RIA, all samples were assayed in duplicate and counted for 1 min in a Wallac Wizard 1470 gamma counter (Perkin Elmer Life Science, USA). The inter-assay variations for plasma analysis were not determined because all samples were measured in the same assay.

2.2.2. Analysis in gonads

Each gonad was sonicated in 0.5 mL of phosphate buffer (0.05 M, pH 7.4) supplemented with sodium azide (NaN_3) using a Microson[™] XL 2000 (USA). Sonicated samples were centrifuged at 20,000 g for 20 min at 4°C and the supernatants stored at -70°C prior to the quantification of E_2 and 11-KT concentrations.

The E_2 concentrations in gonads' extracts were measured using ESTR-CTRIA RIA kit (Cisbio Bioassays, France) according to the method by [Kulczykowska et al. \(2015\)](#). Gonad supernatants (200 μ L) were extracted with 1.6 mL of diethyl ether using the modified methods of [Mori and Kano \(1984\)](#). The recovery rate of the extraction was between 86% and 109%. Dried extracts were stored at -20°C prior to analysis. Extracts were dissolved in phosphate buffer (0.05 M, pH 7.4) supplemented with NaN_3 and samples of 100 μ L were taken for RIA analysis. Further analysis was performed according to the method described for plasma E_2 quantifications (see paragraph 2.2.1.). The detection limit of the assay was 4.25 pg/mL. The intra-assay coefficient of variation was 6.5%.

The 11-KT concentrations in gonads' extracts were determined by EIA using Cayman kit (Ann Arbor, USA) with the extraction procedure described by [Kulczykowska et al. \(2015\)](#). Gonad supernatants (250 μ L) were extracted with 1 mL of ethyl acetate/hexane (50:50 v/v) according to the method recommended in the EIA kit protocol with slight modifications. The recovery of extraction was between 98% and 115%. Dried extracts were stored at -20°C until further analysis. Extracts were dissolved in 0.5 mL of EIA buffer and 50 μ L of the diluted samples were used for the EIA analysis. Further analysis was performed using the methods described for plasma 11-KT quantification (see Section 2.2.1). The detection limit of the assay was 1.092 pg/mL. The intra-assay coefficient of variation was 0.8%. The inter-assay variations for gonads analysis were not determined because there were quantifications for all samples in the same assay.

2.3. Histological analysis of gonads

Gonads were stored in 4% buffered formalin then dehydrated in graded ethanol and embedded in paraffin using standard histological techniques. Sections of 6 μ m were cut using a Leica RM2245 (Germany) microtome and stained with haematoxylin and eosin. Slides from each gonad were examined using a Leica H11210 light microscope (Leica Microsystems, Germany). The developmental stage of gonads was adapted according to [Rocha and Rocha \(2006\)](#) and [Brown-Peterson et al. \(2011\)](#).

2.4. Statistical analysis

Statistical analyses of data were conducted using STATISTICA 13 software. The data were expressed as means $\pm$ SEM. Significance was accepted at $P < 0.05$. For multiple comparisons of hormone concentrations (MEL, T_4 , E_2 and 11-KT), analysis of variance (one-way ANOVA) was performed followed by *post hoc* tests (Sjotvoll and Stoline's test for unequal numbers of cases or LSD




Fig. 1. Concentration of plasma melatonin in round goby females (A) and males (B) in different phases of the reproductive cycle; Data are presented as mean $\pm$ SEM, numbers of individuals are given in circles; Different superscripts indicate statistical differences among phases ($P < 0.05$).

test, and NIR test; as required). For comparisons of all hormone concentrations in plasma and gonads between males and females, Student's *t*-test was used.

2.5. Ethical note

All procedures performed in this study complied with the EC Directive 2010/63/EU for animal experiments and with the guidelines of the Local Ethics Committee on Animal Experimentation (Resolution No. 33/2010).

3. Results

3.1. Analysis of hormones

The pattern of changes in plasma Mel concentrations was similar in both sexes (Fig. 1). In females, there were the greatest concentrations of Mel in the spawning and non-spawning phases (Fig. 1). In males, the greatest concentration of Mel was during the spawning phase (Fig. 1). The differences in Mel concentration between females and males were observed only in the spawning phase ($P < 0.05$; Table 2).

The concentrations of plasma T_4 was of a similar pattern during the various reproductive phases in females and males (Fig. 2). In the non-spawning phase, in both sexes, there were the greatest concentrations of T_4 (Fig. 2). There were greater concentrations of T_4 in males and females in the pre-spawning phase (Fig. 2). In both sexes, there were the least concentrations of T_4 in the late spawning phase (Fig. 2A–B). There were differences in T_4 between females and males only during the non-spawning phase ($P < 0.05$; Table 2).

The patterns of E_2 changes in plasma and gonads were similar in both sexes (Fig. 3A–D). In females, there were the greatest concentrations of plasma E_2 in the pre-spawning and non-spawning phases (Fig. 3A). In ovaries, there was the greatest concentration of E_2 in the pre-spawning phase (Fig. 3B). In males, there were the greatest E_2 concentrations in plasma and gonads in the pre-spawning and non-spawning phases (Fig. 3C–D). The concentrations of E_2 in plasma and gonads of females were greater than those in males in all reproductive phases (Fig. 3A–D, Table 2).

Table 2

Statistical differences (*P*-values) in hormones between males and females of round goby in different phases of the reproductive cycle; significance was accepted at $P < 0.05$.

	Phase			
	pre-spawning	spawning	late spawning	non-spawning
<i>P</i>				
Plasma				
melatonin	0.88	< 0.05	0.93	0.32
thyroxine	0.20	0.13	0.94	< 0.05
17 β -estradiol	< 0.001	< 0.001	< 0.01	< 0.001
11-ketotestosterone	< 0.001	< 0.001	< 0.05	< 0.01
Gonads				
17 β -estradiol	< 0.001	< 0.001	< 0.01	< 0.01
11-ketotestosterone	< 0.01	< 0.001	< 0.05	< 0.001




Fig. 2. Concentration of plasma thyroxine in round goby females (A) and males (B) in different phases of the reproductive cycle; Data are presented as mean $\pm$ SEM, numbers of individuals are given in circles; Different superscripts indicate statistical differences among phases ($P < 0.05$).




Fig. 3. Concentrations of 17 β -estradiol in plasma and gonads of round goby females (A–B) and males (C–D) in different phases of the reproductive cycle Data are presented as mean $\pm$ SEM, numbers of individuals are given in circles; Different superscripts indicate statistical differences among phases ($P < 0.05$).

In females, 11-KT concentrations in plasma and gonads did not differ during the various reproductive phases (Fig. 4A–B). In males, there was a similar pattern of 11-KT in plasma and gonads with the greatest concentrations during spawning (Fig. 4C–D). The concentrations of 11-KT in plasma and gonads of males were greater than those of females during all reproductive phases (Fig. 3A–D; Table 2).




Fig. 4. Concentrations of 11-ketotestosterone in plasma and gonads of round goby females (A–B) and males (C–D) in different phases of the reproductive cycle. Data are presented as mean $\pm$ SEM, numbers of individuals are given in circles; Different superscripts indicate statistical differences among phases ($P < 0.05$).

3.2. Histological analysis

On the basis of histological analysis of ovaries and testes, the following phases of the reproductive cycle in round goby were distinguished: pre-spawning, spawning, late spawning and non-spawning.

Ovaries of round gobies collected during the pre-spawning phase contained mostly follicles at the late stage of vitellogenesis (VTG), before there was germinal vesicle migration (GVM), and oocytes present at the previtellogenic stage (Fig. 5A). Ovaries collected in the spawning phase were characterized by the large number of oocytes at the early stage of GVM. The presence of postovulatory follicles (POFs) indicated these individuals were in the active spawning stage at the time of tissue collection. The ovaries also contained the developing batches of oocytes at the early stage of vitellogenesis (eVTG) and previtellogenic stage (PG), which is characteristic for batch spawners (Fig. 5C). In the late spawning phase, there were large numbers of POFs in the ovaries. Oocytes during GVM, eVTG and PG were also detected (Fig. 5E). During non-spawning phase, ovaries were in regression and contained mostly vitellogenic oocytes in the early atresia stage (A). In this phase, oocytes at the eVTG and PG stages were also present (Fig. 5G).

In the pre-spawning phase, maturing testes contained spermatogonia (SG) - an undifferentiated germ cell (Fig. 5B). In the spawning phase, testes were undergoing active spermatogenesis as indicated by the presence of large numbers of spermatozoa (SZ). Spermatogonial cells, SG and spermatocytes (SC), were also clearly visible during this phase (Fig. 5D). In the late spawning phase, testes were fully matured, filled with SZ (Fig. 5F). In the non-spawning phase, testes were in regression and SGs were the dominant cells. It was also noted that there were residual SZ in the lumen of narrowing seminiferous tubules (ST; Fig. 5H).

4. Discussion

The present study is the first attempt to document the changes in concentrations of plasma Mel and T_4 and plasma and gonadal E_2 and 11-KT as well as with histology of gonads in both sexes of round goby during four different phases of the reproductive cycle: pre-spawning, spawning, late spawning and non-spawning.




Fig. 5. Structures of ovaries and testes in round goby in different phases of the reproductive cycle. Pre-spawning phase: (A) Ovary with developing oocytes in the late stage of vitellogenesis (VTG); (B) Maturing testis with the dominance of spermatogonia (SG). Spawning phase: (C) Partially spent ovary (postovulatory follicles; POFs) with the next developing batches of oocytes in the early stage of maturation (germinal vesicle migration; GVM), early stage of vitellogenesis (eVTG) and previtellogenic stage (PG); (D) Testis under active spermatogenesis with visible spermatocytes (SC) and spermatozoa (SZ) in seminiferous tubules (ST). Late spawning phase: (E) Ovary with the dominance of POFs and the presence of GVM, eVTG and PG; (F) Fully developed testis with seminiferous tubules (ST) filled with SZ. Non-spawning phase: (G) Regressing ovary with the dominance of atretic vitellogenic oocytes (A); (H) regressing testis with the dominance of SG and residual SZ in the lumen of ST. Scale bars correspond to 500 µm.

4.1. Mel

In the present study, the plasma Mel concentrations were of a similar pattern in round goby females and males throughout the four phases where there were assessments. There were similar trends in the batch spawning common carp (*Cyprinus carpio*; [Kezuka et al., 1988](#)) and single spawning pacu (*Piaractus mesopotamicus*; [Iseki et al., 2000](#)). In round goby females, there were greater concentrations of plasma Mel in the pre-spawning phase. These findings are consistent with those of [Maitra et al. \(2013\)](#), where there was a greater concentration of Mel during the pre-spawning phase in the single spawning major carp (*Catla catla*) females. There were the greatest concentrations of Mel in the present study during the spawning phase of round goby females. At this phase, Mel is necessary for maturation processes and to protect the oocytes from free radicals during periods of rapid cell proliferation as has been reported to occur in the batch spawning mummichog (*Fundulus heteroclitus*; [Lombardo et al., 2012](#)) and zebrafish (*Danio rerio*; [Carnevali et al., 2011](#)) as well as in the single spawning major carp ([Maitra and Hasan, 2016](#); [Mondal et al., 2017](#)). In the late-spawning phase, there were the least Mel concentrations in the present study, and the POFs were in predominant numbers compared with the numbers of oocytes that were in relatively greater numbers during other stages of development. The subsequent increase of Mel concentrations occurred in the non-spawning phase while there was ovarian regression occurring. There have been similar previous observations in the batch spawning three-spined stickleback (*Gasterosteus aculeatus*; [Sokołowska et al., 2004](#)), where there was an inhibitory effect on gonads. It resulted in gonadal regression that was attributed to having occurred because of the relatively greater Mel concentrations during this phase. Furthermore, during this phase, there was follicular atresia with the production of free radicals being a potential cause for the increased concentrations of Mel and other antioxidants as previously reported ([Maitra and Hasan, 2016](#); [Mondal et al., 2017](#)) in major carp.

In round goby males of the present study, there was a greater concentration of Mel in the pre-spawning phase when testes were undergoing early maturation. Data from the present study are consistent with previous findings when there was study of the batch spawner three-spined stickleback ([Borg and Ekström, 1981](#)) and single spawner masu salmon (*Oncorhynchus masou*; [Amano et al., 2000](#)), where Mel stimulated gonadal development during the spring of the year. In round goby males, there was the greatest Mel plasma concentration in the spawning phase. Results of studies conducted in fish indicate that relatively greater Mel concentrations enhance the maturation of testes and sperm quality in a single spawning broadhead catfish (*Clarias macrocephalus*; [Aripin et al., 2014](#)). Therefore, the greater concentrations of Mel, in round goby males may also function as a defense against free radicals during the period of rapid cell proliferation that accompanies spermiogenesis, as it was presented in mammals ([Gavella and Lipovac, 2000](#); [Shang et al., 2004](#)). In the present study, in the late spawning phase, when testes were completely matured, there was the least concentration of Mel. The decrease in Mel concentration at the end of spermiogenesis has also been reported in the three-spined stickleback ([Sokołowska et al., 2004](#)). There was a second increase in plasma Mel concentration in round goby males during the non-spawning phase when testes were undergoing regression. There has been a similar increase in Mel concentration observed after spawning in three-spined stickleback ([Sokołowska et al., 2004](#)) and in single spawning major carp ([Bhattacharya et al., 2007](#)). It has been suggested that in fish, relatively greater Mel concentrations may have an inhibitory effect on gonadotropins and induce regression of gonads ([Fenwick, 1970](#); [Maitra and Hasan, 2016](#)). Probably, the increase in Mel concentration during this reproductive phase was also the result of free radical production during the regression of testicular tissue ([Maitra and Hasan, 2016](#)).

4.2. T₄

In the pre-spawning phase, there was a relatively greater concentration of T₄ in plasma of round goby females. Greater plasma concentrations of T₄ have also been reported during the pre-spawning phase in batch spawning of female goldfish (*Carassius auratus*; [Hurlburt, 1977](#)), plaice (*Pleuronectes platessa*; [Osborn and Simpson, 1978](#)), common dentex (*Dentex dentex*; [Pavlidis et al., 2000](#)) and Atlantic cod ([Comeau and Campana, 2006](#)). There have been reports that T₄ increases the growth rate of ovarian follicles and the number of vitellogenic follicles in this phase ([Sage, 1973](#); [Cyr and Eales, 1988](#); [Legler et al., 2000](#); [Supriya et al., 2005](#)). In the spawning and late spawning phase, concentrations of T₄ in plasma of round goby females were basal. There have been similar reports of basal plasma T₄ during this reproductive phase in the batch spawning plaice ([Osborn and Simpson, 1978](#)), common dentex ([Pavlidis et al., 2000](#)) and Atlantic cod ([Comeau and Campana, 2006](#)). There was the greatest concentration of plasma T₄ in round goby females during non-spawning phase when the ovaries contained primarily vitellogenic oocytes undergoing atresia. There has been a similar post-reproductive surge of T₄ observed in such batch spawners as plaice ([Osborn and Simpson, 1978](#)), common dentex ([Pavlidis et al., 2000](#)) and Atlantic cod ([Comeau and Campana, 2006](#)). It was reported that the increase of T₄ in fish occurring after spawning might be associated with the function of T₄ in enhancing development of previtellogenic oocytes still persisting in regressing ovaries ([Pavlidis et al., 2000](#)) and also with ongoing steroidogenesis in the atretic vitellogenic oocytes ([Cyr and Eales, 1988](#); [Rinchard et al., 1993](#)).

In males of round goby, there was a similar trend in changes of plasma T₄ concentration as in females in the present study. The decreasing concentrations of T₄ after the pre-spawning phase, and during the spawning to late spawning phases is a pattern that is consistent with progress in spermatogenesis. There have been reports of similar results for other species such as the batch spawning plaice ([Osborn and Simpson, 1978](#)) and striped bass (*Morone saxatilis*; [Mylonas et al., 1997](#)) or single spawning brown bullhead (*Ictalurus nebulosus*; [Burke and Leatherland, 1983](#)). In the present study, there was the greatest concentration of plasma T₄ during the non-spawning phase when testes were undergoing regression. There was also a similar increase in T₄ concentrations during this reproductive phase in plaice ([Osborn and Simpson, 1978](#)). Furthermore, [Habibi et al. \(2012\)](#) have reported that T₄ inhibits steroidogenesis in the batch spawning goldfish.

4.3. E_2

In the pre-spawning phase of the present study, when a large population of the oocytes commenced active vitellogenesis, there were the greatest concentration of E_2 in plasma and gonads of round goby females. There were relatively greater plasma E_2 concentrations during the pre-spawning phase in batch spawning females of the goldfish (Kagawa et al., 1983a), Atlantic halibut (*Hippoglossus hippoglossus*; Methven et al., 1992), tench (*Tinca tinca*; Pinillos et al., 2003), chub (*Leuciscus cephalus*; Guerriero, 2007), kutum (*Rutilus frisii kutum*; Sabet et al., 2009) and river catfish (*Hemibagrus nemurus*; Adebiyi et al., 2013). Furthermore, there have been *in vivo* and *in vitro* studies reported describing a relatively greater amount of E_2 production by vitellogenic oocytes during the pre-spawning phase in females of the batch spawning tilapia (*Oreochromis mossambicus*; Cornish, 1998), goldfish (Kagawa et al., 1984) and single spawning rainbow trout (*Oncorhynchus mykiss*; van Bohemen and Lambert, 1981). In the spawning phase, E_2 concentrations were less in plasma and ovaries of round goby. There are many batch spawning fish species such as the goldfish (Kagawa et al., 1983a), sea bass (Prat et al., 1990), gudgeon (Rinchard et al., 1993), Jundiá (*Rhamdia quelen*; Barcellos et al., 2001) and gilthead seabream (*Sparus aurata*; Kadmon et al., 1985) in which plasma E_2 concentration is basal during spawning. In round goby, as well as in other batch spawners during the spawning phase, besides oocytes at early stages of vitellogenesis and maturation, the ovaries contain POFs (Kagawa et al., 1984; Rinchard et al., 1993; Barcellos et al., 2001). The POFs, like other oocytes occurring in ovaries during this phase, produce also a very small amount of E_2 (Kagawa et al., 1983b), which could explain the relatively lesser E_2 concentrations in gonads. In the late spawning phase, there were minimal concentrations of E_2 in plasma and there were relatively lesser concentrations in ovaries during this reproductive phase in female round goby. At the end of spawning, there were relatively lesser concentrations of E_2 in other batch spawning teleosts, such as the common dentex (Pavlidis et al., 2000), tench (Pinillos et al., 2003), chub (Guerriero, 2007) and mahseer (*Tor tambroides*; Ismail et al., 2011). Further, the basal level concentration of E_2 in round goby ovaries when there was a predominance of POFs is consistent with results previously reported by Kagawa et al. (1983b) where there were minimal concentrations of E_2 in ovaries following ovulation in the single spawning amago salmon (*Oncorhynchus rhodurus*). During the non-spawning phase in round goby, there was the second greatest concentration of E_2 in plasma and greater concentrations of E_2 in regressed ovaries as compared with the other reproductive phases in the present study. Similarly, during regression of ovaries, there were greater concentrations of E_2 in the plasma of the batch spawning goldfish (Khoo, 1975) and gudgeon (Rinchard et al., 1993). Furthermore, Rinchard et al. (1993) postulated that during regression, atretic vitellogenic oocytes continue to undergo steroidogenesis, which seems to be a sufficient explanation for this observation in the present study.

In round goby males, there were the greatest concentrations of E_2 in plasma and testes in the pre-spawning and non-spawning phases. Relatively greater concentrations of E_2 have been observed in plasma of males of the single spawning Japanese huchen (*Hucho perryi*) (Amer et al., 2001) and the batch spawning gilthead seabream (Chaves-Pozo et al., 2007). The greatest concentration of E_2 in testes of round goby in the pre-spawning phase appears to be associated with the function of this hormone in the early stages of spermatogenesis. Miura et al. (1999) have reported that there are functions of E_2 in the stimulation of spermatogonial stem cell division in the Japanese eel (*Anguilla japonica*). Whereas, the greatest concentration of E_2 observed in testes of the non-spawning round goby is probably associated with the suppression of spermatogenesis. The E_2 inhibits spermatogonial proliferation and induces spermatogonial atresia (Amer et al., 2001; Chaves-Pozo et al., 2005, 2007; Basak et al., 2016). There were lesser concentrations of E_2 in plasma and gonads of round goby males in spawning and late-spawning phases. There were lesser concentrations of E_2 during the spawning season in the serum of the single spawner Japanese huchen (Amer et al., 2001) and in testes in the Japanese eel (Miura et al., 1999; Miura and Miura, 2001). The role of E_2 in spermatogenesis appeared to be less during spawning than in other reproductive phases.

4.4. 11-KT

During the reproductive cycle in round goby females, concentrations of 11-KT in plasma and gonads were less than in males which is consistent with 11-KT being a male-specific hormone in most of the fish species (Borg, 1994; Schulz et al., 2010). Furthermore, the concentrations of 11-KT in plasma and ovaries did not differ during reproductive phases in the present study. There is little information available regarding the functions of 11-KT in the reproductive cycle of female fish. Results from *in vitro* studies in females of the short-finned eel (Lokman et al., 2007), Atlantic cod (Kortner et al., 2008, 2009) and Japanese eel (Endo et al., 2011), however, indicate that 11-KT is one of the factors stimulating the growth of previtellogenic oocytes. In the present study, the previtellogenic oocytes were present in ovaries in similar quantities during all reproductive phases that coincide with the aligned concentrations of 11-KT during the same reproductive phases.

In round goby males, there were the greatest concentrations of 11-KT during the spawning phase in both plasma and testes. The concentrations of 11-KT were associated with the process of active spermatogenesis in the developing testes. These results are consistent with several reports in which concentrations of 11-KT are greatest at the beginning of male spawning and decrease with the advancement of sperm production and testes regression (Borg, 1994; Schulz et al., 2010). For example, a similar pattern was described in batch spawning pejerrey (Elisio et al., 2015) and Atlantic cod (Dahle et al., 2003).

5. Conclusions

There were the greatest Mel concentrations in the spawning and non-spawning phases in the present study, therefore, this hormone appears to have important functions in determining a time frame for spawning in round goby females and males. The greatest concentrations of T_4 and E_2 occurred in the pre-spawning and non-spawning phase, in both females and males, indicating

these hormones are integral in affecting the tissues of the round goby during the active spawning phase and also the timing of initiation of spawning (stimulation of gonad development in pre-spawning phase). The greatest concentrations of 11-KT in males was during the spawning phase and, therefore, this hormone appears to be functional in spermatogenesis in the developing testes. In females, however, there were no changes in 11-KT concentrations during the various reproductive phases. This finding is most likely related to the stimulatory role of 11-KT in the growth of previtellogenic oocytes, which were present in ovaries in similar quantities during all reproductive phases.

Conflict of interests

There is no conflict of interests.

Acknowledgments

The authors would like to thank K.E. Skóra and B. Arciszewski from the Hel Marine Station of the Institute of Oceanography (University of Gdańsk) for providing access to the laboratory facilities, fish maintenance and technical support at the Hel Marine Station. This study was financially supported by the National Science Centre, Poland (grant no. DEC-2012/05/N/NZ8/00739).

References

- Adebisi, F.A., Siraj, S.S., Harmin, S.A., Christianus, A., 2013. Plasma sex steroid hormonal profile and gonad histology during the annual reproductive cycle of river catfish *Hemibagrus nemurus* (Valenciennes, 1840) in captivity. *Fish Physiol. Biochem.* 39, 547–557. <https://doi.org/10.1007/s10695-012-9718-x>.
- Alvarado, M.V., Carrillo, M., Felip, A., 2015. Melatonin-induced changes in kiss/gnrh gene expression patterns in the brain of male sea bass during spermatogenesis. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 185, 69–79. <https://doi.org/10.1016/j.cbpa.2015.03.010>.
- Amano, M., Iigo, M., Ikuta, K., Kitamura, S., Yamada, H., Yamamori, K., 2000. Roles of melatonin in gonadal maturation of underyearling precocious male masu salmon. *Gen. Comp. Endocrinol.* 120, 190–197. <https://doi.org/10.1006/gcen.2000.7547>.
- Amer, M.A., Miura, T., Miura, C., Yamauchi, K., 2001. Involvement of sex steroid hormones in the early stages of spermatogenesis in Japanese huchen (*Hucho perryi*). *Biol. Reprod.* 65, 1057–1066. <https://doi.org/10.1095/biolreprod65.4.1057>.
- Aripin, S.A., Jintasataporn, O., Yoonpundh, R., 2014. Effects of exogenous melatonin in *Clarias macrocephalus* male broodstock first puberty stage. *J. Aquac. Res. Dev.* 6, 307. <https://doi.org/10.4172/2155-9546.1000307>.
- Barcellos, L.J., Wassermann, G.F., Scott, A.P., Woehl, V.M., Quevedo, R.M., Itzès, I., Krieger, M.H., Lulhier, F., 2001. Steroid profiles in cultured female jundiá, the Siluridae *Rhamdia quelen* (Quoy and Gaimard, Pisces Teleostei), during the first reproductive cycle. *Gen. Comp. Endocrinol.* 121, 325–332. <https://doi.org/10.1006/gcen.2001.7603>.
- Basak, R., Roy, A., Rai, U., 2016. Seasonality of reproduction in male spotted murrelet *Channa punctatus*: correlation of environmental variables and plasma sex steroids with histological changes in testis. *Fish Physiol. Biochem.* 42, 1249–1258. <https://doi.org/10.1007/s10695-016-0214-6>.
- Bayarri, M.J., Rodríguez, L., Zanuy, S., Madrid, J.A., Sánchez-Vázquez, F.J., Kagawa, H., Okuzawa, K., Carrillo, M., 2004. Effect of photoperiod manipulation on the daily rhythms of melatonin and reproductive hormones in caged European sea bass (*Dicentrarchus labrax*). *Gen. Comp. Endocrinol.* 136, 72–81. <https://doi.org/10.1016/j.ygcen.2003.12.004>.
- Bhattacharya, S., Chatteraj, A., Maitra, S.K., 2007. Melatonin in the regulation of annual testicular events in carp *Catla catla*: evidence from the studies on the effects of exogenous melatonin, continuous light, and continuous darkness. *Chronobiol. Int.* 24, 629–650. <https://doi.org/10.1080/07420520701534665>.
- Biswas, A., Kundu, S., Roy, S., De, J., Pramanik, M., Ray, A.K., 2006. Thyroid hormone profile during annual reproductive cycle of diploid and triploid catfish, *Heteropneustes fossilis* (Bloch). *Gen. Comp. Endocrinol.* 147, 126–132. <https://doi.org/10.1016/j.ygcen.2005.12.012>.
- Borg, B., 1994. Androgens in teleost fishes. *Comp. Biochem. Physiol. C, Pharmacol. Toxicol. Endocrinol.* 109, 219–245. [https://doi.org/10.1016/0742-8413\(94\)00063-G](https://doi.org/10.1016/0742-8413(94)00063-G).
- Borg, B., Ekström, P., 1981. Gonadal effects of melatonin in the three-spined stickleback, *Gasterosteus aculeatus* L., during different seasons and photoperiods. *Reprod. Nut. Develop.* 6A, 919–927. <https://doi.org/10.1051/rnd:19810704>.
- Bowley, L.A., Yoshikawa, M.A., Wilson, J.Y., 2010. Feminization of male round goby (*Neogobius melanostomus*) from Hamilton Harbour, Lake Ontario. In: Bowley, L.A. (Ed.), *Endocrine disruption in round goby (Neogobius melanostomus) from Hamilton Harbour*. McMaster University, Canada, pp. 62–94.
- Bromage, N., Porter, M., Randall, C., 2001. The environmental regulation of maturation in farmed finfish with special reference to the role of photoperiod and melatonin. *Aquaculture* 197, 63–98. [https://doi.org/10.1016/S0044-8486\(01\)00583-X](https://doi.org/10.1016/S0044-8486(01)00583-X).
- Brown-Peterson, N.J., Wyanski, D.M., Saborido-Rey, F., Macewicz, B.J., Lowerre-Barbieri, S.K., 2011. A standardized terminology for describing reproductive development in fishes. *Mar. Coast. Fish.* 3, 52–70. <https://doi.org/10.1080/19425120.2011.555724>.
- Burke, M., Leatherland, J.F., 1983. Seasonal changes in serum thyroid hormone levels of the feral brown bullhead, *Ictalurus nebulosus* Lesueur. *J. Fish Biol.* 23, 585–593. <https://doi.org/10.1111/j.1095-8649.1983.tb02938.x>.
- Carnevali, O., Gioacchini, G., Maradonna, F., Olivotto, I., Migliarini, B., 2011. Melatonin induces follicle maturation in Danio rerio. *PLoS One* 6, e19978. <https://doi.org/10.1371/journal.pone.0019978>.
- Chaves-Pozo, E., Mulero, V., Meseguer, J., García Ayala, A., 2005. An overview of cell renewal in the testis throughout the reproductive cycle of a seasonal breeding teleost, the gilthead seabream (*Sparus aurata* L.). *Biol. Reprod.* 72, 593–601. <https://doi.org/10.1095/biolreprod.104.036103>.
- Chaves-Pozo, E., Liarte, S., Vargas-Chacoff, L., García-López, A., Mulero, V., Meseguer, J., Mancera, J.M., García-Ayala, A., 2007. 17Beta-estradiol triggers post-spawning in spermatogenically active gilthead seabream (*Sparus aurata* L.) males. *Biol. Reprod.* 76, 142–148. <https://doi.org/10.1095/biolreprod.106.056036>.
- Comeau, L.A., Campana, S.E., 2006. Correlations between thyroidal and reproductive endocrine status in wild Atlantic cod. *Can. Tech. Rep. Fish. Aquat. Sci.* 2682.
- Cornish, D.A., 1998. Seasonal steroid hormone profiles in plasma and gonads of the tilapia, *Oreochromis mossambicus*. *Water S.A.* 24, 257–263.
- Cyr, D.G., Eales, J.G., 1988. *In vitro* effects of thyroid hormones on gonadotropin-induced estradiol-17β secretion by ovarian follicles of rainbow trout, *Salmo gairdneri*. *Gen. Comp. Endocrinol.* 69, 80–87. [https://doi.org/10.1016/0016-6480\(88\)90055-X](https://doi.org/10.1016/0016-6480(88)90055-X).
- Dahle, R., Taranger, G.L., Karlensén, Ø., Kjesbuc, O.S., Norberga, B., 2003. Gonadal development and associated changes in liver size and sexual steroids during the reproductive cycle of captive male and female Atlantic cod (*Gadus morhua* L.). *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* 136, 641–653. [https://doi.org/10.1016/S1095-6433\(03\)00215-0](https://doi.org/10.1016/S1095-6433(03)00215-0).
- Detlaff, T.A., Davydova, S.I., 1979. Differential sensitivity of cells of follicular epithelium and oocytes in the stellate sturgeon to unfavorable conditions, and correlating influence of triiodothyronine. *Gen. Comp. Endocrinol.* 39, 236–243. [https://doi.org/10.1016/0016-6480\(79\)90228-4](https://doi.org/10.1016/0016-6480(79)90228-4).
- Ekström, P., Meissl, H., 1989. Signal processing in a simple vertebrate photoreceptor system: the teleost pineal organ. *Physiol. Bohemoslovaca* 38, 311–326.
- Elisio, M., Chalde, T., Miranda, L.A., 2015. Seasonal changes and endocrine regulation of pejerrey (*Odontesthes bonariensis*) spermatogenesis in the wild. *Gen. Comp. Endocrinol.* 221, 236–243. <https://doi.org/10.1016/j.ygcen.2015.01.011>.
- Endo, T., Todo, T., Lokman, P.M., Kudo, H., Ijiri, S., Adachi, S., Yamauchi, K., 2011. Androgens and very low density lipoprotein are essential for the growth of previtellogenic oocytes from Japanese eel, *Anguilla japonica*, *in vitro*. *Biol. Reprod.* 84, 816–825. <https://doi.org/10.1095/biolreprod.110.087163>.

- Falcón, J., Besseau, L., Sauzet, S., Boeuf, G., 2007. Melatonin effects on the hypothalamo-pituitary axis in fish. *Trends Endocrinol. Metab.* 18, 81–88. <https://doi.org/10.1016/j.tem.2007.01.002>.
- Falcón, J., Migaud, H., Muñoz-Cueto, J.A., Carrillo, M., 2010. Current knowledge on the melatonin system in teleost fish. *Gen. Comp. Endocrinol.* 165, 469–482. <https://doi.org/10.1016/j.ygcen.2009.04.026>.
- Fenwick, J.C., 1970. Demonstration and effect of melatonin in fish. *Gen. Comp. Endocrinol.* 14, 86–97. [https://doi.org/10.1016/0016-6480\(70\)90010-9](https://doi.org/10.1016/0016-6480(70)90010-9).
- Flood, D.E., Fernandino, J.I., Langlois, V.S., 2013. Thyroid hormones in male reproductive development: evidence for direct crosstalk between the androgen and thyroid hormone axes. *Gen. Comp. Endocrinol.* 192, 2–14. <https://doi.org/10.1016/j.ygcen.2013.02.038>.
- Gavella, M., Lipovac, V., 2000. Antioxidative effect of melatonin on human spermatozoa. *Arch. Androl.* 44, 23–27. <https://doi.org/10.1080/014850100262371>.
- Guerriero, G., 2007. Seasonal steroids variations and maturity stages in the female chub, *Leuciscus cephalus* L. (Pisces, Cyprinidae). *Ital. J. Zool.* 74, 317–324. <https://doi.org/10.1080/11250000701448262>.
- Habibi, H.R., Nelson, E.R., Allan, E.R., 2012. New insights into thyroid hormone function and modulation of reproduction in goldfish. *Gen. Comp. Endocrinol.* 175, 19–26. <https://doi.org/10.1016/j.ygcen.2011.11.003>.
- Hurlburt, M.E., 1977. Role of the thyroid gland in ovarian maturation of the goldfish, *Carassius auratus* L. *Can. J. Zool.* 55, 1906–1913. <https://doi.org/10.1139/z77-244>.
- Iseki, K.K., Negrao, J.A., Castrucci, A.M., 2000. Seasonal changes on plasma melatonin levels in the fresh water teleost fish, *Piaractus mesopotamicus* (Characiformes, Characidae). *Comp. Biochem. Physiol. A Mol. Integrative Physiol.* 126, 71. [https://doi.org/10.1016/S1095-6433\(00\)80140-3](https://doi.org/10.1016/S1095-6433(00)80140-3).
- Ismail, M.F., Siraj, S.S., Daud, S.K., Harmin, S.A., 2011. Association of annual hormonal profile with gonad maturity of mahseer (*Tor tambroides*) in captivity. *Gen. Comp. Endocrinol.* 170, 125–130. <https://doi.org/10.1016/j.ygcen.2010.09.021>.
- Janz, D.M., Weber, L.P., 2000. Endocrine system. In: Ostrander, G.K. (Ed.), *The Laboratory Fish: The Handbook of Experimental Animals*. Academic Press, San Diego, pp. 415–449.
- Kadmon, G., Yaron, Z., Gordin, H., 1985. Sequence of gonadal events and oestradiol levels in *Sparus aurata* (L.) under two photoperiod regimes. *J. Fish Biol.* 26, 609–620. <https://doi.org/10.1111/j.1095-8649.1985.tb04301.x>.
- Kagawa, H., Young, G., Nagahama, Y., 1983a. Changes in plasma steroid hormone levels during gonadal maturation in female goldfish *Carassius auratus*. *Bull. Japan Soc. Sci. Fish.* 49, 1783–1787. <https://doi.org/10.2331/suisan.49.1783>.
- Kagawa, H., Young, G., Nagahama, Y., 1983b. Relationship between seasonal plasma estradiol-17 beta and testosterone levels and *in vitro* production by ovarian follicles of amago salmon (*Oncorhynchus rhodurus*). *Biol. Reprod.* 29, 301–309. <https://doi.org/10.1095/biolreprod29.2.301>.
- Kagawa, H., Young, G., Nagahama, Y., 1984. *In vitro* estradiol-17 beta and testosterone production by ovarian follicles of the goldfish, *Carassius auratus*. *Gen. Comp. Endocrinol.* 54, 139–143. [https://doi.org/10.1016/0016-6480\(84\)90209-0](https://doi.org/10.1016/0016-6480(84)90209-0).
- Kalamarz-Kubiak, H., Gozdowska, M., Guellard, T., Kulczykowska, E., 2017. How does oestradiol influence the AVT/IT system in female round gobies during different reproductive phases? *Biol. Open* 6, 1493–1501. <https://doi.org/10.1242/bio.024844>.
- Kezuka, H., Furukawa, K., Aida, K., Hanyu, I., 1988. Daily cycles in plasma melatonin levels under long or short photoperiod in the common carp, *Cyprinus carpio*. *Gen. Comp. Endocrinol.* 72, 296–302. [https://doi.org/10.1016/0016-6480\(88\)90212-2](https://doi.org/10.1016/0016-6480(88)90212-2).
- Khoo, K.H., 1975. The corpus luteum of goldfish (*Carassius auratus* L.) and its functions. *Can. J. Zool.* 53, 1306–1323. <https://doi.org/10.1139/z75-156>.
- Kornis, M.S., Mercado-Silva, N., Vander Zanden, M.J., 2012. Twenty years of invasion: a review of round goby *Neogobius melanostomus* biology, spread and ecological implications. *J. Fish Biol.* 80, 235–285. <https://doi.org/10.1111/j.1095-8649.2011.03157.x>.
- Kortner, T.M., Rocha, E., Silva, P., Filipe, L., Castro, L.F.C., Arukwe, A., 2008. Genomic approach in evaluating the role of androgens on the growth of Atlantic cod (*Gadus morhua*) previtellogenic oocytes. *Comp. Biochem. Physiol. D Genom. Proteom.* 3, 205–218. <https://doi.org/10.1016/j.cbd.2008.04.001>.
- Kortner, T.M., Mortensen, A.S., Hansen, M.D., Arukwe, A., 2009. Neural aromatase transcript and protein levels in Atlantic salmon (*Salmo salar*) are modulated by the ubiquitous water pollutant, 4-nonylphenol. *Gen. Comp. Endocrinol.* 164, 91–99. <https://doi.org/10.1016/j.ygcen.2009.05.009>.
- Kulczykowska, E., Kasprzak, M., Kalamarz, H., Kuriata, M., Nietrzeba, M., Jerzak, L., Kamiński, P., 2007. Melatonin and thyroxine response to pollution in white stork nestlings (*Ciconia ciconia*): aspects of rhythmicity and age. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 146, 392–397. <https://doi.org/10.1016/j.cbpc.2007.04.012>.
- Kulczykowska, E., Kalamarz-Kubiak, H., Nietrzeba, M., Gozdowska, M., 2015. Brain neuropeptide and gonadal steroid responses to deprivation of heterosexual contact in the black molly. *Biol. Open* 4, 69–78. <https://doi.org/10.1242/bio.20149597>.
- Leatherland, J.F., 1994. Reflections on the thyroidology of fishes: from molecules to humankind. *Guelph Ichthyol. Rev.* 2, 1–64.
- Legler, J., Broekhof, J.L.M., Brouwer, A., Lanser, P.H., Murk, A.J., Van Der Saag, P.T., Vethaak, A.D., Wester, P., Zivkovic, D., Van Der Burg, B., 2000. A novel *in vivo* bioassay for (xeno-) estrogens using transgenic zebrafish. *Environ. Sci. Technol.* 34, 4439–4444. <https://doi.org/10.1021/es000605>.
- Lokman, P.M., George, K.A., Divers, S.L., Algie, M., Young, G., 2007. 11-Ketotestosterone and IGF-I increase the size of previtellogenic oocytes from shortfinned eel, *Anguilla australis*, *in vitro*. *Reproduction* 133, 955–967. <https://doi.org/10.1530/REP-06-0229>.
- Lombardo, F., Giorgini, E., Gioacchini, G., Maradonna, F., Ferraris, P., Carnevali, O., 2012. Melatonin effects on *Fundulus heteroclitus* reproduction. *Reprod. Fertil. Dev.* 24, 794–803. <https://doi.org/10.1071/RD11267>.
- Lubzens, E., Young, G., Bobe, J., Cerdà, J., 2010. Oogenesis in teleosts: how eggs are formed. *Gen. Comp. Endocrinol.* 165, 367–389. <https://doi.org/10.1016/j.ygcen.2009.05.022>.
- Maitra, S.K., Hasan, K.N., 2016. The role of melatonin as a hormone and an antioxidant in the control of fish reproduction. *Front. Endocrinol. (Lausanne)* 7, 38. <https://doi.org/10.3389/fendo.2016.00038>.
- Maitra, S.K., Seth, M., Chatteraj, A., 2006. Photoperiod, pineal photoreceptors and melatonin as the signal of photoperiod in the regulation of reproduction in fish. *J. Endocrinol. Reprod.* 10, 73–87. <https://doi.org/10.18311/jer/2006/2125>.
- Maitra, S.K., Chatteraj, A., Mukherjee, S., Moniruzzaman, M., 2013. Melatonin: a potent candidate in the regulation of fish oocyte growth and maturation. *Gen. Comp. Endocrinol.* 181, 215–222. <https://doi.org/10.1016/j.ygcen.2012.09.015>.
- Maitra, S.K., Mukherjee, S., Hasan, K.N., 2015. In: Catalá, Á. (Ed.), *Melatonin: Endogenous Sources and Role in the Regulation of Fish Reproduction*. Nova Science Publishers Inc, Indoleamines, pp. 43–77.
- Marentette, J.R., Fitzpatrick, J.L., Bergera, R.G., Balshine, S., 2009. Multiple male reproductive morphs in the invasive round goby (*Apollonia melanostoma*). *J. Gt. Lakes Res.* 35, 302–308. <https://doi.org/10.1016/j.jglr.2009.01.009>.
- Methven, D.A., Crim, L.W., Norberg, B., Brown, J.A., Goff, G.P., Huse, I., 1992. Seasonal reproduction and plasma levels of sex steroids and vitellogenin in Atlantic halibut (*Hippoglossus hippoglossus*). *Can. J. Fish. Aquat. Sci.* 49, 754–759. <https://doi.org/10.1139/f92-084>.
- Miura, T., Miura, C.I., 2001. Japanese Eel: a model for analysis of spermatogenesis. *Zool. Sci.* 18, 1055–1063. <https://doi.org/10.2108/zsj.18.1055>.
- Miura, T., Miura, C., Ohta, T., Nader, M.R., Todo, T., Yamauchi, K., 1999. Estradiol-17beta stimulates the renewal of spermatogonial stem cells in males. *Biochem. Biophys. Res. Commun.* 264, 230–234. <https://doi.org/10.1006/bbrc.1999.1494>.
- Mondal, P., Hasan, K.N., Pal, P.K., Maitra, S.K., 2017. Influences of exogenous melatonin on the oocyte growth and oxidative status of ovary during different reproductive phases of an annual cycle in carp *Catla catla*. *Theriogenology* 87, 349–359. <https://doi.org/10.1016/j.theriogenology.2016.09.021>.
- Monteverdi, G.H., Di Giulio, R.T., 2000. Vitellogenin association and oocyte accumulation of thyroxine and 3, 5,3'-triiodothyronine in gravid *Fundulus heteroclitus*. *Gen. Comp. Endocrinol.* 120, 198–211. <https://doi.org/10.1006/gcen.2000.7551>.
- Mori, Y., Kano, Y., 1984. Changes in plasma concentrations of LH, progesterone and oestradiol in relation to the occurrence of luteolysis, oestrus and time of ovulation in the Shiba goat (*Capra hircus*). *J. Reprod. Fertil.* 72, 223–230. <https://doi.org/10.1530/jrf.0.0720223>.
- Mylonas, C.C., Woods, L.C., Zohar, Y., 1997. Cyto-histological examination of post-vitellogenesis and final oocyte maturation in captive-reared striped bass. *J. Fish Biol.* 50, 34–49. <https://doi.org/10.1111/j.1095-8649.1997.tb01338.x>.
- Osborn, R.H., Simpson, T.H., 1978. Seasonal changes in thyroidal status in the plaice, *Pleuronectes platessa* L. *J. Fish Biol.* 12, 519–526. <https://doi.org/10.1111/j.1095-8649.1978.tb04197.x>.
- Pavlidis, M., Greenwood, L., Mourot, B., Kokkari, C., Le Menn, F., Divanach, P., Scott, A.P., 2000. Seasonal variations and maturity stages in relation to differences in

- serum levels of gonadal steroids, vitellogenin, and thyroid hormones in the common dentex (*Dentex dentex*). Gen. Comp. Endocrinol. 118, 14–25. <https://doi.org/10.1006/gcen.1999.7440>.
- Pinillos, M.L., Delgado, M.J., Scott, A.P., 2003. Seasonal changes in plasma gonadal steroid concentrations and gonadal morphology of male and female tench (*Tinca tinca*, L.). Aquacul. Res. 34, 1181–1189. <https://doi.org/10.1046/j.1365-2109.2003.00926.x>.
- Popek, W., Bieniarz, K., Epler, P., 1991. Role of pineal gland in the sexual cycle in common carp. In: Surowiak, J.M., Lewandowski (Eds.), Chronobiology and Chronomedicine, pp. 99–102 Frankfurt.
- Popek, W., Drag-Kozak, E., Łuszczek-Trojnar, E., 2010. Influence of melatonin on hypothalamo-pituitary-ovarian axis in carp: mechanisms and actions. In: Kulczykowska, E., Popek, W., Kapoor, B.G. (Eds.), Biological Clock in Fish. Science Publishers, USA, pp. 217–234.
- Prat, F., Zanuy, S., Carrillo, M., de Mones, A., Fostier, A., 1990. Seasonal changes in plasma levels of gonadal steroids of sea bass, *Dicentrarchus labrax* L. Gen. Comp. Endocrinol. 78, 361–373. [https://doi.org/10.1016/0016-6480\(90\)90026-1](https://doi.org/10.1016/0016-6480(90)90026-1).
- Rinchard, J., Kestemont, P., Kühn, E.R., Fostier, A., 1993. Seasonal changes in plasma levels of steroid hormones in an asynchronous fish the gudgeon *Gobio gobio* L. (Teleostei, Cyprinidae). Gen. Comp. Endocrinol. 92, 168–178. <https://doi.org/10.1006/gcen.1993.1153>.
- Rocha, M.J., Rocha, E., 2006. Morphofunctional aspects of reproduction from synchronous to asynchronous fishes - an overview. In: Reinecke, M., Zaccone, G., Kapoor, B.G. (Eds.), Fish Endocrinology. Science Publishers, Enfield, pp. 571–624.
- Rohr, D.H., Lokman, P.M., Davie, P.S., Young, G., 2001. 11-Ketotestosterone induces silvering-related changes in immature female short-finned eels, *Anguilla australis*. Comp. Biochem. Physiol. A Mol. Integr. Physiol. 130, 701–714. [https://doi.org/10.1016/S1095-6433\(01\)00402-0](https://doi.org/10.1016/S1095-6433(01)00402-0).
- Sabet, S.S., Imanpour, M.R., Fatideh, B.A., Saeed, G., 2009. Study on sexual maturity and levels of gonad steroid hormones in female kutum *Rutilus frisii kutum* (Kamenskii, 1901) during spawning season from river Sefid-Rood of the southern Caspian sea. J. Cell. Anim. Biol. 3, 208–215.
- Sage, M., 1973. The evolution of thyroidal function in fishes. Am. Zool. 13, 899–905. <https://doi.org/10.1093/icb/13.3.899>.
- Sapota, M.R., 2012. NOBANIS – Invasive Alien Species Fact Sheet – *Neogobius melanostomus*. Online Database of the European Network on Invasive Alien Species – NOBANIS. (Accessed 04 January 2019). <http://www.nobanis.org/>.
- Schulz, R.W., de França, L.R., Lareyre, J.J., Le Gac, F., Chiarini-Garcia, H., Nobrega, R.H., Miura, T., 2010. Spermatogenesis in fish. Gen. Comp. Endocrinol. 165, 390–411. <https://doi.org/10.1016/j.ygcen.2009.02.013>.
- Shang, X., Huang, Y., Ye, Z., Yu, X., Gu, W., 2004. Protection of melatonin against damage of sperm mitochondrial function induced by reactive oxygen species. Zhonghua Nan Ke Xue. Nat. J. Androl. 10, 604–607.
- Sokołowska, E., Kalamarz, H., Kulczykowska, E., 2004. Seasonal changes in brain melatonin concentration in the three-spined stickleback (*Gasterosteus aculeatus*): towards an endocrine calendar. Comp. Biochem. Physiol. A Mol. Integr. Physiol. 139, 365–369. <https://doi.org/10.1016/j.cbpb.2004.10.001>.
- Sokołowska, E., Kleszczyńska, A., Kalamarz-Kubiak, H., Arciszewski, B., Kulczykowska, E., 2013. Changes in brain arginine vasotocin, isotocin, plasma 11-ketotestosterone and cortisol in round goby, *Neogobius melanostomus*, males subjected to overcrowding stress during the breeding season. Comp. Biochem. Physiol. A Mol. Integr. Physiol. 165, 237–242. <https://doi.org/10.1016/j.cbpa.2013.03.018>.
- Sower, S.A., Schreck, C.B., 1982. Steroid and thyroid hormones during sexual maturation of coho salmon (*Oncorhynchus kisutch*) in seawater or fresh water. Gen. Comp. Endocrinol. 47, 42–53. [https://doi.org/10.1016/0016-6480\(82\)90082-X](https://doi.org/10.1016/0016-6480(82)90082-X).
- Supriya, A., Raghuveer, K., Swapna, I., Rasheeda, M.K., Kobayashi, T., Nagahama, Y., 2005. Thyroid hormone modulation of ovarian recrudescence of air-breathing catfish *Clarias gariepinus*. Fish Physiol. Biochem. 31, 267–270. <https://doi.org/10.1007/s10695-006-0034-1>.
- Swapna, I., Senthilkumaran, B., 2007. Thyroid hormones modulate the hypothalamo–hypophyseal–gonadal axis in teleosts: molecular insights. Fish Physiol. Biochem. 33, 335–345. <https://doi.org/10.1007/s10695-007-9165-2>.
- van Bohemen, C.G., Lambert, J.G., 1981. Estrogen synthesis in relation to estrone, estradiol, and vitellogenin plasma levels during the reproductive cycle of the female rainbow trout, *Salmo gairdneri*. Gen. Comp. Endocrinol. 45, 105–114. [https://doi.org/10.1016/0016-6480\(81\)90175-1](https://doi.org/10.1016/0016-6480(81)90175-1).
- Wiegand, M.D., 1982. Vitellogenesis in fishes. Richter, C.J.J., Goos, H.J.T. (Eds.), Proceedings of the Second International Symposium on Reproductive Physiology of Fish 136–146.
- Zeyl, J.N., Love, O.P., Higgs, D.M., 2014. Evaluating gonadosomatic index as an estimator of reproductive condition in the invasive round goby. J. Gt. Lakes Res. 40, 164–171. <https://doi.org/10.1016/j.jglr.2013.12.004>.