

METHODS

Exploration of peripancreatic lymphatic pathways in a live porcine model



Yohann Renard^{a,b,c,d,*}, Cyril Perrenot^{b,c,1}, Marc Labrousse^{a,c,1}, Claude Avisse^{a,d,1}, Rami Rhaïem^{d,1}, Tullio Piardi^{d,1}, Marc Braun^{b,c,1}, Manuela Perez^{b,c,1}

^a Department of Anatomy, Faculty of Medicine and University Hospital, University of Champagne-Ardenne, Reims, France

^b School of Surgery of Nancy, University of Lorraine, Nancy, France

^c IADI, INSERM U1254, University of Lorraine, Nancy, France

^d Department of Digestive Surgery, Robert-Debre University Hospital, University of Champagne-Ardenne, Reims, France

ARTICLE INFO

Article history:

Received 17 December 2018

Received in revised form 5 June 2019

Accepted 17 June 2019

Keywords:

Lymphatic capillaries

Pancreas

Pancreatic segments

Pancreatic segmental transplantation

Pancreatic neoplasms/surgery

ABSTRACT

Pancreatic cancer is associated with a poor prognosis, mainly due to lymph node invasion and lymph node recurrence after surgical resection, even after extended lymphadenectomy. The peripancreatic lymphatic system is highly complex and the specific lymphatic drainage of each part of the pancreas has not been established.

The aim of this study was to determine the lymphatic drainage pathways specific to each part of the pancreas on live pigs using Patent Blue.

The pancreases of 14 live pigs were injected in different parts of the gland. The technique was efficient and reproducible. The diffusion patterns were similar for each location and were reported. Our results in pigs allowed us to define specific nodal relay stations and lymphatic drainage for each part of the pancreas and confirm that independent anatomical-surgical pancreatic segments can be described. It is interesting to note that lymphatic drainage for the upper part of the proximal part of pancreas (duodenal lobe) occurred on the left side of the portal vein. This suggests that lymph node resection during cephalic duodenopancreatectomy in humans should be extended to the left side of the mesenteric vein, and probably to the right side of the superior mesenteric artery, as recently suggested.

These results could help surgeons perform safe anatomical-segmental pancreatic resections with accurate lymphadenectomies and improve survival in patients with pancreatic cancer. Based on these results we will perform an innovative prospective study. Patent Blue will be injected into different parts of the gland in patients operated for pancreatic resection, and lymphatic diffusion of the dye will be recorded in relation to their origin from the theoretical pancreatic segments (ClinicalTrials.gov Identifier: NCT03597230).

© 2019 Elsevier GmbH. All rights reserved.

1. Introduction

Pancreatic ductal adenocarcinomas (PDAC) are the most frequent pancreatic cancers with a 5-year survival rate of 7% and the poorest prognosis of all cancers (Neuzillet et al., 2015). When possible, surgical resection is the best chance of cure (Siegel et al., 2016). Prognosis is determined by tumor stage, in particular lymph-node status (Fink et al., 2016; Pu et al., 2018) as well as the quality of

lymph node resection (Tol et al., 2014; Zhan et al., 2015). The network of the supra-mesocolic lymphatic vascular system is highly complex (Kanda et al., 2011; Cesmebasi et al., 2015). The high rate of loco-regional lymph-node recurrence after cancer resection, even following extended lymphadenectomy (Riediger et al., 2009) has promoted interest in the anatomical and radiological knowledge of the peripancreatic lymphatic circulation (Akcali et al., 2006; Borghi et al., 1998; Delpero et al., 2017; Pu et al., 2018; Samra et al., 2008; Tol et al., 2014) and the peri-operative detection of lymph node metastases around the pancreas (Beisani et al., 2016; Tomson et al., 2015).

Certain studies, based on child and adult cadaveric dissections as well as lymph-node spreading in PDAC resected specimens, have hypothesized that the pancreas may be divided into several seg-

* Corresponding author at: Department of Anatomy, Faculty of Medicine and University Hospital, University of Champagne-Ardenne, Rue Cognacq-Jay, 51092 Reims Cedex, France.

E-mail address: yrenard@chu-reims.fr (Y. Renard).

¹ These authors contributed equally to this work.




Fig. 1. Anatomy of the pig's pancreas. 1: portal vein. 2: duodenum. 3: aorta. 4: splenic vein. 5: superior mesenteric artery. 6: superior mesenteric vein. 7: inferior mesenteric vein. A: duodenal lobe. B: connecting lobe. C: splenic lobe.

ments (Renard et al., 2017) each with their own lymphatic drainage (Borghini et al., 1998; Cesmebasi et al., 2015; Cubilla et al., 1978; Donatini and Hidden, 1992; Nagakawa et al., 1994; Pissas, 1984). Theoretically, the ability to distinguish independent segments of the pancreatic parenchyma could help surgeons perform safe segmental pancreatic resections and adequate lymphadenectomy to reduce the loco-regional lymph-node recurrence rate. Nevertheless, no independent peripancreatic lymphatic pathways have been demonstrated thus far (Renard et al., 2017).

The goal of this study was to evaluate the feasibility of defining pancreatic segments following lymphatic drainage in a live animal model by mapping the peripancreatic lymphatic pathways using the lymphophil dye, Patent Blue (Beisani et al., 2016).

2. Material and methods

The pig pancreas is composed of three lobes (Fig. 1). The splenic lobe is located posteriorly and to the left, close to the spleen and stomach, and corresponds to the tail and body in the human pancreas. The duodenal lobe is located closest to the duodenum, and corresponds to the head of the pancreas. The connecting lobe is an extension located on the anterior border of the portal vein and should correspond to the uncinate process (Ferrer et al., 2008; Tsuchitani et al., 2016).

Fourteen live pigs were included in this study. They came from the School of Surgery, Lorraine University, Nancy, France. All experiments on pigs were performed in accordance with National Institute of Health recommendations and approved by the corresponding ethics committee. All pigs were initially enrolled for laparoscopic surgical training sessions and were injected 10 min before animal sacrifice (Pissas et al., 1982). The pig was not included in case of surgical dissection of the upper part of the abdomen or in the event of the death of the animal before injection. A median xypho-pubic laparotomy was performed under general anesthesia and precautions were taken to avoid any dissection around the peripancreatic area.

Reproducibility of our protocol was excellent and included an injection of 1 mL of pure Patent Blue (Guerbet[®], Villepinte, France) in the pancreatic parenchyma at a depth of 0.5 to 1 cm from the surface, using a 2 mL syringe (BD Medical[®], Le Pont-de-Claix, France) and a 23 G needle (Fine-Ject, Henke Sass Wolf[®], Tuttlingen, Germany).

We performed the injections to make sure that the results may be applied to human pancreatic surgery. The injections were performed in the upper part of the duodenal lobe and the lower part of this lobe in 4 pigs each, in the distal parts of the splenic lobe in 3 pigs and in the proximal parts of this latter lobe in 3 pigs. The pigs were sacrificed and dissected. Three hours were necessary to make sure that all locations of the blue diffusion were explored. The patterns of blue diffusion were recorded for each pig and pictures were taken.

3. Results

The technique was highly efficient and reproducible. For each location of injection, the diffusion patterns of the blue injection were similar:

- Injection in the superior part of the duodenal lobe (Fig. 2A) resulted in lymphatic diffusion around the hepatic pedicle (Fig. 2B) and towards but not into the splenic lobe until beyond the portal vein (Fig. 2C).
- Injection in the lower part of the duodenal lobe (Fig. 3A) resulted in lymphatic diffusion towards the connecting lobe (Fig. 3C).
- Injection in the distal part of the splenic lobe (Fig. 4A) resulted in lymphatic diffusion around the splenic pedicle and towards the connecting lobe (Fig. 4B).
- Injections in the proximal part of the splenic lobe (Fig. 5A) resulted in lymphatic diffusion around the splenic pedicle and towards the portal vein without crossing the left border (Fig. 5B).

4. Discussion

Most pancreatic tumors (95%) are ductal adenocarcinomas (PDAC) (Amundadottir, 2016; Neuzillet et al., 2015) which is the eleventh cause of cancer and the fourth leading cause of death from cancer worldwide (Ferlay et al., 2016; Siegel et al., 2016). PDAC has the poorest prognosis of all cancers, with a 5-year survival rate of 7% (Neuzillet et al., 2015; Siegel et al., 2016). Lymph node status, including malignant spread and the total number of retrieved lymph nodes as well as the lymph node ratio, is the most important prognostic factor (Fink et al., 2016; Pu et al., 2018; Santi et al., 2011). A prospective multicenter study performed by Delpero et al. (2017)




Fig. 2. Lymphatic diffusion of the Patent Blue after injection in the superior part of the duodenal lobe. A1: Injection in the superior part of the duodenal lobe. A2: Schematic illustration of A1. B1: Diffusion around the hepatic pedicle (white arrow 1). B2: Schematic illustration of B1. C1: Diffusion beyond the portal vein (white arrow 2). C2: Schematic illustration of C1. The blue line corresponds to the portal vein. a, pancreas; b, duodenum; c, portal vein; d, stomach. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

including 147 patients operated for pancreatic cancer showed that lymph node status was the greatest independent prognostic factor (Delpero et al., 2017). Surgical resection with lymphadenectomy seems to be the best curative treatment option (Kayahara et al., 1992; Riediger et al., 2009; Samra et al., 2008; Sauvanet, 2008; Tol et al., 2014; Zhan et al., 2015) in patients with localized cancer (i.e. without spreading to regional and distant lymph node locations, representing up to 10% of cases (Fink et al., 2016; Siegel et al., 2016)). The importance of lymph node status and the high rate of local lymph node recurrence following surgical resection (Delpero et al., 2017; Nimura et al., 2012; Pu et al., 2018; Riediger et al., 2009) have renewed interest in the peripancreatic lymphatic circulation (Akcali et al., 2006; Borghi et al., 1998; Samra et al., 2008). However, because of the absence of strong evidence on the optimal lymphadenectomy to be performed during pancreatic resection, better anatomical understanding of lymphatic spreading around the pancreas, depending on the location of the tumor is still needed (Renard et al., 2017; Tol et al., 2014).

Many reports have described the peripancreatic lymphatic system with various results (Deki and Sato, 1988). Recently, certain studies have focused on the main lymphatic pathways around the

pancreas, with discrepant results on the location of the regional lymph nodes (Cubilla et al., 1978; Deki and Sato, 1988; Donatini and Hidden, 1992; Kanda et al., 2011; Kayahara et al., 1992; Nagakawa et al., 1994; O'Morchoe, 1997; Pissas, 1990, 1984):

- Certain authors have named lymph groups according to their location around the pancreas. Based on 33 pancreatectomy specimens, Cubilla et al. (1978) described 5 main groups: superior, inferior, anterior, posterior and splenic (Cubilla et al., 1978). Borghi et al. (1998) and Pissas (1984) reported a similar regional classification (Borghi et al., 1998; Pissas, 1984).
- Other authors have developed numerical classifications (O'Morchoe, 1997). Based on lymph-node spreading in 44 PDAC resected specimens, Kayahara et al. (1992) and Nagakawa et al. (1994) described 11 nodal areas, forming 4 lymphatic pathways around the head of the pancreas (Kayahara et al., 1992; Nagakawa et al., 1994, 1993). After meticulous macroscopic dissection of 4 cadavers, Deki and Sato (1988) provided a detailed description of 8 lymphatic pathways running towards the abdomino-aortic (para-aortic) lymph-nodes (Deki and Sato, 1988; Pavlidis et al., 2011). They reported 3 lymphatic pathways




Fig. 3. Lymphatic diffusion of the Patent Blue after injection in the lower part of the duodenal lobe. A1: Injection in the lower part of the duodenal lobe. A2: Schematic illustration of A1. B1: Diffusion towards the connecting lobe (white arrow). B2: Schematic illustration of B1. C1 showed no diffusion around the hepatic pedicle. C2: Schematic illustration of C1. a, pancreas; b, duodenum; c, portal vein; d, stomach; e, colon. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

in the anterior surface of the head, 3 in the posterior surface and two major lymphatic routes in the left half of the pancreas (Deki and Sato, 1988).

- Finally, the Japan Pancreatic Society developed an anatomical classification dividing regional and juxtaregional lymph nodes into 35 groups including 23 subgroups based on data from 18,629 cases of carcinoma of the pancreas (Cubilla et al., 1978; Isaji et al., 2004; Nagakawa et al., 1994; Tol et al., 2014).

These authors only reported the location of the lymph nodes around the pancreas and did not try to determine the origin of these pathways from the precise location of the tumors in the pancreatic parenchyma. Further, their works weren't based on a live model.

Very few other studies have evaluated the correlation between these lymphatic pathways and their origin in the different areas of the pancreas, suggesting that each theoretical pancreatic seg-

ment could have its own lymphatic drainage (Borghetti et al., 1998; Cesmebasi et al., 2015; Cubilla et al., 1978; Donatini and Hidden, 1992; Nagakawa et al., 1994; Pissas, 1984). This segmentation would help surgeons perform accurate targeted lymph node resection during segmental pancreatectomy (Fink et al., 2016; Renard et al., 2017). These studies were mainly based on adult, child or embryo cadaveric dissections, or surgical specimens from patients with pancreatic carcinomas. In pilot studies, lymphatic channels were shown to come from two distinct portions of the pancreas, one on the right side and one of the left (head/neck, body/tail, respectively) (Cesmebasi et al., 2015; Fink et al., 2016). Other authors have proposed more precise theories: Cubilla et al. (1978) and Pissas (1984) proposed a division into 5 segments with specific and independent lymphatic drainage (Cubilla et al., 1978; Pissas, 1984), but with significant differences between the 2 theories. Donatini and Hidden (1992) proposed 4 segments with indepen-




Fig. 4. Lymphatic diffusion of the Patent Blue after injection in the distal part of the splenic lobe. A1: Injection in the distal part of the splenic lobe. A2: Schematic illustration of A1. B1: Diffusion around the splenic pedicle (white arrow 1) and towards the connecting lobe (white arrow 2). B2: Schematic illustration of B1. a, pancreas; d, stomach; f, small intestine; g, spleen; h, superior mesenteric vein. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 5. Lymphatic diffusion of the Patent Blue after injection in the proximal part of the splenic lobe. A1: Injection in the proximal part of the splenic lobe. A2: Schematic illustration of A1. B1: Diffusion around the splenic pedicle (white arrow 1) and towards the portal vein but without crossing its left border (white arrow 2). B2: Schematic illustration of B1. The blue line corresponds to the portal vein. a, pancreas; b, duodenum; d, stomach; f, small intestine; g, spleen. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

dent pathways of lymphatic drainage (Donatini and Hidden, 1992), which differed from the two earlier theories.

These complex and non-standardized (Fink et al., 2016) classifications show the difficulty of obtaining a routine description of the peripancreatic lymphatic pathways, and suggest that more accurate and detailed studies are needed to improve staging (Pavlidis et al., 2011).

The goal of the present study was to show the feasibility of defining the origins of the peripancreatic lymphatic pathways in live animals. The pig pancreas includes duodenal, splenic and connecting lobes (Ferrer et al., 2008; Tsuchitani et al., 2016) and is considered a highly suitable model for experimental pancreatic research (Bensley, 2005; Shokouh-Amiri et al., 1989; Troisi et al., 2000), in particular for the lymphatic pathways (Ji and Kato, 1997;

Regoli et al., 2001). To the best of our knowledge, this is the first time that several parts of the pancreas of a live animal have been injected with Patent Blue to describe a cartography of lymphatic drainage for each theoretical segment (Renard et al., 2017). Pissas et al. (1982) reported a similar study in dogs, with a different goal, whose pancreatic anatomy has been already shown to be segmented and highly different from humans (Galindo-Pacheco et al., 1992; Pissas et al., 1982).

The identification of lymphatic vessels is difficult in cadavers and surgical samples (Cesmebasi et al., 2015; Fink et al., 2016; Hirai et al., 2001; O'Morchoe, 1997; Pissas, 1984). Thus several vital staining dyes have been used to distinguish lymphatic vessels from other structures (Evans and Ochsner, 1954; Pissas, 1984; Pissas et al., 1982; Tomson et al., 2015). Although they have been found to be

efficient to achieve this goal when injected in the lung and stomach, they are ineffective in recognizing the lymphatic vessels when injected in the pancreas (Pissas et al., 1982). Pissas (1984) reported a cadaveric study of lymphatic drainage of the pancreas using the previous manufacturing process of Patent Blue dye (Pissas, 1984). More recently, Nagakawa et al. (1994) used carbon particles or radioactive insulin in surgical specimens from patients with pancreatic tumors (Kayahara et al., 1999; Nagakawa et al., 1994, 1993). At present lymphoscintigraphy is the technical reference for the detection of the sentinel node. However, it is difficult to use routinely, in particular during gross dissection and anatomical studies (Niebling et al., 2016; Peek et al., 2017; Tomson et al., 2015).

In the present study we used Patent Blue, first described by Weinberg and Greaney for gastric cancer (Weinberg and Greaney, 1950), which is now a frequently used lymphophil dye, alone or combined with a radiocolloid tracer or lympho-fluoroscopy for sentinel lymph node identification in breast cancer and melanoma (Niebling et al., 2016; Peek et al., 2017; Tomson et al., 2015).

In our study, we were able to describe the peripancreatic lymphatic pathways and show the different pancreatic origins of each. These results confirm the presence of pancreatic segmentation and suggest that independent anatomical-surgical pancreatic parts could be defined with their own specific lymphatic drainage. If these results are confirmed in humans, they could help the surgeon perform limited pancreatic resection procedures safely as well as increase the accuracy of lymph node resections.

For example, there is an ongoing debate on the extent of lymphadenectomy to be performed during pancreatoduodenectomy, in particular around the superior mesenteric artery (Tol et al., 2014). Our results in live pigs show that the upper part of the duodenal lobe drains towards the splenic lobe, beyond the portal vein. If these results are applied to humans this suggests that during surgical resection of a tumor located in the head of pancreas, lymph node resection should be extended to the left, beyond the portal vein up to the right side of the superior mesenteric artery, confirming several recent studies (Borghi et al., 1998; Cesmebasi et al., 2015; Hirai et al., 2001; Kanda et al., 2011; Kayahara et al., 1999; Nimura et al., 2012; Pu et al., 2018; Samra et al., 2008; Tol et al., 2014).

This study presents limitations inherent to the use of live animal. Although pigs are considered as interesting model for experimental pancreatic research, is it not known, to the best of our knowledge, whether the 3 anatomical lobes of the pig's pancreas can be good correlated to human's one. In particular, we did not inject the connecting lobe, probably corresponding the uncinate process, since numerous authors demonstrated that the uncinate process should be an extension of the posterior part of the head (Ferrero et al., 2008; Tsuchitani et al., 2016; Renard et al., 2017). Uchida et al. (1999) and Suda et al. (2006) demonstrated that during embryogenesis, the posterior part of the head and the uncinate process, come from the ventral bud (Uchida et al., 1999; Suda et al., 2006). Moreover, Holt and Varadarajulu (2014) and Rana and Vilmann (2015) reported that the embryological dorsal pancreas is generally more echogenic than the embryological ventral pancreas: a transition area can be distinguished between the darker anterior head to the brighter posterior pancreas (body and tail, including the uncinate process) (Holt and Varadarajulu, 2014; Rana and Vilmann, 2015). In addition, based on cadaveric injections, Donatini and Hidden (1992) showed that the uncinate process was an integral part of the inferior portion of the head (Donatini and Hidden, 1992).

5. Conclusions and perspectives

Mapping the peripancreatic lymphatic pathways with the lymphophil dye, Patent Blue, (Beisani et al., 2016) can offer a reproducible description of pancreatic segments following lymphatic

drainage in a live animal model. If these results are confirmed in the human anatomy, they could help to define patient-specific lymph node dissection during pancreatic adenocarcinoma surgical resections. Based on these results we will perform a prospective study in patients undergoing pancreatic resection. During surgery, different parts of the pancreas will be injected with Patent Blue and the lymphatic pathways will be identified in relation to their origin from the theoretical pancreatic segments (ClinicalTrials.gov Identifier: NCT03597230).

Human and animal rights

Authors declare that the work involves the use of animals, and the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans and EU Directive 2010/63/EU for animal experiments.

Submission declaration and verification

The submitted work has not been published previously anywhere.

Author contributions

Y.R., C.P., M.P. and M.B. conceptualized and validated the research. M.L., C.A. and M.P. performed formal analysis, supervised the project and provided critical review of the manuscript. Y.R., R.R. and M.P. performed the original draft and final writing. All authors approved the manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgements

The authors acknowledge Ms Dale Roche-Lebrec, editor of Medical English (PhD), for editing this manuscript.

References

- Akcali, O., Kiray, A., Ergur, I., Tetik, S., Alici, E., 2006. Thoracic duct variations may complicate the anterior spine procedures. *Eur. Spine J.* 15, 1347–1351, <http://dx.doi.org/10.1007/s00586-006-0082-3>.
- Amundadottir, L.T., 2016. Pancreatic Cancer Genetics. *Int. J. Biol. Sci.* 12, 314–325, <http://dx.doi.org/10.7150/ijbs.15001>.
- Beisani, M., Roca, I., Cardenas, R., Blanco, L., Abu-Suboh, M., Dot, J., Armengol, J.R., Olsina, J.J., Balsells, J., Charco, R., Castell, J., 2016. Initial experience in sentinel lymph node detection in pancreatic cancer. *Rev. Esp. Med. Nucl. Imagen. Mol.* 35, 287–291, <http://dx.doi.org/10.1016/j.remnm.2015.10.006>.
- Bensley, R.R., 2005. Studies on the pancreas of the guinea pig. *Am. J. Anat.* 12, 297–388, <http://dx.doi.org/10.1002/aja.1000120304>.
- Borghi, F., Gattolin, A., Garbossa, D., Bogliatto, F., Garavoglia, M., Levi, A.C., 1998. Embryologic bases of extended radical resection in pancreatic cancer. *Arch. Surg.* 133, 297–301.
- Cesmebasi, A., Malefant, J., Patel, S.D., Du Plessis, M., Renna, S., Tubbs, R.S., Loukas, M., 2015. The surgical anatomy of the lymphatic system of the pancreas. *Clin. Anat.* 28, 527–537, <http://dx.doi.org/10.1002/ca.22461>.
- Cubilla, A.L., Fortner, J., Fitzgerald, P.J., 1978. Lymph node involvement in carcinoma of the head of the pancreas area. *Cancer* 41, 880–887, [http://dx.doi.org/10.1002/1097-0142\(197803\)41:3<880::AID-CNCR2820410315>3.0.CO;2-J](http://dx.doi.org/10.1002/1097-0142(197803)41:3<880::AID-CNCR2820410315>3.0.CO;2-J).
- Deki, H., Sato, T., 1988. An anatomic study of the peripancreatic lymphatics. *Surg. Radiol. Anat.* 10, 121–135.
- Delpero, J.R., Jeune, F., Bachellier, P., Regenet, N., Le Treut, Y.P., Paye, F., Carriere, N., Sauvanet, A., Adham, M., Autret, A., Poizat, F., Turrini, O., Boher, J.M., 2017. prognostic value of resection margin involvement after pancreatoduodenectomy for ductal adenocarcinoma: updates from a French prospective multicenter study. *Ann. Surg.* 266, 787–796, <http://dx.doi.org/10.1097/SLA.0000000000002432>.

- Donatini, B., Hidden, G., 1992. Routes of lymphatic drainage from the pancreas: a suggested segmentation. *Surg. Radiol. Anat.* 14, 35–42.
- Evans, B.P., Ochsner, A., 1954. The gross anatomy of the lymphatics of the human pancreas. *Surgery* 36, 177–191.
- Ferlay, J., Partensky, C., Bray, F., 2016. More deaths from pancreatic cancer than breast cancer in the EU by 2017. *Acta Oncol.* 55, 1158–1160, <http://dx.doi.org/10.1080/0284186X.2016.1197419>.
- Ferrer, J., Scott, W.E., Weegman, B.P., Suszynski, T.M., Sutherland, D.E.R., Hering, B.J., Pappas, K.K., 2008. Pig pancreas anatomy: implications for pancreas procurement, preservation, and islet isolation. *Transplantation* 86, 1503–1510, <http://dx.doi.org/10.1097/TP.0b013e31818bfda1>.
- Fink, D.M., Steele, M.M., Hollingsworth, M.A., 2016. The lymphatic system and pancreatic cancer. *Cancer Lett.* 381, 217–236, <http://dx.doi.org/10.1016/j.canlet.2015.11.048>.
- Galindo-Pacheco, Y., de Santis Prada, L.L., Miglino, M.A., DiDio, L.J., 1992. Pancreatic arteries and anatomicosurgical segments in dogs. *Ital. J. Anat. Embryol.* 97, 273–282.
- Hirai, I., Murakami, G., Kimura, W., Tanuma, K., Ito, H., 2001. Origin of the thoracic duct and pancreaticoduodenal lymphatic pathways to the para-aortic lymph nodes. *J. Hepatobiliary Pancreat. Surg.* 8, 441–448.
- Holt, B.A., Varadarajulu, S., 2014. Features of chronic pancreatitis and associated masses: a focus on endosonography. *Video J. Encycl. G.I. Endosc.* 2, 50–54.
- Isaji, S., Kawarada, Y., Uemoto, S., 2004. Classification of pancreatic cancer: comparison of Japanese and UICC classifications. *Pancreas* 28, 231–234.
- Ji, R.C., Kato, S., 1997. Demonstration of the intralobular lymphatics in the guinea pig pancreas by an enzyme-histochemical method. *J. Anat.* 191 (Pt 1), 15–22.
- Kanda, M., Fujii, T., Nagai, S., Kodera, Y., Kanzaki, A., Sahin, T.T., Hayashi, M., Yamada, S., Sugimoto, H., Nomoto, S., Takeda, S., Morita, S., Nakao, A., 2011. Pattern of lymph node metastasis spread in pancreatic cancer. *Pancreas* 40, 951–955, <http://dx.doi.org/10.1097/MPA.0b013e3182148342>.
- Kayahara, M., Nagakawa, T., Kobayashi, H., Mori, K., Nakano, T., Kadoya, N., Ohta, T., Ueno, K., Miyazaki, I., 1992. Lymphatic flow in carcinoma of the head of the pancreas. *Cancer* 70, 2061–2066.
- Kayahara, M., Nagakawa, T., Ohta, T., Kitagawa, H., Ueno, K., Tajima, H., Elnemr, A., Miwa, K., 1999. Analysis of paraaortic lymph node involvement in pancreatic carcinoma: a significant indication for surgery? *Cancer* 85, 583–590.
- Nagakawa, T., Kobayashi, H., Ueno, K., Ohta, T., Kayahara, M., Miyazaki, I., 1994. Clinical study of lymphatic flow to the paraaortic lymph nodes in carcinoma of the head of the pancreas. *Cancer* 73, 1155–1162.
- Nagakawa, T., Konishi, I., Ueno, K., Ohta, T., Kayahara, M., 1993. A clinical study on lymphatic flow in carcinoma of the pancreatic head area — peripancreatic regional lymph node grouping. *Hepatogastroenterology* 40, 457–462.
- Neuzillet, C., Tijeras-Raballand, A., Bourget, P., Cros, J., Couvelard, A., Sauvanet, A., Vullierme, M.-P., Tournigand, C., Hammel, P., 2015. State of the art and future directions of pancreatic ductal adenocarcinoma therapy. *Pharmacol. Ther.* 155, 80–104, <http://dx.doi.org/10.1016/j.pharmthera.2015.08.006>.
- Niebling, M.G., Pleijhuis, R.G., Bastiaannet, E., Brouwers, A.H., van Dam, G.M., Hoekstra, H.J., 2016. A systematic review and meta-analyses of sentinel lymph node identification in breast cancer and melanoma, a plea for tracer mapping. *Eur. J. Surg. Oncol.* 42, 466–473, <http://dx.doi.org/10.1016/j.ejso.2015.12.007>.
- Nimura, Y., Nagino, M., Takao, S., Takada, T., Miyazaki, K., Kawarada, Y., Miyagawa, S., Yamaguchi, A., Ishiyama, S., Takeda, Y., Sakoda, K., Kinoshita, T., Yasui, K., Shimada, H., Katoh, H., 2012. Standard versus extended lymphadenectomy in radical pancreatoduodenectomy for ductal adenocarcinoma of the head of the pancreas: long-term results of a Japanese multicenter randomized controlled trial. *J. Hepatobiliary Pancreat. Sci.* 19, 230–241, <http://dx.doi.org/10.1007/s00534-011-0466-6>.
- O'Morchoe, C.C., 1997. Lymphatic system of the pancreas. *Microsc. Res. Tech.* 37, 456–477, [http://dx.doi.org/10.1002/\(SICI\)1097-0029\(19970601\)37:5<456::AID-JEMT9>3.0.CO;2-B](http://dx.doi.org/10.1002/(SICI)1097-0029(19970601)37:5<456::AID-JEMT9>3.0.CO;2-B).
- Pavlidis, T.E., Pavlidis, E.T., Sakantamis, A.K., 2011. Current opinion on lymphadenectomy in pancreatic cancer surgery. *Hepatobiliary Pancreat. Dis. Int.* 10, 21–25.
- Peek, M.C., Charalampoudis, P., Anninga, B., Baker, R., Douek, M., 2017. Blue dye for identification of sentinel nodes in breast cancer and malignant melanoma: a systematic review and meta-analysis. *Future Oncol. Lond. Engl.* 13, 455–467, <http://dx.doi.org/10.2217/fo-2016-0255>.
- Pissas, A., 1990. Clinical and surgical anatomy studies of the lymphatic circulation of the pancreas. *Bull. Mem. Acad. R. Med. Belg.* 145, 351–364.
- Pissas, A., 1984. Anatomoclinical and anatomosurgical essay on the lymphatic circulation of the pancreas. *Anat. Clin.* 6, 255–280.
- Pissas, A., Bouchet, Y., Sarrazin, R., 1982. Critical analysis of the different techniques available for the study of the lymphatic drainage of the viscera, and their clinical value. *Anat. Clin.* 4, 315–322, <http://dx.doi.org/10.1007/BF01798561>.
- Pu, N., Li, J., Xu, Y., Lee, W., Fang, Y., Han, X., Zhao, G., Zhang, L., Nuerxiati, A., Yin, H., Wu, W., Lou, W., 2018. Comparison of prognostic prediction between nomogram based on lymph node ratio and AJCC 8th staging system for patients with resected pancreatic head carcinoma: a SEER analysis. *Cancer Manag. Res.* 10, 227–238, <http://dx.doi.org/10.2147/CMAR.S157940>.
- Rana, S.S., Vilmann, P., 2015. Endoscopic ultrasound features of chronic pancreatitis: a pictorial review. *Endosc. Ultrasound* 4, 10–14.
- Regoli, M., Bertelli, E., Orazioli, D., Fonzi, L., Bastianini, A., 2001. Pancreatic lymphatic system in rodents. *Anat. Rec.* 263, 155–160, <http://dx.doi.org/10.1002/ar.1090>.
- Renard, Y., de Mestier, L., Perez, M., Avisse, C., Lévy, P., Kianmanesh, R., 2017. Unraveling pancreatic segmentation. *World J. Surg.*, <http://dx.doi.org/10.1007/s00268-017-4263-5>.
- Riediger, H., Keck, T., Wellner, U., zur Hausen, A., Adam, U., Hopt, U.T., Makowicz, F., 2009. The lymph node ratio is the strongest prognostic factor after resection of pancreatic cancer. *J. Gastrointest. Surg. Tract* 13, 1337–1344, <http://dx.doi.org/10.1007/s11605-009-0919-2>.
- Samra, J.S., Gananadha, S., Hugh, T.J., 2008. Surgical management of carcinoma of the head of pancreas: extended lymphadenectomy or modified en bloc resection? *ANZ J. Surg.* 78, 228–236, <http://dx.doi.org/10.1111/j.1445-2197.2008.04426.x>.
- Santi, I., Brandt, A., Hemminki, K., 2011. What is the major prognostic factor in tumor-node-metastasis staging of pancreatic adenocarcinoma? *Ann. Surg. Oncol.* 18, 300–301, <http://dx.doi.org/10.1245/s10434-010-1189-7>.
- Sauvanet, A., 2008. Lymph node resection for carcinoma of the pancreas. *J. Chir.* 145, 12S31–12S35, Spec no. 4.
- Shokouh-Amiri, M.H., Rahimi-Saber, S., Andersen, H.O., 1989. Segmental pancreatic autotransplantation in the pig. *Transplantation* 47, 42–44.
- Siegel, R.L., Miller, K.D., Jemal, A., 2016. Cancer statistics, 2016. *CA Cancer J. Clin.* 66, 7–30, <http://dx.doi.org/10.3322/caac.21332>.
- Suda, K., Nobukawa, B., Takase, M., Hayashi, T., 2006. Pancreatic segmentation on an embryological and anatomical basis. *J. Hepatobiliary Surg.* 13, 146–148.
- Tol, J.A.M.G., Gouma, D.J., Bassi, C., Dervenis, C., Montorsi, M., Adham, M., Andrén-Sandberg, A., Asbun, H.J., Bockhorn, M., Büchler, M.W., Conlon, K.C., Fernández-Cruz, L., Fingerhut, A., Friess, H., Hartwig, W., Izbicki, J.R., Lillemoe, K.D., Milicevic, M.N., Neoptolemos, J.P., Shrikhande, S.V., Vollmer, C.M., Yeo, C.J., Charnley, R.M., 2014. Definition of a standard lymphadenectomy in surgery for pancreatic ductal adenocarcinoma: a consensus statement by the International Study Group on Pancreatic Surgery (ISGPS). *Surgery* 156, 591–600, <http://dx.doi.org/10.1016/j.surg.2014.06.016>.
- Tomson, D., Lessert, C., Luthi, I., Mazzolai, L., Depairon, M., 2015. Lymphatic vascular pathologies: contribution of lympho-fluoroscopy. *Rev. Med. Suisse* 11, 362–365.
- Troisi, R., Maene, L., Jacobs, B., Berrevoet, F., de Hemptinne, B., Hesse, U.J., 2000. Segmental porcine pancreatic autotransplantation as model for pancreas preservation studies using two different techniques for vascular reconstruction. *J. Invest. Surg.* 13, 203–211.
- Tsushitani, M., Sato, J., Kokoshima, H., 2016. A comparison of the anatomical structure of the pancreas in experimental animals. *J. Toxicol. Pathol.* 29, 147–154, <http://dx.doi.org/10.1293/tox.2016-0016>.
- Uchida, T., Takada, T., Ammori, B.J., Suda, K., Takahashi, T., 1999. Three-dimensional reconstruction of the ventral and dorsal pancreas: a new insight into anatomy and embryonic development. *J. Hepatobiliary Surg.* 6, 176–180.
- Weinberg, J., Greaney, E.M., 1950. Identification of regional lymph nodes by means of a vital staining dye during surgery of gastric cancer. *Surg. Gynecol. Obstet.* 90, 561–567.
- Zhan, H., Xu, J., Wang, L., Zhang, G., Hu, S., 2015. Lymph node ratio is an independent prognostic factor for patients after resection of pancreatic cancer. *World J. Surg. Oncol.* 13, 105, <http://dx.doi.org/10.1186/s12957-015-0510-0>.