



Assessment of the reliability of the motor unit size index (MUSIX) in single subject “round-robin” and multi-centre settings



James J.P. Alix^{a,*,1}, Christoph Neuwirth^{b,1}, Lucy Gelder^c, Christian Burkhardt^b, José Castro^d, Mamede de Carvalho^d, Malgorzata Gawel^e, Stephan Goedee^f, Julian Grosskreutz^g, Timothée Lenglet^h, Cristina Mogliaⁱ, Taha Omer^j, Maarten Schrooten^k, Sanjeev Nandedkar^l, Erik Stalberg^m, Paul E. Barkhausⁿ, Jasna Furtula^o, Johannes P. van Dijk^p, Reto Baldinger^b, Joao Costa^d, Marit Otto^o, Arne Sandberg^m, Markus Weber^b

^aSheffield Institute for Translational Neuroscience, University of Sheffield, 385A Glossop Road, Sheffield S10 2HQ, UK

^bNeuromuscular Diseases Unit/ALS Clinic, Kantonsspital, St. Gallen, Switzerland

^cStatistical Services Unit, University of Sheffield, UK

^dDepartment of Neurosciences, Hospital de Santa Maria, Instituto de Medicina Molecular, Faculty of Medicine, University of Lisbon, Portugal

^eDepartment of Neurology, Medical University of Warsaw, Warsaw, Poland

^fBrain Centre Rudolf Magnus, Department of Neurology and Neurosurgery, UMC Utrecht, Utrecht, The Netherlands

^gHans-Berger Department of Neurology, Jena University Hospital, Jena, Germany

^hDépartement de Neurophysiologie, Groupe hospitalier Pitié-Salpêtrière, APHP, Paris, France

ⁱALS Centre of Torino, Department of Neuroscience “Rita Levi Montalcini”, University of Torino, Torino, Italy

^jTrinity College Biomedical Science Institute (TBSI) and Beaumont Hospital, Dublin, Ireland

^kDepartment of Neurology, University Hospital Leuven, Leuven, Belgium

^lNatus Medical, Inc., 15 Dartantra Drive, Hopewell Junction, NY 12533, USA

^mDepartment of Neuroscience, Clinical Neurophysiology, Uppsala University, Sweden

ⁿMilwaukee Veterans Administration Medical Center and Medical College of Wisconsin, Milwaukee, WI, USA

^oDepartment of Clinical Neurophysiology, Aarhus University Hospital, Aarhus, Denmark

^pDepartment of Orthodontics, University of Ulm, Ulm, Germany

ARTICLE INFO

Article history:

Accepted 3 January 2019

Available online 21 February 2019

Keywords:

Motor unit number index

Motor unit size index

Motor unit

Motor neuron

Amyotrophic lateral sclerosis

Electrophysiology

HIGHLIGHTS

- Motor unit size index (MUSIX) is a rapid, non-invasive means of assessing reinnervation.
- Using healthy volunteers reliable data were provided by both experienced and non-experienced raters.
- Results across single subject and multi-centre settings demonstrate high reliability/reproducibility.

ABSTRACT

Objective: The motor unit size index (MUSIX) is incorporated into the motor unit number index (MUNIX). Our objective was to assess the intra-/inter-rater reliability of MUSIX in healthy volunteers across single subject “round robin” and multi-centre settings.

Methods: Data were obtained from (i) a round-robin assessment in which 12 raters (6 with prior experience and 6 without) assessed six muscles (abductor pollicis brevis, abductor digiti minimi, biceps brachii, tibialis anterior, extensor digitorum brevis and abductor hallucis) and (ii) a multi-centre study with 6 centres studying the same muscles in 66 healthy volunteers. Intra/inter-rater data were provided by 5 centres, 1 centre provided only intra-rater data.

Intra/inter-rater variability was assessed using the coefficient of variation (COV), Bland-Altman plots, bias and 95% limits of agreement.

Abbreviations: ADM, abductor digiti minimi; AH, abductor hallucis; ALS, amyotrophic lateral sclerosis; APB, abductor pollicis brevis; CMAP, compound muscle action potential; COV, coefficient of variation; EDB, extensor digitorum brevis; BB, biceps brachii; EMG, electromyography; LOA, limits of agreement; MOVER, method of variance recovery for ratios; MUNIX, motor unit number index; MUSIX, motor unit size index; SD, standard deviation; SIP, surface interference pattern; TA, tibialis anterior.

* Corresponding author.

E-mail address: j.alix@sheffield.ac.uk (J.J.P. Alix).

¹ These authors contributed equally.

<https://doi.org/10.1016/j.clinph.2019.01.020>

1388-2457/© 2019 International Federation of Clinical Neurophysiology. Published by Elsevier B.V. All rights reserved.

Results: In the round-robin assessment intra-rater COVs for MUSIX ranged from 7.8% to 28.4%. Inter-rater variability was between 7.8% and 16.2%. Prior experience did not impact on MUSIX values. In the multi-centre study MUSIX was more consistent than the MUNIX. Abductor hallucis was the least reliable muscle.

Conclusions: The MUSIX is a reliable neurophysiological biomarker of reinnervation.

Significance: MUSIX could provide insights into the pathophysiology of a range of neuromuscular disorders, providing a quantitative biomarker of reinnervation.

© 2019 International Federation of Clinical Neurophysiology. Published by Elsevier B.V. All rights reserved.

1. Introduction

Motor unit number techniques have been significantly developed since their inception over 30 years ago (McComas et al., 1971). The motor unit number index (MUNIX), first reported in 2004 (Nandedkar et al., 2004), has grown in popularity in recent years due to its ease and speed of application across different muscles (Nandedkar et al., 2010). As a result, it has been employed across a range of neuromuscular conditions, from amyotrophic lateral sclerosis (ALS) (Neuwirth et al., 2015; Escorcio-Bezerra et al., 2016) to demyelinating peripheral neuropathies (Delmont et al., 2016; Philibert et al., 2017). In addition to providing an index of the number of functional motor units, the MUNIX computation also includes a size index, termed the motor unit size index (MUSIX) which is derived from dividing the compound muscle action potential (CMAP) amplitude by the MUNIX value (Nandedkar et al., 2004, 2010). The MUSIX has the possibility of detailing the reinnervation process (e.g. reflecting motor unit size) in neuromuscular disorders which may reveal important information on the natural course of such conditions and the response to new treatments.

Although reports on the MUNIX technique have documented a change in the MUSIX in patients with ALS (Nandedkar et al., 2010; Escorcio-Bezerra et al., 2016), the MUSIX has received little attention. The first step in evaluating its ability to provide novel insights into pathophysiological changes in human subjects is to assess its reliability. We have previously assessed MUNIX in this regard, reporting its reliability in healthy subjects across two settings: a round-robin style assessment of a single healthy subject by multiple raters and a multi-centre study of 66 healthy volunteers (Neuwirth et al., 2011, 2016). In order to ascertain the reliability of the MUSIX, we have analysed data from those studies in order to gauge its utility as a parameter in clinical trials of neuromuscular diseases.

2. Methods

The MUNIX/MUSIX procedure has been described in detail previously (Nandedkar et al., 2004, 2010). A CMAP is recorded and optimised for maximal amplitude using standard neurophysiological techniques and the test subject then asked to contract the given muscle with differing force levels. The computation from these data has been described previously (Nandedkar et al., 2004).

The studies concerning the round-robin MUNIX assessment (Neuwirth et al., 2016) and the multi-centre evaluation of healthy volunteer subjects have been reported previously (Neuwirth et al., 2011). The multi-centre study was approved by the relevant Human Studies Committees in each centre and written consent was obtained for both studies. Briefly, for the round-robin assessment a two-day training course was undertaken by neurophysiologists from across Europe at an ENCALS (European Network for the Cure of ALS) meeting in Dublin 2015. 12 participants were included

in the study, 6 of whom had already completed a prior qualification process for a longitudinal study and were termed “experienced raters”. 6 participants with no prior experience of MUNIX/MUSIX, but participated in the training, were also included and termed “non-experienced raters”. All 12 raters completed two MUNIX/MUSIX assessments of a single healthy subject (M.W) on consecutive days. Assessments were performed using a Keypoint Dantec Focus EMG system. All muscles tested were on the right side: abductor pollicis brevis (APB), abductor digiti minimi (ADM), biceps brachii (BB), tibialis anterior (TA), extensor digitorum brevis (EDB) and abductor hallucis (AH).

In the multi-centre study healthy volunteers were recruited from six centres – Aarhus, Denmark; Lisbon, Portugal; Milwaukee, Wisconsin, USA; Nijmegen, Netherlands; St. Gallen, Switzerland and Uppsala, Sweden (Neuwirth et al., 2011). The demographic details and inclusion/exclusion criteria have been reported previously; briefly potential participants with any medical condition that could impact on MUNIX measurements were excluded (Neuwirth et al., 2011). The study was approved by the relevant Human Studies Committee in each centre and informed consent obtained. Each centre used the same electrodes and measurements were made on Keypoint Classic- and Synergy machines. Recordings were undertaken on right APB, ADM, BB, TA and AH muscles. The EDB muscle was also measured in 32 healthy volunteers across four centres. As documented previously, two investigators performed measurements twice in alternation in four centres (Neuwirth et al., 2011). In one centre only intra-rater variability was assessed (i.e. the same investigator performed measurements twice). In another centre, a first examiner undertook measurements twice and a second examiner performed the recordings once.

2.1. Statistics

The coefficient of variation (COV; standard deviation/mean) was calculated for each muscle. Variability was calculated by: $100 \times (\text{difference of test-retest}) / (\text{mean of test-retest})$. Welch's t-test was used to compare MUSIX values for the experienced and non-experienced rater groups in the round-robin assessment. A hypothetical MUSIX reference value was generated for the round-robin assessment in the same way reported by Neuwirth et al., except on this occasion substituting MUSIX for MUNIX (Neuwirth et al., 2016). Briefly, for each muscle the 6 largest CMAPs taken from the mean of test-retest measurements were determined for each muscle. Of these 6, the 3 test-retest measurements with the smallest variability were used to calculate the “reference” CMAP amplitudes and the corresponding MUSIX values. The difference between the observed measurements and the hypothetical reference was determined and reported as accuracy, expressed as a percentage of the reference value.

For the multi-centre data, inter-rater variability in each study centre was assessed through Bland–Altman plots on log-

transformed data. Displayed are the line of equality (i.e. where the data would lie if both examiners had exact agreement), the mean difference between examiners (bias), and 95% limits of agreement (LOA) for each comparison. The LOA represent the interval of two standard deviations of the measurement differences either side of the mean difference, confidence intervals surrounding these LOA were calculated using the MOVER (method of variance estimates recovery) method (Zou, 2013). Analyses were performed in Statistical Analysis Software, version 9.3.

3. Results

In the single healthy subject round-robin assessment, test-retest data are summarised in Fig. 1 and documented in detail in Tables 1 and 2. Intra-rater COVs for MUSIX ranged from 7.8% in APB to 28.4% in EDB (Table 1). Inter-rater COV ranges were from 7.8% (BB) to 16.2% (AH) (Table 2). Welch's t-test found no evidence of a difference in the values obtained between experienced and non-experienced examiners for any of the muscles examined ($0.19 < p < 0.79$, Table 3). Analysis of the relative mean and standard deviation of the MUSIX measurements of the experienced and non-experienced groups revealed a high degree of accuracy

for all muscles, with the possible exception of ADM and EDB (Fig. 2).

In the multi-centre study of healthy volunteers, MUSIX inter-rater variability for each muscle was examined for all centres, except one where there was only a single examiner (Table 4; complete dataset shown in Supplementary Table 1). AH was the least reliably measured muscle in centres 2–5, and the second least for centre 1. Box plots of the LOA to assess the variability of MUSIX, MUNIX and CMAP over all centres and muscles demonstrated that, on average, the MUNIX measurements had a greater inter-rater variability (Fig. 3; see also Supplementary Table 1). Example inter-rater Bland–Altman plots for APB MUSIX are shown in Fig. 4 (plots for all other muscles can be found in Supplementary Fig. 1), and reveal few points beyond the LOA.

Intra-rater MUSIX variability was assessed in a similar fashion for all examiners, except examiner 8 for whom only data from one visit was available (Table 5; full data set Supplementary Table 2). Example MUSIX intra-rater Bland–Altman plots for the APB muscle for 4 examiners are shown in Fig. 5 (remaining plots for APB and all other muscles available in Supplementary Fig. 2). AH was again the muscle with the most variation between measurements, although EDB was fairly similar. In addition, APB and ADM also showed more variable measurements than BB or TA. As




Fig. 1. Test-retest results for MUSIX in individual muscles in the single subject round-robin study. The dotted line represents a hypothetical reference value (see methods).

Table 1

Intra-rater COV (%) and variability (%) for MUSIX measurements in individual muscles in the single healthy subject study.

Muscle	All raters		Experienced raters		Non-experienced raters	
	COV	Variability	COV	Variability	COV	Variability
APB	8.8	12.4	9.7	13.8	7.8	11.1
ADM	9.7	13.7	10.2	14.5	9.1	12.9
BB	12.5	17.7	16.2	23.0	9.4	13.3
TA	10.1	14.3	8.9	12.6	11.2	15.9
EDB	24.0	33.9	19.6	27.7	28.4	40.1
AH	14.8	20.9	12.0	16.9	17.6	24.9

Table 2

Inter-rater variability for MUSIX expressed as COV (in %) in individual muscles for the first and second measurement and mean of both values (single healthy subject round-robin study).

	Mean of measurements COV	Measurement 1 COV	Measurement 2 COV
<i>Muscle</i>			
APB	11.1	14.6	12.5
ADM	11.2	15.8	12.4
BB	7.8	14.4	13.0
TA	9.3	12.7	12.4
EDB	14.4	21.9	20.7
AH	16.2	14.3	24.8

with inter-rater variation, MUSIX was more variable than either MUSIX or CMAP (Fig. 6). Lastly, MUSIX variability had only a very limited relationship to CMAP, with the CMAP measurement contributing between 0.3% and 3.3% to MUSIX variability (Table 6).

4. Discussion

There is increasing interest in the MUNIX/MUSIX technique as a rapid, non-invasive means to monitor progression in neuromuscular disease. In two previous reports our team detailed the reliability of the MUNIX values (Neuwirth et al., 2011, 2016), but little attention was given to the consistency of the MUSIX. Since the rate of symptom progression in neuromuscular diseases is determined by both the amount of axon loss and effective reinnervation (Gordon et al., 2004; Van Asseldonk et al., 2006), MUSIX could provide valuable information regarding the reinnervation process. Such data could provide important insights into how certain conditions progress and how they respond to new treatments. This paper combines data from two previous studies to investigate whether the MUSIX can be a reliable measurement of motor unit size for future clinical studies.

The dominant finding of the round-robin study is that the coefficient of variability for MUSIX is below 20% for all muscles, except EDB. This finding is similar to the initial report of MUNIX variability, in which the small muscles of the foot (AH and EDB) were also the worst performing (Neuwirth et al., 2016). The reasons for this are multiple. As the MUNIX values obtained for EDB are often smaller than other muscles (which was especially true in the single healthy volunteer study) any variability in the absolute values of EDB measurements will result in a larger relative variability and COV. Furthermore, the peroneal nerve branch to EDB is susceptible to injury and so may changes may not reflect the disease process of interest. Regarding AH, multiple small foot muscles contribute to the recorded measurements of the AH CMAP (Nandedkar and Barkhaus, 2007) and as some individuals find activation of this muscle difficult, the recordings may be susceptible to volume conducted artefact. It is therefore possible that the muscles contributing to the CMAP are different to those activated during voluntary




Fig. 2. Relative mean and standard deviation of MUSIX measurements in individual muscles of the experienced group (filled circles) and non-experienced group (empty circles) compared to the hypothetical reference values, expressed as accuracy (%).

contraction. We would suggest that such technical pitfalls outweigh the benefit of including AH in study protocols and advocate omission from future studies.

Interestingly, in the multi-centre study MUSIX was more consistent, both in intra- and inter-rater assessments, than the MUNIX. Our analyses suggest that one explanation of this observation is that MUSIX is independent of CMAP variability, which is in keeping with prior reports and the opposite of MUNIX, for which variability is highly dependent on the CMAP (Nandedkar et al., 2010). We have previously documented that the relative contribution of the CMAP to MUNIX variation may be as high as 90% which emphasises the importance of CMAP optimisation to the MUNIX measurement (Neuwirth et al., 2011). The underlying cause of MUSIX variation is less clear but may be found in the distribution of surface interference patterns (SIP) across weak and strong contractions.

Overall, comparing the performance MUNIX and MUSIX in the other muscles sampled across both study settings reveals a similar level of performance (Neuwirth et al., 2011, 2016). Encouragingly, there was no significant difference in the values obtained for MUSIX between those with prior experience in the technique and those unfamiliar with it (but with training in standard nerve conduction studies/EMG). This would suggest that that following training, obtaining high quality recordings can be rapidly achieved

Table 3

Mean and standard deviation (SD) of MUSIX values across the groups. All muscles grouped together for Welch's t test.

	MUSIX all raters Mean (SD)	MUSIX experienced raters Mean (SD)	MUSIX non-experienced raters Mean (SD)	Welch's t tests (experience, vs. non-experienced.) p-value
<i>Muscle</i>				
APB	56.12 (7.8)	54.32 (7.16)	57.93 (7.66)	0.61
ADM	67.25 (9.46)	70.28 (9.75)	64.21 (8.5)	0.79
BB	44.16 (5.93)	43.40 (6.64)	44.86 (5.41)	0.32
TA	49.01 (6.02)	48.49 (5.83)	49.53 (6.41)	0.53
EDB	72.04 (17.16)	71.71 (16.60)	72.36 (18.43)	0.45
AH	61.83 (13.57)	66.64 (13.87)	57.02 (11.92)	0.19

Table 4
MUSIX inter-rater variability across all muscles for the different centres.

Centre	Muscle	Mean difference (bias)	Standard deviation difference	Lower LOA	Upper LOA
1	APB	−0.03	0.13	0.21	−0.28
	ADM	−0.01	0.1	0.19	−0.2
	BB	0	0.04	0.08	−0.08
	TA	−0.03	0.09	0.15	−0.21
	AH	0.05	0.015	0.33	−0.24
	EDB	−	−	−	−
2	APB	−0.04	0.09	0.14	−0.21
	ADM	−0.02	0.11	0.2	−0.24
	BB	0.01	0.07	0.14	−0.12
	TA	0.02	0.09	0.18	−0.15
	AH	−0.03	0.14	0.24	−0.3
	EDB	−0.09	0.14	0.18	−0.36
3	APB	−0.06	0.1	0.14	−0.25
	ADM	−0.04	0.09	0.14	−0.22
	BB	0	0.09	0.17	−0.17
	TA	−0.03	0.07	0.1	−0.16
	AH	−0.1	0.18	0.26	−0.46
	EDB	−0.1	0.18	0.26	−0.46
4	APB	0.03	0.1	0.22	−0.16
	ADM	0.01	0.09	0.18	−0.17
	BB	−0.02	0.04	0.07	−0.1
	TA	0	0.05	0.11	−0.11
	AH	−0.01	0.17	0.32	−0.34
	EDB	0.02	0.11	0.22	−0.19
5	APB	0	0.08	0.16	−0.15
	ADM	0	0.08	0.16	−0.17
	BB	−0.02	0.04	0.06	−0.1
	TA	0	0.06	0.12	−0.13
	AH	0.05	0.15	0.35	−0.25
	EDB	0.05	0.17	0.39	−0.13

without prior experience. We would suggest that MUSIX values are an even more reliable than MUNIX values.

However, in keeping with the original multi-centre MUNIX paper, our new analysis demonstrates that there is a degree of vari-

ability between different centres and different muscles (Neuwirth et al., 2011). As noted in the report of Neuwirth et al., training is likely to be important in ensuring consistency of the MUNIX/MUSIX technique across individuals/centres and a training




Fig. 3. Box plots demonstrating the distribution of inter-rater limits of agreement for CMAP/MUNIX/MUSIX across the different centres. The horizontal line represents the median, the larger open circles the mean.




Fig. 4. Example inter-rater MUSIX Bland-Altman plots across participating centres for the APB muscle.

Table 5

MUSIX intra-rater variability across all muscles for the different examiners in the multi-centre study on healthy volunteers.

Examiner	Muscle	Mean difference (bias)	Standard deviation difference	Lower LOA	Upper LOA
1	APB	-0.03	0.12	-0.26	0.21
	ADM	0	0.11	-0.21	0.22
	BB	0	0.03	-0.07	0.07
	TA	0	0.06	-0.12	0.12
	AH	0	0.12	-0.24	0.23
2	EDB	-0.01	0.07	-0.14	0.12
	APB	0.05	0.13	-0.21	0.3
	ADM	0.01	0.12	-0.21	0.24
	BB	0	0.06	-0.13	0.12
	TA	0	0.11	-0.21	0.22
3	AH	0.04	0.12	-0.2	0.28
	EDB	-	-	-	-
	APB	-0.03	0.12	-0.26	0.210
	ADM	0.03	0.12	-0.21	0.27
	BB	-0.02	0.05	-0.12	0.08
4	TA	-0.04	0.08	-0.2	0.11
	AH	-0.03	0.14	-0.3	0.24
	EDB	-0.03	0.11	-0.24	0.17
	APB	0.01	0.08	-0.16	0.17
	ADM	0.05	0.12	-0.18	0.29
5	BB	0.03	0.09	-0.15	0.2
	TA	-0.02	0.09	-0.2	0.15
	AH	-0.05	0.12	-0.28	0.18
	EDB	0.02	0.17	-0.32	0.36
	APB	-0.07	0.07	-0.21	0.08
6	ADM	-0.01	0.11	-0.23	0.2
	BB	0.01	0.07	-0.12	0.14
	TA	-0.01	0.06	-0.12	0.1
	AH	0.09	0.14	-0.19	0.36
	EDB	0.02	0.17	-0.32	0.36
7	APB	-0.02	0.06	-0.14	0.1
	ADM	-0.02	0.08	-0.18	0.14
	BB	0.01	0.06	-0.12	0.13
	TA	0.04	0.09	-0.14	0.21
	AH	0.15	0.21	-0.27	0.58
8	EDB	0.01	0.06	-0.12	0.13
	APB	0.03	0.06	-0.9	0.16
	ADM	0.04	0.09	-0.15	0.22
	BB	0.01	0.04	-0.07	0.08
	TA	0.01	0.04	-0.07	0.09
9	AH	0.04	0.13	-0.21	0.28

(continued on next page)

Table 5 (continued)

Examiner	Muscle	Mean difference (bias)	Standard deviation difference	Lower LOA	Upper LOA
9	EDB	0	0.07	-0.15	0.14
	APB	0	0.08	-0.16	0.15
	ADM	0.01	0.05	-0.08	0.11
	BB	0	0.05	-0.1	0.09
	TA	-0.02	0.07	-0.17	0.12
	AH	0.01	0.1	-0.18	0.21
10	EDB	0.06	0.09	-0.12	0.25
	APB	0.01	0.09	-0.16	0.18
	ADM	-0.01	0.08	-0.17	0.15
	BB	0	0.04	-0.08	0.08
	TA	0.03	0.06	-0.08	0.15
	AH	0.03	0.13	-0.22	0.27
11	EDB	-0.02	0.15	-0.32	0.28
	APB	0.02	0.07	-0.13	0.17
	ADM	0.01	0.07	-0.13	0.15
	BB	-0.04	0.05	-0.15	0.07
	TA	0.01	0.05	-0.09	0.1
	AH	0.02	0.08	-0.14	0.18
	EDB	0.02	0.08	-0.14	0.18




Fig. 5. Example intra-rater MUSIX Bland-Altman plots for 4 examiners and the APB muscle.

experience appeared to negate any effects of unfamiliarity with the technique in our round-robin study (Neuwirth et al., 2016; Nandedkar et al., 2018). This training course based approach was not implemented for the multi-centre study which may explain some of the variation. Recent multi-centre studies (e.g. Sampling and biomarker OPTimization and Harmonization In ALS and other motor neuron diseases (SOPHIA) and the ongoing Biogen trials) have required a training course attendance and completion of test-retest assessments on healthy volunteers prior to commencing recordings on patients, which appears to reduce variability (Neuwirth et al., 2018). Several members of our team have recently published guidelines on performing the MUNIX/MUSIX technique which should provide a further step towards standardisation of the technique (Nandedkar et al., 2018).

Although only limited data are available for MUSIX in conditions such as ALS, studies have shown the expected inverse relationship between MUNIX and MUSIX in such patients (Nandedkar et al., 2010; Escorcio-Bezerra et al., 2016), with one study documenting a 50% increase in MUSIX over 8 months (Boekestein et al., 2012). As emphasised in previous papers focusing on the MUNIX, the MUSIX is not an anatomical measure of motor unit size but an index underpinned by the MUNIX and CMAP measurements and their own relationship to the number of motor axons (Nandedkar et al., 2010; Neuwirth et al., 2011). The relationship to actual motor unit size is therefore unknown. The MUSIX is calculated by dividing the CMAP amplitude by the MUNIX and, as noted many years ago, different methods to derive motor unit size may result in quite different values (Doherty et al., 1993), thus making comparisons, even




Fig. 6. Box plots demonstrating the distribution of intra-rater limits of agreement for CMAP/MUNIX/MUSIX pooled across the examiners. The horizontal line represents the median, the larger open circles the mean.

Table 6

Linear regression analysis (multiple R-Squared). Relative contribution (%) of CMAP variability for total MUSIX variability. The data are pooled from all participating centres. For the EDB muscle this is only centres A, B, D and E.

		All centres
MUSIX ~ CMAP	ABP	0.26
	ADM	0.95
	BB	3.25
	TA	1.80
	AH	1.44
	EDB	7.11

between neurophysiological techniques, difficult. However, a significant positive relationship has been reported between MUSIX and motor unit potential amplitude obtained through macro-EMG, in keeping with the assertion that MUSIX relates to the size of the motor unit (Sandberg et al., 2011). Nonetheless, preliminary studies have documented the expected increase in MUSIX in conditions such as ALS and so the technique may provide valuable insights into which muscles undergo compensatory re-innervation and which do not. Such information may also be of value in other conditions, for example, in spinal muscle atrophy and neuropathies, providing researchers with a further measurement relevant to the pathophysiology of neurogenic diseases. The rapidity and ease of application of MUNIX/MUSIX will hopefully facilitate uptake and enhance our understanding the interplay between denervation and compensatory re-innervation over time in progressive conditions such as ALS.

5. Conclusion

We conclude that the MUSIX measurement can provide a reliable surrogate marker of re-innervation. The index warrants consideration in future longitudinal studies in diseases with either axon or motor neuron loss and compensatory reinnervation, such as ALS.

Acknowledgements

The authors wish to thank Kathleen Baster, Statistical Services Unit, University of Sheffield for statistical advice. Dr. J Alix and Dr. L Gelder were supported by the National Institute for Health Research (NIHR) Sheffield Biomedical Research Centre (Translational Neuroscience). The work was also supported by the Swiss ALS Foundation. Dr. Barkhaus acknowledges funding from the Dolores and Carroll Fund from the Milwaukee Medical Foundation.

Conflict of interest

The authors declare no conflicting interests.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.clinph.2019.01.020>.

References

- Boekestein WA, Schelhaas HJ, van Putten MJ, Stegeman DF, Zwarts MJ, van Dijk JP. Motor unit number index (MUNIX) versus motor unit number estimation (MUNE): a direct comparison in a longitudinal study of ALS patients. *Clin Neurophysiol* 2012;123:1644–9.
- Delmont E, Benvenuto A, Grimaldi S, Duprat L, Philibert M, Pouget J, et al. Motor unit number index (MUNIX): is it relevant in chronic inflammatory demyelinating polyradiculoneuropathy (CIDP)? *Clin Neurophysiol* 2016;127:1891–4.
- Doherty TJ, Stashuk DW, Brown WF. Determinants of mean motor unit size: impact on estimates of motor unit number. *Muscle Nerve* 1993;16:1326–31.
- Escorcion-Bezerra ML, Abrahao A, de Castro I, Chieia MAT, de Azevedo LA, Pinheiro DS, et al. MUNIX: reproducibility and clinical correlations in Amyotrophic Lateral Sclerosis. *Clin Neurophysiol* 2016;127:2979–84.
- Gordon T, Hegedus J, Tam SL. Adaptive and maladaptive motor axonal sprouting in aging and motoneuron disease. *Neurol Res* 2004;26:174–85.
- McComas AJ, Fawcett PR, Campbell MJ, Sica RE. Electrophysiological estimation of the number of motor units within a human muscle. *J Neurol Neurosurg Psychiatry* 1971;34:121–31.

- Nandedkar SD, Barkhaus PE. Contribution of reference electrode to the compound muscle action potential. *Muscle Nerve* 2007;36:87–92.
- Nandedkar SD, Barkhaus PE, Stalberg EV. Motor unit number index (MUNIX): principle, method, and findings in healthy subjects and in patients with motor neuron disease. *Muscle Nerve* 2010;42:798–807.
- Nandedkar SD, Barkhaus PE, Stalberg EV, Neuwirth C, Weber M. Motor unit number index: guidelines for recording signals and their analysis. *Muscle Nerve* 2018. <https://doi.org/10.1002/mus.26099>.
- Nandedkar SD, Nandedkar DS, Barkhaus PE, Stalberg EV. Motor unit number index (MUNIX). *IEEE Trans Biomed Eng* 2004;51:2209–11.
- Neuwirth C, Barkhaus PE, Burkhardt C, Castro J, Czell D, de Carvalho M, et al. Tracking motor neuron loss in a set of six muscles in amyotrophic lateral sclerosis using the Motor Unit Number Index (MUNIX): a 15-month longitudinal multicentre trial. *J Neurol Neurosurg Psychiatry* 2015;86:1172–9.
- Neuwirth C, Braun N, Claeys KG, Bucelli R, Fournier C, Bromberg M, et al. Implementing Motor Unit Number Index (MUNIX) in a large clinical trial: real world experience from 27 centres. *Clin Neurophysiol* 2018;129:1756–62.
- Neuwirth C, Burkhardt C, Alix J, Castro J, de Carvalho M, Gawel M, et al. Quality control of motor unit number index (MUNIX) measurements in 6 muscles in a single-subject “round-robin” setup. *PLoS One* 2016;11 e0153948.
- Neuwirth C, Nandedkar S, Stalberg E, Barkhaus PE, Carvalho M, Furtula J, et al. Motor Unit Number Index (MUNIX): a novel neurophysiological marker for neuromuscular disorders; test-retest reliability in healthy volunteers. *Clin Neurophysiol* 2011;122:1867–72.
- Philibert M, Grapperon AM, Delmont E, Attarian S. Monitoring the short-term effect of intravenous immunoglobulins in multifocal motor neuropathy using motor unit number index. *Clin Neurophysiol* 2017;128:235–40.
- Sandberg A, Nandedkar SD, Stalberg E. Macro electromyography and motor unit number index in the tibialis anterior muscle: differences and similarities in characterizing motor unit properties in prior polio. *Muscle Nerve* 2011;43:335–41.
- Van Asseldonk JT, Van den Berg LH, Kalmijn S, Van den Berg-Vos RM, Polman CH, Wokke JH, et al. Axon loss is an important determinant of weakness in multifocal motor neuropathy. *J Neurol Neurosurg Psychiatry* 2006;77:743–7.
- Zou GY. Confidence interval estimation for the Bland-Altman limits of agreement with multiple observations per individual. *Stat Methods Med Res* 2013;22:630–42.