



Associations of Growth and Body Composition with Brain Size in Preterm Infants

Katherine A. Bell, MD¹, Lillian G. Matthews, PhD¹, Sara Cherkerzian, ScD¹, Caroline Palmer, BSc¹, Kaitlin Drouin, MA, MS¹, Hunter L. Pepin, MS, RDN, LDN², Deirdre Ellard, MS, RDN, LDN², Terrie E. Inder, MBChB, MD¹, Sara E. Ramel, MD³, and Mandy B. Belfort, MD, MPH¹

Objective To assess the association of very preterm infants' brain size at term-equivalent age with physical growth from birth to term and body composition at term.

Study design We studied 62 infants born at <33 weeks of gestation. At birth and term, we measured weight and length and calculated body mass index. At term, infants underwent air displacement plethysmography to determine body composition (fat and fat-free mass) and magnetic resonance imaging to quantify brain size (bifrontal diameter, biparietal diameter, transverse cerebellar distance). We estimated associations of physical growth (Z-score change from birth to term) and body composition with brain size, adjusting for potential confounders using generalized estimating equations.

Results The median gestational age was 29 weeks (range, 24.0-32.9 weeks). Positive gains in weight and body mass index Z-score were associated with increased brain size. Each additional 100 g of fat-free mass at term was associated with larger bifrontal diameter (0.6 mm; 95% CI, 0.2-1.0 mm), biparietal diameter (0.7 mm; 95% CI, 0.3-1.1 mm), and transverse cerebellar distance (0.3 mm; 95% CI, 0.003-0.5 mm). Associations between fat mass and brain metrics were not statistically significant.

Conclusions Weight and body mass index gain from birth to term, and lean mass—but not fat—at term, were associated with larger brain size. Factors that promote lean mass accrual among preterm infants may also promote brain growth. (*J Pediatr* 2019;214:20-6).

For very preterm infants, the period from birth to term represents a critical window for growth of the body and brain. Greater weight gain in the neonatal intensive care unit (NICU) predicts improved neurodevelopment,¹⁻³ but weight alone is a nonspecific marker of nutrient accretion. Distinguishing fat from lean mass accrual characterizes the quality of weight gain and may provide information about brain growth and development. However, little is known about how the composition of weight gain during the NICU hospitalization is related to neonatal brain size and future neurodevelopmental risk.

A few small studies have linked lean mass at term-equivalent age with improved subsequent neurodevelopment, including faster cognitive processing at 4 months and higher cognitive scores at 1 and 4 years. In contrast, greater fat mass in those studies was associated with either neutral or worse cognitive outcomes.⁴⁻⁶ Thus, lean mass at term may be a useful biomarker for brain growth and development. Brain size can be quantified using magnetic resonance imaging (MRI), and larger brain size at term predicts better neurodevelopmental outcomes.⁷⁻⁹ However, studies of body composition and brain size are both sparse and conflicting. One small (n = 42) study found that both lean and fat mass correlated positively with cerebellar volume, whereas another small study (n = 22) found a negative association between fat mass and total brain volume.^{10,11} Both studies are limited by small sample sizes. Further, both assessed brain size only in 1 region or as total brain volume without considering different regions.

From the Departments of ¹Pediatric Newborn Medicine, and ²Nutrition, Brigham & Women's Hospital, Boston, MA; and the ³Division of Neonatology, School of Medicine, University of Minnesota, Minneapolis, MN

Supported by a Marshall Klaus Perinatal Research Award from the American Academy of Pediatrics (to K.B.); the Brigham Research Institute and Brigham and Women's Hospital Stork Fund (to M.B.); Program for Interdisciplinary Neuroscience Traveling Fellowship from the Brigham and Women's Hospital (to L.M.); and by Clinical Translational Science Award UL1RR025758 to Harvard University and Brigham and Women's Hospital from the National Center for Research Resources. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health. The authors declare no conflicts of interest.

Portions of this study were presented at the Pediatric Academic Societies annual meeting, May 4-8, 2018, Toronto, Canada, and at the American Academy of Pediatrics meeting, November 2-6, 2018, Orlando, Florida.

0022-3476/\$ - see front matter. © 2019 Elsevier Inc. All rights reserved.
<https://doi.org/10.1016/j.jpeds.2019.06.062>

BFD	Bifrontal diameter
BPD	Biparietal diameter
BMI	Body mass index
HC	Head circumference
MRI	Magnetic resonance imaging
NICU	Neonatal intensive care unit
SNAPPE-II	Score for Neonatal Acute Physiology-Perinatal Extension-II
TCD	Transverse cerebellar diameter

The primary aim of our study was to assess associations of physical growth and body composition with directly measured brain size among very preterm infants at term-corrected age. Our underlying hypothesis was that lean mass represents the accretion of nutrients, including protein, required for brain growth. For this study, we hypothesized that higher lean mass—but not fat—at term equivalent would be associated with larger brain size. We also hypothesized that greater weight, length and BMI gains from birth to term would be associated with larger brain size.

Methods

We conducted a prospective, longitudinal study of very preterm infants born at a single academic level III NICU from September 2015 to October 2018. Inclusion criteria were gestational age <33 weeks, singleton or twins, and no intention to transfer to another hospital after enrollment. Exclusion criteria were major congenital anomalies, triplets or higher order multiples, and an inability to provide informed consent in English. Of 122 infants enrolled, we excluded 27 who were unexpectedly transferred and 2 who died after enrollment; 18 infants missing MRI or body composition data; 7 infants whose parents withdrew consent; 4 infants diagnosed with a congenital anomaly after enrollment; and 2 infants who could not undergo body composition analysis owing to dependence on continuous respiratory support. Our final sample for this analysis was 62 (Figure 1; available at www.jpeds.com).

This study was approved by the Institutional Review Board of Brigham and Women's Hospital, and all parents of infants who participated in the study provided written consent.

Clinical Data

Clinical data were collected from the electronic health record, including demographics, common comorbidities of prematurity (supplemental oxygen use at 36 weeks postmenstrual age, necrotizing enterocolitis, retinopathy of prematurity, sepsis, patent ductus arteriosus, and intraventricular hemorrhage) defined using standardized definitions, and severity of illness using the Score for Neonatal Acute Physiology-Perinatal Extension-II (SNAPPE-II).^{12,13}

Anthropometrics

Infants were weighed daily by clinical nursing staff on a calibrated infant scale (Scale-Tronix, Inc, White Plains, New York) to the nearest gram. Two trained NICU dietitians performed all length and head circumference (HC) measurements after birth. Length was measured weekly and within 24 hours of the body composition measurement to the nearest 0.1 cm on an infant length board (Ellard Instrumentation Ltd, Monroe, Washington) using the 2-person method.¹⁴ HC was measured weekly to the nearest 0.1 centimeter using a nonstretchable tape. Weight, length, HC, and body mass index (BMI) Z-scores were calculated at birth and at term equivalent using the Olsen intrauterine growth reference.^{15,16}

Body Composition

At term-equivalent age, infants underwent body composition measurement using air displacement plethysmography in the Peapod Infant Body Composition System (COSMED, Concord, California). This device directly measures body mass and volume, then uses a 2-compartment model to determine fat and fat-free mass from infant-specific equations.¹⁷ The accuracy and precision of the device have been validated for preterm infants.¹⁷ We also calculated Z-scores for lean and fat mass at term using recently published reference data, the fat-free mass index using the formula fat-free mass/length² (kg/m²), and the fat mass index as fat mass/length² (kg/m²).¹⁸

Brain Metrics

At term-equivalent age, infants underwent brain MRI without sedation using a Siemens Trio 3 Tesla scanner (Erlangen, Germany).¹⁹ T₂-weighted images were acquired with a sagittal T₂ turbo spin echo sequence, with 1 × 1 × 1-mm isotropic voxels, flip angle of 160°, repetition time of 8630 ms, echo time of 133 ms, field of view of 190 × 190 mm, and matrix of 192 × 192. Total scan time was approximately 1 hour per participant. The images were manually AC-PC aligned using 3DSlicer to standardize their 3-dimensional orientation.²⁰ The aligned images were then used to derive 2-dimensional brain metrics in ITKSNAP, whereby a single observer manually delineated 3 brain metrics—bifrontal diameter (BFD), biparietal diameter (BPD), and transverse cerebellar diameter (TCD)—in the coronal plane (Figure 2; available at www.jpeds.com).^{7,21} In a subset of 29 infants, the reliability of these metrics was assessed by a second observer who was blinded to the body composition data. The interrater reliability of all 3 metrics had intraclass correlation coefficients of >0.9. Intra-rater reliability had intraclass correlation coefficients of 0.83 (BFD), 0.98 (BPD), and 0.93 (TCD). We chose these 3 metrics from among 8 previously described tissue metrics because these 3 have the highest correlation with brain tissue volumes, highest intra-rater and inter-rater reliability, and are all associated with later neurodevelopmental outcomes.^{7,9}

Statistical Analyses

The primary outcome was brain size as determined by brain metrics (BFD, BPD, TCD) at term-equivalent age. The primary exposures were (1) measures of body composition at term-equivalent age (absolute fat-free and fat mass, body fat percentage, fat-free mass index, fat mass index, fat-free mass Z-score and fat mass Z-score) and (2) Z-score change from birth to term-equivalent age of anthropometrics (weight, length, HC, and BMI). We quantified associations between exposure and outcome variables using generalized estimating equations to account for nonindependence of infants born to the same mother. The model included covariables that we identified a priori as determinants of infant growth, body composition, and/or brain size, including gestational age, birthweight Z-score (as a measure of fetal

growth), postmenstrual age at time of outcome measurement, and sex.^{7,22} Further adjusting for severity of illness using the SNAPPE-II score, supplemental oxygen use at 36 weeks, or duration of mechanical ventilation did not substantially change the results, and these variables were not included in the final models. A sensitivity analysis excluding infants with grade 3 or 4 intraventricular hemorrhage or periventricular leukomalacia did not substantially alter the results, so all infants were included in the final analysis.

We used IBM SPSS Statistics version 24.0 (IBM Corp, Armonk, New York) and SAS version 9.4 (SAS Inc, Cary, North Carolina) for all analyses.

Results

Participant characteristics are shown in **Table I**. The median gestational age was 29.1 weeks (range, 24.0–32.9 weeks), and the median birthweight Z-score was 0.05. By term-equivalent age (median postmenstrual age, 39.4 weeks), the median weight Z-score decreased to -0.6 . Length Z-score also declined (median, -0.9 units) from birth to term, whereas the median BMI Z-score change was -0.02 . At term, the average body fat percentage was $17.7\% \pm 5.3\%$ (**Table II**). Infants excluded from the analysis had a similar distribution of sex and race or ethnicity and similar median gestational age, birthweight, and birthweight Z-score, compared with infants who completed the study (data not shown).

Cross-sectional associations of body composition with brain size at term-equivalent age after adjustment for covariables are shown in **Table III**. Lean mass was positively associated with all three metrics. Specifically, each additional 100 g of lean mass was associated with 0.62 mm

Table I. Characteristics of 62 participants

Characteristics	No. (%) or median (range)
Male	35 (56)
Race	
African-American	12 (19)
Asian	3 (5)
Caucasian	36 (58)
Other	11 (18)
Hispanic ethnicity	12 (19)
Gestational age at birth (weeks)	29.1 (24.0 to 32.9)
Birth weight (g)	1088 (410 to 2065)
Birth weight Z-score*	0.05 (-3.1 to 2.3)
Small for gestational age (birthweight <10%)	11 (18)
Multiple gestation	18 (29)
Received antenatal steroids	61 (98)
SNAPPE-II score	8.5 (0 to 48)
Surfactant treatment	40 (65)
Any mechanical ventilation	40 (65)
Duration of mechanical ventilation (days)	1 (0 to 49)
Supplemental oxygen at 36 weeks	23 (37)
Postnatal steroids	7 (11)
Necrotizing enterocolitis Bell stage ≥ 2	3 (5)
Culture-proven sepsis	1 (2)
Patent ductus arteriosus treatment	11 (18)
Intraventricular hemorrhage grade 3 or 4	4 (6)

*From Olsen 2010 reference charts.¹⁵

Table II. Body composition and brain metrics at term-equivalent age (n = 62)

Characteristics	Median (range) or mean \pm SD
Postmenstrual age (weeks)	39.4 (35.9 to 42.1)
Anthropometrics	
Weight at term (g)	2982 \pm 503
Weight Z-score at term*	-0.6 (-3.2 to 0.7)
Δ weight Z-score (birth to term)*	-0.5 (-3.1 to 1.8)
Length at term (cm)	47.8 \pm 2.6
Length Z-score at term*	-0.8 (-4.3 to 0.7)
Δ length Z-score (birth to term)*	-0.9 (-2.0 to 1.4)
HC at term (cm) [†]	34.0 \pm 1.5
HC Z-score at term [†]	-0.1 (-2.0 to 1.9)
Δ HC Z-score (birth to term)*	0.4 (-1.6 to 3.1)
BMI Z-score at term*	0.2 (-1.8 to 1.5)
Δ BMI Z-score (birth to term)*	-0.02 (-2.8 to 2.8)
Body composition	
Fat-free mass (g)	2442 \pm 384
Fat-free mass index (kg/m ²)	10.7 \pm 1.0
Fat mass (g)	539 \pm 207
Fat mass index (kg/m ²)	2.3 \pm 0.8
Body fat percentage	17.7 \pm 5.3
Brain metrics	
BFD (mm)	69.4 \pm 4.7
BPD (mm)	80.4 \pm 5.3
TCD (mm)	51.0 \pm 2.8

*One infant was missing weight, length, and BMI Z-score at term-equivalent age owing to being measured at 42 weeks, beyond the range covered by the Olsen growth charts.¹⁵

[†]Two infants were missing HC data at term-equivalent age.

larger BFD (95% CI, 0.24-1.00), 0.70 mm larger BPD (95% CI, 0.27-1.13), and 0.26 mm larger TCD (95% CI, 0.003-0.51). To express these values in relation to average brain metric values, they represent 1% of average BFD and BPD, and 0.5% of average TCD. Similar to absolute lean mass, the lean mass Z-score and fat-free mass index showed generally positive associations with brain size, although associations with TCD did not meet statistical significance. Scatterplots of lean mass Z-score and brain metrics are shown in **Figure 3** (available at www.jpeds.com). Associations of fat mass with each brain metric were positive but not statistically significant. Whether measured by body fat percentage, body fat percent Z-score, fat mass Z-score, or fat mass index, adiposity was not statistically significantly associated with any measure of brain size (**Table III**).

In a post hoc analysis comparing infants born at <28 weeks of gestation (n = 23) with those born at ≥ 28 weeks of gestation (n = 39), we found no significant difference between groups in mean lean mass (2.4 ± 0.3 kg vs 2.4 ± 0.4 kg; $P = .98$) or lean mass Z-score (-1.5 ± 1.2 vs -1.1 ± 1.4 ; $P = .24$). We also explored effect modification of gestational age category on the association between lean mass and brain metrics by adding an interaction term between gestational age and the exposure variable in each analysis. None of the interaction terms were statistically significant ($P = .2$ -.9).

Longitudinal associations of physical growth from birth to term-equivalent age with brain size at term equivalent are shown in **Table IV**. Weight and BMI Z-score gain were positively associated with all measures of brain size. For each additional Z-score gain in weight, BFD was 2.1 mm

Table III. Cross-sectional associations of body size and composition with brain size at term-equivalent age (n = 62)

Variables	BFD (mm)		BPD (mm)		TCD (mm)	
	Estimate	P value	Estimate	P value	Estimate	P value
Fat-free mass (per 100 g)	0.62 (0.24 to 1.00)	.001	0.70 (0.27 to 1.13)	.001	0.26 (0.003 to 0.51)	.04
Fat-free mass index (kg/m ²)	1.41 (0.56 to 2.27)	.001	1.18 (0.13 to 2.23)	.03	0.34 (−0.20 to 0.88)	.22
Fat-free mass Z-score	1.39 (0.44 to 2.35)	.004	1.40 (0.27 to 2.54)	.02	0.53 (−0.10 to 1.16)	.10
Fat mass (per 100 g)	0.31 (−0.12 to 0.76)	.17	0.50 (−0.16 to 1.16)	.13	0.36 (−0.01 to 0.73)	.06
Fat mass index (kg/m ²)	0.62 (−0.32 to 1.56)	.20	0.96 (−0.61 to 2.54)	.23	0.81 (−0.06 to 1.69)	.07
Fat mass Z-score	0.46 (−0.18 to 1.11)	.16	0.62 (−0.38 to 1.62)	.22	0.39 (−0.11 to 0.89)	.13
Body fat percent	0.04 (−0.09 to 0.16)	.59	0.10 (−0.12 to 0.32)	.39	0.11 (−0.01 to 0.23)	.07
Body fat % Z-score	0.13 (−0.37 to 0.63)	.61	0.33 (−0.52 to 1.19)	.45	0.39 (−0.07 to 0.84)	.09
Weight Z-score	2.1 (0.60 to 3.67)	.006	2.60 (0.90 to 4.31)	.003	1.27 (0.32 to 2.22)	.009
Length Z-score	1.13 (−0.48 to 2.74)	.17	1.48 (−0.32 to 3.27)	.11	0.91 (−0.05 to 1.87)	.06
BMI Z-score	1.79 (0.69 to 2.89)	.001	1.95 (0.45 to 3.44)	.01	0.95 (0.11 to 1.79)	.03
HC Z-score	2.70 (1.74 to 3.67)	<.001	3.10 (2.04 to 4.16)	<.001	1.26 (0.48 to 2.04)	.001

Values are mean (95% CI) increase in brain size. Values in boldface are statistically significant with $P < .05$.

Estimates represent the difference in brain metric measurement associated with one unit difference in each growth or body composition parameter at term-equivalent age, adjusted using generalized estimating equations for gestational age at birth, sex, birthweight Z-score, and postmenstrual age at time of brain MRI, and accounting for nonindependence of infants born to the same mother.

(95% CI, 0.6–3.7) larger, BPD was 2.6 mm (95% CI, 0.9–4.3) larger, and TCD was 1.3 mm (95% CI, 0.3–2.2) larger. To express the associations with weight Z-score gain in relation to average brain metric values, they represent 3% of average BFD and BPD, and 2% of average TCD. Associations between BMI gain and brain metrics were smaller in magnitude but still strongly positive. In contrast, associations of linear growth (Z-score change in body length from birth to term-equivalent age) with brain metrics were not statistically significant.

To assess whether brain metrics provided additional information compared with using HC alone to assess brain size, we calculated Pearson correlation coefficients between HC Z-score at the time of the MRI and each brain metric. HC Z-score was moderately correlated with brain metrics: $r = 0.68$ with BFD, $r = 0.60$ with BPD, and $r = 0.50$ with TCD ($P < .01$ for all correlation coefficients).

Discussion

In a cohort of very preterm infants, we found that more rapid gains in weight and BMI from birth to term-equivalent age were associated with larger brain size at term. Although weight gain in the NICU is a well-established determinant of neurodevelopmental outcomes and predicts brain size at term, relatively little is known about how the composition of weight gain relates to early

brain growth and development.^{1–3,10,23} Using a noninvasive technique to accurately differentiate lean from fat mass, we evaluated relationships of weight composition with brain size at term-equivalent age. Our main finding was that lean mass—whether measured in absolute value, Z-score, or fat-free mass index—was positively associated with brain size, whereas fat mass was not.

Our finding of a positive association between lean mass and brain size complements the limited literature on body composition and neurodevelopment. In the few studies conducted to date, lean mass at term was linked to faster cognitive processing speed in infancy and to improved cognitive scores at 1 and 4 years of age.^{4–6} Taken together, these data suggest that lean mass may be a useful biomarker of brain growth, reflecting factors that facilitate the accretion of nutrients into many tissues, including the brain. One such factor is nutrient intake, particularly protein, which influences lean mass accrual and plays a crucial role in brain developmental processes including neurotransmitter production, cell migration and differentiation, myelination, synaptogenesis, and neuronal growth.^{24–29} Prior observational studies of preterm infants found that greater proportion of total energy intake composed of protein in the first few weeks of life was associated with greater lean mass at term equivalent.²⁶ Similarly, a randomized trial of additional protein and energy intake improved lean mass accretion during the neonatal hospitalization among very low birthweight infants.²⁷ In addition to

Table IV. Longitudinal associations of physical growth from birth to term with brain size at term equivalent (n = 62)

Z-score changes	BFD (mm)		BPD (mm)		TCD (mm)	
	Estimate	P value	Estimate	P value	Estimate	P value
Weight	2.1 (0.6 to 3.7)	.006	2.6 (0.9 to 4.3)	.003	1.3 (0.3 to 2.2)	.008
Length	1.1 (−0.6 to 2.8)	.20	1.7 (−0.01 to 3.5)	.05	0.3 (−0.6 to 1.3)	.50
BMI	1.4 (0.3 to 2.5)	.01	1.4 (0.03 to 2.8)	.04	1.0 (0.3 to 1.7)	.008
HC	1.3 (0.2 to 2.4)	.03	0.8 (−0.7 to 2.3)	.31	1.2 (0.5 to 1.9)	<.001

Values are mean (95% CI) increase in brain size. Values in boldface are statistically significant with $P < .05$.

Estimates represent the difference in brain metric associated with one Z-score difference in physical growth (change from birth to term-equivalent age), adjusted using generalized estimating equations for gestational age at birth, sex, birthweight Z-score, and postmenstrual age at time of brain MRI, and accounting for nonindependence of infants born to the same mother.

nutrient intake, other factors may influence nutrient accretion in the body and brain, including severity of illness, inflammation, postnatal steroid administration, comorbidities such as bronchopulmonary dysplasia, and the insulin-like growth factor axis.^{9,30-37}

In contrast with lean mass, we found little evidence for an association of adiposity—either fat mass or body fat percent—with brain size in any of the three brain regions we studied. In suggesting no benefit of greater adiposity, our results are consistent with a prior study ($n = 22$) in which greater deep subcutaneous fat mass was associated with decreased total brain size.¹¹ Our results are also consistent with prior studies of fat accrual and neurodevelopmental outcomes, which show no benefit of early fat mass gains or fat mass on neurodevelopmental testing from 4 months through 4 years of age.^{4-6,38} Our findings contrast, however, with 1 study ($n = 42$) in which greater fat mass at term equivalent was associated with larger cerebellar size.¹⁰ An important context for our findings is that preterm infants often develop relatively greater adiposity by term-equivalent age than term-born infants.³⁹ Furthermore, preterm infants have an increased risk of adverse cardiometabolic outcomes later in life.⁴⁰ Emerging evidence suggests early fat accrual may exacerbate later cardiometabolic risk in both full-term and preterm infants.⁴¹⁻⁴³ The optimal body composition for preterm infants remains unknown, yet our work adds to the growing body of literature suggesting that excess adiposity is not beneficial for brain growth or development.

Two prior MRI-based studies in preterm infants found that faster weight gain velocity prior to term was associated with larger total brain volume and cerebellum, but neither of those studies addressed the proportionality of weight gain.^{10,23} We noted that gain in BMI—representing weight gain out of proportion to length—predicted larger brain size at term. Unlike older children and adults, BMI in infants is a poor surrogate for fat mass in both preterm infants at birth and full term infants throughout infancy, although the relationship between BMI gain and adiposity gain in preterm infants is unknown.^{44,45} Nevertheless, in settings where body composition analysis is not available, BMI gain may complement weight gain in providing information about nutritional status in relation to brain growth. Routine BMI assessment can be facilitated by published reference charts for preterm infants.¹⁶

Some authors have suggested that gain in body length may be a better index of lean body mass accretion and organ growth than weight gain.^{28,31} In this context, we expected to find an association between linear growth and brain size, but our data showed no association. Similarly, a prior study in the preterm population also found no association between linear growth from birth to term and volume of the total brain, cerebellum, or cortical gray matter.²³ It is notable that the concept of body length as an indicator of lean mass is largely extrapolated from work in the adult population, where height was found to be a better proxy for lean body mass than weight, whereas, studies of preterm infants have shown the opposite, that weight is a better proxy for

lean body mass than length.^{44,46,47} Prior studies linking linear growth from birth to term with neurodevelopmental outcomes have yielded inconsistent results in preterm infants.^{3,30,48} An important caveat is that length measurements are often subject to measurement error, which could obscure true associations between linear growth and outcome; in our study, we minimized measurement error by using an infant length board and having only 2 trained observers perform all measurements.⁴⁹

We used MRI to directly measure brain size at term-equivalent age. Compared with HC, MRI is preferred for measuring brain size because MRI distinguishes brain tissue from extra-axial cerebrospinal fluid—which is increased in preterm infants—and is not influenced by head shape.^{50,51} Brain growth in the NICU is important because early abnormalities in brain size persist throughout childhood and adolescence, and are associated with long-term outcomes.^{34,52,53} In prior studies, the brain metrics we measured at term have been shown to predict later neurodevelopmental outcomes independent of other known risk factors (such as sex, gestational age, severity of illness, maternal education, and overt brain injury).^{54,55} One of those studies compared a contemporary cohort of infants with a historical cohort and found that the contemporary cohort had 4%-10% larger brain metrics and twice as many infants with normal neurodevelopmental outcome at 2 years of age (68% vs 33%).⁵⁵ However, further research is needed to determine the extent of the impact on neurodevelopment that would be expected from the 1%-3% increases in brain size described in our study.

Strengths of our study include direct measurement of brain size using quantitative analysis of high-quality magnetic resonance images, although we acknowledge that a more sophisticated volumetric analysis could further clarify precise regions of the brain where growth is associated with greater lean mass accrual. We also measured body composition directly using a highly accurate device specifically validated for preterm infants.

Our study has several limitations. A large proportion of enrolled infants did not complete the study, primarily owing to early transfer to lower level units for convalescent care, although we did not find any difference in demographic characteristics among those infants and the remaining infants to suggest that this introduced bias in our results. We did not have data on long-term neurodevelopmental outcomes, although prior studies have shown that brain metrics at term are associated with later outcomes. Finally, this is a single-center study and our findings may not be generalizable to other units with different populations or care practices, although our findings regarding weight gain and brain size are consistent with the few prior studies on this topic.

We found that the composition of weight gain from birth to term was associated with brain growth; lean mass was associated with larger brain size whereas fat mass was not. Although the optimal composition of weight gain for preterm infants remains unknown, our work adds to emerging evidence suggesting the importance of early lean mass accrual to optimize long-term neurodevelopment.⁴⁻⁶ The beneficial

effects of lean mass—in contrast with fat mass—are particularly relevant given the striking deficit of lean mass and excess adiposity that preterm infants develop during the NICU hospitalization.³⁹ It is not yet known whether interventions that improve lean mass accrual could also improve brain growth and neurodevelopment. Nevertheless, body composition is an attractive biomarker of brain growth in preterm infants, particularly because recent advances in technology allow rapid noninvasive testing of body composition in the NICU. Attention to the quality and composition of preterm infants' weight gain, rather than quantity alone, is likely important for optimizing brain growth and neurodevelopment as well as mitigating risks to later cardiometabolic health. ■

We thank the infants and their parents who participated in this study. We also thank Danielle Dwyer, RN, and Tina Steele, RN, IBCLC, for their assistance with data collection and Zain Al Tirmizi for his assistance with inter-rater reliability testing of brain metric analysis.

Submitted for publication Feb 22, 2019; last revision received May 8, 2019; accepted Jun 25, 2019.

Reprint requests: Katherine A. Bell, MD, 75 Francis St, Boston, MA 02115.
E-mail: KBell7@partners.org

References

- Ehrenkranz RA, Dusick AM, Vohr BR, Wright LL, Wrage LA, Poole WK. Growth in the neonatal intensive care unit influences neurodevelopmental and growth outcomes of extremely low birth weight infants. *Pediatrics* 2006;117:1253-61.
- Franz AR, Pohlandt F, Bode H, Mihatsch WA, Sander S, Kron M, et al. Intrauterine, early neonatal, and postdischarge growth and neurodevelopmental outcome at 5.4 years in extremely preterm infants after intensive neonatal nutritional support. *Pediatrics* 2009;123:e101-9.
- Belfort MB, Rifas-Shiman SL, Sullivan T, Collins CT, McPhee AJ, Ryan P, et al. Infant growth before and after term: effects on neurodevelopment in preterm infants. *Pediatrics* 2011;128:e899-906.
- Pfister KM, Gray HL, Miller NC, Demerath EW, Georgieff MK, Ramel SE. Exploratory study of the relationship of fat-free mass to speed of brain processing in preterm infants. *Pediatr Res* 2013;74:576-83.
- Ramel SE, Gray HL, Christiansen E, Boys C, Georgieff MK, Demerath EW. Greater early gains in fat-free mass, but not fat mass, are associated with improved neurodevelopment at 1 year corrected age for prematurity in very low birth weight preterm infants. *J Pediatr* 2016;173:108-15.
- Scheurer JM, Zhang L, Plummer EA, Hultgren SA, Demerath EW, Ramel SE. Body composition changes from infancy to 4 years and associations with early childhood cognition in preterm and full-term children. *Neonatology* 2018;114:169-76.
- Nguyen The Tich S, Anderson PJ, Shimony JS, Hunt RW, Doyle LW, Inder TE. A novel quantitative simple brain metric using MR imaging for preterm infants. *AJNR Am J Neuroradiol* 2009;30:125-31.
- Peterson BS, Anderson AW, Ehrenkranz R, Staib LH, Tageldin M, Colson E, et al. Regional brain volumes and their later neurodevelopmental correlates in term and preterm infants. *Pediatrics* 2003;111:939-48.
- Tich SNT, Anderson PJ, Hunt RW, Lee KJ, Doyle LW, Inder TE. Neurodevelopmental and perinatal correlates of simple brain metrics in very preterm infants. *Arch Pediatr Adolesc Med* 2011;165:216-22.
- Paviotti G, De Cunto A, Zennaro F, Boz G, Travan L, Cont G, et al. Higher growth, fat and fat-free masses correlate with larger cerebellar volumes in preterm infants at term. *Acta Paediatr* 2017;106:918-25.
- Vasu V, Durighel G, Thomas EL, Thomas L, Malamateniou C, Bell JD, et al. Preterm nutritional intake and MRI phenotype at term age: a prospective observational study. *BMJ Open* 2014;4:e005390.
- Vermont Oxford Network. Manual of operations: Part 2 data definitions and infant data forms. Burlington (VT): Vermont Oxford Network; 2017.
- Richardson DK, Corcoran JD, Escobar GJ, Lee SK. SNAP-II and SNAPPE-II: simplified newborn illness severity and mortality risk scores. *J Pediatr* 2001;138:92-100.
- Wood AJ, Raynes-Greenow CH, Carberry AE, Jeffery HE. Neonatal length inaccuracies in clinical practice and related percentile discrepancies detected by a simple length-board. *J Paediatr Child Health* 2013;49:199-203.
- Olsen IE, Groveman SA, Lawson ML, Clark RH, Zemel BS. New intrauterine growth curves based on United States data. *Pediatrics* 2010;125:e214-24.
- Olsen IE, Lawson ML, Ferguson AN, Cantrell R, Grabich SC, Zemel BS, et al. BMI curves for preterm infants. *Pediatrics* 2015;135:e572-81.
- Urlando A, Dempster P, Aitkens S. A new air displacement plethysmograph for the measurement of body composition in infants. *Pediatr Res* 2003;53:486-92.
- Norris T, Ramel S, Catalano PM, Ni Caoimh C, Roggero P, Murray D, et al. New charts for the assessment of body composition, according to air-displacement plethysmography, at birth and across the first six months of life. *Am J Clin Nutr* 2019;109:1353-60.
- Mathur AM, Neil JJ, McKinstry RC, Inder TE. Transport, monitoring, and successful brain MR imaging in unsedated neonates. *Pediatr Radiol* 2008;38:260-4.
- Fedorov A, Beichel R, Kalpathy-Cramer J, Finet J, Fillion-Robin J-C, Pujol S, et al. 3D Slicer as an image computing platform for the quantitative imaging network. *Magn Reson Imaging* 2012;30:1323-41.
- Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, et al. User-guided 3D active contour segmentation of anatomical structures: significantly improved efficiency and reliability. *NeuroImage* 2006;31:1116-28.
- Iwata S, Katayama R, Kinoshita M, Saikusa M, Araki Y, Takashima S, et al. Region-specific growth restriction of brain following preterm birth. *Sci Rep* 2016;6:33995.
- Coviello C, Keunen K, Kersbergen KJ, Groenendaal F, Leemans A, Peels B, et al. Effects of early nutrition and growth on brain volumes, white matter microstructure, and neurodevelopmental outcome in preterm newborns. *Pediatr Res* 2018;83:102-10.
- Stephens BE, Walden RV, Gargus RA, Tucker R, McKinley L, Mance M, et al. First-week protein and energy intakes are associated with 18-month developmental outcomes in extremely low birth weight infants. *Pediatrics* 2009;123:1337-43.
- Isaacs EB, Gadian DG, Sabatini S, Chong WK, Quinn BT, Fischl BR, et al. The effect of early human diet on caudate volumes and IQ. *Pediatr Res* 2008;63:308-14.
- Simon L, Frondas-Chauty A, Senterre T, Flamant C, Darmaun D, Rozé J-C. Determinants of body composition in preterm infants at the time of hospital discharge. *Am J Clin Nutr* 2014;100:98-104.
- Costa-Orvay JA, Figueras-Aloy J, Romera G, Closa-Monasterolo R, Carbonell-Estrany X. The effects of varying protein and energy intakes on the growth and body composition of very low birth weight infants. *Nutr J* 2011;10:140.
- Ramel SE, Georgieff MK. Preterm nutrition and the brain. *World Rev Nutr Diet* 2014;110:190-200.
- Morgane PJ, Austin-LaFrance R, Bronzino J, Tonkiss J, Díaz-Cintra S, Cintra L, et al. Prenatal malnutrition and development of the brain. *Neurosci Biobehav Rev* 1993;17:91-128.
- Ramel SE, Demerath EW, Gray HL, Younge N, Boys C, Georgieff MK. The relationship of poor linear growth velocity with neonatal illness and two-year neurodevelopment in preterm infants. *Neonatology* 2012;102:19-24.
- Pfister KM, Ramel SE. Linear growth and neurodevelopmental outcomes. *Clin Perinatol* 2014;41:309-21.
- Kuban KCK, O'Shea TM, Allred EN, Fichorova RN, Heeren T, Paneth N, et al. The breadth and type of systemic inflammation and the risk of

- adverse neurological outcomes in extremely low gestation newborns. *Pediatr Neurol* 2015;52:42-8.
33. Rose J, Vassar R, Cahill-Rowley K, Hintz SR, Stevenson DK. Neonatal biomarkers of inflammation: correlates of early neurodevelopment and gait in very-low-birth-weight preterm children. *Am J Perinatol* 2016;33:71-8.
 34. Matthews LG, Inder TE, Pascoe L, Kapur K, Lee KJ, Monson BB, et al. Longitudinal preterm cerebellar volume: perinatal and neurodevelopmental outcome associations. *Cerebellum* 2018;17:610-27.
 35. Ohlsson A, Calvert SA, Hosking M, Shennan AT. Randomized controlled trial of dexamethasone treatment in very-low-birth-weight infants with ventilator-dependent chronic lung disease. *Acta Paediatr Oslo Nor* 1992;81:751-6.
 36. van de Lagemaat M, Rotteveel J, Heijboer AC, Lafeber HN, van Weissenbruch MM. Growth in preterm infants until six months post-term: the role of insulin and IGF-I. *Horm Res Paediatr* 2013;80:92-9.
 37. Hansen-Pupp I, Hövel H, Löfqvist C, Hellström-Westas L, Fellman V, Hüppi PS, et al. Circulatory insulin-like growth factor-I and brain volumes in relation to neurodevelopmental outcome in very preterm infants. *Pediatr Res* 2013;74:564-9.
 38. Frondas-Chauty A, Simon L, Flamant C, Hanf M, Darmaun D, Rozé J-C. Deficit of fat free mass in very preterm infants at discharge is associated with neurological impairment at age 2 years. *J Pediatr* 2018;196:301-4.
 39. Johnson MJ, Wootton SA, Leaf AA, Jackson AA. Preterm birth and body composition at term equivalent age: a systematic review and meta-analysis. *Pediatrics* 2012;130:e640-9.
 40. Sipola-Leppänen M, Väärasmäki M, Tikanmäki M, Matinelli H-M, Miettola S, Hovi P, et al. Cardiometabolic risk factors in young adults who were born preterm. *Am J Epidemiol* 2015;181:861-73.
 41. Koontz MB, Gunzler DD, Presley L, Catalano PM. Longitudinal changes in infant body composition: association with childhood obesity. *Pediatr Obes* 2014;9:e141-4.
 42. Hawkes CP, Zemel BS, Kiely M, Irvine AD, Kenny LC, Hourihane JO, et al. Body composition within the first 3 months: optimized correction for length and correlation with BMI at 2 years. *Horm Res Paediatr* 2016;86:178-87.
 43. Pfister KM, Zhang L, Miller NC, Ingolfssland EC, Demerath EW, Ramel SE. Early body composition changes are associated with neurodevelopmental and metabolic outcomes at 4 years of age in very preterm infants. *Pediatr Res* 2018;84:713.
 44. Ramel SE, Zhang L, Misra S, Anderson CG, Demerath EW. Do anthropometric measures accurately reflect body composition in preterm infants? *Pediatr Obes* 2017;12(Suppl 1):72-7.
 45. Bell KA, Wagner CL, Perng W, Feldman HA, Shypailo RJ, Belfort MB. Validity of body mass index as a measure of adiposity in infancy. *J Pediatr* 2018;196:168-74.e1.
 46. Skullerud K. Variations in the size of the human brain. Influence of age, sex, body length, body mass index, alcoholism, Alzheimer changes, and cerebral atherosclerosis. *Acta Neurol Scand Suppl* 1985;102:1-94.
 47. Forbes GB. Relation of lean body mass to height in children and adolescents. *Pediatr Res* 1972;6:32-7.
 48. Belfort MB, Gillman MW. Healthy infant growth: what are the trade-offs in the developed world? *Nestle Nutr Inst Workshop Ser* 2013;71:171-84.
 49. Johnson TS, Engstrom JL, Gelhar DK. Intra- and interexaminer reliability of anthropometric measurements of term infants. *J Pediatr Gastroenterol Nutr* 1997;24:497-505.
 50. Keunen K, van Elburg RM, van Bel F, Benders MJNL. Impact of nutrition on brain development and its neuroprotective implications following preterm birth. *Pediatr Res* 2015;77:148-55.
 51. Thompson DK, Warfield SK, Carlin JB, Pavlovic M, Wang HX, Bear M, et al. Perinatal risk factors altering regional brain structure in the preterm infant. *Brain J Neurol* 2007;130:667-77.
 52. de Kieviet JF, Zoetebier L, van Elburg RM, Vermeulen RJ, Oosterlaan J. Brain development of very preterm and very low-birthweight children in childhood and adolescence: a meta-analysis. *Dev Med Child Neurol* 2012;54:313-23.
 53. Cheong JLY, Anderson PJ, Roberts G, Burnett AC, Lee KJ, Thompson DK, et al. Contribution of brain size to IQ and educational underperformance in extremely preterm adolescents. *PLoS One* 2013;8:e77475.
 54. Kidokoro H, Anderson PJ, Doyle LW, Woodward LJ, Neil JJ, Inder TE. Brain injury and altered brain growth in preterm infants: predictors and prognosis. *Pediatrics* 2014;134:e444-53.
 55. Harris SL, Austin NC, Battin M, Broadbent R, Keenan R, Wells S, et al. In-hospital morbidity and brain metrics of preterm neonates born 1998-2009. *N Z Med J* 2016;129:94-107.




Figure 1. Flow diagram of study participants.




Figure 2. A, Measurements of BFD, B, BPD, and C, TCD.




Figure 3. Unadjusted scatterplots of lean mass Z-score at term-equivalent age and brain size measured by 3 brain metrics: **A**, BFD, **B**, BPD, and **C**, TCD.