



Review Article

B cell dysregulation in primary Sjögren's syndrome: A review

Hazim Mahmoud Ibrahim*

Department of Basic Science, Al-Mustansiriyah University, College of Dentistry, Baghdad, Iraq



ARTICLE INFO

Article history:

Received 11 January 2019

Received in revised form 2 July 2019

Accepted 19 September 2019

Keywords:

B cells

Sjögren's syndrome

Peripheral blood

Salivary gland

Germinal center

CD27

ABSTRACT

Primary Sjögren's syndrome is a chronic autoimmune disorder of unknown etiology and is characterized by progressive focal lymphocytic infiltration of the lacrimal and salivary glands. Comparison of B cell subsets from the peripheral blood and salivary glands of patients with primary Sjögren's syndrome and those from healthy individuals shows dysregulation and derangement of B cell subsets in both peripheral circulation and in inflamed glandular tissues. This dysregulation is expressed as a decrease in the percentage of CD27+ memory B cells in peripheral blood and an increase in the CD27+ memory B cells in the affected glands. Further, the overall percentage of long-lived autoantibodies-producing plasma cells within the affected glands is increased. In the last two decades, several studies have shown growing evidences that B cells play multiple roles in primary Sjögren's syndrome pathophysiology, and that dysregulation of these cells may actually play a central role in the disease development.

© 2019 The Author. Published by Elsevier Ltd on behalf of The Japanese Association for Dental Science. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Primary Sjögren's syndrome (pSS) is an autoimmune epithelitis characterized by dry mouth and dry eyes due to the disease-related destruction of the affected salivary and lacrimal glands [1]. Although the pathogenesis of pSS remains unclear, the disease has traditionally been ascribed to T cells [2]. Recent evidences indicate a major contribution of B cells in pSS pathogenesis [3–5]. Patients with pSS demonstrate a decrease in the absolute numbers of circulating CD27+ memory B cells and IgM producing B cell subpopulations accompanied by an increase in circulating naïve CD27– B cells [6]. Furthermore, analysis of B cells in the inflamed salivary gland obtained from a patient with pSS, indicated a striking accumulation of both heavily mutated V_H genes in CD27+ memory B cells and IgM producing plasma cells [7].

2. Primary Sjögren's syndrome

Primary Sjögren's syndrome is a chronic inflammatory autoimmune disease characterized by dry mouth, dry eyes, and sialoadenitis (sialadenitis) with focal periductal lymphocytic infiltration of the lacrimal and salivary glands [8]. The pathogenesis of pSS can virtually be organized in a series of stages. In the **first stage**, environmental factors such as viral infections induce injury

to glandular epithelial cells, thus activating the innate immune system with the release of inflammatory cytokines, chemokines, and autoantigens [9–11]. The release of inflammatory cytokines, chemokines, and autoantigens accompanied by activation of glandular endothelial cells and recruitment of inflammatory cells including macrophages, dendritic cells, and B and T lymphocytes cause an increase in the number of CD27+ memory B cells in the salivary gland [12–14]. In the **second stage**, B cells and T cells are stimulated with the induction of autoantigen-specific autoantibodies (such as anti-SS-A/Ro, anti-SS-B/La, anti-muscarinic receptor, and anti-fodrin receptor antibodies, as well as rheumatoid factor (RF)). These autoantigen-specific autoantibodies react with the corresponding autoantigen resulting in the formation of autoantigen-autoantibody immune complexes that stimulate further activation of inflammatory cells through complement and Fc receptors (FcR), culminating in the production of interferon- α by infiltrating dendritic cells [15,16]. During the **third stage**, further B cell activation and survival occurs, caused mainly by B cell activating factor (BAFF) that is produced by many cell types including B cells, monocytes/macrophages, dendritic cells, neutrophils, epithelial cells and activated T- cells [17]. Moreover, other factors such as IL-2, IFN- γ , IL-10, IL-6, TGF β , IL-4 and IL-5 are released by infiltrating T cells, macrophages and possibly by damaged resident glandular epithelial and mesenchymal cells [18]. During this stage there is a possibility of rearrangement and organization of B-cells within the affected gland resulting in the development of ectopic germinal centers (GCs). These newly formed GCs with a follicular dendritic cell network are found in a subset of pSS patients [19].

* Correspondence to: Current address: 1320 W Somerville Ave, Apt # 703, Philadelphia, PA 19141, United States.

E-mail address: hazimahmoud56@yahoo.com

In pSS, salivary gland hypofunction may occur from the glandular damage caused by the disease-related destruction of glandular tissue and excessive infiltration of inflammatory cells into the gland, or as a result of anti-muscarinic receptor antibodies blocking the parasympathetic stimulation of epithelial cells resulting in reduced saliva production [20,21].

3. B cell biology, development and maturation

In humans, B cells are generated throughout life in the bone marrow [22].

B cells pass through three sequential programmed stages:

First stage: In the bone marrow, B-cell maturation starts from a lymphoid stem cell that differentiates into a progenitor B cell, to a precursor B cell, then to an immature B cell. During this stage B cells randomly rearrange their Ig genes to generate Ag-specific B-cell receptors, which are capable of recognizing a wide variety of antigens [23,24]. **Second stage:** Immature naïve B cells exit the bone marrow and enter the blood stream to complete their maturation in secondary lymphoid tissues, preferentially in the spleen where naïve B cells are generally differentiated into marginal zone (MZ) B cells and follicular B cells [23]. **Third stage:** Follicular B cells proliferate in the germinal center (GC) of lymphoid follicles and differentiate into GC B cells that express high affinity BCR and class-switch isotypes. B cells that leave the GC can develop into memory B cells or plasma cells [23].

Mature B cells recognize various self-antigens and do not react with these self-antigens for the self-reactive or autoreactive B cells must be eliminated and this elimination is part of the immune tolerance process to avoid autoimmunity [25]. B cell tolerance is a mechanism that is important in preventing the development of antibody responses to self-protein antigens. Both central and peripheral mechanisms are implicated in B cell tolerance, and loss of tolerance leads to autoimmune diseases including pSS [25,26].

4. B cells in the immunopathogenesis of primary Sjögren's syndrome

Immunopathogenesis of pSS is a very complex process associated with the innate immune system and adaptive immune system [4]. In patients with pSS, using the indicating B-cell surface markers, CD19 and CD27, several studies have indicated the possible existence of multiple disturbances in B-cell differentiation and maturation; these disturbances include altered B-cell trafficking between circulation and inflamed glands and disturbance in B-cell differentiation with a greater presence of circulating auto-activated plasma B cells [27–30]. B-cell hyperactivity and peripheral B-cell derangement appears to be a hallmark of the disease and might play an important role in the autoimmune/lymphoproliferative processes involved in pSS pathogenesis [27–30]. Furthermore, polarized B lymphocytes possess the capacity to produce a wide range of cytokines such as IL-1, IL-6, IL-7, and TNF- α [31].

5. Disturbances of B-cell subsets in primary Sjögren's syndrome

In patients with pSS, analysis of B-cells shows a major disturbance in the homeostasis of B-cell subsets, mostly the CD27 subset, both in the peripheral blood and in the inflamed salivary glands [7,32–36].

5.1. Peripheral B-cell subsets disturbances in primary Sjögren's syndrome

Immunophenotyping studies indicate that B cell homeostasis is disturbed in patients with pSS and these disturbances present as diminished frequencies and absolute numbers of peripheral CD27+

Table 1

Peripheral B-cell subsets disturbances in primary Sjögren's syndrome.

Disturbances of Peripheral B- cell subsets	References
- Reduced number and frequency of CD27+ memory B cells.	[3,7,22,24,26,30–39]
- Increased level of CD27- naïve B cells.	[3,7,22,24,26,30–39]

Table 2

B cells disturbances within the salivary gland infiltrates in primary Sjögren's Syndrome.

Disturbances of B cells within the exocrine gland infiltrates	References
- Presence of germinal center-like structures.	[12,14,48,66–71]
- High frequency of IgG plasma cells.	[54–61]
- Autoreactive B cells and plasma cells.	[60–63]
- Elevated levels of B cell-associated cytokines and chemokines (e.g., IL-6, IL- 21, BAFF, APRIL, CXCL12, CXCL13).	[12,23,46–48]
- Presence of clonal B cell populations and autoantibody production plasma cells.	[24–26]

memory B cells, in particular reduction in the circulating CD27+ IgM+ subpopulation [8,32–34].

Moreover, pSS patients demonstrate an increase in the number of naïve un-switched peripheral memory B cells (CD19+, CD27–, IgD+) with a decrease in the number of the peripheral memory B cells (CD19+, CD27+, IgD–), possibly because of their retention in inflamed salivary gland tissues [34–37].

Furthermore, analysis of the immunoglobulin repertoire in pSS patients revealed diminished mutational frequencies of the Ig gene rearrangements in individual peripheral CD19+B cells [34,38,39], whereas the vast majority of B cells in the salivary gland expressed heavily mutated IgV_H rearrangements [34].

Recently, an increased frequency of transitional B cells and mature naïve B cells expressing polyreactive antibodies has been demonstrated in the peripheral blood of patients with pSS [40].

Many investigators have focused on the main disturbances of peripheral B cell subsets in pSS [6,12,29,32,34,36,37,41–48], and the results of these studies are summarized in Table 1.

5.2. B cells within salivary gland infiltrates in primary Sjögren's syndrome

The microenvironment of inflamed glandular tissue is rich in chemokines and cytokines that promote B-cell recruitment, homing, survival, activation and plasma cell formation. In patients with pSS, analysis of inflamed glandular tissue revealed polyclonal accumulation of CD27+ memory B cells and CD27^{high} plasma cells [32,34]. The B-cell disturbances within the salivary gland infiltrates in primary Sjögren's syndrome are summarized in Table 2.

5.2.1. Pathogenesis for accumulation of CD27+ memory B cells and plasma cells within the salivary gland infiltrate

The inflamed glands observed in pSS have been shown to express a unique profile of adhesion molecules, cytokines, and chemokines such as B-cell-chemoattracting chemokines CXCL13 and CXCL12 [19,49,50]. CXCL13 is a homeostatic chemokine secreted by follicular dendritic cells, T-helper follicular cells, and antigen-experienced T cells, and interacts with its corresponding receptor CXCR5 on B-cells, to regulate B-cell movement between different tissues [51–54]. CD27+ memory B cells express the corresponding receptors CXCR5 for CXCL13 and CXCR4 for CXCL12 [19,43,49,55]; thus, glandular co-expression of both CXCL12 and CXCL13 seems to attract this subpopulation of peripheral CD27+ memory B cells into the inflamed glands, where they then reside [19,49]. CXCL13 over-expression in inflamed glands of patients




Fig. 1. B cell dysregulation in primary Sjögren's Syndrome.

→ indicates, Take part in.

with pSS plays an active role in the extreme recruitment of circulatory CD27+memory B cells that express the cognate receptor CXCR5 causing their retention in the inflamed glandular tissue, culminating in the formation of ectopic lymphoid tissue composed primarily of B cells [19,49]. Also, CXCR3 expressed on activated B cells/ memory cells will cause the attraction of these cells to the salivary gland due to CXCR3 ligands that are produced locally in the target tissue of pSS [5].

5.2.2. Plasma cells within salivary gland infiltrate

B-cell activating factor (BAFF), a member of the TNF family secreted by inflammatory cells, is necessary for prolonged plasma cell survival and exaggerated immunoglobulin production by these cells [56]. In Sjögren's syndrome, the chronically inflamed glandular tissue microenvironment is rich in BAFF, which causes accumulation of CD27+memory B-cells and increases the numbers of immunoglobulin producing plasma cells in labial salivary glands [57–59]. Analysis of immunoglobulin plasma cells phenotype showed the presence of both IgA and IgG expressing plasma cells in the salivary gland tissue of pSS patients. IgA expressing plasma cells aggregate in small cell clusters in close proximity to ductal and acinar epithelium, IgG expressing plasma cells were found both inside and in the periphery of large focal infiltrates in pSS patient. pSS patients with chronically inflamed salivary glands had significantly higher numbers of interstitial IgG plasma cells compared with healthy tissue [60]. Plasma cells within the salivary gland of patients with pSS comprise autoreactive plasma cells that produce anti-SSA/Ro or anti-SSB/La autoantibodies. These autoreactive plasma cells within the affected gland might be derived either from the differentiation of CD27+memory cells recruited

from peripheral blood or from the differentiation of CD27+ memory B-cells generated in the newly formed ectopic GCs within the affected salivary glands [60–64]. The relative contribution from both sources to memory B-cell formation and subsequent differentiation toward plasma cells is still unknown [5]. Also the heightened expression of CXCR3 and CXCR4 on plasma cells and plasmablasts of patients with pSS, will lead to their migration towards the site of inflammation [46].

5.2.3. Germinal center formation in exocrine glands in Sjögren's syndrome patients

Lymphocytic infiltration of the salivary gland in pSS patients is believed to be a result of chronic inflammation [65]. This infiltrate is a progressive feature which when extensive, may replace large portions of the affected gland [66]. The glandular infiltrates in some patients are organized into tertiary lymphoid tissues with germinal centers resembling the GCs of secondary lymphoid tissues with a central dark and light zone [19,22,49,51,65]. These GCs are found in 18%–59% of pSS patients [67]. In pSS, B cells in the ectopic GCs undergo somatic hypermutation suggesting that these cells react in a manner different from that found in the normal secondary lymphoid organs [68].

5.2.4. Consequences of GC formation within salivary glands

First, extensive lymphocytic organization into GC may lead to increased disruption of the glandular structure, causing increased functional disability [69]. **Second**, the autoantibody production could increase the formation of more immune complexes, contributing to the occurrence of certain extra-glandular manifestations [70]. **Third**, in pSS patients with GCs, special attention is

required for the development of lymphoma because the presence of ectopic GC-like structures in the affected glandular tissue could be associated with an increased risk of lymphoma (Non-Hodgkin and MALT lymphoma) [71,72]. **Fourth**, increased BAFF production may result in continuous B cell activation, leading to B cell lymphoma development [73].

B cell dysregulation in primary Sjögren's Syndrome is summarized in Fig. 1.

6. Possible explanations for dysregulation of B cells in primary Sjögren's syndrome

The mechanism leading to B-cell dysregulation and salivary gland hypofunction in pSS is still unclear, but many explanations are suggested according to the following causes:

First, migration of CD27+ memory B-cells, from peripheral circulation into inflamed salivary glands, results from increased expression of B-cell chemo-attractants CXCL12 and CXCL13 (major B cell attractants mainly produced by follicular dendritic cells) in the inflamed glands [9,30,53]. **Second**, the lower ratio of circulating memory B cells may be explained by the over-expression of chemokine molecules CXCR3 and CXCR4 which guide them into the inflamed tissues [46]. **Third**, shedding of CD27 from the surface of CD27+ memory cells results in an increased percentage of CD27- B-cells [46,68]. Peripheral CD27+ memory B cells tend to differentiate toward plasma cells by ligation of CD27 via CD70 [19]. Consequently, in patients with pSS, elevation of serum CD27 and hypergammaglobulinemia could result from this increased plasma cell differentiation [6]. **Fourth**, alternative B-cell activation pathways via toll-like receptors (TLR) may be used. Studies on pSS patients in the last decade showed that TLR-7 and -9 are expressed at high levels in CD27+ memory B cells, whereas only low to undetectable levels were found in CD27- naïve B cells [74]. When naïve B cells encounter and bind antigens to their B cell TLR, they are activated to develop into plasma or memory B cells, suggesting that the increased level of TLRs detected on memory B cells could be a result of the activation of naïve B cells and that TLRs may be upregulated during this activation [36,75–78]. **Fifth**, the skewing of CD27+ memory B cells towards plasma cells differentiation results in fewer circulatory CD27+ memory B cells. Interaction of CD27 with its ligand CD70 is essential for B-cell differentiation into plasma cell [79]. CD70 is expressed on activated T and B cells [80], and hyperactivation of T and B cells in pSS could result in an elevated differentiation of CD27+ memory B cells into antibody producing plasma cells [33]. **Sixth**, elevated IL-10 production has been reported in pSS patients [81,82]. Interleukin-10 (IL-10) can promote the differentiation of CD27+ memory B cells towards plasma cells [83]. Moreover, IL-10 has been found to down modulate the expression of CD27 on the surface of B cells [84]. Triggering CD27 by CD70 on peripheral blood B cells was found to yield an increase in the number of plasma cells in the presence of IL-10 [79]. **Seventh**, reduced percentage of CD27+ memory B cells may also be caused by a loss of CD27 expression on B cells in pSS [6,33]. Analysis of CD27 expression on peripheral blood B-cell subpopulations revealed that the reduced percentage of CD27+ B cells in pSS is most pronounced in the CD38-IgD+ and CD38-IgD- B-cell subpopulations. These data suggest that some CD27+ B cells die in the early memory B-cell stages and never become memory B cells [6,33,43].

7. Therapies targeting B-cell dysregulation in primary Sjögren's syndrome

The most recent etiopathogenic advances in patients with pSS help evolve new, highly-selective biological therapies targeting molecules and receptors involved in the etiopathogenesis of pSS.

In relation to B cell disturbances, these therapeutics could be classified into two categories [85]:

- 1 Direct B-Cell Blocking: Monoclonal antibodies against B-cell receptors can induce or block B cells from entering the cell cycle; Rituximab blocks the CD20 lymphocytic receptor, leading to complete B-cell depletion in blood [86]; Epratuzumab targets CD22 B-cell antigen receptor signaling, leading to partial B-cell depletion [87]. A study has demonstrated prevention of autoimmunity initiation, slowing of autoimmune disease progression, reduced leukocytic infiltration, and end tissue damage through blockade of CD40 L in murine models [88]. Dysregulated expression of CD40–CD40 L in pSS provides a strong theoretical basis for pursuing blockage of this co-stimulatory pathway. A recent study showed that anti-CD40 L agents revealed promising responses in the treatment of pSS. Anti-CD40 agents, on the other hand, demonstrate more reassuring safety profiles with promising therapeutic benefits [89].
- 2 Indirect B-Cell Blocking: Over-expression of soluble factors, such as antibodies, IFN- α , and BAFF, plays an important role in B-cell differentiation, survival, activation, and the initiation and continuation of pSS. BAFF antagonists, such as anti-BAFF antibodies may have therapeutic efficacy in patients with pSS [87]. The use of BAFF-blocking agent such as Belimumab found to be effective in treatment of pSS [90].

8. Conclusions

The immunopathogenesis of pSS is not clearly understood. The hallmark of this disease is an immunologically mediated inflammatory exocrinopathy that is initially characterized by infiltration of the salivary tissue by lymphocytes, CD27+memory B cells and plasma cells. In pSS, dysregulation of B cell populations is characterized by disturbances in peripheral B-cell homeostasis with depletion of CD27+ memory B cells in peripheral blood accompanied by evidences for the accumulation and retention of autoantibody-producing B cells in the inflamed glands. Moreover, the germinal centers of lymphoid tissue formed within the patients' inflamed glands seem to play a role in these abnormalities. These abnormalities provide new insight into the immunopathogenesis of pSS. The data presented in this review may facilitate an understanding of the underlying pathogenetic mechanisms, involved in this disease, and help in developing improved B cell depletion therapies for this disease.

Conflicts of interest

None to declare.

References

- [1] Moutsopoulos HM. Sjögren's syndrome: autoimmune epithelitis. *Clin Immunol Immunopathol* 1994;72(2):162–5.
- [2] Singh N, Cohen PL. The T cell in Sjögren's syndrome: force majeure, not spectateur. *J Autoimmun* 2012;39(3):229–33.
- [3] Le Pottier L, Devauchelle V, Pers JO, Jamin C, Youinou P. The mosaic of B-cell subsets (with special emphasis on primary Sjögren's syndrome). *Autoimmun Rev* 2007;6(3):149–54.
- [4] Nocturne G, Mariette X. Advances in understanding the pathogenesis of primary Sjögren's syndrome. *Nat Rev Rheumatol* 2013;9(9):544–56.
- [5] Kroese FG, Abdulahad WH, Haacke E, Bos NA, Vissink A, Bootsma H. B-cell hyperactivity in primary Sjögren's syndrome. *Expert Rev Clin Immunol* 2014;10(4):483–99.
- [6] Bohnhorst JØ, Bjørgan MB, Thoen JE, Jonsson R, Natvig JB, Thompson KM. Abnormal B cell differentiation in primary Sjögren's syndrome results in a depressed percentage of circulating memory B cells and elevated levels of soluble CD27 that correlate with Serum IgG concentration. *Clin Immunol* 2002;103(1):79–88.
- [7] Hansen A, Jacobi A, Pruss A, Kaufmann O, Scholze J, Lipsky PE, et al. Comparison of immunoglobulin heavy chain rearrangements between peripheral and glan-

- dular B cells in a patient with primary Sjögren's syndrome. *Scand J Immunol* 2003;57(5):470–9.
- [8] Stefanski AL, Tomiak C, Pleyer U, Dietrich T, Burmester GR, Dörner T. The Diagnosis and Treatment of Sjögren's Syndrome. *Dtsch Arztebl Int* 2017;114(20):354–61.
 - [9] Nagaraju K, Cox A, Casciola-Rosen L, Rosen A. Novel fragments of the Sjögren's syndrome autoantigens *a*-fodrin and type 3 muscarinic acetylcholine receptor generated during cytotoxic lymphocyte granule-induced cell death. *Arthritis Rheum* 2001;44(10):2376–86.
 - [10] Takeda K, Kaisho T, Akira S. Toll-like receptors. *Annu Rev Immunol* 2003;21:335–76.
 - [11] Looney RJ. Will targeting B cells be the answer for Sjögren's syndrome? *Arthritis Rheum* 2007;56(5):1371–7.
 - [12] Aqrawi LA, Skarstein K, Bredholt G, Brun JG, Brokstad KA. Autoantigen-specific memory B cells in primary Sjögren's syndrome. *Scand J Immunol* 2012;75(1):61–8.
 - [13] Sisto M, Lisi S, Ingravalle G, Lofrumento DD, D'Amore M, Ribatti D. Neovascularization is prominent in the chronic inflammatory lesions of Sjögren's syndrome. *Int J Exp Pathol* 2014;95(2):131–7.
 - [14] Carubbi F, Alunno A, Cipriani P, Di Benedetto P, Ruscitti P, Berardicurti O, et al. Is minor salivary gland biopsy more than a diagnostic tool in primary Sjögren's syndrome? Association between clinical, histopathological, and molecular features: a retrospective study. *Semin Arthritis Rheum* 2014;44(3):314–24.
 - [15] Båve U, Nordmark G, Lövgren T, Rönnelid J, Cajander S, Eloranta ML, et al. Activation of the type I interferon system in primary Sjögren's syndrome: a possible etiopathogenic mechanism. *Arthritis Rheum* 2005;52(4):1185–95.
 - [16] Gottenberg JE, Cagnard N, Lucchesi C, Letourneur F, Mistou S, Lazure T, et al. Activation of IFN pathways and plasmacytoid dendritic cell recruitment in target organs of primary Sjögren's syndrome. *Proc Natl Acad Sci U S A* 2006;103(8):2770–5.
 - [17] Mackay F, Schneider P. Cracking the BAFF code. *Nat Rev Immunol* 2009;9:491–502.
 - [18] Moriyama M, Hayashida JN, Toyoshima T, Ohya Y, Shinozaki S, Tanaka A, et al. Cytokine/chemokine profiles contribute to understanding the pathogenesis and diagnosis of primary Sjögren's syndrome. *Clin Exp Immunol* 2012;169(1):17–26.
 - [19] Amft N, Curnow SJ, Scheel-Toellner D, Devadas A, Oates J, Crocker J, et al. Ectopic expression of the B cell-attracting chemokine BCA-1 (CXCL13) on endothelial cells and within lymphoid follicles contributes to the establishment of germinal center-like structures in Sjögren's syndrome. *Arthritis Rheum* 2001;44(11):2633–41.
 - [20] Dawson LJ, Stanbury J, Venn N, Hasdimir B, Rogers SN, Smith PM. Antimutagenic antibodies in primary Sjögren's syndrome reversibly inhibit the mechanism of fluid secretion by human submandibular salivary acinar cells. *Arthritis Rheum* 2006;54(4):1165–73.
 - [21] Dawson LJ, Field EA, Harmer AR, Smith PM. Acetylcholine-evoked calcium mobilization and ion channel activation in human labial gland acinar cells from patients with primary Sjögren's syndrome. *Clin Exp Immunol* 2001;124(3):480–5.
 - [22] Nunez C, Nishimoto N, Gartland GL, Billips LG, Burrows PD, Kubagawa H, et al. B cells are generated throughout life in humans. *J Immunol* 1996;156(2):866–72.
 - [23] Garaud S, Youinou P, Renaudineau Y. DNA methylation and B-cell autoreactivity. *Adv Exp Med Biol* 2011;711:50–60.
 - [24] Tonegawa S. Somatic generation of antibody diversity. *Nature* 1983;302(5909):575–81.
 - [25] Hoffman William, Lakkis Fadi G, Chalasani Geetha. B cells, antibodies, and more. *Clin J Am Soc Nephrol* 2016;11(1):137–54.
 - [26] Scofield RH, Farris AD, Horsfall AC, Harley JB. Fine specificity of the autoimmune response to the Ro/SSA and La/SSB ribonucleoproteins. *Arthritis Rheum* 1999;42:199–209.
 - [27] Ramos-Casals M, Font J. Primary Sjögren's syndrome: current and emergent etiopathogenic concepts. *Rheumatology (Oxford)* 2005;44(11):1354–67.
 - [28] Hansen A, Lipsky PE, Dörner T. B cells in Sjögren's syndrome: indications for disturbed selection and differentiation in ectopic lymphoid tissue. *Arthritis Res Ther* 2007;9(4):218.
 - [29] Nanki T, Takada K, Komano Y, Morio T, Kanegane H, Nakajima A, et al. Chemokine receptor expression and functional effects of chemokines on B cells: implication in the pathogenesis of rheumatoid arthritis. *Arthritis Res Ther* 2009;11(5):R149.
 - [30] Hansen A, Reiter K, Ziprian T, Jacobi A, Hoffmann A, Gosemann M, et al. Dysregulation of chemokine receptor expression and function by B cells of patients with primary Sjögren's syndrome. *Arthritis Rheum* 2005;52(7):2109–19.
 - [31] Bikker A, van Woerkom JM, Kruijs AA, Wenting-van Wijk M, de Jager W, Bijlsma JW, et al. Increased expression of IL-7 in labial salivary glands of patients with primary Sjögren's syndrome correlates with increased inflammation. *Arthritis Rheum* 2010;62:969–77.
 - [32] Bohnhorst JØ, Bjorgan MB, Thoen JE, Natvig JB, Thompson KM. Bm1-bm5 classification of peripheral blood B cells reveals circulating germinal center founder cells in healthy individuals and disturbance in the B cell subpopulations in patients with primary Sjögren's syndrome. *J Immunol* 2001;167(7):3610–8.
 - [33] Bohnhorst JØ, Thoen JE, Natvig JB, Thompson KM. Significantly depressed percentage of CD27 [memory] B cells among peripheral blood B cells in patients with primary Sjögren's syndrome. *Scand J Immunol* 2001;54(4):421–7.
 - [34] Hansen A, Odendahl M, Reiter K, Jacobi AM, Feist E, Scholz J, et al. Diminished peripheral blood memory B cells and accumulation of memory B cells in the salivary glands of patients with Sjögren's syndrome. *Arthritis Rheum* 2002;46(8):2160–71.
 - [35] Binard A, Le Pottier L, Devauchelle-Pensec V, Saraux A, Youinou P, Pers JO. Is the blood B-cell subset profile diagnostic for Sjögren syndrome? *Ann Rheum Dis* 2009;68(9):1447–52.
 - [36] Karlén M, Hansen T, Nordal HH, Brun JG, Jonsson R, Appel S. Expression of toll-like receptor -7 and -9 in B cell subsets from patients with primary Sjögren's syndrome. *PLoS One* 2015;10(3):e0120383.
 - [37] Corsiero E, Sutcliffe N, Pitzalis C, Bombardieri M. Accumulation of self-reactive naive and memory B cell reveals sequential defects in B cell tolerance checkpoints in Sjögren's syndrome. *PLoS One* 2014;9(12):e114575.
 - [38] Brezinschek HP, Brezinschek RI, Lipsky PE. Analysis of the heavy chain repertoire of human peripheral B cells using single-cell polymerase chain reaction. *J Immunol* 1995;155(1):190–202.
 - [39] Brezinschek HP, Foster SJ, Brezinschek RI, Dörner T, Domiati-Saad R, Lipsky PE. Analysis of the human VH gene repertoire. Differential effects of selection and somatic hypermutation on human peripheral CD5(+)IgM+ and CD5(-)IgM+ B cells. *J Clin Invest* 1997;99(10):2488–501.
 - [40] Glauzy S, Sng J, Bannock JM, Gottenberg JE, Korganow AS, Cacoub P, et al. Defective early B cell tolerance checkpoints in Sjögren's syndrome patients. *Arthritis Rheumatol* 2017;69(11):2203–8.
 - [41] Kaschner S, Hansen A, Jacobi A, Reiter K, Monson NL, Odendahl M, et al. Immunoglobulin Vlambda light chain gene usage in patients with Sjögren's syndrome. *Arthritis Rheum* 2001;44(11):2620–32.
 - [42] Heimbacher C, Hansen A, Pruss A, Jacobi A, Reiter K, Lipsky PE, et al. Immunoglobulin Vκ Light Chain Gene Analysis in Patients With Sjögren's Syndrome. *Arthritis Rheum* 2001;44:626–37.
 - [43] d'Arbonneau F, Pers JO, Devauchelle V, Pennec Y, Saraux A, Youinou P. BAFF-induced changes in B cell antigen receptor-containing lipid rafts in Sjögren's syndrome. *Arthritis Rheum* 2006;54(1):115–26.
 - [44] Abdulahad WH, Meijer JM, Kroese FG, Meiners PM, Vissink A, Spijkervet FK, et al. B cell reconstitution and T helper cell balance after rituximab treatment of active primary Sjögren's syndrome: a double-blind, placebo-controlled study. *Arthritis Rheum* 2011;63(4):1116–23.
 - [45] Szyszko EA, Brun JG, Skarstein K, Peck AB, Jonsson R, Brokstad KA. Phenotypic diversity of peripheral blood plasma cells in primary Sjögren's syndrome. *Scand J Immunol* 2011;73(1):18–28.
 - [46] Szabó K, Papp G, Szántó A, Tarr T, Zeher M. A comprehensive investigation on the distribution of circulating follicular T helper cells and B cell subsets in primary Sjögren's syndrome and systemic lupus erythematosus. *Clin Exp Immunol* 2016;183(1):76–89.
 - [47] Mietzner B, Tsuiji M, Scheid J, Velinon K, Tiller T, Abraham K, et al. Autoreactive IgG memory antibodies in patients with systemic lupus erythematosus arise from nonreactive and polyreactive precursors. *Proc Natl Acad Sci U S A* 2008;105(28):9727–32.
 - [48] Dörner T, Jacobi AM, Lee J, Lipsky PE. Abnormalities of B cell subsets in patients with systemic lupus erythematosus. *J Immunol Methods* 2011;363(2):187–97.
 - [49] Salomonsson S, Larsson P, Tengné P, Mellquist E, Hjelmström P, Wahren-Herlenius M. Expression of the B cell-attracting chemokine CXCL13 in the target organ and autoantibody production in ectopic lymphoid tissue in the chronic inflammatory disease Sjögren's syndrome. *Scand J Immunol* 2002;55(4):336–42.
 - [50] Barone F, Bombardieri M, Manzo A, Blades MC, Morgan PR, Challacombe SJ, et al. Association of CXCL13 and CCL21 expression with the progressive organization of lymphoid-like structures in Sjögren's syndrome. *Arthritis Rheum* 2005;52(6):1773–84.
 - [51] Manzo A, Vitolo B, Humby F, Caporali R, Jarroissay D, Dell'accio F, et al. Mature antigen-experienced T helper cells synthesize and secrete the B cell chemoattractant CXCL13 in the inflammatory environment of the rheumatoid joint. *Arthritis Rheum* 2008;58(11):3377–87.
 - [52] Vissers JL, Hartgers FC, Lindhout E, Figdor CG, Adema GJ. BLC (CXCL13) is expressed by different dendritic cell subsets in vitro and in vivo. *Eur J Immunol* 2001;31(5):1544–9.
 - [53] Sáez de Guinoa J, Barrio L, Mellado M, Carrasco YR. CXCL13/CXCR5 signaling enhances BCR-triggered B-cell activation by shaping cell dynamics. *Blood* 2011;118(6):1560–9.
 - [54] Rupprecht TA, Plate A, Adam M, Wick M, Kastenbauer S, Schmidt C, et al. The chemokine CXCL13 is a key regulator of B cell recruitment to the cerebrospinal fluid in acute Lyme neuroborreliosis. *J Neuroinflammation* 2009;6:42–53.
 - [55] Cupedo T, Mebius RE. Role of chemokines in the development of secondary and tertiary lymphoid tissues. *Semin Immunol* 2003;15:243–8.
 - [56] Moisini I, Davidson A. BAFF: a local and systemic target in autoimmune diseases. *Clin Exp Immunol* 2009;158(2):155–63.
 - [57] Daridon C, Devauchelle V, Hutin P, Le Berre R, Martins-Carvalho C, Bendaoud B, et al. Aberrant expression of BAFF by B lymphocytes infiltrating the salivary glands of patients with primary Sjögren's syndrome. *Arthritis Rheum* 2007;56(4):1134–44.
 - [58] Ittah M, Miceli-Richard C, Eric Gottenberg J, Lavie F, Lazure T, Ba N, et al. Mariette X. B cell-activating factor of the tumor necrosis factor family [BAFF] is expressed under stimulation by interferon in salivary gland epithelial cells in primary Sjögren's syndrome. *Arthritis Res Ther* 2006;8(2):R51.
 - [59] Brito-Zerón P, Ramos-Casals M, Nardi N, Cervera R, Yagüe J, Ingelmo M, Font J. Circulating monoclonal immunoglobulins in Sjögren syndrome: prevalence and clinical significance in 237 patients. *Medicine (Baltimore)* 2005;84(2):90–7.

- [60] Szyszko EA, Brokstad KA, Oijordsbakken G, Jonsson MV, Jonsson R, Skarstein K. Salivary glands of primary Sjögren's syndrome patients express factors vital for plasma cell survival. *Arthritis Res Ther* 2011;13(1):R2.
- [61] Tengnér P, Halse AK, Haga HJ, Jonsson R, Wahren-Herlenius M. Detection of anti-Ro/SSA and anti-La/SSB autoantibody-producing cells in salivary glands from patients with Sjögren's syndrome. *Arthritis Rheum* 1998;41(12):2238–48.
- [62] Halse A, Harley JB, Kroneld U, Jonsson R. Ro/SSA-reactive B lymphocytes in salivary glands and peripheral blood of patients with Sjögren's syndrome. *Clin Exp Immunol* 1999;115(1):203–7.
- [63] Aqrawi LA, Skarstein K, Oijordsbakken G, Brokstad KA. Ro52- and Ro60-specific B cell pattern in the salivary glands of patients with primary Sjögren's syndrome. *Clin Exp Immunol* 2013;172(2):228–37.
- [64] Aqrawi LA, Brokstad KA, Jakobsen K, Jonsson R, Skarstein K. Low number of memory B cells in the salivary glands of patients with primary Sjögren's syndrome. *Autoimmunity* 2012;45(7):547–55.
- [65] Theander E, Vasaitis L, Bäcklund E, Nordmark G, Warfvinge G, Liedholm R, et al. Lymphoid organisation in labial salivary gland biopsies is a possible predictor for the development of malignant lymphoma in primary Sjögren's syndrome. *Ann Rheum Dis* 2011;70(8):1363–8.
- [66] Jonsson R, Kroneld U, Bäckman K, Magnusson B, Tarkowski A. Progression of sialoadenitis in Sjögren's syndrome. *Br J Rheumatol* 1993;32(7):578–81.
- [67] Risselada AP, Looije MF, Kruize AA, Bijlsma JW, van Roon JA. The role of ectopic germinal centers in the immunopathology of primary Sjögren's syndrome: a systematic review. *Semin Arthritis Rheum* 2013;42(4):368–76.
- [68] Dörner T, Lipsky PE. Abnormalities of B cell phenotype, immunoglobulin gene expression and the emergence of autoimmunity in Sjögren's syndrome. *Arthritis Res* 2002;4(6):360–71.
- [69] Daniels TE, Cox D, Shiboski CH, Schiødt M, Wu A, Lanfranchi H, et al. Associations between salivary gland histopathologic diagnoses and phenotypic features of Sjögren's syndrome among 1,726 registry participants. *Arthritis Rheum* 2011;63(7):2021–30.
- [70] Voulgarelis M, Tzioufas AG. Pathogenetic mechanisms in the initiation and perpetuation of Sjögren's syndrome. *Nat Rev Rheumatol* 2010;6(9):529–37.
- [71] Sène D, Ismael S, Forien M, Charlotte F, Kaci R, Cacoub P, et al. Ectopic Germinal Center-Like Structures in Minor Salivary Gland Biopsy Tissue Predict Lymphoma Occurrence in Patients With Primary Sjögren's Syndrome. *Arthritis Rheumatol* 2018;70(9):1481–8.
- [72] Alunno A, Leone MC, Giacomelli R, Gerli R, Carubbi F. Lymphoma and Lymphomagenesis in Primary Sjögren's Syndrome. *Front Med (Lausanne)* 2018;5:102.
- [73] Dong L, Chen Y, Masaki Y, Okazaki T, Umehara H. Possible Mechanisms of Lymphoma Development in Sjögren's Syndrome. *Curr Immunol Rev* 2013;9(1):13–22.
- [74] Bernasconi NL, Onai N, Lanzavecchia A. A role for Toll-like receptors in acquired immunity: up-regulation of TLR9 by BCR triggering in naive B cells and constitutive expression in memory B cells. *Blood* 2003;101:4500–4.
- [75] Kiripolsky J, Kramer JM. Current and Emerging Evidence for Toll-Like Receptor Activation in Sjögren's Syndrome. *J Immunol Res* 2018;2018:1246818.
- [76] Browne EP. Regulation of B-cell responses by Toll-like receptors. *Immunology* 2012;136(4):370–9.
- [77] Hua Z, Hou B. TLR signaling in B-cell development and activation. *Cell Mol Immunol* 2013;10(2):103–6.
- [78] Liu YJ, Arpin C. Germinal center development. *Immunol Rev* 1997;156(1):111–26.
- [79] Agematsu K, Nagumo H, Oguti Y, et al. Generation of plasma cells from peripheral blood memory B cells: synergistic effect of interleukin-10 and CD27/CD70 interaction. *Blood* 1998;91:173–80.
- [80] Hintzen RQ, Lens SM, Beckmann MP, Goodwin RG, Lynch D, van Lier RA. Characterization of the human CD27 ligand, a novel member of the TNF gene family. *J Immunol* 1994;152(4):1762–73.
- [81] Halse A, Tengnér P, Wahren-Herlenius M, Haga H, Jonsson R. Increased frequency of cells secreting interleukin-6 and interleukin-10 in peripheral blood of patients with primary Sjögren's syndrome. *Scand J Immunol* 1999;49(5):533–8.
- [82] Llorente L, Richaud-Patin Y, Fior R, Alcocer-Varela J, Wijdenes J, Fourrier BM, et al. In vivo production of interleukin-10 by non-T cells in rheumatoid arthritis, Sjögren's syndrome, and systemic lupus erythematosus. A potential mechanism of B lymphocyte hyperactivity and autoimmunity. *Arthritis Rheum* 1994;37(11):1647–55.
- [83] Avery DT, Ellyard JL, Mackay F, Corcoran LM, Hodgkin PD, Tangye SG. Increased expression of CD27 on activated human memory B cells correlates with their commitment to the plasma cell lineage. *J Immunol* 2005;174(7):4034–42.
- [84] Nagumo H, Agematsu K. Synergistic augmentative effect of interleukin-10 and CD27/CD70 interactions on B-cell immunoglobulin synthesis. *Immunology* 1998;94(3):388–94.
- [85] Brito-Zerón P, Retamozo S, Gheitasi H, Ramos-Casals M. Treating the underlying pathophysiology of primary Sjögren syndrome: recent advances and future prospects. *Drugs* 2016;76(17):1601–23.
- [86] Verstappen GM, van Nimwegen JF, Vissink A, Kroese FGM, Bootsma H. The value of rituximab treatment in primary Sjögren's syndrome. *Clin Immunol* 2017;182:62–71.
- [87] Gottenberg JE, Dörner T, Bootsma H, Devauchelle-Pensec V, Bowman SJ, Mariette X, et al. Efficacy of Epratuzumab, an Anti-CD22 Monoclonal IgG Antibody, in Systemic Lupus Erythematosus Patients With Associated Sjögren's Syndrome: Post Hoc Analyses From the EMBODY Trials. *Arthritis Rheumatol* 2018;70(5):763–73.
- [88] Toubi E, Shoenfeld Y. The role of CD40–CD154 interactions in autoimmunity and the benefit of disrupting this pathway. *Autoimmunity* 2004;37(6–7):457–64.
- [89] Jobling K, Ng WF. CD40 as a therapeutic target in Sjögren's syndrome. *Expert Rev Clin Immunol* 2018;14(7):535–7.
- [90] De Vita S, Quartuccio L, Seror R, Salvin S, Ravaud P, Fabris M, et al. Efficacy and safety of belimumab given for 12 months in primary Sjögren's syndrome: the BELISS open-label phase II study. *Rheumatology (Oxford)* 2015;54(12):2249–56.