



Full Length Article

Cbl-PI3K interaction regulates Cathepsin K secretion in osteoclasts

Jungeun Yu, Naga Suresh Adapala, Laura Doherty, Archana Sanjay*

Department of Orthopaedic Surgery, UConn Health, Farmington, CT 06030, United States of America



ARTICLE INFO

Keywords:

Osteoclast
Vesicular trafficking
Cathepsin K
LAMP2
PI3K and Cbl

ABSTRACT

Effective bone resorption by osteoclasts is critical for balanced bone remodeling. We have previously reported that mice harboring a substitution mutation of tyrosine 737 to phenylalanine in the adapter protein Cbl (Cbl^{Y737F}, YF) have increased bone volume partly due to decreased osteoclast-mediated bone resorption. The Cbl^{Y737F} mutation abrogates interaction between Cbl and the p85 subunit of PI3K. Here, we studied the mechanism for defective resorptive function of YF mutant osteoclasts. The YF osteoclasts had intact actin cytoskeletons and sealing zones. Expression and localization of proteins needed for acidification of the resorptive lacunae were also comparable between the WT and YF osteoclasts. In contrast, secretion of Cathepsin K, a major protease needed to degrade collagen, was diminished in the conditioned media derived from YF osteoclasts. The targeting of Cathepsin K into LAMP2-positive vesicles was also compromised due to decreased number of LAMP2-positive vesicles in YF osteoclasts. Further, we found that in contrast to WT, conditioned media derived from YF osteoclasts promoted increased numbers of alkaline phosphatase positive colonies, and increased expression of osteogenic markers in WT calvarial cultures. Cumulatively, our results suggest that the Cbl-PI3K interaction regulates Cathepsin K secretion required for proper bone resorption, and secretion of factors which promote osteogenesis.

1. Introduction

Osteoclasts are large, multinucleated cells with a unique capacity to degrade the organic and inorganic bone matrices. This function is absolutely necessary for bone modeling bone remodeling and to maintain calcium and phosphate ion homeostasis [1]. The resorptive function of osteoclasts also plays a role in the formation of hematopoietic stem cell niches in the bone marrow [2,3]. Because osteoclasts are a fundamental factor in the pathogenesis of virtually all diseases associated with bone loss, the insights gained into the mechanisms by which they form and degrade bone are clinically significant.

Phosphatidylinositol-3 Kinase (PI3K) is a prominent lipid kinase that phosphorylates the 3'-hydroxyl group on the inositol ring of phosphatidylinositol. PI3K is stimulated by Macrophage colony stimulating factor (M-CSF) and $\alpha_v\beta_3$ integrins, both of which are critical for osteoclast resorptive function. Based on the sequence similarity, composition of subunits and substrate specificity, PI3Ks are divided into three classes: Class I, II, and III. While class II and III PI3Ks are constitutively activated, Class I PI3Ks are activated in response to extracellular stimuli. Class I PI3K is a heterodimer of p110 catalytic and p85 regulatory subunits. The p110 catalytic subunit binds to the p85 regulatory subunit with high affinity and this interaction prevents

inadvertent activation of PI3K [4,5]. Upon engagement of the SH2 domain of the p85 subunit through binding to phosphorylated tyrosines of receptors or signaling proteins, the inhibitory effect of p85 on p110 is released, resulting in increased lipid kinase activity [6].

The E3 ubiquitin ligase Cbl is an adaptor protein that regulates PI3K signaling *via* interaction with the p85 regulatory subunit [7]. In macrophages, Src family kinase-mediated phosphorylation of Cbl is required for Cbl-PI3K association, and for the transfer of this complex to the actin rich cytoskeleton [8]. In the skeletal system, Cbl plays an important role in bone remodeling by regulating bone resorption of osteoclasts [9–11]. While establishing a role for Cbl in bone resorption [9,12,13], we identified a unique function of Cbl through the requirement of a tyrosine at 737 in the C-terminal half, for its interaction with the SH2 domain of the p85 regulatory subunit of PI3K [14]. Substitution mutation of the tyrosine 737 into phenylalanine (Cbl^{Y737F}) resulted in abrogation of Cbl-p85 association, [15] but did not impact the E3 ligase activity of Cbl and the protein level of p85 [16]. We have shown that Src-mediated phosphorylation of Cbl is required for Cbl-p85 interaction, and adenoviral overexpression of Cbl^{Y737F} in WT osteoclasts decreased pit forming activity [10,11,15].

To study the impact of the Cbl-PI3K interaction in the skeletal system, a global knock-in mouse model in which the tyrosine 737 was

* Corresponding author at: Department of Orthopaedic Surgery, UConn Health, 263 Farmington Ave, Farmington, CT 06030, United States of America.
E-mail address: asanjay@uconn.edu (A. Sanjay).

Table 1
Primers used for real-time PCR

Gene	Forward primer	Reverse primer
<i>Atp6v0d2</i>	CAATGAAGCGTCACCTCTGA	TCAGCTATTGAACGCTGGTG
<i>Ctsk</i>	CAGCTTCCCAAGATGTGAT	AAAATGCCCTGTTGTGTCC
<i>Car2</i>	CAAGCACACGGACCAGA	ATGAGCAGAGGCTGTAGG
<i>Clen7</i>	GACTGGCTGTGGAA	TCTCGCTTGAGTGATGTTGAC C
<i>Acp5</i>	GCAGTCCCTAGAAGATGGATT	ATTTGTAGGCCAGCAGCAC
<i>Mtif</i>	TTGATGGATCCGCGCTTGCAAATG	TATGTTGGGAAGGTTGGCTGGACA
<i>Tfeb</i>	GTGATTGACAACATTATGCGCC	GCGTGTAGGCATCTTGCATCT
<i>Tfe3</i>	CCAAGCTGGCTTCCAGGCTCTCAC	GTTAATGTTGAATCGCCTGCGTCG
<i>Wnt10b</i>	AGGCTTCTCCTCCGTTTCAGTTGT	ATTCCCACCTTCTGCTGAAGAA
<i>Pdgfrb</i>	GGGATCCCATTCTGAGGAAC	CGGTCCAGGTGAGAAAGATTG
<i>Sema4d</i>	TCITTTGCTGACGTGATCCAG	CAGATCAGCTGGCCTTTAG
<i>Sphk1</i>	TGAGGTGGTGAATGGGCTAATGGA	AACAGCAGTGTGCAGTTGATGAGC
<i>Sphk2</i>	TGGGCTGTCCITCAACCTCATA	AGTGACAATGCCTTCCCACTCACT
<i>Bmp6</i>	AGAAGGGCACTCTTTCAGGTTCCA	TCACACCACGAGAGTCAACACAA
<i>Osx</i>	GATGGCGTCTCTGCTTG	GCCATAGTGAGCTTCTCTCAA
<i>Alp1</i>	GGAGATGGTATGGGCGTCTC	CTGGCCCTAAGGATTCCGGG
<i>Col1a1</i>	CGATGGATTCCCGTTCGAGT	ACATTAGGCGCAGGAAGGTC
<i>Gapdh</i>	AGGTCGGTGTGAACGGATTG	TGTAGACCATGTAGTTGAGGTCA

substituted to phenylalanine (Cbl^{Y737F/Y737F} mice; hence forth YF mice) was used [17]. Our skeletal characterization of YF mice demonstrated that both bone resorption and formation are perturbed. Increased bone volume resulted due to defective osteoclast-mediated resorption both during homeostasis and upon acute estrogen deprivation [16,18]. In this study, we investigated the molecular mechanism that leads to decreased osteoclast function in YF mice.

2. Materials and methods

2.1. Mice

Generation of Cbl^{Y737F/Y737F} (YF) mice with a point mutation on the p85 regulatory subunit-binding site of Cbl were previously described [17]. Mice were genotyped for WT and YF alleles as previously reported [16]. All mice used in the study were in a mixed C57BL/6JX129SvJ background. For all experiments, 6–8 week old mice were used. All animal procedures were conducted according to protocols approved by the University of Connecticut Health Center Animal Care Committee. Experiments were repeated at least three times, and each time a different cohort of WT and YF mice were used to generate primary osteoclast cultures.

2.2. Culture of bone marrow-derived macrophages and osteoclast formation

To obtain BMMs, marrow was removed by flushing with a 26-gauge needle and erythrocytes were lysed in 150 mM NH₄Cl, 10 mM KHCO₃, and 0.1 mM EDTA (pH 7.4). Cells were centrifuged and the pellet was suspended in Minimum essential medium α -modification (α MEM) in the presence of 10% fetal bovine serum (FBS) (both from Sigma-Aldrich, St. Louis, MO). Cells were seeded at a density of 300,000 cells/cm² and cultured for 3 to 4 days on petri dishes in the presence of recombinant human M-CSF at 30 ng/ml. M-CSF cDNA and expression vectors were obtained from Dr. D. Fremont (St. Louis, MO) and M-CSF purified as previously reported [19]. For osteoclast formation, cells were collected following treatment with 0.25% trypsin/EDTA for 5 min and seeded at a density of 47,000 cells/cm² in α MEM with 10% FBS, M-CSF at 30 ng/ml, and recombinant murine RANKL at 10 ng/ml. RANKL cDNA and expression vectors were obtained from Dr. M. Glogauer (Toronto, Canada), and GST-tagged RANKL was expressed and purified as described [20]. Cultures were carried out until formation of multinucleated TRAP-positive cells. TRAP enzyme histochemistry was conducted using a commercial kit (Sigma-Aldrich, St. Louis, MO), in accordance with manufacturer's instructions. TRAP-positive cells containing > 3 nuclei were considered osteoclasts. To produce conditioned media, BMMs

were either cultured in α MEM with M-CSF (20 ng/ml) for 5 days on petri dishes. To generate osteoclast-conditioned medium, BMMs were cultured on Hydroxyapatite(HA)-coated plates in the presence of M-CSF (20 ng/ml) and RANKL (50 ng/ml) for 6 days. Conditioned media was harvested and centrifuged at 100,000 g for 1 h at 4 °C. The supernatant was filtered through 20- μ m filters and either immediately used or stored at -80 °C. CD1 calvarial osteoprogenitors were derived from 2 to 5 day old CD-1 pups as described previously [16]. After seven days, cultures were stained for toluidine blue and alkaline phosphatase(ALP) staining as per manufacturer's instructions or harvested to isolate mRNA.

2.3. Quantitative reverse transcription PCR

To analyze expression of differentiation markers by real time PCR, cells were lysed with Trizol (Thermo Fisher Scientific, Waltham, MA), and RNAs were extracted as per manufacturer's instructions. 1 μ g of RNA was treated with DNase (Thermo Fisher Scientific), and reverse transcribed to cDNA using the Superscript III cDNA Synthesis Kit (Thermo Fisher Scientific). Quantitative reverse transcription-PCR (qRT-PCR) was performed with 10 ng of cDNA per reaction using SYBR Green Master Mix (Thermo Fisher Scientific) and validated custom designed primers. All primers used in the study are listed in Table 1. ABI Step One Plus Real Time PCR System using the following thermal cycler program: 95 °C for 10 min, 40 cycles of 95 °C for 15 s, and 57 °C for 1 min. Product specificity was confirmed by melt curve analysis, and relative expression was determined using $\Delta\Delta$ CT analysis after normalization to *Gapdh* expression.

2.4. Bone resorption

In vitro bone resorbing activity was assayed as described previously [12,16]. Briefly, osteoclasts were generated in Collagen gel (Nitta Gelatin Co., Osaka, Japan). After 6 days of culture, mature osteoclasts were released from Collagen gel by gentle digestion with 0.1% Collagenase (Calbiochem, San Diego, CA), and cells were seeded onto sterile bovine bone discs in 96-well plates. 48 h later, bone discs were immersed in 1 M Ammonium hydroxide (Sigma-Aldrich) for 5 min, sonicated for 10 s, and then stained for 4 min with 1% toluidine blue in 1% sodium borate (both from Sigma-Aldrich) and briefly rinsed in water. Pit area was quantified with the measuring tool in Adobe Photoshop CS3 Extended Edition, and was normalized to the number of osteoclasts actually present in each sample, determined by counting osteoclasts that were plated on tissue culture plastic.




Fig. 1. Actin organization and sealing zone formation are not altered, but bone-resorbing activity is decreased in YF osteoclasts. BMMs were cultured in collagen gel with M-CSF and RANKL for 6 days. Mature osteoclasts were released by treating cultures with 0.1% Collagenase, and cells were plated on glass cover slips (A) or bovine bone discs (B and C). Cells were stained with rhodamine phalloidin to visualize actin ring (A) and sealing zone (B) formation (orange). Dotted lines indicate boundaries of osteoclasts. (C) Bone discs were stained with toluidine blue to visualize pit formation. (D) Stained area was quantified and normalized for the numbers of osteoclasts. Values are mean ± SD, *n* = 9, *Significantly different between WT and YF, *p* < 0.05. Scale bars represent 100 μM (A), 50 μM f (B), and 100 (C). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

2.5. Inorganic matrix resorption

Mature osteoclasts developed on collagen gel as described above. Cells (10⁵) were then seeded onto HA-coated wells (BD BioCoat™ Osteologic™, BD Bioscience, San Jose, CA). One day after, wells were

rinsed in water and treated with 5% sodium hypochlorite for 5 min. Analysis of resorbed area was performed using bright field microscopy.




Fig. 2. YF osteoclasts can resorb HA-coated surface. (A) Mature osteoclasts were plated on HA-coated surface and resorbed area was visualized. (B) Percentage of total resorbed area is indicated. (C) Cells cultured on coverslips were immunostained for v-ATPase (E subunit, green) using an antibody that was a gift from Dr. Beth Lee, OSU. Similar distribution of v-ATPase was observed for both genotypes. (D) mRNA levels of *Car2*, *Clc7*, and *Atp6v0d2* were analyzed by qRT-PCR and were normalized by mRNA levels of *Gapdh*. Values are mean \pm SD, $n = 5$. Scale bar represents 100 μ m for (A) and (C). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

2.6. Fluorescence microscopy

Cells were plated on sterile FBS-coated glass coverslips (Corning Inc. Corning, NY) and fixed in PBS containing 4% paraformaldehyde for 10 min, then permeabilized with 0.3% Triton X-100 (all from Sigma-Aldrich) for 5 min. Coverslips for Actin labeling were incubated in a 1:40 dilution in PBS of rhodamine phalloidin stock solution (Thermo Fisher Scientific) for 20 min. To detect presences of enzymes, cells were incubated overnight at 4 °C with primary antibodies diluted in PBS supplemented with 3% bovine serum albumin (BSA, Sigma-Aldrich). Cells were rinsed with PBS and then incubated with secondary antibodies conjugated to fluorescein isothiocyanate (FITC) or Phycoerythrin

(PE) (both from Thermo Fisher Scientific) for 1 h at room temperature. The following antibodies were used: anti-vATPase (E subunit) (gift from Dr. Beth Lee, Ohio State University), anti-LAMP2, anti-Cathepsin K, and anti-Cathepsin D (Santa Cruz Biotechnology, Dallas, TX). To visualize nuclei, cells were stained with DAPI (Thermo Fisher Scientific) before mounting. Cells were examined on a Leica fluorescence microscope (Model DMI6000B), and images were collected using the Leica Application Suite X (LASX) CLAS X 1.5.1.1387 (Leica Microsystems, Buffalo Grove, IL). For labeling lysosomes in live cells, cells were stained with LysoTracker® fluorescent dye (Thermo Fisher Scientific) as per manufacturer's instructions. LAMP2-positive vesicles were quantified by using counting tool from LASX after imaging and the number

was normalized by number of nuclei stained with DAPI.

2.7. Cathepsin K and Cathepsin D secretion

Mature osteoclasts were seeded onto 24-well plates (50,000 cells/well). Cells were cultured in the presence of 10% FBS containing medium for 24 h. Conditioned media was collected, and cells were harvested and lysed in mRIPA buffer for SDS-PAGE analysis to detect the presence of proteins.

2.8. Western blotting analysis

Clarified total cell lysate was electrophoresed on 10 or 12% SDS-PAGE as previously described [18]. Western blots were probed with anti-Cathepsin K or anti-Cathepsin D antibodies; the blots were stripped and reprobed with anti- β -Actin antibodies (Cell Signaling Technology, Danvers, MA). The amount of protein in individual bands was quantified by using Odyssey Infrared Imaging Systems software 2.1 (LI-COR Biosciences, Lincoln, NE) as previously reported [16,21].

2.9. Statistical analysis

Experiments conducted in this study were repeated at least three times. Data were expressed, as the mean \pm SD. Significant differences were determined using Student's *t*-test. $p < 0.05$ vs. control was considered significant.

3. Results

3.1. Actin organization and sealing zone formation are intact but bone resorption is decreased in YF osteoclasts

To resorb bone, mature osteoclasts attach to bone *via* integrin-regulated rearrangement of the cytoskeleton, creating a sealing zone that is necessary for activation and polarization of osteoclasts. Osteoclasts were seeded on glass or bovine bone discs, and were stained with rhodamine phalloidin to visualize actin cytoskeleton organization. Both WT and YF osteoclasts formed the typical actin ring on glass cover slips and sealing zone on bone discs, suggesting that the Cbl-PI3K interaction is not required for organization of actin cytoskeleton or the sealing zone formation (Fig. 1A and B). We also determined the *in vitro* pit forming activity of osteoclasts plated on bovine bone discs. In agreement with our previously published results [12,16] we found that compared to WT osteoclast, YF osteoclast formed shallower pits, and there was a 60% decrease in resorbed pit area (Fig. 1C and D) as we have reported previously [16,21].

3.2. Expression of factors needed for acid formation is not altered in YF osteoclasts

Following sealing zone formation, the ruffled border is generated by the simultaneous fusion of numerous secretory lysosomes with the plasma membrane [22]. The ruffled border functions as a site of secretion and externalization of hydrochloric acid and proteases. The acidic environment of osteoclasts dissolves hydroxyapatite (HA), and the proteases hydrolyze the collagen-rich organic matrix [23]. To examine acid secretion, WT and YF osteoclasts were plated on HA-coated surfaces (Fig. 2A). The resorbed area was about 20% decreased in YF osteoclasts, but was not statistically different from WT (Fig. 2B). Proteins implicated in acidification of resorption lacuna include carbonic anhydrase, chloride channel, and vacuolar ATPase. Consistent with no significant difference in resorption of HA-coated slides, which occurs exclusively through release of protons and acidification of the resorptive lacuna, immunofluorescence analysis showed that the cellular localization of vATPase [24] was also comparable between the two genotypes (Fig. 2C). We found that expression of mRNA for *Car2*, *Clc7*,

and *Atp6v0d2* was comparable between WT and YF osteoclasts (Fig. 2D). Together these results suggest that the loss of the Cbl-PI3K interaction in osteoclasts attenuates bone resorption, but does not regulate components of the acid-generating machinery.

3.3. The number of LAMP2-positive vesicles is significantly decreased in YF osteoclasts

Bone resorption by osteoclasts is highly dependent on lysosomes that release proteases in the resorption lacuna. We observed lysosomes in osteoclasts using LysoTracker fluorescent dye. We found that LysoTracker staining was attenuated in YF osteoclasts compared to WT (Fig. 3A). Osteoclasts were also stained for LAMP2, an integral lysosomal membrane protein [25]. We found that the number of LAMP2-positive vesicles in each osteoclast was significantly decreased in YF osteoclasts compared to WT (Fig. 3B and C). In order to understand the reason for decreased LAMP2 positive lysosomes in YF osteoclasts, we determined expression of the MITF/TFE family of transcription factors that is required for lysosomal biogenesis [[26]]. We found that the expression of *Mtif*, *TfeB*, and *Tfe3* was comparable between WT and YF osteoclasts (Supplementary Fig. 1). Together, these results suggest that the loss of Cbl-p85 interaction in osteoclasts results in decreased numbers of LAMP2-positive lysosomes, but this interaction is not involved in regulating the transcription factors needed for lysosomal biogenesis.

3.4. Secretion of Cathepsin K is decreased in YF osteoclasts

Lysosomal proteases required for bone resorption include Tartrate acid resistant phosphatase (TRAP) and Cathepsin K [1]. Most proteases, including Cathepsin D, K, and TRAP are synthesized in the rough endoplasmic reticulum (rER) as a pro-form, and later cleaved into a mature form in the lysosomes [27,28]. The mRNA levels of *Acp5* and *Ctsk* were significantly increased in YF osteoclasts (Fig. 4A). After synthesis in the endoplasmic reticulum, these enzymes are transported from trans Golgi network (TGN) to the secretory lysosomes (SL), which have all the features of lysosomes, but undergo regulated secretion in response to specific stimuli [29]. While a substantial amount of TRAP is secreted directly from TGN, the majority of Cathepsin K is found in SL [30]. We examined the protein expression and secretion of Cathepsin K. For these experiments, equal numbers of mature osteoclasts were plated (Fig. 4B). We found that compared to WT, there was about a 2-fold increase in Cathepsin K protein level in total cell lysates, but the amount of Cathepsin K secreted by YF osteoclasts in the conditioned media was significantly diminished (~75% decrease) (Fig. 4C, left, and 4D). In contrast, protein expression and secretion of Cathepsin D, an aspartic protease mostly present in lysosomes [31], was comparable between WT and YF osteoclasts (Fig. 4C, right). Next we examined the cellular distribution of Cathepsin K. Mature osteoclasts grown on bone slices were stained with rhodamine phalloidin to visualize actin cytoskeleton. As reported by others we also found that in WT osteoclasts, Cathepsin K exclusively localized within actin ring. In contrast, in a majority of YF osteoclasts, Cathepsin K distribution was not within the actin ring (Supplementary Fig. 2A and B). Further, compared to WT, colocalization of Cathepsin K within LAMP2-positive vesicles was also decreased in YF osteoclasts (Supplementary Fig. 2C).

3.5. Conditioned media derived from YF osteoclasts enhances osteogenesis

Independent of their resorptive activity, osteoclasts also regulate bone formation [24,32]. YF mice also have increased numbers of dysfunctional osteoclasts and an increased bone formation rate [16,33]. Therefore, we investigated whether YF osteoclasts can also induce osteogenesis *in vitro*. To test this, bone marrow-derived sorted CD11b⁺ cells were cultured in the presence of M-CSF to generate BMMS, or were seeded onto HA-coated surfaces with M-CSF and RANKL to obtain




Fig. 3. The number of LAMP2-positive vesicles is decreased in YF osteoclasts. Mature osteoclasts seeded on glass cover slips were fixed and stained with LysoTracker® fluorescent dye (green) (A) or anti-LAMP2 antibodies (red) (B) with DAPI to visualize nucleus (blue). (C) Numbers of LAMP2-positive lysosomes. To count LAMP2-positive lysosomes 9 osteoclasts were observed (left) and the lysosome number was normalized to number of nuclei within osteoclast (right). Experiment was performed three times, each time using a different cohort of WT and YF mice. A representative experiment is shown. Values are mean ± SD, *Significance different between WT and YF, $p < 0.05$. Scale bars represent 100 μm and 200 μm for (A) and (B) respectively. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

osteoclasts. Conditioned media from BMMs and mature osteoclast cultures was harvested. WT calvarial osteoblasts were grown in the presence of conditioned media derived from BMMs or osteoclasts of WT or YF mice, and then stained for the formation of CFUs and ALP-stained colonies. We found that cells treated with the YF osteoclast conditioned media had more CFUs and ALP-positive colonies, and expressed increased osteogenic markers than those treated with WT osteoclast conditioned media or either WT or YF BMM conditioned media (Fig. 5 and data not shown for BMM conditioned medium). In osteoclasts

derived from *Ctsk*^{-/-} mice, mRNA expression of sphingosine kinase (SphK1) is upregulated [34]. SphK1 catalyzes the formation of sphingosine-1-phosphate (S1P), and S1P secreted by osteoclasts promotes osteogenesis [34,35]. To determine osteogenic factors generated by osteoclasts, cells were cultured on HA-coated plates and harvested for qRT-PCR analysis. We found that RNA expression for Wnt10B, BMP6, PDGF-BB, and Sphk1 and 2 were comparable between WT and YF cells (Table 2).




Fig. 4. Cathepsin K secretion is perturbed in YF osteoclasts. (A) mRNA expression of *Acp5* and *Ctsk* was examined by qRT-PCR and normalized by mRNA level of *Gapdh*. Values are mean ± SD, *n* = 3, *Significant difference between WT and YF, *p* < 0.05. (B) Equal numbers of purified mature osteoclasts (OC) seeded on tissue culture plastic were stained for TRAP. Scale bar 200 μM (C) Equal numbers of mature osteoclasts seeded in 24-well plates were cultured in medium containing 10% FBS overnight. Total cell lysate (TCL) (50 μg) and conditioned media (CM) (25 μl) were electrophoresed on 10 or 12% SDS-PAGE gels. Western blots were probed with anti-Cathepsin K (CatK) (left) and anti-Cathepsin D (CatD) (right) antibodies. Only the mature CatK form is shown. Blots were stripped and reprobbed with anti-β-Actin antibodies to verify equal loading of TCL. (D) Signal intensity of CatK protein bands in technical duplicates was quantified using LI-COR (Odyssey) system. Experiment was performed three times, each time using a different cohort of WT and YF mice. A representative experiment is shown.

4. Discussion

In osteoclasts, levels of Cathepsin K far exceed that of other proteases [36–38]. It is unique among the mammalian proteases in its ability to cleave both telopeptides and helical region of collagen type I [39]. In humans, mutations in *CTSK* cause pycnodysostosis [37,40]. In *Ctsk*^{-/-} mice, osteoclasts differentiate normally, but degrade the collagenous matrix poorly [39,41]. However, the signaling pathways that regulate Cathepsin K secretion are not completely understood.

In this report, we have demonstrated that abrogation of the Cbl-PI3K interaction in osteoclasts results in decreased lysosome numbers and mis-localization of Cathepsin K, contributing to decreased Cathepsin K secretion, thereby resulting in decreased pit formation. Towards this, our previous work shows increased bone volume and decreased serum level of carboxyterminal collagen crosslinks (CTX)

during homeostasis and upon ovariectomy in YF mice [16,18].

In osteoclasts, a family of v-ATPases function to acidify lysosomes by coupling ATP hydrolysis to proton transport. To maintain electron neutrality, a parallel transport of chloride ions into the lacuna takes place simultaneously. We found that the localization of v-ATPase was also unperturbed in the absence of Cbl-PI3K interaction. While Cathepsin K and v-ATPase are of lysosomal origin, our results suggest that PI3K signaling is preferentially involved in trafficking and secretion of Cathepsin K. These findings suggest that vesicle cargo may be a determining factor in how secretory vesicles are transported to the plasma membrane for exocytosis.

YF osteoclasts are defective in secreting Cathepsin K, but secretion of Cathepsin D is normal. These data suggest that in YF osteoclasts, the cargo present in the secretory lysosomes is not secreted properly, while other secretory mechanisms are not perturbed. The Protein kinase C-δ




Fig. 5. Conditioned media from YF osteoclasts induces osteogenesis. WT calvarial-derived osteoblasts (10^5 cells/well) in 6-well plates were grown in triplicates in conditioned media derived from mature osteoclast cultures. After 7 days, colonies were fixed and stained with toluidine blue to stain CFU fibroblasts (A) or with ALP to stain for CFU-OB (B). Groups of 20 or more cells were counted as a colony. Significant difference between WT and YF, * $p < 0.05$, *** $p < 0.001$. (C) Colonies were harvested after 7 days and expression of osteogenic markers was performed by qRT-PCR and normalized by mRNA level of *Gapdh*. Values are mean \pm SD, and plotted as relative expression to WT. A representation of 2 experiments is shown. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

deficiency in osteoclasts does not affect v-ATPase localization, but attenuates Cathepsin K secretion [42]. Differential secretion and targeting of vesicles is supported by other studies. Disruption of the Mannose-6-phosphate pathway in osteoclasts augmented Cathepsin K secretion without affecting Cathepsin D targeting and secretion [30]. Our results and other findings suggest the existence of distinct/different pools of secretory lysosomes in osteoclasts, each relying on specific signaling cues essential for their exocytosis. This flexibility will allow cells to

secrete cargo based on need.

AKT, a serine/threonine kinase, is the major effector of PI3K in several cell types, including bone cells. We reported that in YF mice, AKT is constitutively phosphorylated, and there is increased phosphorylation of AKT substrates in osteoclasts [12,16]. AKT substrates that are postulated to regulate vesicle include the Rab GTPase-activating proteins AS160/TBC1D4 [43] and TBC1D1 [44]; AKT binding protein ClipR-59 [45]; and CENTB1/ACAP1 and CENTG1 in the

Table 2
Expression analysis of osteogenic factors generated by WT and YF osteoclasts.

	WT	YF
<i>Pdgfbb</i>	1.08 ± 0.150	0.972 ± 0.414
<i>BMP6</i>	1.06 ± 0.138	1.602 ± 0.039
<i>Sema4d</i>	1.017 ± 0.216	1.220 ± 0.174
<i>Sphk-1Sphk1</i>	1.27 ± 0.016	0.945 ± 0.204
<i>Sphk2</i>	0.956 ± 0.03	0.987 ± 0.131
<i>Wnt10b</i>	1.00 ± 0.012	0.823 ± 0.041

centaurin family of GTPases [46,47]. We have also reported that activation of small GTPase is perturbed in YF osteoclasts. Given these findings, we speculate that in YF osteoclasts, decreased lysosome numbers and perturbed AKT activation could impact targeting of Cathepsin K to secretory vesicles, and eventually, bone resorption.

In addition to AKT, other Cbl-related pathways could also regulate vesicular trafficking. A subset of autophagy related proteins were reported to regulate lysosomal secretion in osteoclasts [48]. Interestingly, of the autophagy proteins, microtubule associated protein 1 light chain 3 (LC3) directly binds Cbl via a LC3 interacting motif (WxxL) present in the C-terminal end of Cbl [49]. While binding of mutated Cbl and other signaling proteins such as Src and Grb2 was not affected [16], whether disruption in LC3 binding to mutated Cbl protein, and its impact on Cathepsin K secretion, remain to be tested.

Independent of their resorptive activity, osteoclasts also regulate bone formation. This is especially the case in non-resorptive osteoclasts [24,50]. *Ctsk*^{-/-} mice have increased osteoclast numbers and bone formation rates [37,41,51], and *in vitro* data indicate that osteoclast-specific deletion of *Ctsk* stimulates bone formation [34]. Like *Ctsk*^{-/-} mice, global YF mice also exhibit increased numbers of dysfunctional osteoclasts and an increased bone formation rate [12,16,33]. Additionally, as demonstrated here, Cathepsin K secretion was perturbed in YF osteoclasts. Therefore, we investigated whether YF osteoclasts can influence osteoblast proliferation and differentiation. Our findings show that conditioned media derived from YF osteoclasts enhances osteogenesis. We found that SphK-1 and 2 expression levels were similar between WT and YF samples. mRNA expression levels of other known factors secreted by osteoclasts that modulate osteogenesis were also comparable between WT and YF osteoclasts [24,32,52]. These data suggest that the *Ctsk*^{-/-} osteoclasts and dysfunctional YF osteoclasts secrete different factors that promote osteogenesis. We propose that Cbl-PI3K signaling regulates osteoclast-derived factors that promote osteogenesis, however, experiments to identify the factor(s) are warranted.

The *in vivo* significance of our findings highlights that the selective disruption of Cathepsin K secretion has important implication on bone health. YF mice have increased bone mass under homeostatic conditions [16]. Additionally, YF mice are protected from experimental postmenopausal osteoporosis [18]. Osteoclasts are also responsible for cancer progression [53] and there is a strong correlation between Cathepsin K expression and metastatic tumors [54,55]. > 30 inhibitors of PI3Ks and AKT have been developed for anticancer therapy, and some of them have entered clinical trials [56–58]. Our results suggest that the selective disruption of the Cbl-PI3K interaction can be a strategy to prevent pathological bone loss in several clinical settings, including cancer.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bone.2019.07.009>.

Author contributions

J.Y., N.S.A. and L.D. performed experiments, and analyzed the data. A.S. designed the study, analyzed data, and wrote the manuscript. All authors have reviewed and approved the final submission.

Disclosure

The authors have nothing to disclose.

Acknowledgments

The authors thank D. Fremont for M-CSF cDNA and M. Glogauer for *Trnfsf11* cDNA, B. Lee for the anti vATPase (E subunit) antibody. National Institute of Health Grant (AR0550601) to A.S supported part of the work. Study was supported by United States National Institute of Health grant AR055601 (AS).

References

- [1] S.L. Teitelbaum, Bone resorption by osteoclasts, *Science* 289 (2000) 1504–1508.
- [2] O. Kollet, A. Dar, S. Shvitiel, A. Kalinkovich, K. Lapid, Y. Sztainberg, M. Tesio, R.M. Samstein, P. Goichberg, A. Spiegel, A. Elson, T. Lapidot, Osteoclasts degrade endosteal components and promote mobilization of hematopoietic progenitor cells, *Nat. Med.* 12 (2006) 657–664.
- [3] A. Mansour, A. Wakkach, C. Blin-Wakkach, Role of osteoclasts in the hematopoietic stem cell niche formation, *Cell Cycle* 11 (2012) 2045–2046.
- [4] L.C. Cantley, The phosphoinositide 3-kinase pathway, *Science* 296 (2002) 1655–1657.
- [5] J.A. Engelman, J. Luo, L.C. Cantley, The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism, *Nat. Rev. Genet.* 7 (2006) 606–619.
- [6] L.C. Foukas, K. Okkenhaug, Gene-targeting reveals physiological roles and complex regulation of the phosphoinositide 3-kinases, *Arch. Biochem. Biophys.* 414 (2003) 13–18.
- [7] G. Swaminathan, A.Y. Tsygankov, The Cbl family proteins: ring leaders in regulation of cell signaling, *J. Cell. Physiol.* 209 (2006) 21–43.
- [8] F. Meng, C.A. Lowell, A β_1 integrin signaling pathway involving Src-family kinases, Cbl and PI-3 kinase is required for macrophage spreading and migration, *EMBO J.* 17 (1998) 4391–4403.
- [9] R. Chiusaroli, A. Sanjay, K. Henriksen, M.T. Engsig, W.C. Horne, H. Gu, R. Baron, Deletion of the gene encoding c-Cbl alters the ability of osteoclasts to migrate, delaying resorption and ossification of cartilage during the development of long bones, *Dev. Biol.* 261 (2003) 537–547.
- [10] T. Miyazaki, A. Sanjay, L. Neff, S. Tanaka, W.C. Horne, R. Baron, Src kinase activity is essential for osteoclast function, *J. Biol. Chem.* 279 (2004) 17660–17666.
- [11] A. Sanjay, A. Houghton, L. Neff, E. DiDomenico, C. Bardelay, E. Antoine, J. Levy, J. Gailit, D. Bowtell, W.C. Horne, R. Baron, Cbl associates with Pyk2 and Src to regulate Src kinase activity, $\alpha(v)\beta(3)$ integrin-mediated signaling, cell adhesion, and osteoclast motility, *J. Cell Biol.* 152 (2001) 181–195.
- [12] N.S. Adapala, M.F. Barbe, A.Y. Tsygankov, J.A. Lorenzo, A. Sanjay, Loss of Cbl-PI3K interaction enhances osteoclast survival due to p21-Ras mediated PI3K activation independent of Cbl-b, *J. Cell. Biochem.* 115 (2014) 1277–1289.
- [13] A. Nakajima, A. Sanjay, R. Chiusaroli, N.S. Adapala, L. Neff, C. Itzstein, W.C. Horne, R. Baron, Loss of Cbl-b increases osteoclast bone-resorbing activity and induces osteopenia, *Journal of Bone and Mineral Research: The Official Journal of the American Society for Bone and Mineral Research* 24 (2009) 1162–1172.
- [14] Z. Songyang, L.C. Cantley, Recognition and specificity in protein tyrosine kinase-mediated signaling, *Trends Biochem. Sci.* 20 (1995) 470–475.
- [15] G. DeBlasti, A. Sanjay, T. Miyazaki, W.C. Horne, R. Baron, C-Cbl, but not Cbl-b, requires Src phosphorylation to interact with hte p85 subunit of PI3kinase, *J. Bone Miner. Res.* 17 (Suppl. 1) (2002) S350.
- [16] N.S. Adapala, M.F. Barbe, W.Y. Langdon, M.C. Nakamura, A.Y. Tsygankov, A. Sanjay, The loss of Cbl-phosphatidylinositol 3-kinase interaction perturbs RANKL-mediated signaling, inhibiting bone resorption and promoting osteoclast survival, *J. Biol. Chem.* 285 (2010) 36745–36758.
- [17] J.C. Molero, N. Turner, C.B. Thien, W.Y. Langdon, D.E. James, G.J. Cooney, Genetic ablation of the c-Cbl ubiquitin ligase domain results in increased energy expenditure and improved insulin action, *Diabetes* 55 (2006) 3411–3417.
- [18] N.S. Adapala, D. Holland, V. Scanlon, M.F. Barbe, W.Y. Langdon, A.Y. Tsygankov, J.A. Lorenzo, A. Sanjay, Loss of Cbl-PI3K interaction in mice prevents significant bone loss following ovariectomy, *Bone* 67C (2014) 1–9.
- [19] S.H. Lee, J. Rho, D. Jeong, J.Y. Sul, T. Kim, N. Kim, J.S. Kang, T. Miyamoto, T. Suda, S.K. Lee, R.J. Pignolo, B. Koczon-Jaremko, J. Lorenzo, Y. Choi, V-ATPase V0 subunit d2-deficient mice exhibit impaired osteoclast fusion and increased bone formation, *Nat. Med.* 12 (2006) 1403–1409.
- [20] Y. Wang, D. Lebowitz, C. Sun, H. Thang, M.D. Grynepas, M. Glogauer, Identifying the relative contributions of Rac1 and Rac2 to osteoclastogenesis, *J. Bone Miner. Res.* 23 (2008) 260–270.
- [21] N.S. Adapala, Loss of Cbl-PI3K interaction enhances osteoclast survival due to p21-Ras mediated PI3K activation independent of Cbl-b, *J. Cell. Biochem.* 1 (2014) 1–13.
- [22] R. Baron, L. Neff, W. Brown, P.J. Courtroy, D. Louvard, M.G. Farquhar, Polarized secretion of lysosomal enzymes: co-distribution of cation-independent mannose-6-phosphate receptors and lysosomal enzymes along the osteoclast exocytic pathway, *J. Cell Biol.* 106 (1988) 1863–1872.
- [23] S.L. Teitelbaum, The osteoclast and its unique cytoskeleton, *Ann. N. Y. Acad. Sci.* 1240 (2011) 14–17.

- [24] M.A. Karsdal, T.J. Martin, J. Bollerslev, C. Christiansen, K. Henriksen, Are non-resorbing osteoclasts sources of bone anabolic activity? *J. Bone Miner. Res.* 22 (2007) 487–494.
- [25] H. Zhao, Membrane trafficking in osteoblasts and osteoclasts: new avenues for understanding and treating skeletal diseases, *Traffic* 13 (2012) 1307–1314.
- [26] J. Lacombe, G. Karsenty, M. Ferron, Regulation of lysosome biogenesis and functions in osteoclasts, *Cell Cycle* 12 (2013) 2744–2752.
- [27] M.J. Bossard, T.A. Tomaszek, S.K. Thompson, B.Y. Amegadzie, C.R. Hanning, C. Jones, J.T. Kurdyla, D.E. McNulty, F.H. Drake, M. Gowen, M.A. Levy, Proteolytic activity of human osteoclast cathepsin K. Expression, purification, activation, and substrate identification, *J. Biol. Chem.* 271 (1996) 12517–12524.
- [28] M.S. McQueney, B.Y. Amegadzie, K. D'Alessio, C.R. Hanning, M.M. McLaughlin, D. McNulty, S.A. Carr, C. Ijames, J. Kurdyla, C.S. Jones, Autocatalytic activation of human cathepsin K, *J. Biol. Chem.* 272 (1997) 13955–13960.
- [29] O.J. Holt, F. Gallo, G.M. Griffiths, Regulating secretory lysosomes, *J. Biochem.* 140 (2006) 7–12.
- [30] E. van Meel, M. Boonen, H. Zhao, V. Oorschot, F.P. Ross, S. Kornfeld, J. Klumperman, Disruption of the Man-6-P targeting pathway in mice impairs osteoclast secretory lysosome biogenesis, *Traffic* 12 (2011) 912–924.
- [31] T. Goto, T. Kiyoshima, R. Moroi, T. Tsukuba, Y. Nishimura, M. Himeno, K. Yamamoto, T. Tanaka, Localization of cathepsins B, D, and L in the rat osteoclast by immuno-light and -electron microscopy, *Histochemistry* 101 (1994) 33–40.
- [32] M.A. Karsdal, A.V. Neutzsky-Wulff, M.H. Dziegiel, C. Christiansen, K. Henriksen, Osteoclasts secrete non-bone derived signals that induce bone formation, *Biochem. Biophys. Res. Commun.* 366 (2008) 483–488.
- [33] T. Brennan, N.S. Adapala, M.F. Barbe, V. Yingling, A. Sanjay, Abrogation of Cbl-PI3K interaction increases bone formation and osteoblast proliferation, *Calcif. Tissue Int.* 89 (2011) 396–410.
- [34] S. Lotinun, R. Kiviranta, T. Matsubara, J.A. Alzate, L. Neff, A. Luth, I. Koskivirta, B. Kleuser, J. Vacher, E. Vuorio, W.C. Horne, R. Baron, Osteoclast-specific cathepsin K deletion stimulates S1P-dependent bone formation, *J. Clin. Invest.* 123 (2013) 666–681.
- [35] L. Pederson, M. Ruan, J.J. Westendorf, S. Khosla, M.J. Oursler, Regulation of bone formation by osteoclasts involves Wnt/BMP signaling and the chemokine sphingosine-1-phosphate, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 20764–20769.
- [36] F.H. Drake, R.A. Dodds, I.E. James, J.R. Connor, C. Debouck, S. Richardson, E. Lee-Rykaczewski, L. Coleman, D. Rieman, R. Barthlow, G. Hastings, M. Gowen, Cathepsin K, but not cathepsins B, L, or S, is abundantly expressed in human osteoclasts, *J. Biol. Chem.* 271 (1996) 12511–12516.
- [37] V. Everts, D.C. Aronson, W. Beertsen, Phagocytosis of bone collagen by osteoclasts in two cases of pycnodysostosis, *Calcif. Tissue Int.* 37 (1985) 25–31.
- [38] V. Everts, W. Korper, K.A. Hoeben, I.D. Jansen, D. Bromme, K.B. Cleutjens, S. Heeneman, C. Peters, T. Reinheckel, P. Saftig, W. Beertsen, Osteoclastic bone degradation and the role of different cysteine proteinases and matrix metalloproteinases: differences between calvaria and long bone, *J. Bone Miner. Res.* 21 (2006) 1399–1408.
- [39] P. Saftig, E. Hunziker, O. Wehmeyer, S. Jones, A. Boyde, W. Rommerskirch, J.D. Moritz, P. Schu, K. von Figura, Impaired osteoclastic bone resorption leads to osteopetrosis in cathepsin-K-deficient mice, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 13453–13458.
- [40] B.D. Gelb, G.-P. Shi, H.A. Chapman, R.J. Desnick, Pycnodysostosis, a lysosomal disease caused by cathepsin K deficiency, *Science* 273 (1996) 1236–1238.
- [41] C.Y. Li, K.J. Jepsen, R.J. Majeska, J. Zhang, R. Ni, B.D. Gelb, M.B. Schaffler, Mice lacking cathepsin K maintain bone remodeling but develop bone fragility despite high bone mass, *J. Bone Miner. Res.* 21 (2006) 865–875.
- [42] V. Cremasco, C.E. Decker, D. Stumpo, P.J. Blackshear, K.I. Nakayama, K. Nakayama, T.S. Lupu, D.B. Graham, D.V. Novack, R. Faccio, Protein kinase C-delta deficiency perturbs bone homeostasis by selective uncoupling of cathepsin K secretion and ruffled border formation in osteoclasts, *J. Bone Miner. Res.* 27 (2012) 2452–2463.
- [43] J.C. Hou, J.E. Pessin, Ins (endocytosis) and outs (exocytosis) of GLUT4 trafficking, *Curr. Opin. Cell Biol.* 19 (2007) 466–473.
- [44] S. Chen, J. Murphy, R. Toth, D.G. Campbell, N.A. Morrice, C. Mackintosh, Complementary regulation of TBC1D1 and AS160 by growth factors, insulin and AMPK activators, *Biochem. J.* 409 (2008) 449–459.
- [45] W. Ren, S. Cheema, K. Du, The association of ClipR-59 protein with AS160 modulates AS160 protein phosphorylation and adipocyte Glut4 protein membrane translocation, *J. Biol. Chem.* 287 (2012) 26890–26900.
- [46] T.R. Jackson, B.G. Kearns, A.B. Theibert, Cytohesins and centaurins: mediators of PI 3-kinase-regulated Arf signaling, *Trends Biochem. Sci.* 25 (2000) 489–495.
- [47] M. Soundararajan, X. Yang, J.M. Elkins, F. Sobott, D.A. Doyle, The centaurin gamma-1 GTPase-like domain functions as an NTPase, *Biochem. J.* 401 (2007) 679–688.
- [48] C.J. DeSelm, B.C. Miller, W. Zou, W.L. Beatty, E. van Meel, Y. Takahata, J. Klumperman, S.A. Tooze, S.L. Teitelbaum, H.W. Virgin, Autophagy proteins regulate the secretory component of osteoclastic bone resorption, *Dev. Cell* 21 (2011) 966–974.
- [49] E. Sandilands, B. Serrels, D.G. McEwan, J.P. Morton, J.P. Macagno, K. McLeod, C. Stevens, V.G. Brunton, W.Y. Langdon, M. Vidal, O.J. Sansom, I. Dikic, S. Wilkinson, M.C. Frame, Autophagic targeting of Src promotes cancer cell survival following reduced FAK signalling, *Nat. Cell Biol.* 14 (2011) 51–60.
- [50] T. Segovia-Silvestre, A.V. Neutzsky-Wulff, M.G. Sorensen, C. Christiansen, D. Kimmel, M.A. Karsdal, K. Henriksen, Advances in osteoclast biology resulting from the study of osteopetrotic mutations, *Hum. Genet.* 124 (2009) 561–577.
- [51] B. Pennypacker, M. Shea, Q. Liu, P. Masarachia, P. Saftig, S. Rodan, G. Rodan, D. Kimmel, Bone density, strength, and formation in adult cathepsin K (–/–) mice, *Bone* 44 (2009) 199–207.
- [52] T.J. Martin, N.A. Sims, Osteoclast-derived activity in the coupling of bone formation to resorption, *Trends Mol. Med.* 11 (2005) 76–81.
- [53] K.N. Weilbaecher, T.A. Guise, L.K. McCauley, Cancer to bone: a fatal attraction, *Nat. Rev. Cancer* 11 (2011) 411–425.
- [54] K.D. Brubaker, R.L. Vessella, L.D. True, R. Thomas, E. Corey, Cathepsin K mRNA and protein expression in prostate cancer progression, *J. Bone Miner. Res.* 18 (2003) 222–230.
- [55] A.J. Littlewood-Evans, G. Bilbe, W.B. Bowler, D. Farley, B. Wlodarski, T. Kokubo, T. Inaoka, J. Sloane, D.B. Evans, J.A. Gallagher, The osteoclast-associated protease cathepsin K is expressed in human breast carcinoma, *Cancer Res.* 57 (1997) 5386–5390.
- [56] S. Buonamici, J. Williams, M. Morrissey, A. Wang, R. Guo, A. Vattay, K. Hsiao, J. Yuan, J. Green, B. Ospina, Q. Yu, L. Ostrom, P. Fordjour, D.L. Anderson, J.E. Monahan, J.F. Kelleher, S. Peukert, S. Pan, X. Wu, S.M. Maira, C. Garcia-Echeverria, K.J. Briggs, D.N. Watkins, Y.M. Yao, C. Lengauer, M. Warmuth, W.R. Sellers, M. Dorsch, Interfering with resistance to smoothened antagonists by inhibition of the PI3K pathway in medulloblastoma, *Sci. Transl. Med.* 2 (2010) 51ra70.
- [57] A.J. Folkes, K. Ahmadi, W.K. Alderton, S. Alix, S.J. Baker, G. Box, I.S. Chuckowree, P.A. Clarke, P. Depledge, S.A. Eccles, L.S. Friedman, A. Hayes, T.C. Hancox, A. Kugendradas, L. Lensun, P. Moore, A.G. Olivero, J. Pang, S. Patel, G.H. Pergl-Wilson, F.I. Raynaud, A. Robson, N. Saghiri, L. Salphati, S. Sohal, M.H. Ultsch, M. Valenti, H.J. Wallweber, N.C. Wan, C. Wiesmann, P. Workman, A. Zhyvoloup, M.J. Zvelebil, S.J. Shuttleworth, The identification of 2-(1H-indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-t hieno[3,2-d]pyrimidine (GDC-0941) as a potent, selective, orally bioavailable inhibitor of class I PI3 kinase for the treatment of cancer, *J. Med. Chem.* 51 (2008) 5522–5532.
- [58] S.M. Maira, F. Stauffer, J. Brueggen, P. Furet, C. Schnell, C. Fritsch, S. Brachmann, P. Chene, A. De Pover, K. Schoemaker, D. Fabbro, D. Gabriel, M. Simonen, L. Murphy, P. Finan, W. Sellers, C. Garcia-Echeverria, Identification and characterization of NVP-BEZ235, a new orally available dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor with potent in vivo antitumor activity, *Mol. Cancer Ther.* 7 (2008) 1851–1863.