



Altered resting state functional connectivity of the cognitive control network in fibromyalgia and the modulation effect of mind-body intervention

Jian Kong¹ · Emily Wolcott² · Zengjian Wang¹ · Kristen Jorgenson¹ · William F. Harvey² · Jing Tao¹ · Ramel Rones¹ · Chenchen Wang²

Published online: 2 May 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

This study examines altered resting state functional connectivity (rsFC) of the cognitive control network (CCN) in fibromyalgia patients as compared to healthy controls, as well as how an effective mind-body intervention, Tai Chi, can modulate the altered rsFC of the CCN. Patients with fibromyalgia and matched healthy subjects were recruited in this study. Fibromyalgia patients were scanned 12 weeks before and after intervention. The bilateral dorsolateral prefrontal cortex (DLPFC) was used as a seed to explore the rsFC of the CCN. Data analysis was conducted with 21 patients and 20 healthy subjects. Compared to healthy subjects, fibromyalgia patients exhibited increased rsFC between the DLPFC and the bilateral rostral anterior cingulate cortex (rACC) and medial prefrontal cortex (MPFC) at baseline. The rsFC between the CCN and rACC/MPFC further increased after Tai Chi intervention, and this increase was accompanied by clinical improvements. This rsFC change was also significantly associated with corresponding changes in the Overall Impact domain of the Revised Fibromyalgia Impact Questionnaire (FIQR). Further analysis showed that the rACC/MPFC rsFC with both the PAG and hippocampus significantly decreased following Tai Chi intervention. Our study suggests that fibromyalgia is associated with altered CCN rsFC and that effective mind-body treatment may elicit clinical improvements by further increasing this altered rsFC. Elucidating this mechanism of enhancing the allostasis process will deepen our understanding of the mechanisms underlying mind-body interventions in fibromyalgia patients and facilitate the development of new pain management methods.

Keywords Resting state functional connectivity · Fibromyalgia · Cognitive control network · Dorsal lateral prefrontal cortex · Tai Chi · Anterior cingulate cortex · Mind-body intervention

Introduction

Fibromyalgia (FM) is a multidimensional complex disorder characterized by chronic widespread musculoskeletal pain and physical and psychological limitations (Sumpton and

Moulin 2014). The neuropathology of fibromyalgia remains unclear, but previous studies have suggested that the central nervous system may play an important role in the development and maintenance of fibromyalgia (Jensen et al. 2012b, 2013b; Tracey and Bushnell 2009).

Recently, functional magnetic resonance imaging (fMRI) has been used to investigate the pathophysiology (Jensen et al. 2012b, 2013a; Loggia et al. 2014; Truini et al. 2016) and treatment of fibromyalgia (Cummiford et al. 2016; Flodin et al. 2015). These studies suggest that patients with fibromyalgia exhibit significant changes in brain function and structure, further endorsing the role of the brain in the neuropathology of fibromyalgia. For instance, studies show that fibromyalgia patients exhibit reductions in cortical thickness, brain volume (Jensen et al. 2013b; Kuchinad et al. 2007; Lutz et al. 2008; Robinson et al. 2011), and measures of functional

✉ Jian Kong
kongj@nmr.mgh.harvard.edu

✉ Chenchen Wang
cwang2@tuftsmedicalcenter.org

¹ Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA, USA

² Center For Complementary And Integrative Medicine, Department of Rheumatology, Tufts Medical Center, Tufts University School of Medicine, Boston, MA 02111, USA

connectivity (Jensen et al. 2013b) in the rostral anterior cingulate cortex (rACC) and medial prefrontal cortex (MPFC) compared to healthy controls, demonstrating the important role of the rACC/MPFC in the development of FM.

Although these studies significantly enhanced our understanding of fibromyalgia, the physiological implications of these brain functional and morphometry changes remain unclear. One possible explanation is that some of these changes may reflect an adaptation process of the brain to fibromyalgia, specifically through allostasis, the process of achieving homeostasis through physiological or behavioral changes. Based on the theory of allostasis, the human body, including the brain, adapts to the altered conditions caused by a disorder, such as persistent pain (Borsook et al. 2012). This can be carried out by means of alteration in the hypothalamus – pituitary – adrenal (HPA) axis hormones and the autonomic nervous system, as well as through brain functional and structure changes (McEwen & Wingfield 2003). For example, to adapt and cope with consistent pain, fibromyalgia patients may employ an involuntary distraction strategy through a cognitive control mechanism. In support of this hypothesis, studies have demonstrated that attention can significantly modulate pain perception (Legrain et al. 2009; Lobanov et al. 2013; Torta et al. 2017); that is, focusing on pain may enhance pain responses, while distraction from pain can significantly reduce one's pain experience (Valet et al. 2004). Further studies suggest that the descending pain modulation system, including the periaqueductal gray (PAG), cingulate cortex, and dorsal lateral prefrontal cortex, may play an important role in the attention regulation of pain (Fields 2004; Fields et al. 2005; Kong et al. 2010).

Cognitive control refers to the set of brain processes necessary for goal-directed thought and action (Cole and Schneider 2007), which plays a crucial role in the top-down modulation of attention–memory interactions (Cole and Schneider 2007; Rosen et al. 2015; Sheline et al. 2010), decision-making, and conflict resolution (Sheline et al. 2010). A key factor in cognitive control processing is attention, a mechanism by which sensory input is selected to enter into awareness and which can regulate one's pain experience (Bushnell et al. 2013; Torta et al. 2017; Villemure and Bushnell 2002; Villemure and Schweinhardt 2010; Wiech 2016; Zeidan et al. 2015; Zeidan et al. 2012). Thus, cognitive control may represent an interesting aspect of allostasis in FM patients.

The brain network most closely related to cognitive executive functioning is the cognitive control network (CCN), which includes the frontal gyrus, parietal gyrus, and anterior cingulate cortex (Petersen and Posner 2012; Zanto and Gazzaley 2013). A large body of evidence indicates that the dorsolateral prefrontal cortex (DLPFC) is a key region of the CCN, playing an important role in cognitive control processes (Cieslik et al. 2013; Legrain et al. 2009; Miller and Cohen 2001; Sheline et al. 2010; Tracey and Mantyh 2007). Previous studies have also suggested that using the bilateral DLPFC as a seed can be a valuable tool for exploring the function of the CCN using

resting state fMRI (Cieslik et al. 2013; Hwang et al. 2015; Legrain et al. 2009; Miller and Cohen 2001; Sheline et al. 2010; Tao et al. 2017b; Tracey and Mantyh 2007).

In addition to investigating CCN differences between fibromyalgia patients and healthy controls, we attempted to explore whether effective mind-body treatment (Tai Chi) can modulate the rsFC in fibromyalgia patients. We believe investigating rsFC changes after effective treatment, as well as the association between these rsFC changes and accompanying clinical symptom reductions, will help us understand the significance of brain functional changes observed in FM patients when compared to healthy controls. We chose Tai Chi because pharmacological treatment of FM only achieves limited success with a significant risk of adverse effects that cannot be tolerated by many patients (Goldenberg et al. 2004; Lautenschlager 2000). Previous randomized controlled trials have shown that Tai Chi, a multicomponent intervention that incorporates physical, psychosocial, emotional, spiritual, and behavioral elements, is a potentially useful therapy for patients with fibromyalgia (Wang et al. 2010). Meta-analyses indicate that Tai Chi can enhance cognitive function in older adults, particularly in the realm of executive functioning (Wayne et al. 2014). Imaging studies have also shown that Tai Chi can significantly modulate the function and structure of brain regions associated with cognitive control (Tao et al. 2017b; Wei et al. 2013, 2014, 2017), further endorsing the role of the CCN in Tai Chi intervention.

In this study, we first compared the resting state functional connectivity (rsFC) of the DLPFC between fibromyalgia patients and healthy subjects to characterize the brain pathophysiology of fibromyalgia, and then investigated how the rsFC of the DLPFC are modulated when symptoms are relieved after longitudinal intervention. We hypothesized that fibromyalgia patients would show increased DLPFC-ACC / MPFC rsFC as a distraction/adaptation mechanism. Mind-body interventions may enhance this coping process by increasing DLPFC-ACC/MPFC connectivity, further triggering the descending pain modulation system to relieve clinical symptoms in fibromyalgia patients. This investigation, which combines both healthy controls and an intervention group, allows us to not only illustrate rsFC changes that are sensitive to symptom improvement or treatment, but also highlights the physiological implications of these changes.

Materials and methods

Participants

From May 2015 to September 2015, we recruited 24 patients with fibromyalgia (≥ 21 years old) and 24 healthy controls matched for age, gender, and body mass index (BMI) to participate in the study. After describing the details of the study to interested and eligible subjects, written informed consent was

obtained in accordance with the procedures established by Tufts Medical Center/Tufts University Human Institutional Review Board and the Medical Ethics Committee of Massachusetts General Hospital. The study rheumatologist (WFH) performed clinical examinations and confirmed that participants met the eligibility criteria. All FM patients participated in 12 weeks of Tai Chi training (NCT02407665).

Inclusion criteria were: (1) age 21 years or older, (2) fulfillment of the American College of Rheumatology (ACR) 1990 classification criteria and the ACR 2010 diagnostic criteria for fibromyalgia, (3) willingness to complete a 12-week, twice-a-week Tai Chi exercise program, (4) willingness to undergo an fMRI scan at baseline and follow-up visits, and (5) fluent in English. Exclusion criteria were: (1) diagnosed with medical conditions that are known to contribute to fibromyalgia symptomatology, such as thyroid disease, inflammatory arthritis, systemic lupus erythematosus, rheumatoid arthritis, myositis, vasculitis, or Sjogren's syndrome, (2) inability to pass the Physical Activity Readiness Questionnaire (PAR-Q), (3) score of less than 24 on the Mini-Mental State Examination; (4) plans to relocate from the region during the trial period; (5) verbal confirmation of pregnancy or planned pregnancy during the study period, (6) enrollment in any other clinical trial in the last 30 days, (7) presence of any contraindications to fMRI scanning, including but not limited to: cardiac pacemaker, metal implants, fear of enclosed spaces, pregnancy, and weight > 300 lbs., (8) prior experience with Tai Chi training, or similar types of complementary and alternative medicine (e.g. Qi Gong or yoga) in the past year, and (9) serious medical conditions limiting ability to participate in the Tai Chi training, including dementia, neurological disease, cancer, cardiovascular disease, pulmonary disease, metabolic disease, renal disease, liver disease, or other serious medical conditions, as determined by the study physicians.

Healthy controls were recruited and matched to enrolled fibromyalgia participants. Inclusion criteria were (1) age matched within 5 years (must be ≥ 21 years old), (2) gender and race matched, (3) BMI within 5 kg/m² (must be <300 lbs.), and (4) willingness to attend a single evaluation. Exclusion criteria were (1) prior experience with Tai Chi training or similar types of complementary and alternative medicine, (2) chronic or acute pain (e.g. fibromyalgia, osteoarthritis), and (3) presence of any contraindications to fMRI scanning, including but not limited to: cardiac pacemaker, metal implants, fear of closed spaces, pregnancy, and weight > 300 lbs.

Intervention

All participants in the Tai Chi group attended a 60-min practice session twice a week for 12 weeks at Tufts Medical Center. The instructor (RR), who has extensive experience conducting Tai Chi training programs, followed

a standardized Tai Chi protocol developed for patients with fibromyalgia (Wang et al. 2010). Participants were also provided with printed materials on fibromyalgia and the Tai Chi mind-body program, including Tai Chi principles, practicing techniques, and safety precautions specifically for participants with fibromyalgia. Every session included the following components: (1) warm-up, (2) Tai Chi movement, (3) breathing techniques, and (4) relaxation. Each component of the program was derived from the classical Yang style Tai Chi 108 posture condensed for the 12-week intervention program.

All subjects were encouraged to maintain their usual physical activities and to perform no new additional strength training other than their Tai Chi exercises. Subjects were also allowed to continue taking regular medications and maintain routine visits to their physicians throughout the course of the study. Participants were also instructed to practice at least 30 min a day at home and to complete daily logs indicating the amount of time that they practiced Tai Chi exercises. Data on class attendance and home practice was recorded and verified using standard case report forms.

Outcome measurements

The primary outcome for the study was the resting state functional connectivity (rsFC) of the bilateral DLPFC (a key region of the cognitive control network). The secondary outcomes were: 1) Revised Fibromyalgia Impact Questionnaire (FIQR) including the three domains, i.e., Function, Overall Impact, and Symptom Severity, and 2) Beck Depression Inventory (BDI-II). All outcome measurements were collected at baseline and after 12 weeks of the intervention for the fibromyalgia cohort and at baseline for the healthy subjects.

MRI data acquisition

fMRI scans were performed at Massachusetts General Hospital at the Martinos Center for Biomedical Imaging. Each fibromyalgia subject participated in an identical fMRI scan before and after 12 weeks of Tai Chi exercise, whereas healthy controls were only scanned at baseline. The fMRI brain imaging acquisition was conducted on a 3.0 Tesla Siemens (Skyra syngo) scanner with a 32-channel head coil. Magnetization-prepared rapid gradient echo (MPRAGE) T1-weighted images were collected (flip angle = 7 degree, voxel size 1.0 × 1.0 × 1.0 mm³). The BOLD resting state functional images were obtained with echo-planar imaging (TR = 3000 ms, TE = 30 ms, flip angle = 85 degrees, slice thickness = 2.6 mm, acquisition matrix = 64 × 64, voxel size = 2.6 × 2.6 × 2.6 mm³, 44 axial slices, scan time 8 min 21 s). All patients were required to keep their eyes open during the resting state fMRI scan.

Statistical analysis

Clinical data analysis

Statistical analysis was performed using SPSS 19.0 Software (SPSS Inc., Chicago, IL, USA). A threshold of $p < 0.05$ (2-tailed) was applied. One-way ANOVA and Chi square tests were conducted to compare baseline characteristics of the participants between groups. There were no significant differences in age and gender between the fibromyalgia and matched healthy control groups.

Seed based functional connectivity analysis

Functional BOLD data were preprocessed using SPM 12 (Statistical Parametric Mapping, Wellcome Department of Cognitive Neurology, London, UK; implemented by MATLAB R2012b, Math Works, Inc., Natick, MA, USA). During the preprocessing, images were realigned, segmented, and co-registered to each subject's high-resolution T1 scan, which was used to normalize to the standard Montreal Neurological Institute (MNI) template. Images were also smoothed using an 8 mm full-width at half-maximum (FWHM) Gaussian kernel and filtered with a frequency window of 0.008–0.09 Hz. In addition to these steps, we employed segmentation of gray matter, white matter, and cerebrospinal fluid (CSF) areas for the removal of temporal confounding factors (white matter and CSF) (Whitfield-Gabrieli and Nieto-Castanon 2012). Data were then submitted to motion correction using the artifact detection toolbox (http://www.nitrc.org/projects/artifact_detect/). For each subject, we treated images as outliers if the composite movement from a preceding image exceeded 0.5 mm or if the global mean intensity was greater than 3 standard deviations from the mean image intensity for the entire resting scan. Outliers were included as regressors in the first-level general linear model along with other six regular motion parameters (Redcay et al. 2013).

Resting state functional connectivity analysis was conducted using the CONN toolbox v15.g (Whitfield-Gabrieli and Nieto-Castanon 2012) (<http://www.nitrc.org/projects/conn>). We used an a priori DLPFC seed (peak coordinate: $\pm 36, 27, 29$, with 5 mm radius), which has been used in previous studies (Hwang et al. 2015; Sheline et al. 2010; Tao et al. 2017b). Functional connectivity measures were computed between the seed and every other voxel in the brain. First-level correlation maps were produced by extracting the residual BOLD time course from the DLPFC and by computing Pearson's correlation coefficients between that time course and the time courses of all other voxels in the brain. Correlation coefficients were transformed into Fisher's 'Z' scores to increase normality and allow for improved second-level general linear model analyses.

The baseline DLPFC rsFC of fibromyalgia patients and healthy control subjects were compared using a two-sample t-test. The Tai Chi practice effect (post-practice minus pre-practice) on fibromyalgia patients was compared using a paired t-test. Additionally, we also compared the DLPFC rsFC of fibromyalgia patients after practicing Tai Chi to healthy controls using a two-sample t-test. Age, gender, and BDI scores were included as covariates. A threshold of voxel-wise $p < 0.005$ (uncorrected) and cluster-level $p < 0.05$ (family-wise error correction) were applied for data analyses. Given the important role of the periaqueductal grey in pain modulation, we defined the PAG as a region of interest and used small volume correction to correct the p value at a level of $p < 0.05$. Similar to previous studies (Eippert et al. 2009; Kong et al. 2013), correction was based on peak coordinates (x, y, z: 1, -25, -12) with a 4 mm radius obtained from a previous PAG meta-analysis (Linnman et al. 2012).

Results

The study was completed with 21 fibromyalgia patients and 20 healthy controls. One fibromyalgia patient was excluded from the rsFC analysis due to excessive head movement during scanning. Additionally, one healthy control was excluded due to brain atrophy.

The mean age of fibromyalgia subjects was 53.10 ± 11.58 (mean \pm SD) and 52.90 ± 11.12 for control subjects (Table 1). There were no significant differences in age, gender, race, and BMI between the fibromyalgia and healthy control groups. FIQR scores demonstrated moderate to severe fibromyalgia in the majority of the fibromyalgia subjects with an average score of 45.1 ± 18.6 . BDI-II scores revealed moderate depression with an average score of 19.71 ± 11.12 in the fibromyalgia group, and a two-sample t-test showed significant differences between the fibromyalgia and healthy control groups in BDI-II scores (2.75 ± 3.77) ($p < 0.0001$). Paired t-tests showed significant pre- and post-Tai Chi differences in general FIQR scores (mean \pm SD, pre: 45.1 ± 18.6 , post: 35.8 ± 21.4 , $p = 0.003$), as well as in the three FIQR domains: Function (pre: 12.1 ± 6.3 , post: 8.5 ± 6.4 , $p = 0.001$), Overall Impact (pre: 8.8 ± 6.4 , post: 7.0 ± 5.7 , $p = 0.05$), and Symptoms (pre: 25.1 ± 8.4 , post: 20.8 ± 12 , $p = 0.017$). Analysis of BDI-II scores showed a significant difference between pre- and post-treatment scores in the fibromyalgia patients (pre: 19.71 ± 11.12 , post: 9.95 ± 8.55 , $p = 0.0027$).

Functional connectivity results

rsFC analysis indicated that fibromyalgia patients showed significantly greater rsFC between the DLPFC and both the bilateral rostral anterior cingulate cortex (rACC) and medial prefrontal cortex (MPFC) compared to healthy

Table 1 Baseline characteristics and health outcome variables of study participants, all values are mean (SD) unless otherwise noted

Variable	Tai Chi (<i>n</i> = 21)	Healthy controls (<i>n</i> = 20)
Female, no. (%)	20 (95.24)	19 (95.00)
Age, yr.	53.10 (11.58)	52.9 (11.12)
Race, no. (%)		
White	13 (61.90)	12 (60.00)
Black	6 (28.57)	6 (30.00)
Asian	2 (9.52)	2 (10.00)
Body mass index, kg/m ²	29.37 (6.96)	27.28 (3.58)
Beck depression inventory-II	19.71 (11.12)	2.75 (3.77)

controls at baseline. There were no significant DLPFC rsFC decreases in fibromyalgia patients compared to healthy subjects at baseline. The comparison between healthy controls and fibromyalgia patients after intervention showed significant CCN rsFC increases at the bilateral rACC MPFC and temporal pole and rsFC decreases at the left superior parietal lobule and postcentral gyrus in fibromyalgia patients (Table 2).

A direct comparison between the DLPFC before and after intervention in FM patients indicates a significant rsFC increase with the bilateral rACC/MPFC, frontal gyrus, precuneus, occipital gyrus, cerebellum; left temporal lobe,

angular/supramarginal gyrus; and right operculum and precentral gyrus and a significant rsFC decrease at the brain stem after Tai Chi.

To explore the association between the CCN rsFC changes after intervention and the corresponding Overall Impact domain score changes (a domain reflecting the cognitive aspect of the FIQR), we extracted the Fisher *z* value of the rACC/MPFC cluster and applied a regression analysis, including age, gender, and BDI-II score as covariates. We found a significant positive association between CCN and rACC/MPFC rsFC changes and corresponding Overall Impact domain score changes ($p = 0.01$) (Fig. 1).

Table 2 Regions with significantly different connectivity between the DLPFC and other brain regions in fibromyalgia patients (FM) and healthy controls before and after treatment

Conditions	Brain regions	mm ³	Cluster centroid (MNI)			
			x	y	z	Z
FM (pre) > HC	Bilateral rACC/ MPFC	540	18	48	6	3.91
HC > FM (pre)	NA					
FM (post) > FM (pre)	Bilateral rACC/MPFC	312	-10	58	-4	3.89
	Bilateral middle frontal gyrus /superior frontal gyrus	1380	-16	28	38	5.61
	Left middle frontal gyrus	768	-36	26	28	4.68
	Left middle frontal /medial frontal gyrus	255	-20	42	18	4.59
	Right operculum	246	46	24	-2	4.22
	Left angular / supramarginal gyrus	235	-54	-58	22	4.27
	Right precentral gyrus	211	46	-8	40	4.71
	Left temporal lobe	202	-50	12	-30	3.96
	Bilateral middle occipital gyrus	1561	34	-72	22	5.57
	Bilateral precuneus	804	28	-72	-4	5.18
	Right cerebellum	378	40	-48	-40	4.92
	Left cerebellum	308	-22	-60	-54	4.73
	Left cerebellum/fusiform gyrus	209	-44	-46	-28	4.63
FM (pre) > FM (post)	Brainstem	253	8	-24	-56	4.52
FM(post) > HC	Bilateral rACC/MPFC	2650	18	42	26	4.76
	Left temporal pole	251	-52	16	-14	3.96
	Right temporal pole	190	42	24	-22	4.05
HC > FM(post)	Left superior parietal lobule / postcentral gyrus	449	-42	-44	56	4.22

FM(pre), Fibromyalgia pre-Tai Chi intervention; FM(post), Fibromyalgia post-Tai Chi intervention; HC, Healthy control; DLPFC, dorsolateral prefrontal cortex; rACC, rostral anterior cingulate cortex; MPFC, medial prefrontal cortex




Fig. 1 Comparison of dorsal lateral prefrontal cortex (DLPFC) connectivity maps in fibromyalgia patients and healthy controls. The partial correlation scatter indicates a regression analysis between the Overall Impact domain score and DLPFC-rACC/MPFC rsFC changes after Tai Chi

Given the importance of the rACC/MPFC in pain modulation and pathophysiology of chronic pain (Fields 2004; Li et al. 2016; Yu et al. 2014), we extracted the overlapping areas of the rACC/MPFC (contrasts of fibromyalgia post-intervention > healthy and fibromyalgia post-intervention > fibromyalgia pre-intervention) as an ROI. We defined the ROI using the overlapping areas so that we could target the areas reflecting both the neuropathology of fibromyalgia and the areas that would be modulated by Tai Chi intervention. We then compared the rsFC of the rACC/MPFC before and after Tai Chi using the same method as the ROI of the DLPFC and found that after Tai Chi practice, FM patients showed significant rsFC decreases between the rACC/MPFC and the left hippocampus, parahippocampus, fusiform, cerebellum; right PAG, and caudate; and bilateral inferior temporal gyrus (Table 3, Fig. 2). There were also significant rsFC increases between the rACC/MPFC and bilateral MPFC, precuneus, middle frontal gyrus, middle

cingulate cortex, supplementary motor cortex, operculum; left middle frontal gyrus, postcentral gyrus/supramarginal gyrus; and right precentral gyrus (Table 3).

Discussion


In this study, we found that compared to matched healthy controls, the cognitive control network (CCN) of fibromyalgia patients showed increased rsFC with the bilateral rACC/MPFC prior to Tai Chi practice. After intervention, fibromyalgia patients demonstrated further increased rsFC between the CCN and the bilateral rACC and MPFC, which was significantly associated with a decrease in Overall Impact domain scores on the FIQR. Further rsFC analyses using the rACC/MPFC as a seed found that, after Tai Chi, the rsFCs between the rACC/MPFC and both the PAG and hippocampus were significantly decreased.

Table 3 Regions with significantly different connectivity between the rACC/MPFC and other brain regions in FM and healthy controls before and after treatment

Conditions	Brain regions	mm ³	Cluster centroid (MNI)			Z
			x	y	z	
FM (post) > FM (pre)	Right middle frontal gyrus/precentral gyrus	811	42	12	38	6.10
	Left postcentral gyrus/supramarginal gyrus/operculum	1113	-44	-24	40	5.54
	Right middle frontal gyrus	227	40	54	2	4.39
	Bilateral middle cingulate cortex/MPFC/supplementary motor cortex	361	-4	26	36	4.07
	Left middle frontal gyrus	230	-36	40	46	4.41
	Left middle frontal gyrus	804	-38	48	8	5.65
	Bilateral precuneus	1238	-6	-68	46	5.03
FM (pre) > FM (post)	Right operculum	239	66	-16	16	3.76
	Left inferior temporal gyrus/fusiform gyrus/ hippocampus/parahippocampus	212	-48	-18	-44	4.40
	Right PAG (small volume correction)	6	22-8	-22	-8	2.88
	Right caudate	264	22	-4	36	5.58
	Right inferior temporal gyrus	498	52	-24	-38	5.02
	Left cerebellum gyrus	419	-20	-62	-24	5.28

FM (pre), Fibromyalgia pre-Tai Chi intervention; FM (post), Fibromyalgia post-Tai Chi intervention; HC, Healthy control; DLPFC, dorsolateral prefrontal cortex; rACC, rostral anterior cingulate cortex; MPFC, medial prefrontal cortex

Fig. 2 A representative result of resting state functional connectivity using the rostral anterior cingulate cortex (rACC)/medial prefrontal cortex (MPFC) as a seed (ROI). **a** seed region used in rsFC analysis **b** periaqueductal grey (PAG) showed significant rsFC decreases with rACC/MPFC after Tai Chi exercise



Our findings are consistent with a previous study demonstrating the beneficial effects of Tai Chi (Wang et al. 2010) and exercise (Flodin et al. 2015) in fibromyalgia patients. In addition, our results are consistent with previous brain imaging studies, which demonstrated a significant modulation effect of Tai Chi on brain function and structure in healthy human subjects (Fong et al. 2014; Tao et al. 2016, 2017a, b, c; Wei et al. 2013, 2014, 2017) and of exercise on brain responses to heat pain (Ellingson et al. 2016). Our results are also similar with other studies that explore resting state functional connectivity of pain-related brain regions (Flodin et al. 2015) in patients with fibromyalgia.

The DLPFC is a brain region in the dorsal pathway (stream) that is associated with reactions to various stimuli, including pain. Lorenz and colleagues (Lorenz et al. 2003) suggest that the DLPFC exerts active control on pain perception by modulating cortico-subcortical and cortico-cortical pathways. Napadow et al. (2010) found a negative association between the right frontoparietal network and both the PAG and hippocampus (greater right frontoparietal network connectivity in relation to lower pain levels) in fibromyalgia patients with greater spontaneous pain. In one of our previous studies, we found significant fMRI signal increases in the bilateral DLPFC and MPFC/ACC during both anticipation and the application of pain within a placebo analgesia conditioning paradigm (Kong et al. 2013). We also found that the baseline rsFC between the right frontoparietal network and the rACC could significantly predict placebo analgesia.

Both the DLPFC and ACC/MPFC play an important role in executive control processing (Petersen and Posner 2012) and non-pharmacological therapeutic modalities such as mind-body intervention (Tang et al. 2015) and acupuncture (Chen et al. 2014, 2015; Wang et al. 2016, 2017), as well as placebo effects (Gollub et al. 2018; Kong et al. 2006; Petrovic et al. 2002; Wager et al. 2004; Zubieta et al. 2005). Studies also suggest that the rACC/MPFC is a key region in the descending pain modulatory system (Eippert and Tracey 2014; Fields 2004; Kong et al. 2013; Kucyi et al. 2013; Tracey et al. 2002; Yu et al. 2014), forming a core network with the PAG and rostral ventral medulla (RVM) for pain modulation (Kong

et al. 2010). Patients with fibromyalgia have been shown to exhibit reduced connectivity during repeated pressure pain stimuli between the rACC and the amygdala, hippocampus, and brainstem (PAG) compared to healthy controls (Jensen et al. 2012b). Thus, we postulate that enhanced rsFC between the DLPFC and the rACC/MPFC in fibromyalgia patients at baseline may indicate that the brain employs a self-regulation mechanism to cope with repeated pain experiences.

Consistent with the aforementioned studies and our hypothesis, we found that after 12 weeks of Tai Chi intervention, the rsFC between the CCN and the rACC/MPFC rsFC was further enhanced, and this increase was associated with decreases in the Overall Impact domain score of the FIQR, suggesting that Tai Chi practice can further amplify this self-regulation process (Tang et al. 2015). Our finding is consistent with a previous study (Jensen et al. 2012a) investigating the effects of cognitive behavioral therapy on fibromyalgia patients. Investigators found that rather than reducing pain responses in patients with fibromyalgia, cognitive behavioral therapy increased access to executive regions for the reappraisal of pain. These results are further supported by another brain imaging study (Li et al. 2017) on the effect of acupuncture treatment in patients with migraines, in which the investigators discovered decreased rsFC between the right frontoparietal network and bilateral precuneus in migraine patients when compared to healthy controls. After acupuncture treatment, the rsFC between the frontoparietal network and precuneus was further reduced, and this reduction was associated with headache intensity relief.

Taken together, we believe our results suggest the existence of allostasis in patients with fibromyalgia. Based on the allostasis theory proposed by McEwen and colleagues (McEwen and Stellar 1993), physiological systems within the body fluctuate to meet demands from external forces to maintain homeostasis. In this context, allostasis represents the ability to protect the body through the altered activity of mediators that normally promote adaptation (Borsook et al. 2012). Most importantly, it seems that different treatment modalities, such as behavioral therapy, mind-body interventions, and acupuncture,

can further enhance allostasis to promote self-regulation and adaptation, and this may represent a common mechanism underlying non-pharmacological treatments. Elucidating this mechanism might shed light on the development of new pain management methods.

To further explore the role of the rACC/MPFC in the development of FM and the associated clinical improvements, we compared the rsFC changes of the rACC/MPFC before and after intervention. We found that after 12 weeks of Tai Chi practice, the rsFCs between the rACC/MPFC and both the hippocampus and PAG were significantly reduced. Recently, investigators have conceptualized chronic pain as a type of long-term learning (Apkarian et al. 2011), characterized by an accumulation of nociceptive memory (Yi and Zhang 2011) and an inability to extinguish negative emotional associations and anxiety (Mutso et al. 2014; Ploghaus et al. 2001; Vachon-Preseau et al. 2013, 2016). A recently proposed model of aversive prediction error in pain learning has suggested that the PAG is the primary hub in the pain learning neural network (Roy et al. 2014), relying on the rACC/MPFC and the hippocampus. We (Kong et al. 2018) found that the rsFC of the nucleus accumbens (NAc) showed significant increase with both the rACC/MPFC and dorsolateral prefrontal cortex in a boosted acupuncture group using an expectancy manipulation (Kong et al. 2006) as compared to a standard acupuncture group without such a manipulation. In another study, we found that acupuncture significantly increased the rsFC between the hippocampus and PAG (Egorova et al. 2015) and improved pain intensity in knee osteoarthritis patients, providing further support for the role of the hippocampus in pain modulation.

We also found that after the intervention, fibromyalgia patients showed increased rsFC between the rACC/MPFC and the middle/inferior prefrontal gyrus, middle cingulate, and postcentral gyrus/operculum. It is well known that these regions are involved in the cognitive, affective, and sensory aspects of pain, respectively (Tracey and Mantyh 2007). The increased rsFC between the rACC/MPFC and these regions may reflect the enhanced modulation effects and therapeutic benefits of Tai Chi.

Although our results may provide some novel insights into the mechanisms underlying the effectiveness of Tai Chi, the lack of a control condition has significantly limited our ability to identify the precise mechanism behind Tai Chi's therapeutic effects. We would like to emphasize here that the aim of this study was to investigate the rsFC alterations of the CCN in FM patients and how these rsFC changes following effective treatment were associated with symptom relief. These findings have the potential to shed light on our understanding of the pathophysiology and development of FM. We acknowledge that pharmacological medication (all participants were allowed to keep their regular medication during experiment) and other unknown factors may have also contributed to the clinical improvements observed. Nevertheless, this paradigm

has been used in previous studies (Rodriguez-Raecke et al. 2009). Additional controls should be applied in future studies to elucidate the specific effect of Tai Chi.

In summary, we found that fibromyalgia patients showed increased resting state functional connectivity between the cognitive control network and the rACC/MPFC, which may reflect an adaptive brain response to fibromyalgia. Effective interventions such as Tai Chi can further enhance the cognitive control network and rACC/MPFC rsFC and relieve clinical symptoms in fibromyalgia patients. Finally, we found that rACC/MPFC rsFCs with both the PAG and hippocampus were significantly reduced after the intervention. Our findings suggest the existence of allostasis in patients with fibromyalgia. A novel, effective pain management intervention such as Tai Chi may further enhance allostasis to promote self-regulation and adaptation mechanisms, and ultimately, provide significant therapeutic benefit to FM patients.

Acknowledgements Dr. Chenchen Wang is supported by R01 AT005521, R01AT006367 and K24AT007323, the National Center for Research Resources, National Institutes of Health (UL1 RR025752) and the National Center for Advancing Translational Sciences, National Institutes of Health (UL1TR000073 and UL1TR001064). Dr. Jian Kong is supported by R01AT006364, R01AT008563, R61 AT009310, R21AT008707, and P01 AT006663 from NIH/NCCIH. We also thank Courtney Lang for her help in manuscript revision and proof reading.

Author contributions Experimental design: Chenchen Wang, Jian Kong, Ramel Rones, William F Harvey.

Data collection: Jian Kong, Emily Wolcott, Jing Tao, Kristen Jorgenson, William F Harvey.

Data analysis: Zengjian Wang, Jian Kong.

Manuscript preparation: Jian Kong, Zengjian Wang, Chenchen Wang, Kristen Jorgenson, William F Harvey.

Funding This study was supported by the National Center for Complementary and Integrative Health of the National Institutes of Health (R01 AT005521, R01 AT006367 and K24 AT007323, R01AT006364, R01 AT008563, R21AT008707, R61 AT009310, and P01 AT006663 from NIH/NCCIH), the National Center for Research Resources, National Institutes of Health (UL1 RR025752) and the National Center for Advancing Translational Sciences, National Institutes of Health (UL1TR000073 and UL1TR001064).

Compliance with ethical standards

Conflict of interest Jian Kong holds equity of a start-up Inc. (MNT). All authors declare that they have no competing financial interests.

References

- Apkarian, A. V., Hashmi, J. A., & Baliki, M. N. (2011). Pain and the brain: specificity and plasticity of the brain in clinical chronic pain. *Pain*, *152*(3 Suppl), S49–S64. <https://doi.org/10.1016/j.pain.2010.11.010>.
- Borsook, D., Maleki, N., Becerra, L., & McEwen, B. (2012). Understanding migraine through the lens of maladaptive stress

- responses: a model disease of allostatic load. *Neuron*, 73(2), 219–234. <https://doi.org/10.1016/j.neuron.2012.01.001>.
- Bushnell, M. C., Ceko, M., & Low, L. A. (2013). Cognitive and emotional control of pain and its disruption in chronic pain. *Nature Reviews. Neuroscience*, 14(7), 502–511. <https://doi.org/10.1038/nrn3516>.
- Chen, X., Spaeth, R. B., Freeman, S. G., Scarborough, D. M., Hashmi, J. A., Wey, H. Y., et al. (2015). The modulation effect of longitudinal acupuncture on resting state functional connectivity in knee osteoarthritis patients. *Molecular Pain*, 11(1), 67. <https://doi.org/10.1186/s12990-015-0071-9>.
- Chen, X., Spaeth, R. B., Retzepi, K., Ott, D., & Kong, J. (2014). Acupuncture modulates cortical thickness and functional connectivity in knee osteoarthritis patients. *Scientific Reports*, 4, 6482. <https://doi.org/10.1038/srep06482>.
- Cieslik, E. C., Zilles, K., Caspers, S., Roski, C., Kellermann, T. S., Jakobs, O., et al. (2013). Is there "one" DLPFC in cognitive action control? Evidence for heterogeneity from co-activation-based parcellation. *Cerebral Cortex*, 23(11), 2677–2689. <https://doi.org/10.1093/cercor/bhs256>.
- Cole, M. W., & Schneider, W. (2007). The cognitive control network: Integrated cortical regions with dissociable functions. *NeuroImage*, 37(1), 343–360. <https://doi.org/10.1016/j.neuroimage.2007.03.071>.
- Cummiford, C. M., Nascimento, T. D., Foerster, B. R., Clauw, D. J., Zubieta, J. K., Harris, R. E., & DaSilva, A. F. (2016). Changes in resting state functional connectivity after repetitive transcranial direct current stimulation applied to motor cortex in fibromyalgia patients. *Arthritis Research & Therapy*, 18, 40. <https://doi.org/10.1186/s13075-016-0934-0>.
- Egorova, N., Gollub, R. L., & Kong, J. (2015). Repeated verum but not placebo acupuncture normalizes connectivity in brain regions dysregulated in chronic pain. *NeuroImage Clinical*, 9, 430–435. <https://doi.org/10.1016/j.nicl.2015.09.012>.
- Eippert, F., Bingel, U., Schoell, E. D., Yacubian, J., Klinger, R., Lorenz, J., & Buchel, C. (2009). Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron*, 63(4), 533–543 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19709634.
- Eippert, F., & Tracey, I. (2014). Pain and the PAG: learning from painful mistakes. *Nature Neuroscience*, 17(11), 1438–1439. <https://doi.org/10.1038/nn.3844>.
- Ellingson, L. D., Stegner, A. J., Schwabacher, I. J., Koltyn, K. F., & Cook, D. B. (2016). Exercise Strengthens Central Nervous System Modulation of Pain in Fibromyalgia. *Brain Sciences*, 6(1). <https://doi.org/10.3390/brainsci6010008>.
- Fields, H. (2004). State-dependent opioid control of pain. *Nature Reviews. Neuroscience*, 5(7), 565–575 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15208698.
- Fields, H. L., Basbaum, A. I., & Heinricher, M. M. (2005). Central nervous system mechanisms of pain modulation. In S. B. McMahon & M. Koltzenburg (Eds.), *Wall and Melzack's Textbook of Pain* (pp. 125–142). London: Elsevier - Churchill Livingstone.
- Flodin, P., Martinsen, S., Mannerkorpi, K., Lofgren, M., Bileviciute-Ljungar, I., Kosek, E., & Fransson, P. (2015). Normalization of aberrant resting state functional connectivity in fibromyalgia patients following a three month physical exercise therapy. *NeuroImage Clinical*, 9, 134–139. <https://doi.org/10.1016/j.nicl.2015.08.004>.
- Fong, D. Y., Chi, L. K., Li, F., & Chang, Y. K. (2014). The benefits of endurance exercise and Tai Chi Chuan for the task-switching aspect of executive function in older adults: an ERP study. *Frontiers in Aging Neuroscience*, 6, 295. <https://doi.org/10.3389/fnagi.2014.00295>.
- Goldenberg, D. L., Burckhardt, C., & Crofford, L. (2004). Management of fibromyalgia syndrome. *JAMA*, 292(19), 2388–2395. <https://doi.org/10.1001/jama.292.19.2388>.
- Gollub, R. L., Kirsch, I., Maleki, N., Wasan, A. D., Edwards, R. R., Tu, Y., et al. (2018). A functional neuroimaging study of expectancy effects on pain response in patients with knee osteoarthritis. *The Journal of Pain*. <https://doi.org/10.1016/j.jpain.2017.12.260>.
- Hwang, J., Egorova, N., Yang, X. Q., Zhang, W. Y., Chen, J., Yang, X. Y., et al. (2015). Subthreshold depression is associated with impaired resting state functional connectivity of the cognitive control network. *Translational Psychiatry*, 5(e683). <https://doi.org/10.1038/tp.2015.1174>.
- Jensen, K., Srinivasan, P., Spaeth, R., Tan, Y., Kosek, E., Petzke, F., et al. (2013a). Overlapping structural and functional brain changes in patients with long-term exposure to fibromyalgia. *Arthritis and Rheumatism*, 65(12), 3293–3303.
- Jensen, K. B., Kosek, E., Wicksell, R., Kemani, M., Olsson, G., Merle, J. V., et al. (2012a). Cognitive behavioral therapy increases pain-evoked activation of the prefrontal cortex in patients with fibromyalgia. *Pain*, 153(7), 1495–1503. <https://doi.org/10.1016/j.pain.2012.04.010>.
- Jensen, K. B., Loitole, R., Kosek, E., Petzke, F., Carville, S., Fransson, P., et al. (2012b). Patients with fibromyalgia display less functional connectivity in the brain's pain inhibitory network. *Molecular Pain*, 8(1), 32 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=22537768.
- Jensen, K. B., Srinivasan, P., Spaeth, R., Tan, Y., Kosek, E., Petzke, F., et al. (2013b). Overlapping structural and functional brain changes in patients with long-term exposure to fibromyalgia pain. *Arthritis and Rheumatism*, 65(12), 3293–3303. <https://doi.org/10.1002/art.38170>.
- Kong, J., Gollub, R. L., Rosman, I. S., Webb, J. M., Vangel, M. G., Kirsch, I., & Kaptchuk, T. J. (2006). Brain activity associated with expectancy-enhanced placebo analgesia as measured by functional magnetic resonance imaging. *The Journal of Neuroscience*, 26(2), 381–388 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16407533.
- Kong, J., Jensen, K., Loitole, R., Cheetham, A., Wey, H. Y., Tan, T., et al. (2013). Functional connectivity of frontoparietal network predicts cognitive modulation of pain. *Pain*, 154(3), 459–467.
- Kong, J., Tu, P. C., Zyloney, C., & Su, T. P. (2010). Intrinsic functional connectivity of the periaqueductal gray, a resting fMRI study. *Behavioural Brain Research*, 211(2), 215–219 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20347878.
- Kong, J., Wang, Z. J., Leiser, J., Minicucci, D., Edwards, R., Kirsch, I., et al. (2018). Enhancing treatment of osteoarthritis knee pain by boosting expectancy: A functional neuroimaging study. *NeuroImage Clinical*, 18, 325–334.
- Kuchinad, A., Schweinhardt, P., Seminowicz, D. A., Wood, P. B., Chizh, B. A., & Bushnell, M. C. (2007). Accelerated brain gray matter loss in fibromyalgia patients: premature aging of the brain? *The Journal of Neuroscience*, 27(15), 4004–4007. <https://doi.org/10.1523/jneurosci.0098-07.2007>.
- Kucyi, A., Salomons, T. V., & Davis, K. D. (2013). Mind wandering away from pain dynamically engages antinociceptive and default mode brain networks. *Proceedings of the National Academy of Sciences of the United States of America*, 110(46), 18692–18697.
- Lautenschlager, J. (2000). Present state of medication therapy in fibromyalgia syndrome. *Scandinavian Journal of Rheumatology. Supplement*, 113, 32–36 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11028829>.
- Legrain, V., Damme, S. V., Eccleston, C., Davis, K. D., Seminowicz, D. A., & Crombez, G. (2009). A neurocognitive model of attention to

- pain: behavioral and neuroimaging evidence. *Pain*, 144(3), 230–232. <https://doi.org/10.1016/j.pain.2009.03.020>.
- Li, Z., Lan, L., Zeng, F., Makris, N., Hwang, J., Guo, T., et al. (2017). The altered right frontoparietal network functional connectivity in migraine and the modulation effect of treatment. *Cephalalgia*, 37(2), 161–176. <https://doi.org/10.1177/0333102416641665>.
- Li, Z., Liu, M., Lan, L., Zeng, F., Makris, N., Liang, Y., et al. (2016). Altered periaqueductal gray resting state functional connectivity in migraine and the modulation effect of treatment. *Scientific Reports*, 6, 20298. <https://doi.org/10.1038/srep20298>.
- Linnman, C., Moulton, E. A., Barmettler, G., Becerra, L., & Borsook, D. (2012). Neuroimaging of the periaqueductal gray: state of the field. *NeuroImage*, 60(1), 505–522. <https://doi.org/10.1016/j.neuroimage.2011.11.095>.
- Lobanov, O. V., Quevedo, A. S., Hadsel, M. S., Kraft, R. A., & Coghill, R. C. (2013). Frontoparietal mechanisms supporting attention to location and intensity of painful stimuli. *Pain*, 154(9), 1758–1768. <https://doi.org/10.1016/j.pain.2013.05.030>.
- Loggia, M. L., Berna, C., Kim, J., Cahalan, C. M., Gollub, R. L., Wasan, A. D., et al. (2014). Disrupted brain circuitry for pain-related reward/punishment in fibromyalgia. *Arthritis & Rheumatology*, 66(1), 203–212. <https://doi.org/10.1002/art.38191>.
- Lorenz, J., Minoshima, S., & Casey, K. L. (2003). Keeping pain out of mind: the role of the dorsolateral prefrontal cortex in pain modulation. *Brain*, 126, 1079–1091.
- Lutz, J., Jager, L., de Quervain, D., Krauseneck, T., Padberg, F., Wichnalek, M., et al. (2008). White and gray matter abnormalities in the brain of patients with fibromyalgia: a diffusion-tensor and volumetric imaging study. *Arthritis and Rheumatism*, 58(12), 3960–3969. <https://doi.org/10.1002/art.24070>.
- McEwen, B. S., & Stellar, E. (1993). Stress and the individual. Mechanisms leading to disease. *Archives of Internal Medicine*, 153(18), 2093–2101 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8379800>.
- McEwen, B. S., & Wingfield, J. C. (2003). The concept of allostasis in biology and biomedicine. *Hormones and Behavior*, 43(1), 2–15 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12614627>
- Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, 24, 167–202 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11283309.
- Mutso, A. A., Petre, B., Huang, L., Baliki, M. N., Torbey, S., Herrmann, K. M., et al. (2014). Reorganization of hippocampal functional connectivity with transition to chronic back pain. *Journal of Neurophysiology*, 111(5), 1065–1076. <https://doi.org/10.1152/jn.00611.2013>.
- Napadow, V., LaCount, L., Park, K., As-Sanie, S., Clauw, D. J., & Harris, R. E. (2010). Intrinsic brain connectivity in fibromyalgia is associated with chronic pain intensity. *Arthritis and Rheumatism*, 62(8), 2545–2555 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20506181.
- Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years after. *Annual Review of Neuroscience*, 35, 73–89. <https://doi.org/10.1146/annurev-neuro-062111-150525>.
- Petrovic, P., Kalso, E., Petersson, K. M., & Ingvar, M. (2002). Placebo and opioid analgesia—imaging a shared neuronal network. *Science*, 295(5560), 1737–1740.
- Ploghaus, A., Narain, C., Beckmann, C. F., Clare, S., Bantick, S., Wise, R., et al. (2001). Exacerbation of pain by anxiety is associated with activity in a hippocampal network. *The Journal of Neuroscience*, 21(24), 9896–9903 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11739597.
- Redcay, E., Moran, J. M., Mavros, P. L., Tager-Flusberg, H., Gabrieli, J. D., & Whitfield-Gabrieli, S. (2013). Intrinsic functional network organization in high-functioning adolescents with autism spectrum disorder. *Frontiers in Human Neuroscience*, 7, 573. <https://doi.org/10.3389/fnhum.2013.00573>.
- Robinson, M. E., Craggs, J. G., Price, D. D., Perlstein, W. M., & Staud, R. (2011). Gray matter volumes of pain-related brain areas are decreased in fibromyalgia syndrome. *The Journal of Pain*, 12(4), 436–443. <https://doi.org/10.1016/j.jpain.2010.10.003>.
- Rodriguez-Raecke, R., Niemeier, A., Ihle, K., Ruether, W., & May, A. (2009). Brain gray matter decrease in chronic pain is the consequence and not the cause of pain. *The Journal of Neuroscience*, 29(44), 13746–13750. <https://doi.org/10.1523/JNEUROSCI.3687-09.2009>.
- Rosen, M. L., Stern, C. E., Michalka, S. W., Devaney, K. J., & Somers, D. C. (2015). Cognitive control network contributions to memory-guided visual attention. *Cerebral Cortex*. <https://doi.org/10.1093/cercor/bhv028>.
- Roy, M., Shohamy, D., Daw, N., Jepma, M., Wimmer, G. E., & Wager, T. D. (2014). Representation of aversive prediction errors in the human periaqueductal gray. *Nature Neuroscience*, 17(11), 1607–1612. <https://doi.org/10.1038/nm.3832>.
- Sheline, Y. I., Price, J. L., Yan, Z., & Mintun, M. A. (2010). Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proceedings of the National Academy of Sciences of the United States of America*, 107(24), 11020–11025. <https://doi.org/10.1073/pnas.1000446107>.
- Sumpton, J. E., & Moulin, D. E. (2014). Fibromyalgia. *Handbook of Clinical Neurology*, 119, 513–527. <https://doi.org/10.1016/B978-0-7020-4086-3.00033-3>.
- Tang, Y. Y., Holzel, B. K., & Posner, M. I. (2015). The neuroscience of mindfulness meditation. *Nature Reviews. Neuroscience*, 16(4), 213–225. <https://doi.org/10.1038/nrn3916>.
- Tao, J., Chen, X., Liu, J., Egorova, N., Xue, X., Liu, W., et al. (2017a). Tai Chi Chuan and Baduanjin mind-body training changes resting-state low-frequency fluctuations in the frontal lobe of older adults: a resting-state fMRI study. *Frontiers in Human Neuroscience*, 11, 514. <https://doi.org/10.3389/fnhum.2017.00514>.
- Tao, J., Chen, X. L., Egorova, N., Liu, J., Xue, X. H., Huang, J., et al. (2017b). Tai Chi Chuan and Baduanjin practice modulates functional connectivity of the cognitive control network in older adults. *Scientific Reports*, 7(41581). <https://doi.org/10.1038/srep41581>.
- Tao, J., Liu, J., Egorova, N., Chen, X., Sun, S., Xue, X., et al. (2016). Increased hippocampus-medial prefrontal cortex resting-state functional connectivity and memory function after Tai Chi Chuan practice in elder adults. *Frontiers in Aging Neuroscience*, 8, 25. <https://doi.org/10.3389/fnagi.2016.00025>.
- Tao, J., Liu, J., Liu, W., Huang, J., Xue, X., Chen, X., et al. (2017c). Tai Chi Chuan and Baduanjin increase grey matter volume in older adults: a brain imaging study. *Journal of Alzheimer's Disease*. <https://doi.org/10.3233/JAD-170477>.
- Torta, D. M., Legrain, V., Mouraux, A., & Valentini, E. (2017). Attention to pain! A neurocognitive perspective on attentional modulation of pain in neuroimaging studies. *Cortex*, 89, 120–134. <https://doi.org/10.1016/j.cortex.2017.01.010>.
- Tracey, I., & Bushnell, M. C. (2009). How neuroimaging studies have challenged us to rethink: is chronic pain a disease? *The Journal of Pain*, 10(11), 1113–1120. <https://doi.org/10.1016/j.jpain.2009.09.001>.
- Tracey, I., & Mantyh, P. W. (2007). The cerebral signature for pain perception and its modulation. *Neuron*, 55(3), 377–391 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17678852.
- Tracey, I., Ploghaus, A., Gati, J. S., Clare, S., Smith, S., Menon, R. S., & Matthews, P. M. (2002). Imaging attentional modulation of pain in the periaqueductal gray in humans. *The Journal of Neuroscience*, 22(7), 2748–2752 Retrieved from <http://www.ncbi.nlm.nih.gov/>

- entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11923440.
- Truini, A., Tinelli, E., Gerardi, M. C., Calistri, V., Iannuccelli, C., La Cesa, S., et al. (2016). Abnormal resting state functional connectivity of the periaqueductal grey in patients with fibromyalgia. *Clinical and Experimental Rheumatology*, *34*(2 Suppl 96), S129–S133 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/27157397>.
- Vachon-Preseau, E., Roy, M., Martel, M. O., Caron, E., Marin, M. F., Chen, J., et al. (2013). The stress model of chronic pain: evidence from basal cortisol and hippocampal structure and function in humans. *Brain*, *136*(Pt 3), 815–827. <https://doi.org/10.1093/brain/aws371>.
- Vachon-Preseau, E., Tetreault, P., Petre, B., Huang, L., Berger, S. E., Torbey, S., et al. (2016). Corticolimbic anatomical characteristics predetermine risk for chronic pain. *Brain*. <https://doi.org/10.1093/brain/aww100>.
- Valet, M., Sprenger, T., Boecker, H., Willloch, F., Rummeny, E., Conrad, B., et al. (2004). Distraction modulates connectivity of the cingulo-frontal cortex and the midbrain during pain—an fMRI analysis. *Pain*, *109*(3), 399–408 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15157701.
- Villemure, C., & Bushnell, M. C. (2002). Cognitive modulation of pain: how do attention and emotion influence pain processing? *Pain*, *95*, 195–199.
- Villemure, C., & Schweinhardt, P. (2010). Supraspinal pain processing: distinct roles of emotion and attention. *The Neuroscientist*, *16*(3), 276–284. <https://doi.org/10.1177/1073858409359200>.
- Wager, T. D., Rilling, J. K., Smith, E. E., Sokolik, A., Casey, K. L., Davidson, R. J., et al. (2004). Placebo-induced changes in FMRI in the anticipation and experience of pain. *Science*, *303*, 1162–1167.
- Wang, C., Schmid, C. H., Rones, R., Kalish, R., Yinh, J., Goldenberg, D. L., et al. (2010). A randomized trial of tai chi for fibromyalgia. *The New England Journal of Medicine*, *363*(8), 743–754 Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20818876.
- Wang, X., Wang, Z., Liu, J., Chen, J., Liu, X., Nie, G., et al. (2016). Repeated acupuncture treatments modulate amygdala resting state functional connectivity of depressive patients. *Neuroimage Clinical*, *12*, 746–752. <https://doi.org/10.1016/j.nicl.2016.07.011>.
- Wang, Z., Wang, X., Liu, J., Chen, J., Liu, X., Nie, G., et al. (2017). Acupuncture treatment modulates the corticostriatal reward circuitry in major depressive disorder. *Journal of Psychiatric Research*, *84*, 18–26. <https://doi.org/10.1016/j.jpsychires.2016.09.014>.
- Wayne, P. M., Walsh, J. N., Taylor-Piliae, R. E., Wells, R. E., Papp, K. V., Donovan, N. J., & Yeh, G. Y. (2014). Effect of tai chi on cognitive performance in older adults: systematic review and meta-analysis. *Journal of the American Geriatrics Society*, *62*(1), 25–39. <https://doi.org/10.1111/jgs.12611>.
- Wei, G. X., Dong, H. M., Yang, Z., Luo, J., & Zuo, X. N. (2014). Tai Chi Chuan optimizes the functional organization of the intrinsic human brain architecture in older adults. *Frontiers in Aging Neuroscience*, *6*, 74. <https://doi.org/10.3389/fnagi.2014.00074>.
- Wei, G. X., Gong, Z. Q., Yang, Z., & Zuo, X. N. (2017). Mind-body practice changes fractional amplitude of low frequency fluctuations in intrinsic control networks. *Frontiers in Psychology*, *8*, 1049. <https://doi.org/10.3389/fpsyg.2017.01049>.
- Wei, G. X., Xu, T., Fan, F. M., Dong, H. M., Jiang, L. L., Li, H., et al. (2013). Can Taichi reshape the brain? A brain morphometry study. *PLoS One*, *8*(4), e61038. <https://doi.org/10.1371/journal.pone.0061038>.
- Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connectivity*, *2*(3), 125–141. <https://doi.org/10.1089/brain.2012.0073>.
- Wiech, K. (2016). Deconstructing the sensation of pain: The influence of cognitive processes on pain perception. *Science*, *354*(6312), 584–587. <https://doi.org/10.1126/science.aaf8934>.
- Yi, M., & Zhang, H. (2011). Nociceptive memory in the brain: cortical mechanisms of chronic pain. *The Journal of Neuroscience*, *31*(38), 13343–13345. <https://doi.org/10.1523/JNEUROSCI.3279-11.2011>.
- Yu, R., Gollub, R., Spaetha, R., Napadowa, V., Wasana, A., & Kong, J. (2014). Disrupted functional connectivity of the periaqueductal gray in chronic low back pain. *NeuroImage Clinical*, *6*, 100–108.
- Zanto, T. P., & Gazzaley, A. (2013). Fronto-parietal network: flexible hub of cognitive control. *Trends in Cognitive Sciences*, *17*(12), 602–603. <https://doi.org/10.1016/j.tics.2013.10.001>.
- Zeidan, F., Emerson, N. M., Farris, S. R., Ray, J. N., Jung, Y., McHaffie, J. G., & Coghill, R. C. (2015). Mindfulness meditation-based pain relief employs different neural mechanisms than placebo and sham mindfulness meditation-induced analgesia. *The Journal of Neuroscience*, *35*(46), 15307–15325. <https://doi.org/10.1523/JNEUROSCI.2542-15.2015>.
- Zeidan, F., Grant, J. A., Brown, C. A., McHaffie, J. G., & Coghill, R. C. (2012). Mindfulness meditation-related pain relief: evidence for unique brain mechanisms in the regulation of pain. *Neuroscience Letters*, *520*(2), 165–173. <https://doi.org/10.1016/j.neulet.2012.03.082>.
- Zubieta, J. K., Bueller, J. A., Jackson, L. R., Scott, D. J., Xu, Y., Koeppe, R. A., et al. (2005). Placebo effects mediated by endogenous opioid activity on mu-opioid receptors. *The Journal of Neuroscience*, *25*(34), 7754–7762.