



Brazilian Journal of OTORHINOLARYNGOLOGY

www.bjorl.org



REVIEW ARTICLE

The efficacy and safety of ciclesonide for the treatment of perennial allergic rhinitis: a systematic review and meta-analysis[☆]



Qi Yang^{a,b}, Fei Wang^{a,b}, Bin Li^a, Wenbin Wu^c, Dengpiao Xie^a, Li He^a,
Nan Xiang^d, Yan Dong^{a,c,*}

^a Chengdu University of Traditional Chinese Medicine, Chengdu, China

^b Sichuan College of Traditional Chinese Medicine, Mianyang, China

^c Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu, China

^d Chengdu First People's Hospital, Chengdu, China

Received 19 August 2018; accepted 24 October 2018

Available online 22 November 2018

KEYWORDS

Meta-analysis;
Ciclesonide;
Rhinitis;
Allergic;
Perennial

Abstract

Introduction: Allergic rhinitis is a chronic inflammatory disease which affects 1 out of 6 individuals. Perennial allergic rhinitis accounts for 40% of AR cases. Ciclesonide is one of the relatively new intranasal steroid for allergic rhinitis.

Objective: The purpose of this study was to evaluate the efficacy and safety of ciclesonide in the treatment of perennial allergic rhinitis.

Methods: We searched Pubmed, Scientific Citation Index, Embase, Clinical Trial Registries for randomized controlled trials and Cochrane Central Register of Controlled Trials to find out the randomized controlled Trial comparing ciclesonide with placebo for PAR.

Results: Eight studies were included. In comparison with placebo groups, ciclesonide groups significantly decreased Reflective Total Nasal Symptom Score (MD = -0.56; 95% CI -0.72 to -0.39, $p < 0.00001$) with heterogeneity ($p = 0.19$, $I^2 = 24\%$), Instantaneous Total Nasal Symptom Score (MD = -0.57; 95% CI -0.75 to -0.39, $p < 0.00001$) with heterogeneity ($p = 0.34$, $I^2 = 11\%$). A significant effect for Reflective Nasal Symptom Score Subtotal (MD = -0.15; 95% CI -0.18 to -0.13, $p < 0.00001$) with heterogeneity ($p = 0.12$, $I^2 = 24\%$) was also demonstrated. Rhinoconjunctivitis quality of life questionnaire score (RQLQs) (MD = -0.27; 95% CI -0.39 to -0.15, $p < 0.00001$) with heterogeneity ($p = 0.58$, $I^2 = 0\%$) in the treatment of ciclesonide was also significantly reduced. In addition, the difference in Treatment-Emergent Adverse Events between the two groups was not significant.

[☆] Please cite this article as: Yang Q, Wang F, Li B, Wu W, Xie D, He L, et al. The efficacy and safety of ciclesonide for the treatment of perennial allergic rhinitis: a systematic review and meta-analysis. Braz J Otorhinolaryngol. 2019;85:371–8.

* Corresponding author.

E-mail: dongyancd@163.com (Y. Dong).

PALAVRAS-CHAVE

Metanálise;
Ciclesonida;
Rinite;
Alérgica;
Perene

Conclusion: Ciclesonide can improve perennial allergic rhinitis without increasing adverse events. Ciclesonide may be another valuable choice for perennial allergic rhinitis in the future. © 2018 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Published by Elsevier Editora Ltda. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Eficácia e segurança da ciclesonida no tratamento da rinite alérgica perene: uma revisão sistemática e metanálise

Resumo

Introdução: A rinite alérgica é uma doença inflamatória crônica que afeta um a cada seis indivíduos. A rinite alérgica perene é responsável por 40% dos casos de rinite alérgica. A ciclesonida é um dos corticosteroides intranasais mais novos para o tratamento dessa condição clínica.

Objetivo: Avaliar a eficácia e segurança da ciclesonida no tratamento da rinite alérgica perene.

Método: Uma busca foi feita nos bancos de dados Pubmed, *Scientific Citation Index*, Embase e *Clinical Trial Registries* por ensaios clínicos randomizados e *Cochrane Central Register of Controlled Trials* por estudos controlados randomizados que comparassem ciclesonida com placebo no tratamento da rinite alérgica perene.

Resultados: Oito estudos foram incluídos. Em comparação com os grupos placebo, os grupos ciclesonida mostraram diminuição significativa no escore do *Reflective Total Nasal Symptom Score* (DM = -0,56; IC 95%: -0,72 a -0,39, $p < 0,00001$) com heterogeneidade ($p = 0,19$, $I^2 = 24\%$), do *Instantaneous Total Nasal Symptom Score* (DM = -0,57; IC95%: -0,75 a -0,39, $p < 0,00001$) com heterogeneidade ($p = 0,34$, $I^2 = 11\%$). Um efeito significativo no escore do *Reflective Nasal Symptom Score Subtotal* (DM = -0,15; IC 95%: -0,18 a -0,13, $p < 0,00001$) com heterogeneidade ($p = 0,12$, $I^2 = 24\%$) também foi demonstrado. O escore do *Rhinoconjunctivitis Quality of Life Questionnaire score* (RQLQs) (DM = -0,27; IC 95%: -0,39 a -0,15, $p < 0,00001$) com heterogeneidade ($p = 0,58$, $I^2 = 0\%$) também foi significativamente reduzido no tratamento com ciclesonida. Além disso, a diferença em relação aos eventos adversos emergentes do tratamento entre os dois grupos não foi significativa.

Conclusão: A ciclesonida pode melhorar a rinite alérgica perene sem aumentar os eventos adversos. Esse fármaco pode ser outra opção valiosa para a rinite alérgica perene no futuro.

© 2018 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Publicado por Elsevier Editora Ltda. Este é um artigo Open Access sob uma licença CC BY (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Allergic Rhinitis (AR), a chronic inflammatory disease, is characterized by nasal itching, sneezing, runny nose and congestion.¹ As a highly prevalent condition, AR affects 1 out of 6 individuals. The symptoms of AR interfere with all aspects of daily life that are associated with decreased sleep quality and performance at work.²

Despite currently available treatment options, the incidence of AR is increasing. It remains the leading cause of morbidity, absenteeism and restricted activities and is related to considerable cost pressures in the health care system.^{3,4} AR can be divided into seasonal and perennial forms. Perennial allergic rhinitis (PAR) accounts for 40% of AR cases.⁵ AR is a Type 1 IgE-mediated hypersensitivity reaction.⁶

Intranasal corticosteroids (INS) represent the standard treatment for AR of all severities owing to their anti-inflammatory activity.^{7,8} Systematic reviews and meta-analyses revealed topical corticosteroids are superior to

antihistamines in putting nasal symptoms of AR under control.^{9,10} Ciclesonide was approved by the FDA as one of the relatively new INS additions to the AR armamentarium in 2006.¹¹

Ciclesonide in PAR have been evaluated in several randomized controlled trials (RCT). However, the evidence from the currently available individual randomized trials concerning ciclesonide in PAR is not convincing. Whether ciclesonide has an effect on PAR and whether it plays a role in prevention and treatment remains to be seen. To figure out these issues, we engaged in a systematic review with meta-analysis of randomized controlled trials to analyze the effect of ciclesonide in the treatment of PAR.

Methods

Data sources

We searched Pubmed, Scientific Citation Index, Embase, Clinical Trial Registries for randomized controlled trials and




Figure 1 Selection of studies.

Cochrane Central Register of Controlled Trials, with a search deadline of July 2018. We used the following keywords: "Rhinitis, Allergic, Perennial", "Rhinitis, Allergic, Non-seasonal", "Ciclesonide" and "random* controlled trial" (Fig. 1). In order to identify potentially pertinent studies, we scanned the citations of the included studies.

Study selection

Two independent reviewers assessed the title and abstract of relevant papers. If the study was randomized trials and contrasted ciclesonide with placebo for patients with "perennial allergic rhinitis", the study was included.

Data extraction and quality assessment

The data information of characteristics of methods, participants, interventions and results were extracted independently by two reviewers. The Cochrane Handbook for Systematic Reviews of Interventions¹² was used to assess the quality of included studies by evaluating the risk of bias. Any discrepancy was resolved by the third author.

Outcome definition

The primary outcome was the change in average A.M. and P.M. reflective Total Nasal Symptom Score (rTNSS), A.M. instantaneous Total Nasal Symptom Score (iTNSS) and the

second outcome included changes in average A.M. and P.M. reflective Nasal Symptom Score (rNSS), Rhinoconjunctivitis Quality of Life Questionnaire score (RQLQs). Treatment-Emergent Adverse Events (TEAEs) were used to monitor the safety.

Data synthesis and analysis

The effect size of continuous outcomes was evaluated by weighted mean difference (WMD) and dichotomous outcomes assessed by Risk Ratio (RR) with 95% Confidence Interval (CI). Heterogeneity was evaluated with I^2 statistics. A random-effects model was applied regardless of the heterogeneity of the results. Statistical assessments were performed using Review Manager, version 5.3.

Results

Study selection and study characteristics

In the initial search 53 related publications were identified in total. 14 duplicates were removed afterward. 29 studies were excluded by reading the title or abstract. The remaining 10 full-text articles were reviewed and 2 studies were excluded. Finally, 8 trials¹³⁻¹⁷ (NCT01451541, NCT01033825, NCT01378429) enrolling 4039 patients were included (Table 1). The selection process of studies

Table 1 Summary of trials included in the meta-analysis.

Study	Year	No. of patients (ciclesonide/placebo)	Interventions	Duration (weeks)	Outcomes
NCT01451541	2014	758 (511/247)	Ciclesonide 37 µg	12	rTNSS iTNSS
NCT01033825	2012	160 (105/55)	Ciclesonide 74 µg	6	TEAE
			Ciclesonide 160 µg		rTNSS iTNSS
			Ciclesonide 200 µg		rNSS
			Ciclesonide 320 µg		
Eli (13)	2007	471 (238/233)	Ciclesonide 200 µg	6	rTNSS rNSS
					RQLQ TEAE
Paul (14)	2007	663 (441/222)	Ciclesonide 200 µg	52	rTNSS rNSS
					RQLQ TEAE
William (15)	2012	1110 (803/307)	Ciclesonide 74 µg	26	rTNSS iTNSS
			Ciclesonide 148 µg		rNSS RQLQ
					TEAE
Kenneth (16)	2007	123 (81/42)	Ciclesonide 200 µg	12	rTNSS rNSS
					TEAE
NCT01378429	2014	89 (47/42)	Ciclesonide 74 µg	6	rTNSS
William (17)	2008	665 (500/165)	Ciclesonide 25 µg	12	rTNSS TEAE
			Ciclesonide 100 µg		
			Ciclesonide 200 µg		

was showed in Fig. 1. Duration of treatment was 6–52 weeks.

Quality assessment of included studies

All 8 studies^{13–17} (NCT01451541, NCT01033825, NCT01378429) were randomized and double-blind. Seven studies^{13–15,17} (NCT01451541, NCT01033825, NCT01378429) were multicentre trials. However, all studies did not provide concrete randomization methods. All studies^{13–17} (NCT01451541, NCT01033825, NCT01378429) reported blinding of participants and personnel. All studies did not have reporting bias. Five studies reported withdrawals and one study¹⁵ was analyzed on an intention-to-treat basis (Fig. 2).

Effects on rTNSS

Eight studies included the comparison of the change in rTNSS between eight groups^{13–17} (NCT01451541, NCT01033825, NCT01378429). The pooled result showed that there was significant difference between the two groups (MD = −0.56; 95% CI −0.72 to −0.39, $p < 0.00001$) (Fig. 3) with heterogeneity ($p = 0.19$, $I^2 = 24\%$) (Fig. 3).

Effects on iTNSS

Three studies included comparison of iTNSS between three groups¹⁵ (NCT01451541, NCT01033825). Pooled results showed that there was a significant difference between the two groups (MD = −0.57; 95% CI −0.75 to −0.39, $p < 0.00001$) (Fig. 4) with heterogeneity ($p = 0.34$, $I^2 = 11\%$) (Fig. 4).

Effects on rNSS

We compared rNSS in five trials^{14–17} (NCT01033825). There was significant difference between the two groups, sneezing (MD = −0.15; 95% CI −0.21 to −0.10, $p < 0.00001$) with heterogeneity ($p = 0.29$, $I^2 = 18\%$) (Fig. 5), runny nose (MD = −0.16; 95% CI −0.22 to −0.10, $p < 0.00001$) with heterogeneity ($p = 0.28$, $I^2 = 19\%$) (Fig. 5), nasal itching (MD = −0.14; 95% CI −0.20 to −0.09, $p < 0.00001$) with heterogeneity ($p = 0.37$, $I^2 = 8\%$) (Fig. 5), nasal congestion (MD = −0.17; 95% CI −0.25 to −0.09, $p < 0.00001$) with heterogeneity ($p = 0.03$, $I^2 = 55\%$) (Fig. 5), Subtotal (MD = −0.15; 95% CI −0.18 to −0.13, $p < 0.00001$) with heterogeneity ($p = 0.12$, $I^2 = 24\%$) (Fig. 5).

Effects on RQLQs

RQLQs was compared in the three trials.^{13–15} Compared with placebo, ciclesonide significantly reduced RQLQs (MD = −0.27; 95% CI −0.39 to −0.15, $p < 0.00001$) with heterogeneity ($p = 0.58$, $I^2 = 0\%$) (Fig. 6).

Safety

Safety was assessed by monitoring TEAEs. For TEAEs, six trials^{13–17} (NCT01451541) reported complete data. There was no significant difference between the two groups (RR = 1.02; 95% CI 0.94–1.10, $p = 0.61$) with heterogeneity ($p = 0.17$, $I^2 = 36\%$) (Fig. 7).

Publication bias and sensitivity analysis

There was no evidence of significant publication bias by Egger's test for rTNSS ($t = -0.52$, $p = 0.609$).




Figure 2 Risk of bias graph according to recommendations from the Cochrane collaboration.

Discussion and conclusions

Antihistamines and corticosteroids are current treatments for controlling AR symptoms. INS are the most effective available drug suppressing all rhinitis symptoms which include nasal blockage.¹⁸ However, although widely used, ciclesonide for AR still is lacking in clear evidence to make decisive recommendations for a therapeutic option. In the present study, we performed a search to evaluate the efficacy and safety of ciclesonide in patients with PAR. In this review, we found that the ciclesonide might be able to decrease rTNSS, iTNSS, rNSS, RQLQs without increasing TEAEs in the short term.

Ciclesonide is the latest inhaled glucocorticosteroid to treat symptoms of asthma and AR.¹⁹ The anti-inflammatory effect of ciclesonide is seen solely at the bronchial level, only a fraction of the drug reaching the gastrointestinal tract and becoming inactive.^{14,20}

In our review, ciclesonide produced significant relief in rTNSS and iTNSS. In the study by Eli,¹³ it was suggested that improvement in the rTNSS continued to increase throughout the 6 weeks of treatment. These continued improvements in nasal symptoms are associated with AR and may encourage patients to stick to treatment.

All individual rNSS declined in all patients treated with ciclesonide, especially nasal congestion, which is the most difficult symptom to treat. The overall change in rTNSS was driven by all four nasal symptoms, suggesting that all individual rNSS contributed to the overall difference between two groups. Owing to a variable degree of heterogeneity in these studies, we performed sensitivity analysis. We should treat the results cautiously, although the results did not change.

Ciclesonide produced a statistically significant reduction in RQLQs. However, studies on the clinical relevance of questionnaires showed that only 0.5 or more of the changes were clinically relevant.²¹

For TEAEs, there were no significant differences between the two groups. In our review, most of adverse events of ciclesonide were mild or moderate and well tolerated. Rates of discontinuation were similar to placebo. Typically seen Adverse Events (AEs) with INS are usually topical in nature and include nasal discomfort and nosebleeds.²² In the study by William,¹⁵ epistaxis and upper respiratory tract infection were the most commonly reported AEs. It may be as a consequence of negligible oral bioavailability (1%), high protein




Figure 3 Forest plots of rTNSS of patients treated with ciclesonide.




Figure 4 Forest plots of iTNSS of patients treated with ciclesonide.




Figure 5 Forest plots of rNSS of patients treated with ciclesonide.




Figure 6 Forest plots of RQLQs of patients treated with ciclesonide.




Figure 7 Forest plots of TEAEs of patients treated with ciclesonide.

binding (99%) of ciclesonide and the active metabolite, with negligible impact on the hypothalamic–pituitary–adrenal axis.^{23–25}

There were several limitations in our meta-analysis. First, several notable areas of variability existed in the data. The duration of intervention varied between 2 and 52 weeks and the baseline severity of the disease had some differences. Second, there is a possibility of study selection bias. Third, four of the eight studies¹⁵ (NCT01451541, NCT01033825, NCT01378429) were sponsored by pharmaceutical companies. We conduct a subgroup analysis by excluding these data and the results did not change.

In conclusion, ciclesonide can improve PAR without increasing adverse events. Ciclesonide may be another valuable choice for patients with PAR in the future.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Informed consent

Informed consent was obtained from all individual participants included the study.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Greiner AN, Hellings PW, Rotiroti G, Scadding GK. Allergic rhinitis. *Lancet*. 2011;378:2112–22.
- Dykewicz MS, Hamilos DL. Rhinitis and sinusitis. *J Allergy Clin Immunol*. 2010;125 Suppl. 2:S103–15.
- Benninger MS, Benninger RM. The impact of allergic rhinitis on sexual activity, sleep, and fatigue. *Allergy Asthma Proc*. 2009;30:358–65.
- Meltzer EO, Nathan R, Derebery J, Stang PE, Campbell UB, Yeh WS, et al. Sleep, quality of life, and productivity impact of nasal symptoms in the United States: findings from the Burden of Rhinitis in America survey. *Allergy Asthma Proc*. 2009;30:244–54.
- Skoner DP. Allergic rhinitis: definition, epidemiology, pathophysiology, detection, and diagnosis. *J Allergy Clin Immunol*. 2001;108 Suppl. 1:S2–8.
- Kakli HA, Riley TD. Allergic rhinitis. *Prim Care*. 2016;43:465–75.
- Wallace DV, Dykewicz MS, Bernstein DI, Blessing-Moore J, Cox L, Khan DA, et al. The diagnosis and management of rhinitis: an updated practice parameter. *J Allergy Clin Immunol*. 2008;122 Suppl. 2:S1–84.
- Sur DK, Scandale S. Treatment of allergic rhinitis. *Am Fam Phys*. 2010;81:1440–6.
- Weiner JM, Abramson MJ, Puy RM. Intranasal corticosteroids versus oral H1 receptor antagonists in allergic rhinitis: systematic review of randomised controlled trials. *BMJ*. 1998;317:1624–9.
- Yáñez A, Rodrigo GJ. Intranasal corticosteroids versus topical H1 receptor antagonists for the treatment of allergic rhinitis: a systematic review with meta-analysis. *Ann Allergy Asthma Immunol*. 2002;89:479–84.
- Meltzer EO, Bensch GW, Storms WW. New intranasal formulations for the treatment of allergic rhinitis. *Allergy Asthma Proc*. 2014;35 Suppl. 1.
- Higgins JPT, Altman DG, Sterne JAC. Assessing risk of bias in included studies. In: Higgins JPT, Green S. editors. *Cochrane handbook for systematic reviews of interventions* version 5.1.0.2011.
- Meltzer EO, Kunjibettu S, Hall N, Wingertzahn MA, Murcia C, Berger W, et al. Efficacy and safety of ciclesonide, 200 microg once daily, for the treatment of perennial allergic rhinitis. *Ann Allergy Asthma Immunol*. 2007;98:175–81.
- Chervinsky P, Kunjibettu S, Miller DL, Prenner BM, Raphael G, Hall N, et al. Long-term safety and efficacy of intranasal ciclesonide in adult and adolescent patients with perennial allergic rhinitis. *Ann Allergy Asthma Immunol*. 2007;99:69–76.
- Berger WE, Mohar DE, LaForce C, Raphael G, Desai SY, Huang H, et al. A 26 week tolerability study of ciclesonide nasal aerosol

- in patients with perennial allergic rhinitis. *Am J Rhinol Allergy*. 2012;26:302–7.
16. Kim K, Weiswasser M, Nave R, Ratner P, Nayak A, Herron J, et al. Safety of once-daily ciclesonide nasal spray in children 2 to 5 years of age with perennial allergic rhinitis. *Pediatr Asthma Allergy Immunol*. 2007;20:229–42.
 17. Berger WE, Nayak A, Lanier BQ, Kaiser HB, Laforce C, Darken P, et al. Efficacy and safety of once-daily ciclesonide nasal spray in children with allergic rhinitis. *Pediatr Asthma Allergy Immunol*. 2008;21:73–82.
 18. Eifan AO, Durham SR. Pathogenesis of rhinitis. *Clin Exp Allergy*. 2016;46:1139–51.
 19. Jacobs RL. Ciclesonide for the treatment of seasonal allergic rhinitis. *Exp Rev Clin Immunol*. 2011;7:735–41.
 20. Passalacqua G, Ciprandi G. Novel therapeutic interventions for allergic rhinitis. *Exp Opin Invest Drugs*. 2006;15:1615–25.
 21. Juniper EF, Guyatt GH. Development and testing of a new measure of health status for clinical trials in rhinoconjunctivitis. *Clin Exp Allergy*. 1991;21:77–83.
 22. Salib RJ, Howarth PH. Safety and tolerability profiles of intranasal antihistamines and intranasal corticosteroids in the treatment of allergic rhinitis. *Drug Saf*. 2003;26:863–93.
 23. Rohatagi S, Luo Y, Shen L, Guo Z, Schemm C, Huang Y, et al. Protein binding and its potential for eliciting minimal systemic side effects with a novel inhaled corticosteroid, ciclesonide. *Am J Ther*. 2005;12:201–9.
 24. Rohatagi S, Krishnaswami S, Pfister M, Sahasranaman S. Model-based covariate pharmacokinetic analysis and lack of cortisol suppression by the new inhaled corticosteroid ciclesonide using a novel cortisol release model. *Am J Ther*. 2005;12:385–97.
 25. Dietzel K, Engelstätter R, Keller A. Ciclesonide: an on-site-activated steroid. Karger; 2004. p. 31.