



Endogenous opioid signalling in the brain during pregnancy and lactation

Paula J. Brunton^{1,2} 

Received: 11 September 2018 / Accepted: 10 October 2018 / Published online: 10 November 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

During pregnancy, the regulation of several neuroendocrine systems is altered to support the pregnancy and facilitate the transition to motherhood. These adaptations are organised by the mother's brain and include those that serve to optimise foetal growth, protect the foetus(es) from adverse prenatal programming by maternal stress, facilitate timely parturition and ensure the offspring are nourished and cared for after birth. Although pregnancy hormones are important in inducing and maintaining many of these adaptations, their effects are often mediated via interactions with neuropeptide systems in the brain. In particular, endogenous opioids in the maternal brain play key roles in altered regulation of the stress axis, the oxytocin system, the prolactin system and the neural circuits controlling maternal behaviour. Together, these adaptations maximise the likelihood of a successful pregnancy outcome.

Keywords Enkephalin · HPA axis · Oxytocin · Prolactin · Maternal behaviour

Introduction

During pregnancy, altered neuropeptide signalling in the maternal brain allows physiological adaptations that maximise the likelihood of a successful pregnancy outcome. Several neuroendocrine systems are modified in pregnancy. For example, oxytocin stores in the posterior pituitary are enlarged so there are sufficient reserves to meet the demand at parturition and during lactation. At the same time, oxytocin neurones have to be kept in check to prevent premature activation and minimise the risk of preterm birth. Exposure of the developing foetus to maternal stress hormones can lead to intra-uterine growth restriction and have long-term adverse effects on the foetal brain; hence, maternal stress responses are restrained in pregnancy to conserve energy and protect the foetus. The central control of prolactin secretion is altered so a state of hyperprolactinemia can develop to support milk production

and secretion, to sustain lactation and nourish the offspring. To facilitate the transition to motherhood, the neural circuits governing maternal behaviour are primed to ensure the rapid expression of appropriate maternal care after birth. These adaptations are driven primarily by the actions of pregnancy hormones on the brain; however, endogenous opioid mechanisms play important regulatory roles in these systems. Here, focusing on the rodent literature, we consider the contribution of endogenous opioids in mediating critical changes in the maternal brain that support the pregnancy and prepare females for birth and motherhood.

Endogenous opioids

There are three families of endogenous opioids—the enkephalins, dynorphins and endorphins—each derived from different precursors, proenkephalin (pENK), prodynorphin (pDYN) and proopiomelanocortin (POMC), respectively (Drolet et al. 2001) and three main types of opioid receptor, referred to as mu (μ), delta (δ) and kappa (κ) (Mansour et al. 1994a). Enkephalins bind to μ - and δ -receptors, dynorphins to κ -receptors and endorphins to μ - and κ -receptors (Russell and Douglas 2000). Endogenous opioids and their receptors are widely distributed throughout the CNS (Le Merrer et al. 2009) where they play an important role in regulating responses to

✉ Paula J. Brunton
p.j.brunton@ed.ac.uk

¹ Centre for Discovery Brain Sciences, University of Edinburgh, Hugh Robson Building, George Square, Edinburgh, Scotland EH8 9XD, UK

² Zhejiang University–University of Edinburgh Institute, Haining, Zhejiang, People's Republic of China

stress and modulating pain, mood and emotionality, food intake, reward and attachment (Bali et al. 2015; Benarroch 2012; Bodnar 2011; Drolet et al. 2001). During pregnancy, endogenous opioids play key roles in promoting adaptations in the maternal brain, particularly in the arcuate, paraventricular and supraoptic nuclei (Fig. 1), which ultimately serve to maximise the likelihood of a successful pregnancy outcome.

The HPA axis

The hypothalamo-pituitary-adrenal (HPA) axis is the major neuroendocrine system that responds to stress. Stress exposure results in activation of a population of neurosecretory neurones in the parvocellular region of the paraventricular nucleus (pPVN) in the hypothalamus, which synthesise two neuropeptides: corticotropin-releasing hormone (CRH) and arginine vasopressin (AVP). CRH and AVP are released from nerve terminals at the median eminence into the hypophysial portal blood system and act synergistically on anterior pituitary corticotrophs to stimulate ACTH secretion, which in turn triggers secretion of glucocorticoids from the adrenal cortex. Glucocorticoids play a vital role in restoring homeostasis in the face of stress by mediating appropriate behavioural and metabolic adaptations and also switch off the HPA axis, via a negative feedback loop, when the stress no longer poses a threat.

Attenuated HPA axis responses in pregnancy

In late-pregnant rodents, the HPA axis is less responsive to a host of psychological and physical stressors, such as exposure to a novel environment, air-puff, the elevated plus maze, noise, social stress, forced swimming, immune challenge, as well as to stimulation by metabolic peptides or CRH; and this hyporesponsive state is maintained through parturition and lactation, until weaning (Brunton et al. 2006a; Brunton et al. 2005; Brunton and Russell 2003; Brunton and Russell 2010; da Costa et al. 1997; Douglas et al. 2003; Ma et al. 2005; Neumann et al. 2003; Neumann et al. 1998; Shanks et al. 1999; Toufexis et al. 1999; Windle et al. 1997). Hence, stress-induced ACTH and corticosterone secretion is markedly reduced or completely suppressed in late pregnancy and during lactation, as a result of adaptations at the level of both the anterior pituitary and the brain, though those in the hypothalamus seem to be of key importance (Brunton 2010). For example, stress is less effective in activating the parvocellular CRH/AVP neurones in the PVN, meaning activity of the HPA axis is restrained at the apex of the system (Brunton et al. 2005; da Costa et al. 2001; da Costa et al. 1996; Lightman and Young 1989). Importantly, this phenomenon of reduced glucocorticoid responses to stress is also observed in pregnant women (Kammerer et al. 2002; Schulte et al. 1990). It is

thought that attenuated HPA axis responses to stress in pregnancy function to conserve energy supplies for the pregnancy and minimise exposure of the foetus(es) to excessive levels of maternal glucocorticoids, which can have long-term adverse programming effects (Maccari et al. 2014).

Endogenous opioids and regulation of the HPA axis

Opioid-synthesising neurones are present in stress-related regions of the brain, including the cerebral cortex, amygdala, lateral septum, bed nucleus of the stria terminalis (BNST), preoptic area, hypothalamus, nucleus tractus solitarius (NTS) and ventrolateral medulla (VLM) (Harlan et al. 1987; Khachaturian et al. 1983). Several studies provide evidence for endogenous opioids in modulating behavioural and HPA axis responses to stress (Barfield et al. 2010; Brunton et al. 2005; Buckingham and Cooper 1986; Buller et al. 2005; Douglas et al. 1998a; Drolet et al. 2001; Grisel et al. 2008; Sanders et al. 2005) and acute stress results in an increase in gene expression for enkephalin in brain regions involved in stress processing, including the PVN and the brainstem (Ceccatelli and Orazzo 1993; Harbuz et al. 1994; Harbuz and Lightman 1989; Mansi et al. 2000). The effects of endogenous opioids in the brain on stress-related behaviours and neuroendocrine stress responses are complex and region- and opioid-receptor type-dependent. For example, there is evidence that endogenous endorphins and enkephalins aid stress-coping, particularly under chronic stress conditions and have anxiolytic actions, whereas dynorphin has been demonstrated to have anxiogenic actions (Barfield et al. 2010; Berube et al. 2013; Berube et al. 2014; Grisel et al. 2008; Henry et al. 2018; Kang et al. 2000; Katoh et al. 1990; Konig et al. 1996), though these effects depend on the site of action in the brain.

In male rats (and non-pregnant females), administration of an exogenous opioid receptor agonist (e.g., morphine) enhances, while an opioid receptor antagonist (naloxone) suppresses the HPA axis responses to acute stress (Buckingham and Cooper 1984, 1986; Buller et al. 2005). This is evidently the result of a central action, as morphine, met-enkephalin and leu-enkephalin stimulate CRH secretion from hypothalamic preparations but not ACTH release from the pituitary (Buckingham and Cooper 1984) and centrally administered naloxone reduces stress-induced activation of CRH neurones in the pPVN (Buller et al. 2005). However, in pregnancy, endogenous opioids switch to having a net inhibitory role over HPA axis function.

Endogenous opioids and HPA axis regulation during late pregnancy and parturition

Attenuated HPA axis responses to stress in pregnancy are a consequence of the emergence of a central inhibitory opioid




Fig. 1 Anatomical distribution of opioid peptides and receptors in the arcuate, paraventricular and supraoptic nuclei of the rat hypothalamus. The left-hand side of the diagram summarises the distribution of immunoreactivity for dynorphin (dyn), enkephalin (enk) and β -endorphin (β end) in cell bodies (stars) or in neuronal fibres (circles). The right-hand side shows the distribution pattern for mu (μ)-, delta (δ)- and kappa (κ)-opioid receptors. For cell bodies (stars), receptors were identified by in situ hybridisation; while receptor binding (triangles) was identified by autoradiography. In both cases, the relative expression levels are

indicated by the number of a given symbol i.e., one symbol = low; 2 symbols = moderate; and 3 symbols = high expression. 3V, 3rd ventricle; Arc, arcuate nucleus; OT, optic tract; PVN, paraventricular nucleus; SON, supraoptic nucleus. Based on data from Abe et al. (1987), Beaulieu et al. (1996), Desjardins et al. (1990), Fallon and Leslie (1986), Khachaturian et al. (1983), Khachaturian et al. (1982), Mansour et al. (1994a), Mansour et al. (1994b), Mansour et al. (1994c), Mansour et al. (1987), Merchenthaler et al. (1986), Minami et al. (1994) and Wamsley et al. (1980)

mechanism. Hence, systemic administration of naloxone reinstates HPA axis responses to stressful stimuli, such as forced swimming and immune challenge, in late-pregnant rats (Brunton et al. 2005; Douglas et al. 1998a) and to parturition-related stimuli at birth (Wigger et al. 1999), reflected by increased ACTH and corticosterone secretion, together with increased activation of pPVN neurones and increased gene expression for *Crh* in the pPVN (Brunton et al. 2005).

In late pregnancy, gene expression for pENK-A (*Penk*) and μ -opioid receptor (MOR; *Oprm1*) are up-regulated in the NTS of the brainstem and *Pomc* mRNA and β -endorphin immunoreactivity is increased in the arcuate nucleus compared with non-pregnant rats (Brunton et al. 2005; Douglas et al. 2002). However, it is likely that the NTS is the source of the opioids that are involved in restraining HPA axis responses in late-pregnant rats, at least for those stressors where activation of the CRH neurones is mediated via brainstem afferents. The evidence is as follows: immune challenge with the cytokine, interleukin-1 β (IL-1 β), activates hypothalamic-projecting noradrenergic neurones in the A2 region of the NTS. This results in increased noradrenaline release from nerve terminals in the PVN that stimulates CRH neurones and, hence, ACTH and corticosterone secretion. In late pregnancy, despite a similar activation of the NTS neurones, systemic IL-1 β fails to

result in increased noradrenaline release in the PVN. Similarly, noradrenaline release in the PVN evoked by forced swim stress is significantly lower in late-pregnant rats compared with virgin females (Douglas et al. 1995). Naloxone infused directly into the PVN restores IL-1 β -induced noradrenaline release in the PVN (Brunton et al. 2005), indicating that endogenous opioids act presynaptically on noradrenergic inputs to the PVN to inhibit evoked noradrenaline release. If the noradrenergic NTS neurones are the same population that display increased gene expression for MORs and enkephalin in pregnancy, this would provide a mechanism whereby endogenous opioids (i.e., enkephalin) act on up-regulated presynaptic MORs on noradrenergic nerve terminals to auto-inhibit noradrenaline release in the PVN (Fig. 2), hence reducing stimulation of the CRH neurones in late pregnancy (Brunton et al. 2005). The same mechanism is hypothesised to limit activation of the magnocellular oxytocin neurones in late pregnancy (discussed below) (Brunton et al. 2006b; Leng et al. 1995).

A reduction in the effectiveness of the noradrenergic input to the pPVN CRH neurones has also been implicated in underlying reduced HPA axis responses to stress during lactation (Toufexis et al. 1998; Toufexis and Walker 1996); though the precise mechanisms are not known. However, in contrast to pregnancy and parturition, HPA axis hyporesponsiveness




Fig. 2 Endogenous opioid inhibition of CRH/oxytocin neurones in late pregnancy. Noradrenergic (NAergic) A2 neurones in the nucleus tractus solitarius (NTS) of the brainstem project to parvocellular corticotropin-releasing hormone (CRH) and magnocellular oxytocin (OT) neurones in the paraventricular nucleus (PVN). In non-pregnant rats, stress (e.g., interleukin-1 β , forced swimming) activates these noradrenergic A2 neurones triggering noradrenaline (NA) release from the nerve terminals in the PVN. Noradrenaline acts via $\alpha 1$ adrenoceptors ($\alpha 1R$) to stimulate (+) the CRH/OT neurones, resulting in CRH release at the median eminence (ME) and OT release from the posterior pituitary (PP). In late pregnancy, systemic interleukin-1 β fails to stimulate noradrenaline release in the PVN (despite similar activation of the noradrenergic cell bodies in the NTS); hence, the CRH and OT neurones are less excited and ACTH/corticoesterone and OT secretion is markedly suppressed (\downarrow) or completely absent. This is a consequence of increased inhibition (\ominus) by a central endogenous opioid mechanism. It is proposed that enkephalin (enk) acts presynaptically on up-regulated MOR on the noradrenergic terminals to inhibit stimulated noradrenaline release, since the NTS A2 neurones display increased (\uparrow) expression of proenkephalin-A and MOR mRNA in late pregnancy and infusion of the opioid receptor antagonist naloxone directly into the PVN restores IL-1 β -evoked noradrenaline release in the PVN. Moreover, naloxone activates the CRH and oxytocin neurones leading to increased ACTH, corticoesterone and oxytocin secretion into the blood

during lactation is dependent upon the suckling stimulus provided by the pups (Lightman and Young 1989) and is likely to involve central prolactin actions (Torner and Neumann 2002; Torner et al. 2002).

Oxytocin

Oxytocin is synthesised by magnocellular neurones in the PVN and the supraoptic nuclei (SON), whose axons project to the posterior pituitary gland (Russell and Brunton 2009). Upon activation, oxytocin is secreted from the nerve terminals in the posterior pituitary into the systemic circulation. Oxytocin plays an important role in promoting uterine contractions during birth and is vital in stimulating milk ejection during lactation (for review, see Burbach et al. 2006). Oxytocin is also released within the brain from the dendrites of the magnocellular neurones and also from the terminals of centrally projecting neurones, where it plays an important role in promoting the expression of maternal behaviour after birth (Knobloch et al. 2012; Lee et al. 2009; Ludwig and Leng 2006; Pedersen et al. 1994). In the rat, oxytocin is also considered to be a “stress hormone”; hence, stress exposure activates oxytocin neurones (reflected by increased expression of the protein product of the immediate early gene, c-fos) and stimulates oxytocin secretion (Brunton et al. 2006b; Douglas et al. 2000; Lang et al. 1983).

During pregnancy, the oxytocin neurones are maintained in a quiescent state that limits oxytocin secretion and permits the accumulation of oxytocin stores in the posterior pituitary (which expand by around 75% in rats) (Douglas et al. 1993b; Fuchs and Saito 1971), for when it is required at parturition and subsequent lactation. Accruing these stores is important as oxytocin synthesis is evidently not increased in pregnancy (Douglas et al. 1998b). Critically, limiting activity of the oxytocin neurones in pregnancy also minimises the likelihood of premature oxytocin secretion that could lead to uterine contractions and preterm birth (Elovitz and Mrinalini 2004; Goodwin et al. 1994). There are several mechanisms that contribute to restraining oxytocin secretion during pregnancy (Brunton et al. 2014). These include auto-inhibition by nitric oxide (Srisawat et al. 2000) and the potentiating effects of allopregnanolone on GABA actions at GABA_A receptors (Brussaard and Herbison 2000); however, here, the focus is on endogenous opioid mechanisms.

Oxytocin and parturition

A2 noradrenergic neurones in the NTS of the brainstem project to magnocellular oxytocin neurones in the SON and PVN (Ericsson et al. 1994; Meddle et al. 2000). At parturition,

noradrenaline release is increased in the SON and PVN and acts on $\alpha 1$ adrenoceptors to activate the oxytocin neurones (Douglas et al. 1998b; Douglas et al. 2001; Herbison et al. 1997; Meddle et al. 2000; Randle et al. 1986). Activated oxytocin neurones fire in synchronised bursts, secreting pulses of oxytocin into the blood, which acts on myometrium oxytocin receptors (OTR), up-regulated by increased levels of estradiol acting on oestrogen receptor- α (ER α) (Mesiano et al. 2002; Murata et al. 2000; Murata et al. 2003; Welsh et al. 2012), to stimulate uterine contractions. Signals from the contracting uterus and distended birth canal are relayed by spinal and vagal afferents (Higuchi et al. 1986; Peters et al. 1987) to the A2 noradrenergic NTS neurones, resulting in a positive feedback loop that drives further oxytocin release, known as the Ferguson reflex (Ferguson 1941). Oxytocin release from the dendrites of activated magnocellular neurones acts locally to synchronise burst firing of the oxytocin neurones, necessary for pulsatile oxytocin secretion at parturition and during milk ejection (Dyball and Leng 1986; Moos et al. 1998; Rossoni et al. 2008).

Endogenous opioids and control of oxytocin secretion

Oxytocin secretion is restrained by endogenous opioids. Opioids act both centrally, on the oxytocin neurones or their inputs, as well as on axon terminals in the posterior pituitary (Bicknell et al. 1988; Zhao et al. 1988a, b). In non-pregnant rats and during early pregnancy, endogenous opioids acting via κ -opioid receptors on axon terminals in the posterior pituitary inhibit oxytocin secretion and this is the principal mechanism involved in regulating oxytocin secretion (Douglas et al. 1993b; Leng et al. 1994). However, during the last week of pregnancy, a κ -agonist is less effective in inhibiting electrically stimulated oxytocin release from isolated neural lobes, indicating a desensitisation of this inhibitory opioid mechanism in the pituitary (Douglas et al. 1993b), allowing for enhanced oxytocin secretion at parturition in response to excitation of the oxytocin cell bodies. At this time, a central inhibitory opioid system becomes activated (described in the subsequent text) and this mechanism predominates in restraining activity of the oxytocin neurones in response to excitatory stimuli in late pregnancy (Douglas et al. 1993b; Douglas et al. 1995).

Stress-induced oxytocin responses in pregnancy

In non-pregnant rats, in addition to activating the HPA axis, systemic IL-1 β also activates oxytocin neurones, increasing their firing rate and resulting in increased oxytocin secretion into the blood (Brunton et al. 2012; Brunton et al. 2006b). In contrast, IL-1 β has no effect on the electrical or secretory activity of oxytocin neurones in late-pregnant rats (Brunton

et al. 2012; Brunton et al. 2006b). This is the result of a central endogenous opioid mechanism, since naloxone increases the firing rate of oxytocin neurones and unmasks a much exaggerated oxytocin secretory response to IL-1 β in late-pregnant rats (Brunton et al. 2006b). A similar response is seen following forced swimming or CCK administration, where oxytocin secretion, Fos induction in the SON and the firing rate of SON oxytocin neurones are potentiated by naloxone to a far greater extent in late-pregnant rats compared with virgin females (Douglas et al. 1998a; Douglas et al. 1995). Forced swimming, CCK and IL-1 β mediate their excitatory effect on oxytocin neurones via noradrenergic neurones in the NTS. As mentioned previously, in late pregnancy, these stimuli fail to trigger noradrenaline release in the PVN (Brunton et al. 2005; Douglas et al. 2005). It is proposed that the same endogenous opioid mechanism that prevents excitation of the CRH neurones in the PVN (described previously) by inhibiting noradrenaline release from afferent inputs to the PVN (mediated by enkephalin actions on presynaptic MORs) also serves to keep the oxytocin neurones in the SON and PVN under control in late pregnancy (Fig. 2). Given that A2 noradrenergic NTS neurones also relay stimuli from the birth canal to drive oxytocin secretion at parturition, it is hypothesised that maintaining quiescence of this pathway during pregnancy plays an important role in preventing premature activation of the oxytocin neurones and in ensuring birth is not triggered prematurely.

While this inhibitory opioid mechanism is anticipated to prevent inappropriate activation of the oxytocin neurones by stimuli such as stress in late pregnancy, central opioid inhibition of the oxytocin neurones persists during parturition (Leng et al. 1988; Neumann et al. 1991) but it is withdrawn soon after birth (Leng et al. 1988) and is not detected in lactation (Douglas et al. 1993b; Neumann et al. 1993). μ - and κ -receptor agonists inhibit the electrical activity of oxytocin neurones, inhibit oxytocin secretion and delay established parturition (Bicknell et al. 1988; Douglas et al. 1993a; Leng et al. 1988; Pumford et al. 1991; Pumford et al. 1993; Russell et al. 1989). Conversely, naloxone administration during parturition increases oxytocin secretion and reduces the interval between pup births (Leng et al. 1988). Similarly, naloxone can prevent the reduction in oxytocin secretion and the interruption to parturition induced by stress (Leng et al. 1988). Together, these data indicate potent opioid inhibition over oxytocin neurones at parturition, which may serve to regulate the timing of births so inter-birth intervals are optimal to allow for appropriate maternal behaviour between deliveries.

Endogenous opioids also restrain oxytocin release within the SON and PVN in late pregnancy; however, this mechanism is down-regulated at parturition (Douglas et al. 1995; Neumann et al. 1993). Given local oxytocin release within the magnocellular nuclei plays an important role in positive

feedback regulation of oxytocin neuronal activity (Neumann et al. 1996), withdrawal of this opioid inhibition permits potent excitation of the oxytocin neurones and promotes synchronised burst firing of oxytocin neurones during parturition and suckling for effective uterine contractions and milk ejection, respectively.

Induction of inhibitory endogenous opioid tone over HPA axis and oxytocin responses in pregnancy

The endogenous opioid mechanism that restrains activity of the HPA axis and magnocellular oxytocin system in pregnancy is induced by the neuroactive metabolite of progesterone, allopregnanolone (Brunton et al. 2012; Brunton et al. 2009). Allopregnanolone concentrations progressively increase in the blood and in the brain during pregnancy (Concas et al. 1998). In the brain, allopregnanolone concentrations reach maximal levels in late pregnancy, before declining dramatically before parturition (Concas et al. 1998). Gene expression for 5 α -reductase (*Srd5a*) and 3 α -hydroxysteroid dehydrogenase (*Akr1c4*), the enzymes responsible for the conversion of progesterone to dihydroprogesterone and then allopregnanolone, are up-regulated in the NTS in late pregnancy (Brunton et al. 2009). The activity of these enzymes is also increased in the hypothalamus of late pregnant rats (Brunton et al. 2009), presumably leading to increased local allopregnanolone synthesis. Blocking allopregnanolone synthesis in pregnant rats with the 5 α -reductase inhibitor, finasteride, not only restores ACTH and oxytocin secretory responses to immune challenge but also reduces *Penk* gene expression in the NTS to levels similar to those found in non-pregnant animals (Brunton et al. 2012; Brunton et al. 2009). Moreover, administration of allopregnanolone to virgin rats (to simulate pregnancy levels) rapidly up-regulates *Penk* gene expression in the NTS to levels equivalent to those observed in late gestation and induces opioid inhibition over stress-induced ACTH and oxytocin secretion (Brunton et al. 2012; Brunton et al. 2009). Hence, increased concentrations of allopregnanolone in the NTS in pregnancy evidently induce and maintain an inhibitory opioid mechanism that tightly controls both the oxytocin and the CRH neurones. This is important since inhibiting 5 α -reductase activity (and hence allopregnanolone production) during the last week of gestation leads to preterm birth and increased neonatal mortality (Paris et al. 2011). It remains to be established whether this is a result of premature activation of the oxytocin neurones as a result of withdrawal of endogenous opioid tone or is due to altered GABA signalling to oxytocin neurones, since allopregnanolone has also been demonstrated to potentiate the inhibitory influence of GABA on oxytocin neurones in late pregnancy (Koksmas et al. 2003).

Mimicking pregnancy levels of sex steroids in virgin females with chronic estradiol and progesterone administration induces opioid inhibition over the neurohypophysial oxytocin

system but not the HPA axis (Douglas et al. 2000). Hence, naloxone administration potentiates stress-induced oxytocin secretion but not ACTH or corticosterone in these rats (Douglas et al. 2000). It is unclear why this treatment induces opioid tone over the oxytocin system but not the HPA axis. It is not known whether exogenously administered progesterone is converted to allopregnanolone to exert these effects, though this seems unlikely as this would be expected to result in similar effects on both systems (Brunton et al. 2012; Brunton et al. 2009). Rather, it is possible that the sex steroid treatment expands pituitary stores of oxytocin (as in pregnancy), meaning a greater pool is available for release when opioid restraint is blocked by naloxone.

Central control of prolactin secretion

In non-pregnant animals, prolactin secretion by anterior pituitary lactotrophs is under tonic inhibitory control by dopamine, produced and secreted by hypothalamic tuberoinfundibular dopamine (TIDA) neurones in the arcuate nucleus neurones that project to the median eminence and secrete dopamine into the hypothalamo-hypophysial portal blood system (Reymond and Porter 1985). In the anterior pituitary, dopamine acts on D2 receptors expressed on lactotrophs to inhibit prolactin production (Caron et al. 1978; Guillou et al. 2015; Mansour et al. 1990). Prolactin can enter the brain (Walsh et al. 1987); however, the process by which it crosses the blood-brain barrier is unclear (Augustine et al. 2003), although, evidently, this does not depend upon prolactin receptors (PRL-R) (Brown et al. 2016). Once in the brain, prolactin acts directly on TIDA neurones, which express PRL-Rs, to stimulate activity of the TIDA neurones and up-regulate tyrosine hydroxylase (*Th*; the rate-limiting enzyme in dopamine synthesis) gene expression in TIDA neurones (Arbogast and Voogt 1991; Lerant and Freeman 1998). This results in increased dopamine synthesis and secretion into the portal blood generating a short-loop negative-feedback system whereby prolactin controls its own secretion (Ben-Jonathan et al. 1980).

Prolactin secretion during pregnancy

In the rat, prolactin plays an essential role in preventing regression of the corpus luteum (luteolysis) in early pregnancy, ensuring progesterone secretion to sustain the pregnancy and in preparing the mammary alveoli for milk production (Gibori and Richards 1978; Ormandy et al. 1997). In addition, prolactin (together with oestrogen, progesterone and oxytocin) primes the brain to promote maternal behaviour after birth and plays an essential role in lactation by stimulating milk production and secretion (Bridges et al. 1985; Bridges et al. 1990; Bridges and Ronsheim 1990; Lucas et al. 1998;

Ormandy et al. 1997). Moreover, increased prolactin in the maternal brain in the perinatal period contributes to restraining basal and stress-induced activity of the HPA axis and oxytocin system (Gustafson et al. 2017; Torner et al. 2002). Accordingly, in pregnancy, there are important adaptations in the hypothalamic mechanisms controlling the prolactin system that must occur in order to facilitate increased prolactin secretion.


In rodents, mating stimulates the onset of twice daily surges in prolactin secretion, which persist for the first 8–10 days of pregnancy (Gunnert and Freeman 1983). Around mid-pregnancy, these surges are suppressed when the secretion of placental lactogen begins to increase, which acts in a similar fashion to prolactin to stimulate dopamine release from TIDA neurones and thereby inhibit prolactin secretion from the lactotrophs (Demarest et al. 1983a; Lee and Voogt 1999; Smith and Neill 1976; Voogt et al. 1982; Voogt 1984). The night before parturition and following progesterone withdrawal, the TIDA neurones become seemingly insensitive to negative feedback by prolactin and placental lactogen—despite increasing their firing rate, they fail to secrete dopamine in response to prolactin (Anderson et al. 2006a; Romano et al. 2013) and this persists into lactation. This ineffective negative feedback permits a large nocturnal surge in prolactin secretion at the end of pregnancy, facilitating the transition from gestation to motherhood and allowing a state of hyperprolactinemia to develop to sustain lactation (Grattan and Averill 1990; Grattan et al. 2008). The mechanism involved is not fully understood but it is evidently induced by progesterone withdrawal (Grattan and Averill 1990; Soaje et al. 2006) and probably involves changes in prolactin receptor intracellular signalling and reduced TH activity (Anderson et al. 2006a; Anderson et al. 2006b; Steyn et al. 2008). Following parturition, suckling by the pups reduces activity of the TIDA neurones, down-regulates *Th* gene expression and suppresses dopamine release at the median eminence, hence increasing prolactin secretion until weaning when tonic inhibition of prolactin secretion by TIDA neurones is reinstated (Ben-Jonathan et al. 1980; Demarest et al. 1983b; Feher et al. 2010; Selmanoff and Gregerson 1985; Selmanoff and Wise 1981; Wang et al. 1993).

Endogenous opioids and regulation of prolactin secretion during pregnancy and lactation

Endogenous opioids play a key role in regulating prolactin secretion by the pituitary through actions on TIDA neurones. Endogenous opioid peptides inhibit TIDA neurones and, hence, dopamine secretion, promoting the prolactin surges in early pregnancy and pre-partum and suckling-induced prolactin secretion during lactation (Andrews and Grattan 2002, 2003; Callahan et al. 2000; Sagrillo and Voogt 1991). Hence, naloxone administration during the night before

parturition increases dopamine turnover and inhibits the pre-partum prolactin surge (Andrews and Grattan 2002). Interestingly, treatment with naloxone a couple of days earlier in pregnancy (on day 19) has no effect on prolactin secretion or dopaminergic activity in the mediobasal hypothalamus (Soaje et al. 2006), unless progesterone actions are blocked, which decreases TIDA neurone activity and permits an increase in prolactin secretion (Soaje et al. 2006), supporting the idea that progesterone withdrawal at the end of pregnancy plays a key role in reducing activity of the TIDA neurones and facilitating increased prolactin secretion (Andrews and Grattan 2002; Soaje and Deis 1994). However, on day 19 of pregnancy, despite facilitating increased prolactin secretion, naloxone has no greater effect on the activity of the dopamine neurones than the anti-progesterone given alone (Soaje et al. 2006), suggesting that in addition to the inhibition of TIDA neurones, endogenous opioids may regulate prolactin secretion through another mechanism in the presence of high levels of progesterone, possibly via inhibition of putative prolactin-releasing factor neurones. One such candidate is oxytocin, since oxytocin stimulates prolactin secretion (Gonzalez-Iglesias et al. 2015; Samson et al. 1986) and as described previously, oxytocin neurones are inhibited by endogenous opioids in late pregnancy (Brunton et al. 2006b; Douglas et al. 2000). Indeed, recent findings support a role for oxytocin in facilitating prolactin secretion in late pregnancy but this effect is only apparent when progesterone action is blocked (Villegas-Gabutti et al. 2018).

Immunoreactivity for dynorphin, enkephalin and β -endorphin is present in the arcuate nucleus (Fitzsimmons et al. 1992; Merchenthaler 1994) and contact between opioid peptide-containing nerve terminals and TIDA neurones has been described (Fitzsimmons et al. 1992; Horvath et al. 1992). However, β -endorphin innervation of dopaminergic neurones only represents a small proportion of opioidergic connections in the arcuate nucleus, with the majority expressing dynorphin or enkephalin (Fitzsimmons et al. 1992). During pregnancy and lactation, POMC mRNA expression and β -endorphin immunoreactivity are only modestly increased in the arcuate nucleus in late pregnancy (Douglas et al. 2002); however, immunoreactivity and gene expression for enkephalin are markedly up-regulated, such that all TIDA neurones co-express enkephalin (Merchenthaler 1994; Merchenthaler et al. 1995), which is likely driven by increased prolactin (Nahi and Arbogast 2003). Given the lack of δ -receptors in the arcuate nucleus (Mansour et al. 1994a; Mansour et al. 1987), it is reasonable to assume that enkephalin released from TIDA neurones acts on μ -opioid receptors in the arcuate nucleus, perhaps in an auto-inhibitory manner (Fig. 3). Indeed, the prolactin surges in early pregnancy, late pregnancy and lactation are inhibited and the activity of TIDA neurones increased, by administration of selective μ - and κ - but not δ -receptor



antagonists (Andrews and Grattan 2003), indicating these receptors play more important roles.

In late pregnancy, in addition to attenuating HPA axis and oxytocin responses to stress, endogenous opioids also restrain stress-induced prolactin secretion (Valdez et al. 2014). This

effect is mediated through μ -opioid receptors and estradiol and progesterone actions are involved in the induction and maintenance of this inhibitory opioid tone (Valdez et al. 2014). Hence, in pregnancy, endogenous opioids, through their inhibitory actions on TIDA neurones, play a key role in

◀ **Fig. 3 Endogenous opioid control of TIDA neurones in late pregnancy.** In virgin rats, prolactin secretion is under tonic inhibition (–) by dopamine (DA) secreted by tubero-infundibular dopamine (TIDA) neurones in the arcuate nucleus and secreted into the portal blood system to act on dopamine D2 receptors expressed on lactotrophs in the anterior pituitary. Prolactin acts via prolactin receptors (PRL-R) on TIDA neurons to stimulate (+) dopamine release, producing a short-loop negative-feedback system. Endogenous opioids inhibit TIDA neurones and hence stimulate prolactin secretion; this facilitates the surges in prolactin secretion in early pregnancy and pre-partum. Enkephalin (enk) is markedly up-regulated (↑) in arcuate nucleus TIDA neurones in late pregnancy and it is proposed that enkephalin released from TIDA neurones acts on μ - κ -opioid receptors to inhibit the dopamine neurones and hence dopamine release (possibly via auto-inhibition). Dynorphin (dyn)-containing nerve terminals also make contact with arcuate TIDA neurones and dynorphin acts via κ -opioid receptors to inhibit TIDA neurones. In addition, before parturition, the TIDA neurones become insensitive to negative feedback by prolactin and placental lactogen (PL). These adaptations permit a state of hyperprolactinemia to develop that supports lactation and promotes maternal behaviour

permitting a state of hyperprolactinemia, which maintains the pregnancy and facilitates lactation. In addition, prolactin actions on the brain also promote maternal behaviour post-partum (discussed in the subsequent section).

Maternal behaviour

Following the birth of the offspring, maternal behaviour must be expressed swiftly if the young are to be reared successfully. Maternal care consists of a complex set of behaviours including nest-building, retrieving and gathering the young, nursing, grooming and protection from the environment (e.g., cold) and other threats (e.g., unfamiliar conspecifics). The maternal brain is primed for this transition by prolonged exposure to pregnancy levels of ovarian and lactogenic hormones during gestation (for review, see Bridges 2015).

Neural circuitry and hormonal priming of maternal behaviour

The different elements of maternal behaviour involve distinct, though interconnected circuits in the brain (for review, see Numan 2007). Of particular importance in the onset and integration of maternal behavioural responses is the medial preoptic area (mPOA); hence, lesions to this region disrupt or abolish maternal behaviour (Cohn and Gerall 1989; Numan et al. 1977; Numan and Smith 1984; Stack et al. 2002; Tsuneoka et al. 2013). Hormone priming mediated by the decline in circulating progesterone at the end of pregnancy, concurrent with increased estradiol, activates neurones in the mPOA and induces maternal behaviour (Bridges et al. 1978; Sheehan and Numan 2002). Moreover, maternal behaviour can be induced in nulliparous rats treated with sex steroids to mimic

the concentrations and pattern of secretion seen in pregnancy (Bridges 1984). Activation of the mPOA and the neighbouring ventral BNST results in inhibition of neural circuits involving the anterior hypothalamic nucleus and the periaqueductal grey that normally mediate aversive responses to newborn pups and pup odour in virgin females, thus eliminating avoidance behaviour (Bridges et al. 1999; Numan and Numan 1996; Sukikara et al. 2006) while activation of dopaminergic neurones in the ventral tegmental area (VTA) that project to the nucleus accumbens (NAc) stimulate motivation and reward, making new mothers more responsive to pup-related stimuli (Champagne et al. 2004; Hansen et al. 1991; Keer and Stern 1999; Numan and Smith 1984; Parada et al. 2008; Stolzenberg et al. 2007).

Prolactin and oxytocin also play key roles in stimulating maternal care. Central prolactin infusions stimulate maternal behaviour towards foster pups in steroid-primed virgin rats and prolactin is more effective when administered directly into the MPOA (Bridges et al. 1990); while suppression of endogenous prolactin secretion with bromocriptine delays the onset of maternal behaviour (Bridges and Ronsheim 1990). Moreover, a recent study employing conditional PRL-R knockout mice demonstrated that prolactin action, specifically in the MPOA, is vital for the normal expression of maternal behaviour after birth (Brown et al. 2017). At parturition, oxytocin receptors are up-regulated in several brain regions linked to maternal behaviour, including in the MPOA, BNST and VTA (Meddle et al. 2007; Pedersen et al. 1994). Administration of an oxytocin receptor antagonist into the VTA or the MPOA of parturient rats impairs maternal behaviour (Pedersen et al. 1994) and centrally-administered exogenous oxytocin is able to induce maternal care in hormonally primed nulliparous rats (Pedersen et al. 1982).

Endogenous opioids and maternal behaviour

At parturition, the withdrawal of up-regulated central endogenous opioid mechanisms, in particular in the mPOA, is necessary to facilitate maternal behaviour (Byrnes and Bridges 2000; Byrnes et al. 2000). Levels of opioids and opioid receptor binding sites in the MPOA are elevated during pregnancy but decrease in lactation, evidently as a result of changes in ovarian sex steroids (Hammer Jr and Bridges 1987). Administration of morphine or the μ -receptor agonist, β -endorphin (but not δ - or κ -receptor agonists), into the MPOA disrupts maternal behaviour in lactating or hormone-primed-sensitised virgin rats (Mann and Bridges 1992; Mann et al. 1991; Rubin and Bridges 1984). The role of physiological levels of endogenous opioids in the maternal brain post-partum is unclear; however, there is evidence to indicate that they may act to regulate nursing bout duration, at least in early lactation (Byrnes et al. 2000) and are involved in regulating the switch from nursing behaviour to predatory/hunting behaviour (Sukikara et al. 2007).

Large-scale gene expression studies of the MPOA, medial prefrontal cortex, lateral septum and NAc have identified over 1000 genes/region that are differentially expressed in the maternal brain compared with non-pregnant females (Driessen et al. 2014; Eisinger et al. 2014; Eisinger et al. 2013; Zhao et al. 2014). As well as identifying potential new candidate genes involved in regulating maternal behaviour, studies such as these are paving the way for exploring the networks and interactions between novel genes and the already well-characterised “maternal” genes. Post-partum dams find pups rewarding (Ferris et al. 2005; Lee et al. 2000); therefore, it is particularly interesting that the top 700 of the identified “maternal genes” are significantly enriched for genes involved in reward and addiction, including genes for the opioid peptides, *Penk* and prodynorphin (*Pdyn*) and *Oprm1* (Gammie et al. 2016), further supporting the hypothesis that endogenous opioid mechanisms play a role in maternal reward and reinforcement (Nelson and Panksepp 1998).

Summary

In summary, endogenous opioids play key roles in mediating adaptations in the maternal brain. Here, the focus was on adaptations in the regulation of the HPA axis, oxytocin system, prolactin system and the circuits governing maternal behaviour. These changes are important for conserving energy, protecting the foetus from harmful levels of maternal glucocorticoids, providing sufficient oxytocin for parturition and lactation, ensuring that the onset of parturition occurs in a timely manner and that after birth the offspring are suckled and properly cared for. Appropriate withdrawal of up-regulated central endogenous opioid mechanisms is equally important to support the transition from pregnancy to motherhood.

Acknowledgements The research performed in the laboratory of PJB described here was funded by the Biotechnology and Biological Sciences Research Council.

Compliance with ethical standards

Conflict of interest The author declares that she has no conflict of interest.

References

- Abe J, Okamura H, Makino S, Yanaihara N, Iyata Y (1987) Immunocytochemical distribution of [Met]enkephalin-Arg-Gly-Leu immunoreactivity in the rat diencephalon. *Brain Res Bull* 19: 735–741
- Anderson GM, Beijer P, Bang AS, Fenwick MA, Bunn SJ, Grattan DR (2006a) Suppression of prolactin-induced signal transducer and activator of transcription 5b signaling and induction of suppressors of cytokine signaling messenger ribonucleic acid in the hypothalamic arcuate nucleus of the rat during late pregnancy and lactation. *Endocrinology* 147:4996–5005
- Anderson ST, Barclay JL, Fanning KJ, Kusters DH, Waters MJ, Curlewis JD (2006b) Mechanisms underlying the diminished sensitivity to prolactin negative feedback during lactation: reduced STAT5 signaling and up-regulation of cytokine-inducible SH2 domain-containing protein (CIS) expression in tuberoinfundibular dopaminergic neurons. *Endocrinology* 147:1195–1202
- Andrews ZB, Grattan DR (2002) Opioid control of prolactin secretion in late pregnant rats is mediated by tuberoinfundibular dopamine neurons. *Neurosci Lett* 328:60–64
- Andrews ZB, Grattan DR (2003) Opioid receptor subtypes involved in the regulation of prolactin secretion during pregnancy and lactation. *J Neuroendocrinol* 15:227–236
- Arbogast LA, Voogt JL (1991) Hyperprolactinemia increases and hypoprolactinemia decreases tyrosine hydroxylase messenger ribonucleic acid levels in the arcuate nuclei, but not the substantia nigra or zona incerta. *Endocrinology* 128:997–1005
- Augustine RA, Kokay IC, Andrews ZB, Ladyman SR, Grattan DR (2003) Quantitation of prolactin receptor mRNA in the maternal rat brain during pregnancy and lactation. *J Mol Endocrinol* 31: 221–232
- Bali A, Randhawa PK, Jaggi AS (2015) Stress and opioids: role of opioids in modulating stress-related behavior and effect of stress on morphine conditioned place preference. *Neurosci Biobehav Rev* 51:138–150
- Barfield ET, Barry SM, Hodgins HB, Thompson BM, Allen SS, Grisel JE (2010) Beta-endorphin mediates behavioral despair and the effect of ethanol on the tail suspension test in mice. *Alcohol Clin Exp Res* 34: 1066–1072
- Beaulieu J, Champagne D, Drolet G (1996) Enkephalin innervation of the paraventricular nucleus of the hypothalamus: distribution of fibers and origins of input. *J Chem Neuroanat* 10:79–92
- Benarroch EE (2012) Endogenous opioid systems: current concepts and clinical correlations. *Neurology* 79:807–814
- Ben-Jonathan N, Neill MA, Arbogast LA, Peters LL, Hoefler MT (1980) Dopamine in hypophysial portal blood: relationship to circulating prolactin in pregnant and lactating rats. *Endocrinology* 106:690–696
- Berube P, Laforest S, Bhatnagar S, Drolet G (2013) Enkephalin and dynorphin mRNA expression are associated with resilience or vulnerability to chronic social defeat stress. *Physiol Behav* 122:237–245
- Berube P, Poulin JF, Laforest S, Drolet G (2014) Enkephalin knockdown in the basolateral amygdala reproduces vulnerable anxiety-like responses to chronic unpredictable stress. *Neuropsychopharmacology* 39:1159–1168
- Bicknell RJ, Leng G, Russell JA, Dyer RG, Mansfield S, Zhao BG (1988) Hypothalamic opioid mechanisms controlling oxytocin neurons during parturition. *Brain Res Bull* 20:743–749
- Bodnar RJ (2011) Endogenous opiates and behavior: 2010. *Peptides* 32: 2522–2552
- Bridges RS (1984) A quantitative analysis of the roles of dosage, sequence, and duration of estradiol and progesterone exposure in the regulation of maternal behavior in the rat. *Endocrinology* 114:930–940
- Bridges RS (2015) Neuroendocrine regulation of maternal behavior. *Front Neuroendocrinol* 36:178–196
- Bridges RS, Ronsheim PM (1990) Prolactin (PRL) regulation of maternal behavior in rats: bromocriptine treatment delays and PRL promotes the rapid onset of behavior. *Endocrinology* 126:837–848
- Bridges RS, Rosenblatt JS, Feder HH (1978) Serum progesterone concentrations and maternal behavior in rats after pregnancy termination: behavioral stimulation after progesterone withdrawal and inhibition by progesterone maintenance. *Endocrinology* 102:258–267

- Bridges RS, DiBiase R, Loundes DD, Doherty PC (1985) Prolactin stimulation of maternal behavior in female rats. *Science* 227:782–784
- Bridges RS, Numan M, Ronsheim PM, Mann PE, Lupini CE (1990) Central prolactin infusions stimulate maternal behavior in steroid-treated, nulliparous female rats. *Proc Natl Acad Sci U S A* 87:8003–8007
- Bridges RS, Mann PE, Coppeta JS (1999) Hypothalamic involvement in the regulation of maternal behaviour in the rat: inhibitory roles for the ventromedial hypothalamus and the dorsal/anterior hypothalamic areas. *J Neuroendocrinol* 11:259–266
- Brown RS, Wyatt AK, Herbison RE, Knowles PJ, Ladyman SR, Binart N, Banks WA, Grattan DR (2016) Prolactin transport into mouse brain is independent of prolactin receptor. *FASEB J* 30:1002–1010
- Brown RSE, Aoki M, Ladyman SR, Phillipps HR, Wyatt A, Boehm U, Grattan DR (2017) Prolactin action in the medial preoptic area is necessary for postpartum maternal nursing behavior. *Proc Natl Acad Sci U S A* 114:10779–10784
- Brunton PJ (2010) Resetting the dynamic range of hypothalamo-pituitary-adrenal axis stress responses through pregnancy. *J Neuroendocrinol* 22:1198–1213
- Brunton PJ, Russell JA (2003) Hypothalamic-pituitary-adrenal responses to centrally administered orexin-A are suppressed in pregnant rats. *J Neuroendocrinol* 15:633–637
- Brunton PJ, Russell JA (2010) Prenatal social stress in the rat programmes neuroendocrine and behavioural responses to stress in the adult offspring: sex specific effects. *J Neuroendocrinol* 22:258–271
- Brunton PJ, Meddle SL, Ma S, Ochedalski T, Douglas AJ, Russell JA (2005) Endogenous opioids and attenuated hypothalamic-pituitary-adrenal axis responses to immune challenge in pregnant rats. *J Neurosci* 25:5117–5126
- Brunton PJ, Bales J, Russell JA (2006a) Neuroendocrine stress but not feeding responses to centrally administered neuropeptide Y are suppressed in pregnant rats. *Endocrinology* 147:3737–3745
- Brunton PJ, Sabatier N, Leng G, Russell JA (2006b) Suppressed oxytocin neuron responses to immune challenge in late pregnant rats: a role for endogenous opioids. *Eur J Neurosci* 23:1241–1247
- Brunton PJ, McKay AJ, Ochedalski T, Piastowska A, Rebas E, Lachowicz A, Russell JA (2009) Central opioid inhibition of neuroendocrine stress responses in pregnancy in the rat is induced by the neurosteroid allopregnanolone. *J Neurosci* 29:6449–6460
- Brunton PJ, Bales J, Russell JA (2012) Allopregnanolone and induction of endogenous opioid inhibition of oxytocin responses to immune stress in pregnant rats. *J Neuroendocrinol* 24:690–700
- Brunton PJ, Russell JA, Hirst JJ (2014) Allopregnanolone in the brain: protecting pregnancy and birth outcomes. *Prog Neurobiol* 113:106–136
- Brussaard AB, Herbison AE (2000) Long-term plasticity of postsynaptic GABA(A)-receptor function in the adult brain: insights from the oxytocin neurone. *Trends Neurosci* 23:190–195
- Buckingham JC, Cooper TA (1984) Differences in hypothalamo-pituitary-adrenocortical activity in the rat after acute and prolonged treatment with morphine. *Neuroendocrinology* 38:411–417
- Buckingham JC, Cooper TA (1986) Effects of naloxone on hypothalamo-pituitary-adrenocortical activity in the rat. *Neuroendocrinology* 42:421–426
- Buller KM, Hamlin AS, Osborne PB (2005) Dissection of peripheral and central endogenous opioid modulation of systemic interleukin-1beta responses using c-fos expression in the rat brain. *Neuropharmacology* 49:230–242
- Burbach JPH, Young LJ, Russell JA (2006) Oxytocin: synthesis, secretion, and reproductive functions. In: Neill JD (ed) *Knobil and Neill's physiology of reproduction*. Elsevier, Amsterdam, pp 3055–3128
- Byrnes EM, Bridges RS (2000) Endogenous opioid facilitation of maternal memory in rats. *Behav Neurosci* 114:797–804
- Byrnes EM, Riger BA, Bridges RS (2000) Opioid receptor antagonism during early lactation results in the increased duration of nursing bouts. *Physiol Behav* 70:211–216
- Callahan P, Klosterman S, Prunty D, Tompkins J, Janik J (2000) Immunoneutralization of endogenous opioid peptides prevents the suckling-induced prolactin increase and the inhibition of tuberoinfundibular dopaminergic neurons. *Neuroendocrinology* 71:268–276
- Caron MG, Beaulieu M, Raymond V, Gagne B, Drouin J, Lefkowitz RJ, Labrie F (1978) Dopaminergic receptors in the anterior pituitary gland. Correlation of [³H]dihydroergocryptine binding with the dopaminergic control of prolactin release. *J Biol Chem* 253:2244–2253
- Ceccatelli S, Orazzo C (1993) Effect of different types of stressors on peptide messenger ribonucleic acids in the hypothalamic paraventricular nucleus. *Acta Endocrinol* 128:485–492
- Champagne FA, Chretien P, Stevenson CW, Zhang TY, Gratton A, Meaney MJ (2004) Variations in nucleus accumbens dopamine associated with individual differences in maternal behavior in the rat. *J Neurosci* 24:4113–4123
- Cohn J, Gerall AA (1989) Pre- and postpuberal medial preoptic area lesions and maternal behavior in the rat. *Physiol Behav* 46:333–336
- Concas A, Mostallino MC, Porcu P, Follesa P, Barbaccia ML, Trabucchi M, Purdy RH, Grisenti P, Biggio G (1998) Role of brain allopregnanolone in the plasticity of gamma-aminobutyric acid type A receptor in rat brain during pregnancy and after delivery. *Proc Natl Acad Sci U S A* 95:13284–13289
- da Costa APC, Wood S, Ingram CD, Lightman SL (1996) Region-specific reduction in stress-induced c-fos mRNA expression during pregnancy and lactation. *Brain Res* 742:177–184
- da Costa APC, Kampa RJ, Windle RJ, Ingram CD, Lightman SL (1997) Region-specific immediate-early gene expression following the administration of corticotropin-releasing hormone in virgin and lactating rats. *Brain Res* 770:151–162
- da Costa AP, Ma X, Ingram CD, Lightman SL, Aguilera G (2001) Hypothalamic and amygdaloid corticotropin-releasing hormone (CRH) and CRH receptor-1 mRNA expression in the stress-hyporesponsive late pregnant and early lactating rat. *Brain Res Mol Brain Res* 91:119–130
- Demarest KT, Duda NJ, Riegler GD, Moore KE (1983a) Placental lactogen mimics prolactin in activating tuberoinfundibular dopaminergic neurons. *Brain Res* 272:175–178
- Demarest KT, McKay DW, Riegler GD, Moore KE (1983b) Biochemical indices of tuberoinfundibular dopaminergic neuronal activity during lactation: a lack of response to prolactin. *Neuroendocrinology* 36:130–137
- Desjardins GC, Brawer JR, Beaudet A (1990) Distribution of mu, delta, and kappa opioid receptors in the hypothalamus of the rat. *Brain Res* 536:114–123
- Douglas AJ, Clarke G, MacMillan SJ, Bull PM, Neumann I, Way SA, Wright DM, McGrory BG, Russell JA (1993a) Effects of the kappa-opioid agonist U50,488 on parturition in rats. *Br J Pharmacol* 109:251–258
- Douglas AJ, Dye S, Leng G, Russell JA, Bicknell RJ (1993b) Endogenous opioid regulation of oxytocin secretion through pregnancy in the rat. *J Neuroendocrinol* 5:307–314
- Douglas AJ, Neumann I, Meeren HKM, Leng G, Johnstone LE, Munro G, Russell JA (1995) Central endogenous opioid inhibition of supraoptic oxytocin neurons in pregnant rats. *J Neurosci* 15:5049–5057
- Douglas AJ, Johnstone HA, Wigger A, Landgraf R, Neumann ID (1998a) The role of endogenous opioids in neurohypophysial and hypothalamo-pituitary-adrenal axis hormone secretory responses to stress in pregnant rats. *J Endocrinol* 158:285–293
- Douglas AJ, Meeren HK, Johnstone LE, Pfaff DW, Russell JA, Brooks PJ (1998b) Stimulation of expression of the oxytocin gene in rat supraoptic neurons at parturition. *Brain Res* 782:167–174

- Douglas AJ, Johnstone H, Brunton P, Russell JA (2000) Sex-steroid induction of endogenous opioid inhibition on oxytocin secretory responses to stress. *J Neuroendocrinol* 12:343–350
- Douglas AJ, Scullion S, Antonijevic IA, Brown D, Russell JA, Leng G (2001) Uterine contractile activity stimulates supraoptic neurons in term pregnant rats via a noradrenergic pathway. *Endocrinology* 142:633–644
- Douglas AJ, Bicknell RJ, Leng G, Russell JA, Meddle SL (2002) Beta-endorphin cells in the arcuate nucleus: projections to the supraoptic nucleus and changes in expression during pregnancy and parturition. *J Neuroendocrinol* 14:768–777
- Douglas AJ, Brunton PJ, Bosch OJ, Russell JA, Neumann ID (2003) Neuroendocrine responses to stress in mice: hyporesponsiveness in pregnancy and parturition. *Endocrinology* 144:5268–5276
- Douglas A, Meddle S, Toschi N, Bosch O, Neumann I (2005) Reduced activity of the noradrenergic system in the paraventricular nucleus at the end of pregnancy: implications for stress hyporesponsiveness. *J Neuroendocrinol* 17:40–48
- Driessen TM, Eisinger BE, Zhao C, Stevenson SA, Saul MC, Gammie SC (2014) Genes showing altered expression in the medial preoptic area in the highly social maternal phenotype are related to autism and other disorders with social deficits. *BMC Neurosci* 15:11
- Drolet G, Dumont EC, Gosselin I, Kinkead R, Laforest S, Trottier JF (2001) Role of endogenous opioid system in the regulation of the stress response. *Prog Neuro-Psychopharmacol Biol Psychiatry* 25:729–741
- Dyball RE, Leng G (1986) Regulation of the milk ejection reflex in the rat. *J Physiol* 380:239–256
- Eisinger BE, Zhao C, Driessen TM, Saul MC, Gammie SC (2013) Large scale expression changes of genes related to neuronal signaling and developmental processes found in lateral septum of postpartum outbred mice. *PLoS One* 8:e63824
- Eisinger BE, Driessen TM, Zhao C, Gammie SC (2014) Medial prefrontal cortex: genes linked to bipolar disorder and schizophrenia have altered expression in the highly social maternal phenotype. *Front Behav Neurosci* 8:110
- Elovitz MA, Mrinalini C (2004) Animal models of preterm birth. *Trends Endocrinol Metab* 15:479–487
- Ericsson A, Kovacs KJ, Sawchenko PE (1994) A functional anatomical analysis of central pathways subserving the effects of interleukin-1 on stress-related neuroendocrine neurons. *J Neurosci* 14:897–913
- Fallon JH, Leslie FM (1986) Distribution of dynorphin and enkephalin peptides in the rat brain. *J Comp Neurol* 249:293–336
- Feher P, Olah M, Bodnar I, Hechtel D, Bacskay I, Juhasz B, Nagy GM, Vecsernyes M (2010) Dephosphorylation/inactivation of tyrosine hydroxylase at the median eminence of the hypothalamus is required for suckling-induced prolactin and adrenocorticotrop hormone responses. *Brain Res Bull* 82:141–145
- Ferguson JKW (1941) A study of the motility of the intact uterus at term. *Surg Gynecol Obstet* 73:359–366
- Ferris CF, Kulkarni P, Sullivan JM Jr, Harder JA, Messenger TL, Febo M (2005) Pup suckling is more rewarding than cocaine: evidence from functional magnetic resonance imaging and three-dimensional computational analysis. *J Neurosci* 25:149–156
- Fitzsimmons MD, Olschowka JA, Wiegand SJ, Hoffman GE (1992) Interaction of opioid peptide-containing terminals with dopaminergic perikarya in the rat hypothalamus. *Brain Res* 581:10–18
- Fuchs AR, Saito S (1971) Pituitary oxytocin and vasopressin content of pregnant rats before, during and after parturition. *Endocrinology* 88:574–578
- Gammie SC, Driessen TM, Zhao C, Saul MC, Eisinger BE (2016) Genetic and neuroendocrine regulation of the postpartum brain. *Front Neuroendocrinol* 42:1–17
- Gibori G, Richards JS (1978) Dissociation of two distinct luteotropic effects of prolactin: regulation of luteinizing hormone-receptor content and progesterone secretion during pregnancy. *Endocrinology* 102:767–774
- Gonzalez-Iglesias AE, Fletcher PA, Arias-Cristancho JA, Cristancho-Gordo R, Helena CV, Bertram R, Tabak J (2015) Direct stimulatory effects of oxytocin in female rat gonadotrophs and somatotrophs in vitro: comparison with lactotrophs. *Endocrinology* 156:600–612
- Goodwin TM, Paul R, Silver H, Spellacy W, Parsons M, Chez R, Hayashi R, Valenzuela G, Creasy GW, Merriman R (1994) The effect of the oxytocin antagonist atosiban on preterm uterine activity in the human. *Am J Obstet Gynecol* 170:474–478
- Grattan DR, Averill RL (1990) Effect of ovarian steroids on a nocturnal surge of prolactin secretion that precedes parturition in the rat. *Endocrinology* 126:1199–1205
- Grattan DR, Steyn FJ, Kokay IC, Anderson GM, Bunn SJ (2008) Pregnancy-induced adaptation in the neuroendocrine control of prolactin secretion. *J Neuroendocrinol* 20:497–507
- Grisel JE, Bartels JL, Allen SA, Turgeon VL (2008) Influence of beta-endorphin on anxious behavior in mice: interaction with EtOH. *Psychopharmacology* 200:105–115
- Guillou A, Romano N, Steyn F, Abitbol K, Le Tissier P, Bonnefont X, Chen C, Mollard P, Martin AO (2015) Assessment of lactotroph axis functionality in mice: longitudinal monitoring of PRL secretion by ultrasensitive-ELISA. *Endocrinology* 156:1924–1930
- Gunnet JW, Freeman ME (1983) The mating-induced release of prolactin: a unique neuroendocrine response. *Endocr Rev* 4:44–61
- Gustafson P, Bunn SJ, Grattan DR (2017) The role of prolactin in the suppression of Crh mRNA expression during pregnancy and lactation in the mouse. *J Neuroendocrinol* 29
- Hammer RP Jr, Bridges RS (1987) Preoptic area opioids and opiate receptors increase during pregnancy and decrease during lactation. *Brain Res* 420:48–56
- Hansen S, Harthorn C, Wallin E, Lofberg L, Svensson K (1991) Mesotelencephalic dopamine system and reproductive behavior in the female rat: effects of ventral tegmental 6-hydroxydopamine lesions on maternal and sexual responsiveness. *Behav Neurosci* 105:588–598
- Harbuz MS, Lightman SL (1989) Responses of hypothalamic and pituitary mRNA to physical and psychological stress in the rat. *J Endocrinol* 122:705–711
- Harbuz MS, Jessop DS, Lightman SL, Chowdrey SL (1994) The effects of restraint or hypertonic saline stress on corticotropin-releasing factor, arginine-vasopressin, and proenkephalin-a messenger-RNAs in the CFY, Sprague-Dawley and Wistar strains of rat. *Brain Res* 667:6–12
- Harlan RE, Shivers BD, Romano GJ et al (1987) Localization of preproenkephalin mRNA in the rat brain and spinal cord by in situ hybridization. *J Comp Neurol* 258:159–184
- Henry MS, Bisht K, Vernoux N, Gendron L, Torres-Berrio A, Drolet G, Tremblay ME (2018) Delta opioid receptor signaling promotes resilience to stress under the repeated social defeat paradigm in mice. *Front Mol Neurosci* 11:100
- Herbison AE, Voisin DL, Douglas AJ, Chapman C (1997) Profile of monoamine and excitatory amino acid release in rat supraoptic nucleus over parturition. *Endocrinology* 138:33–40
- Higuchi T, Uchida K, Honda K, Negoro H (1986) Oxytocin release during parturition in the pelvic-neurectomized rat. *J Endocrinol* 109:149–154
- Horvath TL, Naftolin F, Leranath C (1992) Beta-endorphin innervation of dopamine neurons in the rat hypothalamus: a light and electron microscopic double immunostaining study. *Endocrinology* 131:1547–1555
- Kammerer M, Adams D, Castelberg Bv B, Glover V (2002) Pregnant women become insensitive to cold stress. *BMC Pregnancy Childbirth* 2:8
- Kang W, Wilson SP, Wilson MA (2000) Overexpression of proenkephalin in the amygdala potentiates the anxiolytic effects of benzodiazepines. *Neuropsychopharmacology* 22:77–88

- Katoh A, Nabeshima T, Kameyama T (1990) Behavioral changes induced by stressful situations: effects of enkephalins, dynorphin, and their interactions. *J Pharmacol Exp Ther* 253:600–607
- Keer SE, Stern JM (1999) Dopamine receptor blockade in the nucleus accumbens inhibits maternal retrieval and licking, but enhances nursing behavior in lactating rats. *Physiol Behav* 67:659–669
- Khachaturian H, Watson SJ, Lewis ME, Coy D, Goldstein A, Akil H (1982) Dynorphin immunocytochemistry in the rat central nervous system. *Peptides* 3:941–954
- Khachaturian H, Lewis ME, Watson SJ (1983) Enkephalin systems in diencephalon and brainstem of the rat. *J Comp Neurol* 220:310–320
- Knobloch HS, Charlet A, Hoffmann LC, Eliava M, Khrulev S, Cetin AH, Osten P, Schwarz MK, Seeburg PH, Stoop R, Grinevich V (2012) Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73:553–566
- Kokma JJ, van Kesteren RE, Rosahl TW, Zwart R, Smit AB, Luddens H, Brussaard AB (2003) Oxytocin regulates neurosteroid modulation of GABA(A) receptors in supraoptic nucleus around parturition. *J Neurosci* 23:788–797
- Konig M, Zimmer AM, Steiner H, Holmes PV, Crawley JN, Brownstein MJ, Zimmer A (1996) Pain responses, anxiety and aggression in mice deficient in pre-proenkephalin. *Nature* 383:535–538
- Lang RE, Heil JWE, Ganten D, Hermann K, Unger T, Rascher W (1983) Oxytocin unlike vasopressin is a stress hormone in the rat. *Neuroendocrinology* 37:314–316
- Le Merrer J, Becker JA, Befort K, Kieffer BL (2009) Reward processing by the opioid system in the brain. *Physiol Rev* 89:1379–1412
- Lee Y, Voogt JL (1999) Feedback effects of placental lactogens on prolactin levels and Fos-related antigen immunoreactivity of tuberoinfundibular dopaminergic neurons in the arcuate nucleus during pregnancy in the rat. *Endocrinology* 140:2159–2166
- Lee A, Clancy S, Fleming AS (2000) Mother rats bar-press for pups: effects of lesions of the MPOA and limbic sites on maternal behavior and operant responding for pup-reinforcement. *Behav Brain Res* 108:215–231
- Lee HJ, Macbeth AH, Pagani JH, WSr Y (2009) Oxytocin: the great facilitator of life. *Prog Neurobiol* 88:127–151
- Leng G, Mansfield S, Bicknell RJ, Blackburn RE, Brown D, Chapman C, Dyer RG, Hollingsworth S, Shibuki K, Yates JO, Way SA (1988) Endogenous opioid actions and effects of environmental disturbance on parturition and oxytocin secretion in rats. *J Reprod Fertil* 84:345–356
- Leng G, Bicknell R, Brown D, Bowden C, Chapman C, Russell J (1994) Stimulus-induced depletion of pro-enkephalins, oxytocin and vasopressin and pro-enkephalin interaction with posterior pituitary hormone release in vitro. *Neuroendocrinology* 60:559–566
- Leng G, Brown CH, Murphy NP, Onaka T, Russell JA (1995) Opioid-noradrenergic interactions in the control of oxytocin cells. *Adv Exp Med Biol* 395:95–104
- Lerant A, Freeman ME (1998) Ovarian steroids differentially regulate the expression of PRL-R in neuroendocrine dopaminergic neuron populations: a double label confocal microscopic study. *Brain Res* 802:141–154
- Lightman SL, Young WSI (1989) Lactation inhibits stress-mediated secretion of corticosterone and oxytocin and hypothalamic accumulation of corticotropin-releasing factor and enkephalin messenger ribonucleic acids. *Endocrinology* 124:2358–2364
- Lucas BK, Ormandy CJ, Binart N, Bridges RS, Kelly PA (1998) Null mutation of the prolactin receptor gene produces a defect in maternal behavior. *Endocrinology* 139:4102–4107
- Ludwig M, Leng G (2006) Dendritic peptide release and peptide-dependent behaviours. *Nat Rev Neurosci* 7:126–136
- Ma S, Shipston MJ, Morilak D, Russell JA (2005) Reduced hypothalamic vasopressin secretion underlies attenuated adrenocorticotropin stress responses in pregnant rats. *Endocrinology* 146:1626–1637
- Maccari S, Krugers HJ, Morley-Fletcher S, Szyf M, Brunton PJ (2014) The consequences of early-life adversity: neurobiological, behavioural and epigenetic adaptations. *J Neuroendocrinol* 26:707–723
- Mann PE, Bridges RS (1992) Neural and endocrine sensitivities to opioids decline as a function of multiparity in the rat. *Brain Res* 580:241–248
- Mann PE, Kinsley CH, Bridges RS (1991) Opioid receptor subtype involvement in maternal behavior in lactating rats. *Neuroendocrinology* 53:487–492
- Mansi JA, Laforest S, Drolet G (2000) Effect of stress exposure on the activation pattern of enkephalin-containing perikarya in the rat ventral medulla. *J Neurochem* 74:2568–2575
- Mansour A, Khachaturian H, Lewis ME, Akil H, Watson SJ (1987) Autoradiographic differentiation of mu, delta, and kappa opioid receptors in the rat forebrain and midbrain. *J Neurosci* 7:2445–2464
- Mansour A, Meador-Woodruff JH, Bunzow JR, Civelli O, Akil H, Watson SJ (1990) Localization of dopamine D2 receptor mRNA and D1 and D2 receptor binding in the rat brain and pituitary: an in situ hybridization-receptor autoradiographic analysis. *J Neurosci* 10:2587–2600
- Mansour A, Fox CA, Burke S, Meng F, Thompson RC, Akil H, Watson SJ (1994a) Mu, delta, and kappa opioid receptor mRNA expression in the rat CNS: an in situ hybridization study. *J Comp Neurol* 350:412–438
- Mansour A, Fox CA, Meng F, Akil H, Watson SJ (1994b) Kappa 1 receptor mRNA distribution in the rat CNS: comparison to kappa receptor binding and prodynorphin mRNA. *Mol Cell Neurosci* 5:124–144
- Mansour A, Fox CA, Thompson RC, Akil H, Watson SJ (1994c) Mu-opioid receptor mRNA expression in the rat CNS: comparison to mu-receptor binding. *Brain Res* 643:245–265
- Meddle SL, Leng G, Selvarajah JR, Bicknell RJ, Russell JA (2000) Direct pathways to the supraoptic nucleus from the brainstem and the main olfactory bulb are activated at parturition in the rat. *Neuroscience* 101:1013–1021
- Meddle SL, Bishop VR, Gkoumassi E, van Leeuwen FW, Douglas AJ (2007) Dynamic changes in oxytocin receptor expression and activation at parturition in the rat brain. *Endocrinology* 148:5095–5104
- Merchenthaler I (1994) Induction of enkephalin in tuberoinfundibular dopaminergic neurons of pregnant, pseudopregnant, lactating and aged female rats. *Neuroendocrinology* 60:185–193
- Merchenthaler I, Maderdrut JL, Altschuler RA, Petrusz P (1986) Immunocytochemical localization of proenkephalin-derived peptides in the central nervous system of the rat. *Neuroscience* 17:325–348
- Merchenthaler I, Lennard DE, Cianchetta P, Merchenthaler A, Bronstein D (1995) Induction of proenkephalin in tuberoinfundibular dopaminergic neurons by hyperprolactinemia: the role of sex steroids. *Endocrinology* 136:2442–2450
- Mesiano S, Chan EC, Fitter JT, Kwek K, Yeo G, Smith R (2002) Progesterone withdrawal and estrogen activation in human parturition are coordinated by progesterone receptor A expression in the myometrium. *J Clin Endocrinol Metab* 87:2924–2930
- Minami M, Onogi T, Taya T, Katao Y, Hosoi Y, Maekawa K, Katsumata S, Yabuuchi K, Satoh M (1994) Molecular cloning and in situ hybridization histochemistry for rat mu-opioid receptor. *Neurosci Res* 18:315–322
- Moos F, Gouzenes L, Brown D, Dayanithi G, Sabatier N, Boissin L, Rabie A, Richard P (1998) New aspects of firing pattern autocontrol in oxytocin and vasopressin neurones. *Adv Exp Med Biol* 449:153–162
- Murata T, Murata E, Liu CX, Narita K, Honda K, Higuchi T (2000) Oxytocin receptor gene expression in rat uterus: regulation by ovarian steroids. *J Endocrinol* 166:45–52
- Murata T, Narita K, Honda K, Matsukawa S, Higuchi T (2003) Differential regulation of estrogen receptor alpha and beta mRNAs

- in the rat uterus during pregnancy and labor: possible involvement of estrogen receptors in oxytocin receptor regulation. *Endocr J* 50: 579–587
- Nahi F, Arbogast LA (2003) Prolactin modulates hypothalamic preproenkephalin, but not proopiomelanocortin, gene expression during lactation. *Endocrine* 20:115–122
- Nelson EE, Panksepp J (1998) Brain substrates of infant-mother attachment: contributions of opioids, oxytocin, and norepinephrine. *Neurosci Biobehav Rev* 22:437–452
- Neumann I, Russell JA, Wolff B, Landgraf R (1991) Naloxone increases the release of oxytocin, but not vasopressin, within limbic brain areas of conscious parturient rats: a push-pull perfusion study. *Neuroendocrinology* 54:545–551
- Neumann I, Russell JA, Landgraf R (1993) Oxytocin and vasopressin release within the supraoptic and paraventricular nuclei of pregnant, parturient and lactating rats: a microdialysis study. *Neuroscience* 53: 65–75
- Neumann I, Douglas AJ, Pittman QJ, Russell JA, Landgraf R (1996) Oxytocin released within the supraoptic nucleus of the rat brain by positive feedback action is involved in parturition-related events. *J Neuroendocrinol* 8:227–233
- Neuman ID, Johnstone HA, Hatzinger M, Liebsch G, Shipston M, Russell JA, Landgraf R, Douglas AJ (1998) Attenuated neuroendocrine responses to emotional and physical stressors in pregnant rats involve adeno-hypophysial changes. *J Physiol* 508:289–300
- Neumann ID, Bosch OJ, Toschi N, Torner L, Douglas AJ (2003) No stress response of the hypothalamo-pituitary-adrenal axis in parturient rats: lack of involvement of brain oxytocin. *Endocrinology* 144: 2473–2479
- Numan M (2007) Motivational systems and the neural circuitry of maternal behavior in the rat. *Dev Psychobiol* 49:12–21
- Numan M, Numan M (1996) A lesion and neuroanatomical tract-tracing analysis of the role of the bed nucleus of the stria terminalis in retrieval behavior and other aspects of maternal responsiveness in rats. *Dev Psychobiol* 29:23–51
- Numan M, Smith HG (1984) Maternal behavior in rats: evidence for the involvement of preoptic projections to the ventral tegmental area. *Behav Neurosci* 98:712–727
- Numan M, Rosenblatt JS, Komisaruk BR (1977) Medial preoptic area and onset of maternal behavior in the rat. *J Comp Physiol Psychol* 91:146–164
- Ormandy CJ, Binart N, Kelly PA (1997) Mammary gland development in prolactin receptor knockout mice. *J Mammary Gland Biol Neoplasia* 2:355–364
- Parada M, King S, Li M, Fleming AS (2008) The roles of accumbal dopamine D1 and D2 receptors in maternal memory in rats. *Behav Neurosci* 122:368–376
- Paris JJ, Brunton PJ, Russell JA, Walf AA, Frye CA (2011) Inhibition of 5 α -reductase activity in late pregnancy decreases gestational length and fecundity and impairs object memory and central progesterone milieu of juvenile rat offspring. *J Neuroendocrinol* 23:1079–1090
- Pedersen CA, Ascher JA, Monroe YL, Prange AJ Jr (1982) Oxytocin induces maternal behavior in virgin female rats. *Science* 216:648–650
- Pedersen CA, Caldwell JD, Walker C, Ayers G, Mason GA (1994) Oxytocin activates the postpartum onset of rat maternal behavior in the ventral tegmental and medial preoptic areas. *Behav Neurosci* 108:1163–1171
- Peters LC, Kristal MB, Komisaruk BR (1987) Sensory innervation of the external and internal genitalia of the female rat. *Brain Res* 408:199–204
- Pumford KM, Leng G, Russell JA (1991) Morphine actions on supraoptic oxytocin neurones in anaesthetized rats: tolerance after i.c.v. morphine infusion. *J Physiol* 440:437–454
- Pumford KM, Russell JA, Leng G (1993) Effects of the selective kappa-opioid agonist U50,488 upon the electrical activity of supraoptic neurones in morphine-tolerant and morphine-naive rats. *Exp Brain Res* 94:237–246
- Randle JC, Mazurek M, Kneifel D, Dufresne J, Renaud LP (1986) Alpha 1-adrenergic receptor activation releases vasopressin and oxytocin from perfused rat hypothalamic explants. *Neurosci Lett* 65:219–223
- Reymond MJ, Porter JC (1985) Involvement of hypothalamic dopamine in the regulation of prolactin secretion. *Horm Res* 22:142–152
- Romano N, Yip SH, Hodson DJ, Guillou A, Parnaudeau S, Kirk S, Tronche F, Bonnefont X, Le Tissier P, Bunn SJ, Grattan DR, Mollard P, Martin AO (2013) Plasticity of hypothalamic dopamine neurons during lactation results in dissociation of electrical activity and release. *J Neurosci* 33:4424–4433
- Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F (2008) Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput Biol* 4:e1000123
- Rubin BS, Bridges RS (1984) Disruption of ongoing maternal responsiveness in rats by central administration of morphine sulfate. *Brain Res* 307:91–97
- Russell JA, Brunton PJ (2009) Oxytocin. In: Squire LR (ed) *Encyclopedia of neuroscience*, vol 7. Oxford Academic Press, pp 337–347
- Russell JA, Douglas AJ (2000) Opioids. In: Fink G (ed) *Encyclopedia of Stress*, vol 3. Academic Press, San Diego, pp 86–98
- Russell JA, Gosden RG, Humphreys EM, Cutting R, Fitzsimons N, Johnston V, Liddle S, Scott S, Stirland JA (1989) Interruption of parturition in rats by morphine: a result of inhibition of oxytocin secretion. *J Endocrinol* 121:521–536
- Sagrillo CA, Voogt JL (1991) Endogenous opioids mediate the nocturnal prolactin surge in the pregnant rat. *Endocrinology* 129:925–930
- Samson WK, Lumpkin MD, McCann SM (1986) Evidence for a physiological role for oxytocin in the control of prolactin secretion. *Endocrinology* 119:554–560
- Sanders MJ, Kieffer BL, Fanselow MS (2005) Deletion of the mu opioid receptor results in impaired acquisition of Pavlovian context fear. *Neurobiol Learn Mem* 84:33–41
- Schulte HM, Weisner D, Allolio B (1990) The corticotrophin releasing hormone test in late pregnancy: lack of adrenocorticotrophin and cortisol response. *Clin Endocrinol* 33:99–106
- Selmanoff M, Gregerson KA (1985) Suckling decreases dopamine turnover in both medial and lateral aspects of the median eminence in the rat. *Neurosci Lett* 57:25–30
- Selmanoff M, Wise PM (1981) Decreased dopamine turnover in the median eminence in response to suckling in the lactating rat. *Brain Res* 212:101–115
- Shanks N, Windle RJ, Perks P, Wood S, Ingram CD, Lightman SL (1999) The hypothalamic-pituitary-adrenal axis response to endotoxin is attenuated during lactation. *J Neuroendocrinol* 11:857–865
- Sheehan T, Numan M (2002) Estrogen, progesterone, and pregnancy termination alter neural activity in brain regions that control maternal behavior in rats. *Neuroendocrinology* 75:12–23
- Smith MS, Neill JD (1976) Termination at midpregnancy of the two daily surges of plasma prolactin initiated by mating in the rat. *Endocrinology* 98:696–701
- Soaje M, Deis RP (1994) A modulatory role of endogenous opioids on prolactin secretion at the end of pregnancy in the rat. *J Endocrinol* 140:97–102
- Soaje M, Valdez S, Bregonzio C, Penissi A, Deis RP (2006) Dopaminergic mechanisms involved in prolactin release after mifepristone and naloxone treatment during late pregnancy in the rat. *Neuroendocrinology* 84:58–67
- Srisawat R, Ludwig M, Bull PM, Douglas AJ, Russell JA, Leng G (2000) Nitric oxide and the oxytocin system in pregnancy. *J Neurosci* 20: 6721–6727

- Stack EC, Balakrishnan R, Numan MJ, Numan M (2002) A functional neuroanatomical investigation of the role of the medial preoptic area in neural circuits regulating maternal behavior. *Behav Brain Res* 131:17–36
- Steyn FJ, Anderson GM, Grattan DR (2008) Hormonal regulation of suppressors of cytokine signaling (SOCS) messenger ribonucleic acid in the arcuate nucleus during late pregnancy. *Endocrinology* 149:3206–3214
- Stolzenberg DS, McKenna JB, Keough S, Hancock R, Numan MJ, Numan M (2007) Dopamine D1 receptor stimulation of the nucleus accumbens or the medial preoptic area promotes the onset of maternal behavior in pregnancy-terminated rats. *Behav Neurosci* 121: 907–919
- Sukikara MH, Mota-Ortiz SR, Baldo MV, Felicio LF, Canteras NS (2006) A role for the periaqueductal gray in switching adaptive behavioral responses. *J Neurosci* 26:2583–2589
- Sukikara MH, Platero MD, Canteras NS, Felicio LF (2007) Opiate regulation of behavioral selection during lactation. *Pharmacol Biochem Behav* 87:315–320
- Tomer L, Neumann ID (2002) The brain prolactin system: involvement in stress response adaptations in lactation. *Stress* 5:249–257
- Tomer L, Toschi N, Nava G, Clapp C, Neumann ID (2002) Increased hypothalamic expression of prolactin in lactation: involvement in behavioural and neuroendocrine stress responses. *Eur J Neurosci* 15:1381–1389
- Toufexis DJ, Walker C-D (1996) Noradrenergic facilitation of the adrenocorticotropin response to stress is absent during lactation in the rat. *Brain Res* 737:71–77
- Toufexis DJ, Thirivikraman KV, Plotsky PM, Morilak DA, Huang N, Walker C-D (1998) Reduced noradrenergic tone to the hypothalamic paraventricular nucleus contributes to the stress hyporesponsiveness of lactation. *J Neuroendocrinol* 10:417–427
- Toufexis DJ, Tesolin S, Huang N, Walker C (1999) Altered pituitary sensitivity to corticotropin-releasing factor and arginine vasopressin participates in the stress hyporesponsiveness of lactation in the rat. *J Neuroendocrinol* 11:757–764
- Tsuneoka Y, Maruyama T, Yoshida S, Nishimori K, Kato T, Numan M, Kuroda KO (2013) Functional, anatomical, and neurochemical differentiation of medial preoptic area subregions in relation to maternal behavior in the mouse. *J Comp Neurol* 521:1633–1663
- Valdez SR, Pennacchio GE, Gamboa DF, de Di Nasso EG, Bregonzio C, Soaje M (2014) Opioid modulation of prolactin secretion induced by stress during late pregnancy. Role of ovarian steroids. *Pharmacol Rep* 66:386–393
- Villegas-Gabutti CM, Pennacchio GE, Vivas L, Jahn GA, Soaje M (2018) Role of oxytocin in prolactin secretion during late pregnancy. *Neuroendocrinology* 106:324–334
- Voogt JL (1984) Evidence for an inhibitory influence of rat placental lactogen on prolactin release in vitro. *Biol Reprod* 31:141–147
- Voogt J, Robertson M, Friesen H (1982) Inverse relationship of prolactin and rat placental lactogen during pregnancy. *Biol Reprod* 26:800–805
- Walsh RJ, Slaby FJ, Posner BI (1987) A receptor-mediated mechanism for the transport of prolactin from blood to cerebrospinal fluid. *Endocrinology* 120:1846–1850
- Wamsley JK, Young WS 3rd, Kuhar MJ (1980) Immunohistochemical localization of enkephalin in rat forebrain. *Brain Res* 190:153–174
- Wang HJ, Hoffman GE, Smith MS (1993) Suppressed tyrosine hydroxylase gene expression in the tuberoinfundibular dopaminergic system during lactation. *Endocrinology* 133:1657–1663
- Welsh T, Johnson M, Yi L, Tan H, Rahman R, Merlino A, Zakar T, Mesiano S (2012) Estrogen receptor (ER) expression and function in the pregnant human myometrium: estradiol via ER α activates ERK1/2 signaling in term myometrium. *J Endocrinol* 212:227–238
- Wigger A, Lorsch P, Oehler I, Keck ME, Naruo T, Neumann ID (1999) Nonresponsiveness of the rat hypothalamo-pituitary-adrenocortical axis to parturition-related events: inhibitory action of endogenous opioids. *Endocrinology* 140:2843–2849
- Windle RJ, Wood S, Shanks N, Perks P, Conde GL, da Costa APC, Ingram CD, Lightman SL (1997) Endocrine and behavioural responses to noise stress: comparison of virgin and lactating female rats during non-disrupted maternal activity. *J Neuroendocrinol* 9: 407–414
- Zhao BG, Chapman C, Bicknell RJ (1988a) Functional kappa-opioid receptors on oxytocin and vasopressin nerve terminals isolated from the rat neurohypophysis. *Brain Res* 462:62–66
- Zhao BG, Chapman C, Bicknell RJ (1988b) Opioid-noradrenergic interactions in the neurohypophysis. I. Differential opioid receptor regulation of oxytocin, vasopressin, and noradrenaline release. *Neuroendocrinology* 48:16–24
- Zhao C, Eisinger BE, Driessen TM, Gammie SC (2014) Addiction and reward-related genes show altered expression in the postpartum nucleus accumbens. *Front Behav Neurosci* 8:388