



## Corrigendum

## Corrigendum to “Unexpectedly higher diazoxon hydrolysis by serum paraoxonase-1 in coronary heart disease” [Clinical Biochemistry 65 (2019) 21–23]



Bharti Mackness, Mike Mackness\*

Avda. Princip D'Espanya, 43892 Miami Playa, Tarragona, Spain

The authors regret that there were two numerical errors in Table 2 of this article. In Table 2 CONTROL 192QQ should be 11.04 and CHD

192RR should be 12.99. The correct version of the table is reproduced below.

The authors would like to apologise for any inconvenience caused.

**Table 2**

DIAZ activity of PON1 according to 192 and 55 genotypes.

	CONTROL		CHD	
	n	Activity <sup>a</sup>	n	Activity <sup>a</sup>
192QQ	155	11.04 ± 0.36 <sup>+</sup>	206	17.60 ± 0.37**
QR	97	9.54 ± 0.93	172	16.45 ± 0.43**
RR	22	8.81 ± 0.41	32	12.99 ± 0.87** <sup>+</sup>
55MM	26	8.73 ± 0.76	45	13.79 ± 0.66** <sup>+</sup>
LM	146	9.67 ± 0.36	219	16.28 ± 0.37**
LL	102	11.14 ± 0.45 <sup>+</sup>	146	16.95 ± 0.47**

Significantly different from control \*\**P* < .0001 or other genotypes +*P* < .0001.

<sup>a</sup> DIAZ activity = μmol/min/ml.

DOI of original article: <https://doi.org/10.1016/j.clinbiochem.2019.01.003>

\* Corresponding author.

E-mail address: [mike.mackness@gmail.com](mailto:mike.mackness@gmail.com) (M. Mackness).

<https://doi.org/10.1016/j.clinbiochem.2019.03.001>

Available online 06 March 2019

0009-9120/© 2019 The Canadian Society of Clinical Chemists. Published by Elsevier Inc. All rights reserved.