

Review article

# Clinical implications of intravenous drug incompatibilities in critically ill patients



Malik Benlabed <sup>a</sup>, Maxime Perez <sup>a,b</sup>, Romain Gaudy <sup>a</sup>, Stéphanie Genay <sup>a,b</sup>,  
 Damien Lannoy <sup>a,b</sup>, Christine Barthélémy <sup>a</sup>, Pascal Odou <sup>a,b</sup>, Gilles Lebuffle <sup>a,c</sup>,  
 Bertrand Décaudin <sup>a,b,\*</sup>

<sup>a</sup> Groupe de recherche sur les formes injectables et les technologies associées (GRITA), University Lille, EA 7365, 59000 Lille, France

<sup>b</sup> Pharmacie, CHU de Lille, 59000 Lille, France

<sup>c</sup> Pôle d'anesthésie-réanimation, CHU de Lille, 59000 Lille, France

ARTICLE INFO

Article history:  
 Available online 19 April 2018

Keywords:  
 Parenteral nutrition  
 Intravenous  
 Infusion pumps  
 Filters  
 Drug incompatibility  
 Critical care

ABSTRACT

**Objective:** The aim of this review is to analyse the clinical consequences of intravenous drug incompatibilities in critically ill patients, especially the incidence of organ dysfunctions and mortality.  
**Methods:** A review of literature was conducted according to the PRISMA statement in June 2017, using Medline, ISI Web of Science and Clinicaltrials.gov.

**Data extraction:** Eligible studies were case reports and randomised controlled trials (RCTs) that assessed the effects of drug incompatibilities in critically ill patients on morbidity or mortality as primary or secondary outcomes, or adverse events. Two investigators independently reviewed the eligibility of the study from abstracts or manuscript data.

**Data synthesis:** Twelve articles met the selection criteria. The six articles reporting RCTs concern only four RCTs. RCTs were single-centre studies comparing infusion with or without filter. One of them included adult patients. The others included paediatric and neonatal intensive care unit patients. Primary endpoints were SIRS, organ failure, overall complication rate, bacteraemia, sepsis, phlebitis and length of stay. The results are mixed with one RCT reporting a reduction in SIRS, organ failure and overall complication rate, two studies in disagreement over the occurrence of sepsis and one study reporting no impact on length of hospital stay. The six articles on case reports show different drug incompatibility situations. They report pulmonary toxicity.

**Conclusion:** Little data is available on this topic. Infused particles may induce organ failure, in particular pulmonary toxicity and SIRS. Further studies are needed to establish a link between the level of exposure to drug incompatibilities and clinical implication.

© 2018 Société française d'anesthésie et de réanimation (Sfar). Published by Elsevier Masson SAS. All rights reserved.

Contents

1. Introduction	174
2. Method	174
2.1. Data sources and search strategy	174
2.2. Study selection	174
3. Results	174
3.1. Case reports	174
3.1.1. Total parenteral nutrition	174
3.1.2. Other drugs	175
3.2. Randomised controlled trials	175

\* Corresponding author. Laboratoire de biopharmacie, pharmacie Galénique, hospitalière, 3, rue du Professeur-Laguesse, BP 83, 59006 Lille cedex, France.  
 E-mail address: bertrand.decaudin@univ-lille2.fr (B. Décaudin).

4. Discussion .....	178
5. Conclusion .....	179
References .....	179

## 1. Introduction

In intensive care units (ICUs), patients require numerous intravenous (IV) drugs administered via a central venous catheter and this parenteral drug infusion is often complicated by the fact that the number of concurrently administered drugs exceeds the number of available venous lines. Drug incompatibility, particularly between two drugs, is sometimes visible and recognisable through precipitate, especially in multidrug IV therapy [1]. The mechanisms of drug incompatibility were well described by Newton [2] who drew a distinction between physical and chemical reactions. Physical incompatibilities include visible (precipitation, turbidity and colour change) and sub-visible (pH change, sub-visible particles, decrease in drug concentration) reactions. Chemical drug incompatibilities are often sub-visible and lead to reactions such as redox, complexation or racemisation, which reduce the effectiveness of the administered drug or create toxicity. The frequency of drug incompatibility has been evaluated in different studies. For example, one observational study, performed in two adult ICUs and a medical unit, showed that 3.7% of drug combinations used were incompatible [3]. Another showed that physicochemical incompatibilities represented 19 out of 102 (18.6%) recorded nursing acts during a 30-day period in an adult ICU [4].

Drug incompatibilities increase particle formation. In animal studies, this particulate contamination induces thrombogenesis, impairs microcirculation and modulates the immune response [5]. In addition to particles arising from drug incompatibilities, particulate contamination may be introduced into the patient from the IV solutions. X-ray analyses have shown that particulate matter consists mainly of glass from ampoules, rubber from stoppers of infusion bottles, and plastic from infusion sets [6]. To improve the detection of drug incompatibility, various decision-supporting tools are available such as Trissel's handbook, the King guide to parenteral admixtures, the cross-table on the compatibility of injectable Y-delivered drugs from the University Hospital of Geneva and some databases, the most essential being the French Stabilis database [7,8]. In ICUs, the use of a multi-lumen catheter improves efficacy in terms of drug delivery and safety by minimising the risk of incompatibility reactions. To decrease the particulate contamination generated by drug incompatibility, the use of filters and in-line filtration is recommended [9] particularly during parenteral nutrition, to reduce inflammatory response and microcirculation impairment [10]. The clinical impact of drug incompatibilities might be particularly deleterious in critically ill patients. This indicates how important it is to clarify and synthesise the relative strengths and weaknesses of literature on this subject.

The objective of this review was to evaluate the clinical consequences of intravenous drug incompatibilities in critically ill patients.

## 2. Method

### 2.1. Data sources and search strategy

A review of literature was conducted according to the PRISMA statement in June 2017, using Medline, ISI Web of Science and Clinicaltrials.gov.

The following keywords were used: "parenteral nutrition" OR "parenteral nutrition, total" AND "crystallisation" OR "granuloma, foreign body"; "parenteral nutrition" OR "parenteral nutrition, total" AND "calcium phosphates"; "critical care" OR "care, critical" OR "critical ill" OR "intensive care units" OR "intensive care units, neonatal" OR "intensive care units, pediatric" AND "infusion, intravenous" AND "particulate matter" AND "filtration"; "drug incompatibility" AND "particulate matter"; "\*phlebitis" AND "filtration" AND "infusion intravenous"; "drug contamination\*" AND "infusion, intravenous" AND "filtration; "parenteral nutrition" OR "parenteral nutrition, total" AND "infusion intravenous" AND "drug therapy, combination".

### 2.2. Study selection

Included studies were human case reports and randomised controlled trials (RCTs) published after 1990 that evaluated the effects of drug incompatibility in critically ill patients on morbidity or mortality as primary or secondary outcomes, or adverse events. Adult and paediatric populations were concerned. We included studies investigating the effects of IV in-line filtration on organ functions in intensive care patients. Animal and in vitro studies were not included. Reviewed articles were restricted to those in English and French, and published as full-text articles. Reference lists of all included studies were also scanned. Two investigators (MB and MP) independently reviewed the identified abstracts or manuscripts to determine which studies were eligible. Discrepancies were resolved through the intervention of two other authors (BD and GL).

## 3. Results

A total of 187 studies were screened for eligibility (Fig. 1). Twelve publications (six case reports and six RCT) met the selection criteria.

### 3.1. Case reports

The six articles on case reports show different drug incompatibility situations (i.e. total parenteral nutrition (TPN), ceftriaxone with calcium gluconate and various medications) (Table 1). These articles report pulmonary toxicity (diffuse interstitial pneumonitis, occlusions in pulmonary artery branches and pulmonary granulomata).

#### 3.1.1. Total parenteral nutrition

Four articles concern drug incompatibilities involving TPN [9–12]. All of them highlight vascular crystal precipitation in the lung. Three articles contain clinical data on adults [9–11].

Hill et al. reported sudden cardio-respiratory arrest in two previously healthy young women receiving peripheral vein parenteral nutrition [11]. Autopsies documented amorphous material containing calcium and obstructing the pulmonary microvasculature of each patient. Mc Nearney et al. reported the history of a young woman with Crohn's disease who, six weeks after starting TPN, developed fever, shortness of breath, a dry cough, diffuse crackles in all pulmonary fields and milary nodules visible on a high-resolution computed tomography (HRCT)-scan [12]. Reedy et al. reported the case of a surgical patient who




Fig. 1. Flow chart.

developed chest tightness, shortness of breath and fever while receiving TPN [13]. Investigations undertaken by the authors of clinical cases focused mainly on lung biopsy and chest HRCT-scan. The clinical symptoms presented by these patients ranged from respiratory failure including shortness of breath, chest tightness, and dry cough with or without fever to sudden death or cardio-respiratory arrest. The persistence of severe respiratory status or persistent hypoxemia resulted in a high-resolution CT scan being taken, which showed either pulmonary miliary nodules or something resembling ground glass. Trans-bronchial or open-lung biopsies were needed to document TPN-associated pulmonary complications i.e. thrombi or granulomas containing crystalline precipitates. The last case reported post-mortem examinations of 41 parenterally-fed infants [14].

### 3.1.2. Other drugs

Bradley et al. [15] reported the deaths of seven neonates from cardio-respiratory arrest after receiving the concomitant IV infusion of ceftriaxone and calcium. This data was drawn from the Food and Drug Administration adverse event reporting system database [15]. The autopsy findings in four of the five infants for whom information was available revealed the presence of crystalline material or white precipitates in vascular beds and most often in the lungs. Precipitate formation occurred when both ceftriaxone and calcium were infused simultaneously, as well as when given at different time intervals. These observations suggest that for some infants the serum concentration of ceftriaxone may be sufficiently high for hours after infusion to interact with subsequently administered IV calcium.

Felton et al. observed life-threatening pulmonary hypertension and right ventricular failure in a 43 year-old man, a complication due to calcium and phosphate replacement in an ICU [16].

### 3.2. Randomised controlled trials

The six articles reporting RCTs concern four RCTs (different aspects of one RCT reported in three articles) (Table 2). The RCTs, concerning 1906 patients, were single-centre studies comparing infusion with or without filter. One of them included adult patients. The others included paediatric ICU and neonatal ICU patients. Primary endpoints were systemic inflammatory response syndrome (SIRS), organ failure, overall complication rate, bacteraemia, sepsis, phlebitis and length of stay. The results (Table 3) are heterogeneous with one RCT showing a reduction in SIRS, organ failure and overall complication rate, two studies in contradiction over the occurrence of sepsis and one reporting no consequence on length of stay.

One RCT was conducted in a PICU. This randomised single-centre, prospective trial, including 807 patients, aimed to assess the effects of in-line filtration on reducing the incidence of overall complications (sepsis, organ failure and thrombosis) and particularly SIRS, in critically ill children. Jack et al. reported a reduction in the incidence of SIRS and overall complication rate in the filter group [17]. The duration of mechanical ventilation was also reduced, as were both PICU and length of hospital stay. Boehne et al. assessed the influence of in-line filtration on different organ functions on the same cohort of patients [18]. The incidence of respiratory, renal and haematological dysfunction was lower in the filter group. No difference was observed for cardiovascular, hepatic or neurological morbidities between groups. Sasse et al. performed a sub-group analysis on 305 critically ill cardiac children [19]. The risk of SIRS, renal and haematological dysfunction was significantly lower within the filter group.

Two RCTs were conducted in a NICU. Van Lingen et al. reported a significant reduction in major complications through a composite

**Table 1**  
Data from case reports.

Reference	Patient(s)	Treatment involved	Detectability (biopsy, autopsy, clinical observation)	Treatment undertaken	Time between treatment and occurrence	Context of care	Clinical data reported
Bradley et al., 2009 [15]	Nine neonates and young infants, mostly younger than 2 months	Ceftriaxone + calcium gluconate with concurrent IV drugs (metronidazole, midazolam, furosemide, etc)	Autopsy findings	–	From a few minutes after ceftriaxone infusion to around 2 hours	–	Presence of crystalline material or white precipitate in vascular beds (in the lungs) for 4 infants
Felton et al., 2006 [16]	43-year-old man	5 L of crystalloid fluids, furosemide 10 mg, calcium gluconate 10% 10 mL and potassium phosphate 20 mmol/14 mmol TPN	Electrocardiogram, transthoracic echocardiography white precipitate in the drip chamber Autopsy	Anaesthesia with fentanyl, propofol and succinylcholine, intubation and mechanical ventilation	90 minutes	Ischiorectal abscess	Acute severe deterioration by flooding pulmonary oedema, severe pulmonary hypertension and right heart failure Death of the two patients. Amorphous material containing calcium obstructing the pulmonary microvasculature revealed by the autopsy Vascular pulmonary precipitates with occlusions in pulmonary artery branches containing crystalline precipitates visible with X-RAY analysis. Alveolar foreign body giant cell granulomas containing crystalline precipitates
Hill et al., 1996 [11]	Two young women with pelvic infections	Home TPN + manual addition of selenium 40 µg/mL 1.05 mL. No phosphorus salt in the TPN	Chest CT scan, bronchoscopy, BAL, open lung biopsy of the lingula (crystalline precipitates visible in energy dispersive X-Ray analysis)	Reintroduction of TPN on hospital day 2. Well tolerated until day 8 (spike fevers). IV antibiotherapies. Discontinuous TPN on day 11 with oral intake. Supplemental oxygen	Fever, shortness of breath and exercise intolerance 2 weeks after TPN. Hospitalisation of dehydration secondary to decreased oral intake, abdominal pain and fistulae 3 weeks after	Crohn's disease receiving home TPN	2 cases of pulmonary granulomata in the pulmonary arterial system. No identification of specific foreign material, but occasionally fragments and visible cotton fibers
McNearney et al., 2003 [12]	26-year-old woman	Parenteral nutrition for prematurity, according to a standard protocol with a fluid intake of 150 mL/kg	Necropsy findings of tissue sections from each lobe of each lung	–	–	Prematurity and association with respiratory distress requiring ventilatory support ( $n=38$ ) or necrotising or ischaemic enterocolitis ( $n=3$ )	Amorphous material obstructing the pulmonary microvasculature revealed by transbronchial biopsy
Puntis et al., 1992 [14]	Postmortem material from 41 parenterally fed infants from ICU	Home TPN + hydrocortisone, trimethoprim-sulfamethoxazole and ganciclovir during hospitalisation	Chest X-ray and CT scan. Biopsy during hospitalisation and autopsy of the lungs	Reduction in calcium gluconate from 30 to 20 mEq, and magnesium from 45 to 35 mEq per bag	Patient received regimens of TPN from the age of 6 weeks	Complex medical history notable for short gut secondary to mesenteric ischemia (intestinal malrotation)	
Reedy et al., 1999 [13]	21-year-old man						

**Table 2**  
RCT study design.

Reference	Design	Participants n	Treatment (n)	Study period (months)	Inclusion criteria	Exclusion criteria	Primary Endpoints	Secondary Endpoints
Van Lingen et al., 2004 [20]	Single-centre, prospective, randomised, controlled study	88	Control (44)/ Filter (44)	–	Premature infants > 26 weeks and < 37 weeks of gestation with respiratory distress syndrome, asphyxia or pneumonia/septicaemia, and patients on antibiotic therapy	Patients with congenital malformations, preterm infants < 26 weeks	Bacteraemia, phlebitis, extravasation, thrombosis, septicaemia and necrosis	Costs attributable to patients
Van den Hoogen et al., 2006 [21]	Single-centre, randomised trial	507	Control (214)/ Filter group (228)	12	All infants admitted in NICU who needed a CVC	–	CVC insertion site, number of catheter days, occurrence of sepsis or phlebitis, costs for disposable materials and nursing time	–
Jack et al., 2012 [17]	Single-centre, prospective, randomised, controlled trial	807	Control (406)/ Filter group (401)	43	Patients < 18 years	Expected death within 48 h of admission, absence of IV infusion therapy	Overall complication rate of major events (SIRS, sepsis, organ failure and thrombosis)	Length of stay on PICU and overall hospital stay
Boehne et al., 2013 [18]	Single-centre, prospective, randomised, controlled trial	807	Control (406)/ Filter group (401)	43	Patients < 18 years	Expected death within 48 h of admission, recruitment for other trials, absence of IV infusion therapy, discharge from PICU within 6 h, discontinuation of intervention	Overall complication rate of major events (incidence of SIRS, sepsis, thrombosis, acute liver failure, ARDS, acute renal and circulatory failure)	–
Sasse et al., 2015 [19]	Single-centre, prospective, randomised, controlled trial	305	Control (150)/ Filter group (155)	43	Patients < 18 years, Patients with cardiac diseases	Expected death within 48 h of admission, recruitment for other trials, absence of IV infusion therapy, discharge from PICU within 6 h, discontinuation of intervention	SIRS, sepsis, mortality, organ failure (circulation, lung, liver and kidney) or organ dysfunction (cardiovascular, respiratory, neurological, hematological)	Mortality rates, length of stay on PICU and duration of mechanical ventilation
Gradwohl-Matis et al., 2015 [22]	Single-centre, prospective, randomised, controlled, open-label trial	504	Control (252)/ Filter group (252)	16	All critically ill patients > 18 years, expected length of stay > 24 h and a CVC in place	Age < 18 years, pregnancy, neutropenia (< 1,5 G/L) or known immunosuppression, limited intensive care	Number of ICU days with SIRS	Incidence of SIRS, number of SIRS criteria per day with the syndrome, the duration of invasive mechanical ventilation, length of stay in the ICU, incidence of acute lung injury/acute respiratory distress syndrome, maximum C-reactive protein serum concentration, maximum white blood cell count, incidence of new candida and/or central-line-associated bloodstream infections, incidence of new thromboembolic complications, cumulative insulin requirements and presence of hyper- or hypoglycaemia

**Table 3**  
RCT main results.

Reference	Study selection	Study characteristics	Primary endpoints (control vs filter group)	Secondary Endpoints
Van Lingen et al., 2004 [20]	Exclusion: 4 in control group (death)	Age 30.7 weeks	Reduction in major complications [21 vs. 8; $P < 0.05$ ]	Reduction in costs/infant for the study group with filter, but no statistical data
Van den Hoogen et al., 2006 [21]	Exclusion: 39 in control group, 25 in filter group	Age 33 weeks, male 57%	No significant difference in the incidence of sepsis (also sepsis per 1000 catheter days), on catheter days. No phlebitis during the study	
Jacket et al., 2012 [17]	–	Age 5.8 years old, male 57%	Reduction of the overall complication rate [40.9% vs. 30.9%; 95% CI 0.484 to 0.865], SIRS [30.3% vs. 22.4%; 95% 0.485 to 0.913]	Reduction in length of stay [3.89 (2.96 to 4.81) vs. 2.98 (2.33 to 3.63)] and duration of mechanical ventilation [14.0 (5.6 to 22.4) vs. 11.0 (7.1 to 14.9)]
Boehne et al., 2013 [18]	–	Age 5.8 years old, male 57%	Reduction in respiratory [14.5% vs. 9.5%; CI –9.52 to –0.59%], renal [9.9% vs. 6.0%; CI –7.58 to –0.15%] and hematological dysfunction [8.4% vs. 4.5%; CI –7.26 to –0.51%]	–
Sasse et al., 2015 [17]	Exclusion: 32 in control group, 35 in filter group	Age 3.4 years old, male 63%	Reduction in SIRS [36.0% vs. 25.2%; 95% CI –21.8 to –0.5%], renal dysfunction [10.3% vs. 19.3%; CI –17.0 to –3.0%] and hematological dysfunction [5.8% vs. 12.7%; CI –14.2 to –0.2%]	No significant reduction
Gradwohl-Matis et al., 2015 [20]	No exclusion	Age 67.3 years old, male 59%	No significant difference	Higher incidence of SIRS in in-line microfilter group [96.8% vs. 99.6%]

criterion including the occurrence of thrombosis, sepsis and necrotising enterocolitis (NEC) [20]. Van den Hoogen et al. evaluated the incidence of nosocomial sepsis and did not find any statistical difference between infants infused with or without filters [21].

One RCT was performed on adult ICU patients. Gradwohl-Matis et al. carried out a prospective randomised, controlled, open-label study that evaluated the influence of in-line filters on systemic immune activation in critically ill adults [22]. The author observed that the number of ICU days with SIRS, the incidence of SIRS, acute lung injury, the number of new infections and thromboembolic complications and the duration of mechanical ventilation were no different between filter and control group.

#### 4. Discussion

This review of literature shows the potentially deleterious clinical effects of intravenous drug incompatibilities, which may induce or aggravate organ dysfunctions in critically ill patients. Pulmonary morbidities seem to be the main complications observed. These have been reported by authors cited above and are mainly caused by the micro-emboli of crystal precipitates obstructing pulmonary vessels and generating granulomatous pulmonary arteritis and granulomatous interstitial pneumonitis. Severe arterial pulmonary hypertension associated with cardiac arrest can occur during a diffuse and multiple emboli procedure. One of the first clinical complications was described by the FDA who reported that the ceftriaxone-calcium complexation formed during the concomitant IV administration of ceftriaxone and calcium led to the cardiac arrest and death of seven neonates due to pulmonary emboli of white precipitate [15]. This report is one of the best examples of pulmonary complication induced by drug incompatibility in a paediatric population. In contrast, Dalton et al. [23] showed that, in critically ill adult patients, the simultaneous IV administration of high concentrations of calcium and ceftriaxone was not significantly associated with greater mortality or adverse events compared to matched unexposed patients. Although Dalton's study is limited by its low-powered capacity for detecting effects i.e. only severe cardiorespiratory events or death, discrepancies in results may be explained by differences in calcium concentration and infusion method, ceftriaxone dose and respiratory function.

Similar pulmonary complications have been observed with TPN [10,12–14]. The origin of precipitation remains to be elucidated. One of the mechanisms could be related to the proportion or mixing sequence of electrolytes in the solution and exposure to extremes of temperature in storage or transfer. Such pulmonary complications occur frequently from a few days up to six weeks after the start of TPN. Vascular occlusion by precipitates should be hypothesised in patients on TPN who experience unexplainable fever, cough and shortness of breath. Diagnosis may require a CT scan showing diffuse pulmonary nodular opacities or even a lung biopsy in search of micro-emboli precipitates. In this context, Mc Nearney et al. suggested a prompt discontinuation of parenteral nutrition [12].

There are many arguments suggesting that the proportion of electrolytes in the TPN solution may be the cause of certain pulmonary complications. The high concentration of di- or trivalent cations leads to the degradation of lipid emulsions and the aggregation of lipid particles. High concentrations of calcium, phosphate and magnesium in the TPN solution contribute to the formation of calcium phosphate crystals.

As a result of drug incompatibilities, particles can activate the immune system and lead to the development of SIRS, one of the major risks for organ failure and death in critically ill patients. The incidence of SIRS, studied by Jack et al. in a randomised controlled trial evaluating the effect of in-line filtration, was significantly lower in the group of critically ill children using in-line filter [17]. Based on this data, in-line filtration seems effective and represents a potent strategy for preventing SIRS but is in contrast to the negative results in adults shown by Gradwohl-Matis et al. [22].

The effects of drug incompatibility on organ dysfunction constitute the essential part of this review. Mechanisms of particle-induced organ damage are probably accounted for by a mechanical blockage of micro-vessels, an activation of platelets and neutrophil granulocytes with the generation of occlusive micro-thrombi and the formation of foreign-body granulomas [8,23–25]. To elucidate the pathophysiological effects of particle infusion, Boehne et al. [18] and Sasse et al. [19] evaluated, in critically ill children, the influence of in-line filtration on different organ functions and observed a significant reduction in the incidence of SIRS, respiratory, renal and haematological dysfunction in the filter group. However, the absence of any difference

between groups concerning other organ dysfunctions is questionable.

This raises the problem of the effectiveness of in-line filtration and reliability of different studies, particularly a meta-analysis of four randomised trials performed by Forster et al., including 704 neonates, which did not demonstrate any difference in mortality or morbidity between filtration group and control group [24]. Nevertheless, the positive effects of filtration on organ function need to be considered. Renal and respiratory dysfunctions induced by particles are probably due to microcirculation impairment. Lehr et al. underlined the important role of microcirculation [10]. They investigated in an animal model whether particles from different antibiotic solutions might affect the blood supply in a micro vascular bed that has been compromised by prior injury induced experimentally by exposure of muscle tissue to ischemia and reperfusion. The loss of capillary perfusion due to particle infusion was dependent on the extent of ischaemia-induced muscle injury with more capillaries lost in the more severely compromised muscle areas. These findings suggest that particles may severely compromise tissue perfusion in patients with prior micro vascular compromise of vital organs (i.e. sepsis, trauma) and thus predispose to complications such as acute respiratory distress syndrome, acute renal failure and multi-organ failure. Therefore, prior microcirculation impairment in pathological conditions such as sepsis can induce or worsen organ dysfunction. Arima et al. investigated whether the injection of polystyrene bead as an artificial contaminant of intravenous fluids after lipopolysaccharide (LPS) administration affected mortality and organ damage in mice [25]. The mice were divided into four groups and received injections of: polystyrene beads only, LPS only, polystyrene beads 30 mn after LPS or saline. The survival rate after LPS injection associated with polystyrene bead injection was significantly lower than that of the other three groups. In kidney sections, injured glomeruli were significantly higher with LPS and polystyrene bead injection than in the other three groups. LPS and polystyrene bead injection decreased the glomerular filtration rate and led to renal failure. Orbegozo et al. observed that micro vascular reactivity is soon affected in patients with ARDS and this impairment is directly related to the severity of the disease. We can therefore easily understand that particles may aggravate this pathology [26]. Robinson et al. used an isolated rat heart model to assess whether unfiltered direct intra-arterial administration of cardioplegic solutions, as in open-heart surgery, could be damaging to the myocardium [27]. This led to a significant reduction in coronary blood flow, which could be largely prevented by filtration.

Other studies have also shown the benefit of a multi-lumen strategy to prevent particle contamination. Perez et al. underlined the advantage of a multi-lumen infusion set in reducing particulate contamination [28]. In an in vitro study, the authors reproduced the parenteral multidrug infusion used in a paediatric haematology unit. With a multi-lumen infusion set, particulate contamination was significantly reduced by 68% compared to the standard infusion set. The same team, in a controlled in vitro study, tested the compatibility of medications during multi-infusion therapy and confirmed the impact of a new multi-lumen infusion access device on the occurrence and prevention of known physical drug incompatibilities [29]. Foinard et al. evaluated in vitro the impact of multi-lumen infusion devices on the occurrence of known physical drug incompatibilities [30]. They observed that physical incompatibilities between two drugs could lead to a significant reduction in drug delivery to the patient, even in the absence of visible particles. Indeed, furosemide precipitation resulting in the formation of visible and/or sub-visible particles led to a drug loss to the patient estimated at between 10% and 15% in the presence of midazolam. This experimental data requires to be confirmed in

clinical practice so as to determine to what extent the choice of device affects the efficiency and safety of IV therapeutics administered to the patient.

Our analysis focused mainly on the clinical implications of the administration to patient of particles resulting from intravenous drug incompatibilities. This is a limitation to our study as other consequences of drug incompatibilities are not included. Chemical incompatibilities induce drug degradation that may lead to a decrease in drug delivery and/or the formation of toxic substances. For example, diluting epinephrine in a sodium bicarbonate solution induces its oxidation and decreases its concentration in solution [31]. The clinical relevance of such incompatibility depends on the contact time between the two chemically incompatible drugs [1,31,32]. It is therefore important to read carefully the results of drug stability studies to know how long the contact time is between mixed drugs in the same container [1].

## 5. Conclusion

This review shows that drug incompatibilities via particulate contamination may increase the risk of severe complications and damage to different organs. Drug incompatibilities can increase morbidity in critically ill patients particularly during prolonged TPN. Prevention of drug incompatibilities therefore appears to be a safety procedure to be implemented for critically ill patients by using multi-lumen infusion lines and catheters and/or in-line filtration. In critically ill neonates and children, in-line filtration reduces and even prevents some organ dysfunctions with no effect on mortality. Randomised controlled trials with a larger patient population are required to confirm the deleterious prognosis of particle contamination and to evaluate the preventive effect of in-line filtration and/or a multi-lumen strategy on morbidity.

## Funding

No financial support was obtained for the writing of this manuscript.

## Disclosure of interest

Dr. Décaudin reports non-financial support from Doran International, outside the submitted work. The other authors declare that they have no competing interest.

## References

- [1] Kanji S, Lam J, Johanson C, Singh A, Goddard R, Fairbairn J, et al. Systematic review of physical and chemical compatibility of commonly used medications administered by continuous infusion in intensive care units. *Crit Care Med* 2010;38:1890–8.
- [2] Newton DW. Y-site Compatibility of intravenous drugs with parenteral nutrition. *J Parenter Enter Nutr* 2012;37:297–9.
- [3] Gikic M, Di Paolo ER, Pannatier A, Cotting J. Evaluation of physicochemical incompatibilities during parenteral drug administration in a paediatric intensive care unit. *Pharm World Sci* 2000;22(3):88–9.
- [4] Tissot E, Cornette C, Demoly P, Jacquet M, Barale F, Capellier G. Medication errors at the administration stage in an intensive care unit. *Intensive Care Med* 1999;25(4):353–9.
- [5] Walpot H, Franke RP, Burchard WG, Agternkamp C, Müller FG, Mittermayer C, et al. [Particulate contamination of infusion solutions and drug additives in the framework of long-term intensive therapy. 2. An animal model]. *Anaesthesist* 1989;38:617–21.
- [6] Machotka O, Manak J, Kubena A, Vlcek J. Incidence of intravenous drug incompatibilities in intensive care units. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;159:652–6.
- [7] De Giorgi I, Guignard B, Fonzo-Christe C, Bonnabry P. Evaluation of tools to prevent drug incompatibilities in paediatric and neonatal intensive care units. *Pharm World Sci PWS* 2010;32:520–9.
- [8] Fonzo-Christe C, Bouchoud L, Pfister R. Incompatibilités médicamenteuses et nutrition parentérale en néonatalogie. *Nutr Clin Metabol* 2017;31:24–7.

- [9] Ball PA, Bethune K, Fox J, Ledger R, Barnett M. Particulate contamination in parenteral nutrition solutions: still a cause for concern? *Nutrition* 2001;17:926–9.
- [10] Lehr H-A, Brunner J, Rangoonwala R, James Kirkpatrick C. Particulate matter contamination of intravenous antibiotics aggravates loss of functional capillary density in postischemic striated muscle. *Am J Respir Crit Care Med* 2002;165:514–20.
- [11] Hill SE, Heldman LS, Goo ED, Whippe PE, Perkinson JC. Fatal microvascular pulmonary emboli from precipitation of a total nutrient admixture solution. *JPEN J Parenter Enteral Nutr* 1996;20:81–7.
- [12] McNearney T, Bajaj C, Boyars M, Cottingham J, Haque A. Total parenteral nutrition associated crystalline precipitates resulting in pulmonary artery occlusions and alveolar granulomas. *Dig Sci* 2003;48:1352–4.
- [13] Reedy JS, Kuhlman JE, Voytovich M. Microvascular pulmonary emboli secondary to precipitated crystals in a patient receiving total parenteral nutrition: a case report and description of the high-resolution ct findings. *Chest* 1999;115:892–5.
- [14] Puntis JW, Wilkins KM, Ball PA, Rushton DI, Booth IW. Hazards of parenteral treatment: do particles count? *Arch Dis Child* 1992;67:1475–7.
- [15] Bradley J, Wassel RT, Lee L, Nambiar S. Intravenous ceftriaxone and calcium in the neonate: assessing the risk for cardiopulmonary adverse events. *Pediatrics* 2009;123:609–13.
- [16] Felton TW, McCormick BA, Finfer SR, Fisher MM. Life-threatening pulmonary hypertension and right ventricular failure complicating calcium and phosphate replacement in the intensive care unit. *Anaesthesia* 2006;61:49–53.
- [17] Jack T, Boehne M, Brent BE, Hoy L, Köditz H, Wessel A, et al. In-line filtration reduces severe complications and length of stay on pediatric intensive care unit: a prospective randomized controlled trial. 38. 2012;p. 1008–16.
- [18] Boehne M, Jack T, Köditz H, Seidemann K, Schmidt F, Abura M, et al. In-line filtration minimizes organ dysfunction: new aspects from a prospective, randomized, controlled trial. *BMC Pediatr* 2013;13:21.
- [19] Sasse M, Dziuba F, Jack T, Koditz H, Kaussen T, Bertram H, et al. In-line filtration decreases systemic inflammatory response syndrome. Renal and hematologic dysfunction in pediatric cardiac intensive care patients. *Pediatr Cardiol* 2015;1270–8.
- [20] van Lingen R, Baerts W, Marquering A, Ruijs G. The use of in-line intravenous filters in sick newborn infants. *Acta Paediatr* 2004;93:658–62.
- [21] Van den Hoogen A, Krediet T, Uiterwaal C, Bolenius J, Gerards L, Fleer A. In-line filters in central venous catheters in a neonatal intensive care unit. *J Perinat Med* 2006;71–4.
- [22] Gradwohl-Matis I, Brunauer A, Dankl D, Wirthel E, Meburger I, Bayer A. Influence of in-line microfilters on systemic inflammation in adult critically ill patients: a prospective, randomized, controlled open label trial. *Ann Intensive Care* 2015;5:1–8.
- [23] Dalton BR, Zuege DJ, Shahpori R, Laupland KB. Concomitant ceftriaxone and high-concentration intravenous calcium therapy in adult critical care patients: a matched cohort study. *Ann Pharmacother* 2010;44:1158–63.
- [24] Foster JP, Richards R, Showell M, Jones LJ. Intravenous in-line filters for preventing morbidity and mortality in neonates. *Cochrane Database Syst Rev* 2015;CD005248.
- [25] Arima H, Hirate H, Sugiura T, Suzuki S, Takahashi S, Sobue K. IV injection of polystyrene beads for mouse model of sepsis causes severe glomerular injury. *J Intensive Care* 2014;2(1):21.
- [26] Orbegozo Cortés D, Rahmania L, Irazabal M, Santacruz C, Fontana V, De Backer D, et al. Microvascular reactivity is altered early in patients with acute respiratory distress syndrome. *Respir Res* 2016;17:59.
- [27] Robinson LA, Braimbridge MV, Hearse DJ. Impairment of myocardial function induced by particulate contamination of cardioplegic solutions. *Adv Myocardiol* 1985;6:503–12.
- [28] Perez M, Décaudin B, Abou Chahla W, Nelken B, Barthélémy C, Lebuffe G, et al. In vitro analysis of overall particulate contamination exposure during multidrug IV therapy: Impact of infusion sets. *Pediatr Blood Cancer* 2015;62:1042–7.
- [29] Perez M, Décaudin B, Foinard A, Barthélémy C, Debaene B, Lebuffe G, et al. Compatibility of medications during multi-infusion therapy: a controlled in vitro study on a multilumen infusion device. *Anaesth Crit Care Pain Med* 2015;34:83–8.
- [30] Foinard A, Décaudin B, Barthélémy C, Debaene B, Odou P. The impact of multilumen infusion devices on the occurrence of known physical drug incompatibility: a controlled in vitro study. *Anesth Analg* 2013;116:101–6.
- [31] Bonhomme L, Benhamou D, Comoy E, Preaux N. Stability of epinephrine in alkalized solutions. *Ann Emerg Med* 1990;19:1242–4.
- [32] Gedeberg R, Wiklund L. The biological activity of adrenaline after injection through an intravenous cannula containing alkaline buffer. *Resuscitation* 1989;18:49–58.