

LETTERS TO THE EDITOR

Comments regarding: Kim KA, Oh WS, Lee JJ. Recording an accurate maxillomandibular relationship by adding vertical stops to the occlusal rims. J Prosthet Dent 2019;121:242-245.

Dear Editor:



We read the article entitled “Recording an accurate maxillomandibular relationship by adding vertical stops to the occlusal rims” by Kim et al¹ with great interest. Different methods of registering centric relation have been described²⁻⁴; the most common method being the wax record procedure.² However, there have been contradictory reports on recording centric relation by this technique.⁵⁻⁷ Uneven contact in the occlusal rims can result in uneven pressure on the denture-supporting tissues and subsequent errors in centric relation recording as stated by Gysi.⁶ Therefore, equalization of contacts on the denture-supporting areas or, more precisely, equalization of vertical contacts has been considered as a basic requirement for recording centric relation.⁸

In the technique proposed by Kim et al,¹ a nonuniform contact between the wax occlusal rims was corrected by placing dome-shaped baseplate wax as vertical stops on the occlusal surface of the mandibular occlusal rims. However, we feel that closing on 4 isolated wax vertical stops will create a nonuniform distribution of pressure on the underlying denture bearing mucosa, thereby defeating the concept of obtaining equalized and uniform contact. Moreover, the length of time to complete this action and relative difference in softness of wax on both sides may also affect the accuracy and reliability in recording centric relation.²

In our opinion, centric relation records should be made with accurately fitted and stable trial denture bases⁸ having uniform contact of wax occlusal rim throughout the occlusal table under minimal pressure.² Attempts to make uniform occlusal contact by introducing noncontinuous wax interferences in the form of 4 dome-shaped elevations that contact only in the posterior region of the rims further complicates the basic concept of recording centric relation. We suggest that the authors objectively validate their technique in terms of precision achieved in recording

centric relation by this method before it can be advocated.

Himanshi Aggarwal, MDS
Lucknow, India

Arvind Tripathi, MDS, MNAMS
Lucknow, India

Ashutosh Gupta, MDS
Lucknow, India

REFERENCES

1. Kim KA, Oh WS, Lee JJ. Recording an accurate maxillomandibular relationship by adding vertical stops to the occlusal rims. J Prosthet Dent 2019;121:242-5.
2. Yurkstas AA, Kapur KK. Factors influencing centric relation records in edentulous mouths. J Prosthet Dent 1964;14:1054-65.
3. Shanahan TEJ. Physiologic jaw relations and occlusion of complete dentures. J Prosthet Dent 1955;5:319-24.
4. Dixon DL. Overview of articulation materials and methods for the prosthodontic patient. J Prosthet Dent 2000;83:235-47.
5. Wright WH. Use of intra-oral jaw relation wax records in completed denture prosthesis. J Am Dent Assoc 1939;26:542-55.
6. Gysi A. Practical application of research results in denture construction. J Am Dent Assoc 1929;16:199-223.
7. Trapozzano VR. Occlusal records. J Prosthet Dent 1955;5:325-32.
8. Kingery RH. A review of some of the problems associated with centric relation. J Prosthet Dent 1952;307-19.

Reply:



The occlusal rim should ideally demonstrate a uniform contact for the registration of maxillomandibular record. Even contact should be applied during the recording procedure to avoid displacement (or distortion) of the record base.¹ However, attainment of the ideal pattern of uniform contact is a challenge.

One common clinical procedure includes creation of space in the posterior segment between the maxillary and mandibular occlusal rims.² The vertical stop is established in the anterior segment of occlusal rims to maintain the occlusal vertical dimension. However, the record base can be displaced unless the stability of the record base is ensured during the closure of the