



Contents lists available at ScienceDirect

Comparative Immunology, Microbiology and Infectious Diseases

journal homepage: www.elsevier.com/locate/cimid

Corrigendum

Corrigendum to “A survey on *Toxocara cati* eggs on the hair of stray cats: A potential risk factor for human toxocariasis in Northeastern Iran” [Comp. Immunol. Microbiol. Infect. Dis. 64 (2019) 10–13]

Amin Bakhshani, Mohsen Maleki, Alireza Haghparast, Sima Parande Shirvan, Hassan Borji*

Department of Pathobiology, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran

The authors regret < As we have used EndNote program for reference list, some of reference listed only in Tables 1 and 4 have missed from reference list >

The authors would like to apologise for any inconvenience caused.

Table 1 Prevalence of *Toxocara cati* in stray cats in different regions of Iran and other country by necropsy and fecal analysis.

Region	Place	Prevalence (%)	Reference
Iran	Northeastern Iran	28 ^a	Borji et al. [19]
	Other countries		
	Northwestern Iran	78 ^a	Hajipour et al. [20]
	South of Iran	26.7 ^a	Mikaeili et al. [21]
	Central of Iran	13.3 ^a	Mohsen and Hossein [22]
	Northern Iran	44 ^a	Sharif et al. [23]
	Northwestern Iran	86.3 ^a	Yakhchali et al. [24]
	South of Iran	42.6 ^a	Zibaei et al. [25]
	Northern Germany	27.1 ^b	Becker et al. [26]
	Central Spain	11.7 ^b	Montoya et al. [27]
	Netherlands	27 ^b	Nijse et al. [28]
	Argentina	61.2 ^b	Sommerfelt et al. [29]
	Northern Italy	33.1 ^b	Spada et al. [30]
	Greece	24.2 ^b	Symeonidou et al. [31]
	Poland	27.9 ^b	Szwabe and Blaszkowska [32]
	Portugal	38.3 ^b	Waap et al. [33]

a-necropsy, b-Fecal analysis.

Table 4 A comparison of numbers of *Toxocara* spp. eggs recovered from hair samples from stray cats and those recovered from dog hairs.

Host	Place	Prevalence (%)	Total number eggs/gr hair	Reference
Stray cats	Turkey	22	28	Oge et al. [5]
	Netherlands	3.4	28	Overgaaauw et al. [4]
	Mexico	41.7	nd	Rojas et al. [10]
Stray dogs	Brazil	24	614.8	Amaral et al. [34]
	Vietnam	56.3	nd	Anh et al. [35]
	Turkey	21.56	62	Aydenizöz-Ozkayhan et al. [36]
	Egypt	26.6	77.6	El-Tras et al. [37]
	Ireland	8.8	26	Keegan and Holland [38]
	Brazil	6.7	57.5	Merigueti et al. [39]
	Turkey	14	58	Oge et al. [5]
	Netherlands	12.2	17	Overgaaauw et al. [4]
	Italy	2.9	nd	Paoletti et al. [40]
	Ireland	67	583.88	Roddie et al. [7]
Mexico	41.7	nd	Rojas et al. [10]	
north-west of Iran	36.2	nd	Tavassoli et al. [41]	
Ireland and Britain	25	71	Wolfe and Wright [6]	

Nd. Not determined.

References

- [1] D. Despommier, Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular aspects, Clin. Microbiol. Rev. 16 (April (2)) (2003) 265–272.
- [2] H. Smith, et al., How common is human toxocariasis? Towards standardizing our knowledge, Trends Parasitol. 25 (April (4)) (2009) 182–1828, <https://doi.org/10.1016/j.pt.2009.01.006>. Epub 2009 Mar 5.
- [3] P.A. Overgaaauw, F. van Knapen, Veterinary and public health aspects of *Toxocara* spp, Vet. Parasitol. 193 (April (4)) (2013) 398–403, <https://doi.org/10.1016/j.vetpar.2012.12.035>. Epub 2012 Dec 20.
- [4] P.A. Overgaaauw, et al., Zoonotic parasites in fecal samples and fur from dogs and cats in the Netherlands, Vet. Parasitol. 163 (July (1-

DOI of original article: <https://doi.org/10.1016/j.cimid.2019.02.002>

* Corresponding author.

E-mail address: hborji@um.ac.ir (H. Borji).

<https://doi.org/10.1016/j.cimid.2019.02.008>

- 2)) (2009) 115–122, <https://doi.org/10.1016/j.vetpar.2009.03.044>. Epub 2009 Apr 5.
- [5] H. Oge, et al., Comparison of *Toxocara* eggs in hair and faecal samples from owned dogs and cats collected in Ankara, Turkey, *Vet. Parasitol.* 206 (December (3-4)) (2014) 227–231, <https://doi.org/10.1016/j.vetpar.2014.10.005>.
- [6] A. Wolfe, I.P. Wright, Parasitic nematode eggs in fur samples from dogs, *Vet. Rec.* 154 (March (13)) (2004) 408.
- [7] G. Roddie, et al., Contamination of dog hair with eggs of *Toxocara canis*, *Vet. Parasitol.* 152 (March (1-2)) (2008) 85–93, <https://doi.org/10.1016/j.vetpar.2007.12.008>. Epub 2007 Dec 15.
- [8] N. Yamaguti, *Systema helminthum, Nematodes of Vertebrates Vol. III* Inter Science Publisher Inc., New York, 1961.
- [9] E.J.L. Soulsby, *Helminths, Arthropods and Protozoa of Domesticated Animals*, 6 ed., (1977) Philadelphia.
- [10] T.O. Rojas, et al., Identification of *Toxocara* spp. eggs in dog hair and associated risk factors, *Vet. World* 10 (July (7)) (2017) 798–802, <https://doi.org/10.14202/vetworld.2017.798-802>. Epub 2017 Jul 20.
- [11] A. Mohsen, H. Hossein, Gastrointestinal parasites of stray cats in Kashan, Iran, *Trop. Biomed.* 26 (April (1)) (2009) 16–22.
- [12] D. Barutzki, R. Schaper, Results of parasitological examinations of faecal samples from cats and dogs in Germany between 2003 and 2010, *Parasitol. Res.* 109 (August (Suppl 1)) (2011) S45–S60, <https://doi.org/10.1007/s00436-011-2402-8>.
- [13] A.C. Becker, et al., Prevalence of endoparasites in stray and fostered dogs and cats in Northern Germany, *Parasitol. Res.* 111 (August (2)) (2012) 849–857, <https://doi.org/10.1007/s00436-012-2909-7>. Epub 2012 Apr 13.
- [14] A. Kleine, A. Springer, C. Strube, Seasonal variation in the prevalence of *Toxocara* eggs on children's playgrounds in the city of Hanover, Germany, *Parasite Vectors* 10 (May (1)) (2017) 248, <https://doi.org/10.1186/s13071-017-2193-6>.
- [15] J. Gawor, A. Borecka, Quantifying the risk of zoonotic geohelminth infections for rural household inhabitants in Central Poland, *Ann. Agric. Environ. Med.* 24 (March (1)) (2017) 44–48, <https://doi.org/10.5604/12321966.1230679>.
- [16] H. Mizgajska-Wiktor, et al., Distribution and dynamics of soil contamination with *Toxocara canis* and *Toxocara cati* eggs in Poland and prevention measures proposed after 20 years of study, *Vet. Parasitol.* 30 (January (234)) (2017) 1–9, <https://doi.org/10.1016/j.vetpar.2016.12.011>. Epub 2016 Dec 15.
- [17] H. Avcioglu, I. Balkaya, The relationship of public park accessibility to dogs to the presence of *Toxocara* species ova in the soil, *Vector Borne Zoonotic Dis.* 11 (February (2)) (2011) 177–180, <https://doi.org/10.1089/vbz.2009.0244>. Epub 2010 Jun 23.
- [18] M. Zibaei, et al., Soil contamination with *Toxocara* spp. eggs in the public parks from three areas of Khorram Abad, Iran, *Nepal Med. Coll. J.* 12 (June (2)) (2010) 63–65.
- [19] H. Borji, et al., A survey on endoparasites and ectoparasites of stray cats from Mashhad (Iran) and association with risk factors, *J. Parasit. Dis.* 35 (2) (2011) 202–6. Epub 2011 Aug 2.
- [20] N. Hajipour, et al. A survey study on gastrointestinal parasites of stray cats in Azarshahr, (East Azerbaijan province, Iran), *J. Parasit. Dis.* 40(4) (2016) 1255–1260. Epub 2015 Feb 28.
- [21] F. Mikaeili et al., *Toxocara* nematodes in stray cats from Shiraz, southern Iran: intensity of infection and molecular identification of the isolates, *Iran. J. Parasitol.* 8 (4) (2013) 593–600.
- [22] A. Mohsen, H. Hossein H., Gastrointestinal parasites of stray cats in Kashan, Iran, *Trop. Biomed.* 26 (1) (2009) 6–22.
- [23] M. Sharif, et al., *Toxocara cati* infections in stray cats in northern Iran, *J. Helminthol.* 1 (1) (2007) 63–6.
- [24] M. Yakhchali, et al. Gastrointestinal Helminths and Ectoparasites in the Stray Cats (Felidae: *Felis catus*) of Ahar Municipality, Northwestern Iran, Iran. *J. Parasitol.* 12 (2) (2017) 298–304.
- [25] M. Zibaei et al. Prevalence of *Toxocara cati* and other intestinal helminths in stray cats in Shiraz, Iran, *Trop. Biomed.* 24 (2) (2007) 39–43.
- [26] A.C. Becker, et al. Prevalence of endoparasites in stray and fostered dogs and cats in Northern Germany. *Parasitol. Res.* 111 (2) (2012) 849–57, <http://doi.org/10.1007/s00436-012-2909-7>. Epub 2012 Apr 13.
- [27] A. Montoya, et al. Implications of zoonotic and vector-borne parasites to free-roaming cats in central Spain, *Vet. Parasitol.* 15 (251) (2018) 125–130, <http://doi.org/10.1016/j.vetpar.2018.01.009>.
- [28] R. Nijse, et al. Environmental contamination with *Toxocara* eggs: a quantitative approach to estimate the relative contributions of dogs, cats and foxes, and to assess the efficacy of advised interventions in dogs, *Parasit. Vectors.* 8 (2015) 397, <http://doi.org/10.1186/s13071-015-1009-9>.
- [29] I.E. Sommerfelt et al., Prevalence of *Toxocara cati* and other parasites in cats' faeces collected from the open spaces of public institutions: Buenos Aires, Argentina, *Vet. Parasitol.* 140 (3-4) (2006) 296–301. Epub 2006 May 2.
- [30] E. Spada et al., Prevalence of faecal-borne parasites in colony stray cats in northern Italy, *J. Feline Med. Surg.*; 15 (8): (2013) 672–7, <http://doi.org/10.1177/1098612X12473467>. Epub 2013 Jan 17.
- [31] I. Symeonidou et al., Feline gastrointestinal parasitism in Greece: emergent zoonotic species and associated risk factors, *Parasit. Vectors.* 11 (1) (2018) 227, <http://doi.org/10.1186/s13071-018-2812-x>.
- [32] K. Szwabe, J. Blaszkowska, Stray dogs and cats as potential sources of soil contamination with zoonotic parasites, *Ann. Agric. Environ. Med.* 24 (1) (2017) 39–43, <http://doi.org/10.5604/12321966.1234003>.
- [33] H. Waap et al., Parasite communities in stray cat populations from Lisbon, Portugal, *J. Helminthol.* 88 (4) (2014) 389–95, <http://doi.org/10.1017/S0022149X1300031X>. Epub 2013 May 30.
- [34] H.L. Amaral, et al., Presence of *Toxocara canis* eggs on the hair of dogs: a risk factor for Visceral Larva Migrants, *Vet. Parasitol.* 174 (1-2) (2010) 115–8, <https://doi.org/10.1016/j.vetpar.2010.07.016>. Epub 2010 Aug 6. Epub 2010 Aug 6.
- [35] N.T. Anh, et al., Levels of *Toxocara* infections in dogs and cats from urban Vietnam together with associated risk factors for transmission, *J. Helminthol.* 4 (Jul 90) (2016) 508–10, <https://doi.org/10.1017/S0022149X15000619>. Epub 2015 Jul 30.
- [36] M. Aydenizöz-Ozkayhan, et al., The investigation of *Toxocara canis* eggs in coats of different dog breeds as a potential transmission route in human toxocarosis. *Vet. Parasitol.* 25; 152 (1-2) (2008) 94–100. <http://doi.org/10.1016/j.vetpar.2007.12.002>. Epub 2007 Dec 8.
- [37] W.F.El-Tras, et al., Risk of *Toxocara canis* eggs in stray and domestic dog hair in Egypt, *Vet. Parasitol.* 10; 178 (3-4) (2010) 319–23, <http://doi.org/10.1016/j.vetpar.2010.12.051>. Epub 2011 Jan 26.
- [38] J.D. Keegan, C.V. Holland, Contamination of the hair of owned dogs with the eggs of *Toxocara* spp., *Vet. Parasitol.* 11; 173 (1-2) (2010) 161–4, <http://doi.org/10.1016/j.vetpar.2010.06.010>. Epub 2010 Jun 15.
- [39] Y.F.F.B Meriguetti et al., Protective and risk factors associated with the presence of *Toxocara* spp. eggs in dog hair, *Vet. Parasitol.*, 15; 244 (2017) 39–43, <http://doi.org/10.1016/j.vetpar.2017.07.020>. Epub 2017 Jul 21.
- [40] B. Paoletti, et al. Zoonotic parasites in feces and fur of stray and private dogs from Italy. *Parasitol. Res.* 114 (6) (2015) 2135–41, <http://doi.org/10.1007/s00436-015-4402-6>. Epub 2015 Mar 17.
- [41] M. Tavassoli et al., Hair Contamination of Sheepdog and Pet Dogs with *Toxocara canis* Eggs, Iran. *J. Parasitol.* 7 (4) (2012) 110–5.