



Comparison of adverse events between cluster and conventional immunotherapy for allergic rhinitis patients with or without asthma: A systematic review and meta-analysis

Zihan Jiang, Hao Xiao, Hongting Zhang, Shixi Liu, Juan Meng*

Sichuan University (四川大学), West China Hospital, Department of Otorhinolaryngology, Chengdu, China

ARTICLE INFO

Keywords:

Rhinitis, allergic
Desensitization, immunologic/method
Desensitization, immunotherapy/adverse effects
Allergens/administration and dosage
Allergens/adverse effects
Allergen/therapeutic use
Humans

ABSTRACT

Background: Cluster schedule of allergen-specific immunotherapy (AIT) is a cost-effective choice for allergic rhinitis (AR) patients, but its safety has been questioned due to the greater dosages required at each treatment compared with conventional immunotherapy. It remains a question that whether cluster schedule leads to a higher risk of side effects.

Objective: This study was designed to update the evidence and investigate whether cluster schedule leads to a higher risk of local adverse reactions (LARs) and systemic adverse reactions (SARs) than cluster schedule does.

Methods: We searched the Cochrane Central Register of Controlled Trials, EMBASE and Medline thoroughly and included studies comparing cluster and conventional schedules. A meta-analysis of 5 outcomes related to adverse events was performed after bias and heterogeneity assessments. And as a result of language limitations, we considered only articles in Chinese and English.

Results: 5 observational studies and 6 interventional studies were included in the meta-analysis. There were no differences between cluster and conventional schedules when analyzing SARs by the number of patients, delayed SARs, grade 2 SARs and LARs. Analyses of SARs by injection, grade 1 SARs and LARs by injection in observational studies showed that cluster schedule had a lower risk of adverse events than did conventional schedule.

Conclusion: Our data suggest that cluster schedule is as safe as or even safer than conventional schedule for AR patients with or without asthma (AS).

1. Introduction

Allergen-specific immunotherapy (AIT) is an important treatment strategy for allergic rhinitis (AR) patients with or without asthma (AS). AIT is currently the only curative intervention that potentially rebalances the immune system and thus affects the natural course of allergic disease. The long-term efficacy and safety of AIT have been confirmed by many clinical trials and studies. Moreover, AIT can prevent the development of new sensitization [1] and prevent development of AR into AS [2,3].

Subcutaneous immunotherapy (SCIT) is the most universal route and its efficacy has been widely confirmed in clinical trials. The standard procedure of SCIT contains two phases: a dose-increase phase and a maintenance phase. Depending on different dose-increase regimens, SCIT can be divided into conventional schedule, cluster schedule and

rush schedule. With conventional schedule, one injection is administered per week, and a period of approximately 15 weeks is needed before reaching the maintenance dose. With cluster schedule, 2–3 injections are administered on a weekly visit, so the duration to reach the maintenance dose is shortened. Compared with conventional schedule, the cluster schedule cuts the time of the dose-increase phase nearly in half, so it is more convenient and more cost-effective for patients. However, this raises a question that whether cluster schedule leads to a higher risk of side effects and patients who are susceptible to adverse reactions should choose conventional schedule for safety concern. This systematic review aims to analyze the relevant literature comparing the risk of adverse reactions between these 2 schedules and provide evidence for clinical practice.

Abbreviations: AIT, allergen-specific immunotherapy; AR, allergic rhinitis; AS, asthma; LARs, local adverse reactions; SARs, systemic adverse reactions; SCIT, subcutaneous immunotherapy; RCTs, randomized controlled trials; PAR, perennial(or persistent) allergic rhinitis; SAR, seasonal allergic rhinitis; IAR, intermittent allergic rhinitis

* Corresponding author at: Sichuan University, Department of Otorhinolaryngology, West China Hospital, 37 Guoxue Alley, Chengdu 610041, China.

E-mail address: mjmelinda@163.com (J. Meng).

<https://doi.org/10.1016/j.amjoto.2019.07.013>

Received 24 May 2019

0196-0709/ © 2019 Elsevier Inc. All rights reserved.

2. Material and methods

2.1. Criteria of studies for this review

2.1.1. Types of studies

Considering that adverse reactions during allergen immunotherapy are rare, we decided to include both randomized controlled trials (RCTs) and other studies such as nonrandomized controlled trials, case-control studies, cross-sectional studies and both prospective and retrospective cohort studies, provided that they included a comparison of adverse events between cluster and conventional schedule therapies.

2.1.2. Types of participants

Patients suffering AR with or without AS were qualified regardless of their age, gender, kinds of allergens, duration and severity of disease and other coexisting conditions. All patients underwent objective examinations such as positive skin prick tests and elevated serum allergen-specific IgE tests to confirm their allergies.

2.1.3. Types of interventions

Conventional immunotherapy involves a dose accumulation phase of 8–16 weeks, and one injection is given on a weekly basis; on the other hand, cluster immunotherapy involves 2–3 injections given during one visit at intervals of 30 min, and the dose maintenance phase is reached within 8 weeks. After the build-up phase, patients under both schedules receive weekly injections until the end of the treatment. Studies involving only the dose accumulation phase or both phases of these 2 schedules were considered.

2.1.4. Types of outcome measures

Five aspects of adverse events related to AIT were analyzed: systemic adverse reactions (SARs), delayed SARs (happened 30 min after treatment), severity of SARs on the basis of a grade system in accordance with grading guidelines, local adverse reactions (LARs), delayed LARs (happened 30 min after treatment).

2.2. Search methods for the identification of studies

Published and unpublished articles in the EMBASE (1974 to 2019 February 01), Medline (1946 to February 01, 2019) and Cochrane Central Register of Controlled Trials (December 2018) databases were searched via OvidSP. The Medline search strategy is presented below, and the strategy when querying other databases was reorganized on the basis of the following example:

Search Strategy:

- 1 1 exp. Rhinitis, Allergic/ or exp Rhinitis, Allergic, Seasonal/ or exp Rhinitis, Allergic, Perennial/ (20374)
- 2 (allergic rhinitis or atopic rhinitis or (SAR or IAR or PAR)).af. (163365)
- 3 (hayfever or pollinosis or pollenosis or (hay and fever)).af. (5810)
- 4 ((allergen* or aeroallergen*) and rhinitis).af. (10767)
- 5 1 or 2 or 3 or 4 (177152)
- 6 exp immunotherapy/ or exp desensitization, immunologic/ (258009)
- 7 (immunotherapy or immunomodula*).af. (153720)
- 8 (immune or allergen* or antigen* or desensiti*).af. (1475094)
- 9 (therapy or vaccin* or inject*).af. (4559197)
- 10 8 and 9 (383754)
- 11 (cluster or accelerat*).af. (451277)
- 12 6 or 7 or 10 (634209)
- 13 5 and 11 and 12 (123)

2.3. Data extraction and analysis

2.3.1. Data extraction

Each study was viewed by 2 researchers. Four aspects of the data, including details of study design and methodology, characteristics of the population, description of the interventions and the statistics of the outcome measures, were collected with a prepared extraction sheet.

2.3.2. Quality assessment

The risk of bias of studies was assessed according to the Cochrane Collaboration tool and other scales recommended in the Cochrane handbook, such as the Newcastle-Ottawa Scale (NOS) for cohort studies and case-control studies. A total score more than or equal to 6 for NOS is qualified.

2.3.3. Data analysis

RevMan 5.3 was used for data processing. Trials and observational studies were analyzed separately. Chi-squared tests and the I^2 statistic were performed to assess heterogeneity, with $P < 0.1$ and $I^2 > 50\%$ indicating significant differences between studies and the necessity to develop critical explanations. The outcome measures mentioned above were all dichotomous. The Mantel-Haenszel method under fixed-effect model was used for analysis, and relative risks (RR) with 95% confidence intervals (CIs) were used to evaluate the safety of the schedules. All the tests were two-tailed. Sensitivity analysis and subgroup analysis would be applied when necessary.

3. Results

3.1. Description of studies

A total of 423 reports were identified through database searching, and 4 additional reports were identified from the reference lists. After removing duplicates, 359 records remained, of which 28 records were further assessed. Six studies were excluded because of the absence of full texts [4–9], 1 was excluded for repeated data [10]; therefore, 21 full-text articles were examined. Finally, 11 studies, including 5 RCTs [11–15] (1 full text was obtained by sending an e-mail request to the author [15]), 1 nonrandomized controlled trial [16] and 5 observational studies [17–21], met the eligibility criteria. The flow diagram of the study selection is illustrated in Fig. 1. The reasons for excluding the other 10 studies are as follows:

Five studies did not use a standardized conventional schedule [22–24] or cluster schedule [25,26],

One study compared a cluster schedule with a placebo [27],

Two were excluded for insufficient data [28,29], and

Two investigated only patients under the cluster schedule [30,31].

A total of 643 patients were investigated in the included trials: 319 were administered AIT under a cluster schedule, and 324 were administered AIT under a conventional schedule as a control (Table 1a). Five observational studies involved a population of 1907 participants: 966 were administered AIT under a cluster schedule, and 941 were administered AIT under a conventional schedule. Additional details about the characteristics of the studies are shown in Table 1b.

3.2. Risk of bias in the included studies

The risk of bias evaluation for trials was performed as recommended by the Cochrane handbook [32] and the accredited Newcastle-Ottawa Scale was used to evaluate the observational studies (Fig. 2 and Table 2). The result showed acceptable bias in all included studies.

3.3. Effects of interventions

Considering design heterogeneity, trials and observational studies were analyzed separately. SARs and LARs were assessed by counting




Fig. 1. Preferred reporting items for systematic reviews and meta-analyses (PRISMA) 2009 flow diagram.

the number of injections and participants.

3.3.1. Systemic adverse reactions in the build-up phase (by injection and by patient)

4 trials [11–13,16] reported systemic reactions during the initial phase by injection. A total of 3514 doses were administered to patients under the cluster schedule, and 3293 doses were administered under the conventional schedule, with incidence rates of systemic reactions of 19 and 22, respectively. No difference (RR = 0.89; 95% CI 0.49–1.64; $P = 0.72$) and low heterogeneity ($P = 0.77$; $I^2 = 0\%$) was discovered between groups. 139 patients received cluster regimen and 142 patients received the conventional regimen in 4 trials [12,13,15,16], with 20 and 17 patients, respectively, having systemic reactions. Analysis showed low heterogeneity between studies ($P = 0.76$; $I^2 = 0\%$) and no differences in the effects of the 2 regimens (RR = 1.23; 95% CI 0.69–2.19; $P = 0.49$).

3 observational studies [17,19,20] reported SARs data by injection during the initial phase including 4830 and 4006 injections in the cluster and conventional groups, with 17 and 29 total cases of SAR, respectively. The cluster schedule group was at a lower risk of SARs (RR = 0.47; 95% CI 0.25–0.88; $P = 0.02$) than was the conventional schedule group although significant heterogeneity was observed ($P = 0.01$; $I^2 = 77\%$; (Fig. 3). 19 out of 914 patients in the cluster group and 29 out of 899 patients in the conventional group experienced SARs in 4 observational studies [17,19–21]. The tests for overall effect

showed no significant differences between 2 groups (RR = 0.59; 95% CI 0.33–1.06).

3.3.2. Severity of systemic reactions in the build-up phase

4 trials [11–13,16] classified SARs by injection according to the guidelines of the Immunotherapy Subcommittee of the European Academy of Allergy and Clinical Immunology (EAACI) [33]. No grade 3 or grade 4 SARs were reported. There was no significant difference in the SAR incidence between the cluster and conventional schedule groups for grade 1 SARs (RR = 0.90; 95% CI 0.37–2.16; $P = 0.81$) and grade 2 SARs (RR = 0.89; 95% CI 0.38–2.07; $P = 0.79$) with accredited heterogeneity.

Analysis for 3 observational studies [17,19,20] indicated that the cluster schedule group was less likely to present grade 1 SARs (RR = 0.28; 95% CI 0.13–0.62; $P = 0.002$), although heterogeneity was observed between studies ($P = 0.09$, $I^2 = 58\%$) (Fig. 4a). For grade 2 SARs, the test revealed no difference (Fig. 4b).

3.3.3. Delayed systemic reactions

Two trials [12,14] reported no late-phase systemic reactions, while another trial reported that all SARs occurred after 30 min [16]. Thus, analysis was not applicable.

No difference was observed between cluster and conventional schedules according to the result for observational studies (Fig. 5).

Table 1a
Characteristics of the trials.

| Study design | Schedule | No. of patients | Age | Patients withdrawn | Patients with AR and AS | Vaccine | Build-up phase | Premedication | Outcomes |
|-----------------|------------------|-----------------|-----------------|--------------------|-------------------------|--|------------------------------|---------------|--|
| Fan, Q 2017 | Controlled trial | 30 | 23.5 (8.5-20) | 2 | 5 | <i>D. pteronyssinus</i> extract (Alutard SQ, ALK-Abelló) | 6w 14w | - | SAR, clinical efficacy |
| Tabar, A.I 2005 | Double-blind RCT | 30 | 22.0 (5.5-32.5) | 1 | 4 | Pangramin Depot UM D. | 6w | - | SAR, LAR, PEF, VAS, SPT, sigE, IgG, IgG4, etc. |
| Wang, C.S 2011 | RCT | 120 | 19.34 ± 9.8* | 13 | 94 | pteronysinus extract (Alutard SQ, ALK-Abelló) | 6w | - | LAR, SAR, T5SS, medication score, sigE, IgG4 |
| Zhang, L. 2009 | RCT | 119 | 18.47 ± 9.49* | 9 | 82 | <i>D. pteronyssinus</i> extract (Alutard SQ, ALK-Abelló) | 12w | - | LAR, SAR, medication score, QOL, VAS, SPT, IgE, etc. |
| Luo, Z 2015 | RCT | 40 | 26 (18,38) | 2 | 0 | <i>D. pteronyssinus</i> extract (Alutard SQ, ALK-Abelló) | 6w | - | SAR, RQLQ, medication score, clinical efficacy, etc. |
| Sandlos, C 2016 | RCT | 48 | 25 (15-36) | 1 | 0 | Stallergenes (France) and Immunotek (Spain) | 6w 14w 15w 3w 7w | - | SAR, clinical efficacy |

Note: “*”: Data in these cells is described by the mean ± standard deviation, while others is presented as median (25–75th percentiles). “-”: Data is not available. AR: allergic rhinitis. SAR: systemic adverse reaction. LAR: local adverse reaction. PEF: peak expiratory flow. VAS: visual analog scale. SPT: subcutaneous prick test. T5SS: total 5 symptom score. QOL: quality of life. RQLQ: mini rhinoconjunctive quality of life questionnaire.

Table 1b
Characteristics of the observational studies.

| Study design | Schedule | No. of patients | Age | Patients withdrawn | Patients with AR and AS | Vaccine | Build-up phase | Premedication | Outcome |
|----------------------------|---------------|-----------------|-------------|--------------------|-------------------------|---|----------------|---------------|----------|
| Garde, J. 2005 | Prospective | 66 | 28 ± 11.6 | 4 | 19 | <i>Salsola kali</i> (25 BU/ml, Pangramin® Depot BU, ALK-Abelló, S.A.) | 4w 13w | No | LAR, SAR |
| Martines-Canavate, A. 2005 | Retrospective | 22 | 9.6 ± 3.1 | - | 60 | <i>Alternaria alternata</i> 100% (Pangramin® Depot-UM, ALK-ABELLÓ) | 4w 7w | - | LAR, SAR |
| Barth, C. 2010 | Retrospective | 42 | 31 (6-71)* | 11 | 43 | Allergopharma-Joachim Ganzer, ALK-Scherax, Bencard Allergie, HAL Allergie, Leti Pharma and Stallergenes | - | - | LAR, SAR |
| Quiralte, J. 2013 | Retrospective | 75 | 26.2 ± 13.3 | - | 164 | Alustal Rapid®, Stallergenes Ibérica SA | - | - | LAR, SAR |
| Rodriguez Del Rio 2017 | Prospective | 339 | 26.7 ± 13.8 | - | 143 | Mites, pollen, altermaria, dander | 4w 8w | - | LAR, SAR |
| | Conventional | 437 | 11.7 ± 3.9 | - | - | | - | - | SAR |
| | Conventional | 483 | | | | | | | |

Note: “*”: Data in this cell is presented as mean(range), while others are presented as mean ± standard deviations. “-”: Data is not available. AR: allergic rhinitis. LAR: local adverse reaction. SAR: systemic adverse reaction.

| | | | | | | |
|----------------|----------------|------------------|-------------------|--------------|-------------|---|
| Zhang, L. 2009 | Wang, CS. 2011 | Tabar, A.I. 2005 | Sandlios, C. 2016 | Lou, Z. 2015 | Fan, Q 2017 | |
| + | ? | ? | ? | + | - | Random sequence generation (selection bias) |
| - | - | + | - | - | - | Allocation concealment (selection bias) |
| + | + | + | - | + | + | Blinding of participants and personnel performance (performance bias) |
| + | + | + | + | + | + | Blinding of outcome assessment (detection bias) |
| + | ? | - | + | + | ? | Incomplete outcome data (attrition bias) |
| + | + | - | + | + | + | Selective reporting (reporting bias) |
| + | + | + | + | + | + | Other bias |

Fig. 2. Risk of bias summary for the included trials.

Table 2
Newcastle-Ottawa Scale results for the observational studies.

| | Garde, J 2005 | Martines Canavate, A. 2005 | Barth, C.2010 | Quiralte, J. 2013 | Rodriguez Del Rio, P 2017 |
|--|---------------|----------------------------|---------------|-------------------|---------------------------|
| Selection | | | | | |
| Representativeness of the exposed cohort | 1 | 1 | 1 | 1 | 1 |
| Selection of the nonexposed cohort | 1 | 1 | 1 | 1 | 1 |
| Ascertainment of exposure | 1 | 1 | 1 | 1 | 1 |
| Demonstration that outcome of interest was not present at start of study | 1 | 0 | 0 | 0 | 1 |
| Comparability | | | | | |
| Comparability of cohorts on the basis of the design or analysis | 0 | 0 | 0 | 2 | 1 |
| Outcome | | | | | |
| Assessment of outcome | 1 | 1 | 1 | 1 | 1 |
| Was follow-up long enough for outcomes to occur | 1 | 1 | 1 | 1 | 1 |
| Adequacy of follow up of cohorts | 1 | 1 | 1 | 1 | 1 |
| Total | 7 | 6 | 6 | 8 | 8 |

| Study or Subgroup | Cluster schedule | | Conventional schedule | | Weight | Risk Ratio M-H, Fixed, 95% CI | Year | Risk Ratio M-H, Fixed, 95% CI |
|---|------------------|-------------|-----------------------|-------------|---------------|----------------------------------|------|----------------------------------|
| | Events | Total | Events | Total | | | | |
| Garde, J. 2005 | 4 | 528 | 2 | 286 | 8.4% | 1.08 [0.20, 5.88] | 2005 | |
| Barth, C.2010 | 8 | 1590 | 3 | 1168 | 11.2% | 1.96 [0.52, 7.37] | 2010 | |
| Quiralte, J. 2013 | 5 | 2712 | 24 | 2552 | 80.3% | 0.20 [0.07, 0.51] | 2013 | |
| Total (95% CI) | | 4830 | | 4006 | 100.0% | 0.47 [0.25, 0.88] | | |
| Total events | 17 | | 29 | | | | | |
| Heterogeneity: Chi ² = 8.57, df = 2 (P = 0.01); I ² = 77% | | | | | | | | |
| Test for overall effect: Z = 2.38 (P = 0.02) | | | | | | | | |

Fig. 3. SARs in the build-up phase by injection in observational studies.

3.3.4. Local adverse reactions in the build-up phase (by injection and by patient)

Only 2 trials [12,16] reported LARs, including 22 LAR cases out of 1026 injections in the cluster schedule group and 24 LAR cases out of 1112 injections in the conventional schedule group, during the build-up phase. Test showed no significant difference (RR = 1.01; 95% CI 0.57–1.78; P = 0.99) with a I² value of 0%. There was not enough data for analysis by patient.

Three observational studies [17,19,20] reported LARs, including 219 LAR cases out of 4830 injections in the cluster schedule group and 303 LAR cases out of 4006 injections in the conventional schedule group. Test indicated that cluster schedule caused a lower risk of LARs


in the build-up phase (RR = 0.64; 95% CI 0.54–0.76; P < 0.0001) (Fig. 6) with heterogeneity (I² = 67%; P = 0.05). No difference in LARs by patient between 2 groups (RR = 1.00; 95% CI 0.79–1.28; P = 0.98).

3.3.5. Delayed local adverse reactions


The data were insufficient for analysis.

3.3.6. Sensitivity analysis and publication bias

We made a sensitivity analysis by adding studies which was excluded for lacking full texts. Data relevant to our outcomes were acquired from the abstracts of 2 studies [6,8] and pooled into analysis for SARs by patient. The result illustrated that the risk of SARs by patient in



(a)



(b)

Fig. 4. a. Grade 1 SARs in observational studies.
b. Grade 2 SARs in the observational studies.




Fig. 5. Delayed SARs in observational studies.




Fig. 6. LARs in the build-up phase by injection in observational studies.




Fig. 7. SARs by patients in included observational studies and 2 observational studies lacking full texts.

cluster group is significantly lower (RR = 0.55; 95% CI 0.35–0.86; $P = 0.009$) with an acceptable heterogeneity ($I^2 = 42\%$; $P = 0.13$) (Fig. 7). The number of studies was inadequate for analyzing publication bias.

4. Discussion

As presented above, we analyzed statistics from trials and observational studies separately. Among the included studies, we detected statistical heterogeneity in some observational studies thus the result should be interpreted critically.

The tests for all outcomes in included trials showed a similar risk of adverse events between 2 schedules. However, significant lower RRs of cluster schedule were revealed in observational studies with respect to SARs in the build-up phase by injection, grade 1 SARs and LARs in the build-up phase by injection because of high statistical heterogeneity between Quiralte, J's study and others, which may indicate some publication bias.

Here, we attribute the variance to 4 main reasons. First, rare adverse events are less likely to be observed in clinical trials when the sample size is small. Apart from the double-blind RCT by Tabar [11], which included 239 individuals, the trials all included a relatively small sample size, although analyses by the number of injections could enlarge the sample size and eliminate some errors.

Second, every participant was given an antihistamine drug before the injections as premedication in 2 trials [12,14], 1 cohort study [17] reported no pretreatment, and relevant information was not available in the other studies. Previous studies suggested that premedication such as antihistamines and leukotriene antagonists could significantly reduce the incidence of adverse reactions [34,35]; however, further studies are necessary to confirm this idea.

Third, the duration of the dose-increasing phase and the number of doses administered at each visit were well controlled in the trials, except for one [15], but were variable in the observational studies. Finally, different types of allergens, including pollen, fungi, dander and mites were considered in the observational studies. In contrast, all but one trial applied vaccines of dust mite only.

Beyond the aforementioned factors, there are still some other varying elements, such as age and sex distribution and the number of patients with asthma, which may contribute to heterogeneity.

A 10-year retrospective study demonstrated that half of the systemic reactions to AIT were delayed and 10% occurred after 90 min when patients had already left the hospital [36]. Most of these reactions required medical treatment. Therefore, more attention should be paid to delayed reactions when assessing the risk of adverse reactions to immunotherapy.

Sensitivity analysis for included studies was not necessary since only one study showed inconsistent result, while analysis by adding 2 observational studies lacking full texts showed cluster schedule had lower hazard to develop SARs. This could be reliable because of acceptable statistical heterogeneity between studies.

5. Conclusion

Based on the above outcomes analyzed, our study suggests that cluster schedule is as safe as or even safer than conventional schedule for AR patients with or without AS. Large-scale and well-controlled RCTs are needed to address whether the exact duration of the build-up phase and the number of injections per visit for each schedule influence the occurrence of adverse reactions.

Acknowledgements

None.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] DesRoches A, Paradis L, Menardo JL, Bouges S, Daures JP, Bousquet J. Immunotherapy with a standardized Dermatophagoides pteronyssinus extract .6. Specific immunotherapy prevents the onset of new sensitizations in children. *J Allergy Clin Immunol* 1997;99:450–3. [https://doi.org/10.1016/s0091-6749\(97\)70069-1](https://doi.org/10.1016/s0091-6749(97)70069-1).
- [2] Jacobsen L, Niggemann B, Dreborg S, et al. Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy* 2007;62:943–8. <https://doi.org/10.1111/j.1398-9995.2007.01451.x>.
- [3] Niggemann B, Jacobsen L, Dreborg S, et al. Five-year follow-up on the PAT study: specific immunotherapy and long-term prevention of asthma in children. *Allergy* 2006;61:855–9. <https://doi.org/10.1111/j.1398-9995.2006.01068.x>.
- [4] Muro MD, Tabar AI, Garcia BE, Lizaso MT, Olaguibel JM. Cluster versus conventional immunotherapy in allergic to Dermatophagoides pteronyssinus. A controlled study of safety and efficacy. *Journal of Investigational Allergology and Clinical Immunology* 1997;7:472.
- [5] Jacobs J, Somani S, Turner R. Comparison of the safety of cluster vs. conventional immunotherapy schedules in a community based allergy clinic. *Ann Allergy Asthma Immunol* 2012;5:A122.
- [6] Minocha R, Detjen PF, Masood D, Story RE, Schafer E. Comparing systemic reaction rates and time to maintenance for inhalant aeroallergens in cluster versus standard immunotherapy in an academic affiliated hospital: a preliminary analysis. *Ann Allergy Asthma Immunol* 2013;1:A102–3.
- [7] Roger A, Depreux N, Jurgens Y, Heath MD, Garcia G, Skinner MA. Acarovac Plus development and clinical report of a novel tyrosine-adsorbed, modified house dust-mite allergen product. *Allergy: European Journal of Allergy and Clinical Immunology* 2014;99:561.
- [8] Penas-Aguilera A, Mascaró-Casanovas I, Feliu-Alsina M, Egues-Cachau G, Villalobos-Arebalo P, Basagana-Toronto M. Tolerance of a high-dose allergoid subcutaneous immunotherapy treatment in children. *Allergy: European Journal of Allergy and Clinical Immunology* 2017;72(Supplement 103):490.
- [9] Oskouei YM, Hosseini RF, Azad FJ, Rafatpanah H, Rezaei A, Esmaili H. Comparing cluster subcutaneous immunotherapy with conventional protocol in allergic rhinitis patients based on clinical and immunologic data. *Iranian journal of allergy, asthma and immunology* 2018; Conference: 3rd international congress of immunology, asthma and allergy. Iran: ICIAAI; 2017. p. 78–9. Islamic Republic of 17.
- [10] Quiralte J, Cardona V, Davila I, Moreno E, Ruiz B, Garcia M. Short conventional versus cluster immunotherapy for respiratory allergies: a retrospective safety comparison. *Allergy: European Journal of Allergy and Clinical Immunology* 2010;92:265.
- [11] Tabar AI, Echechipia S, Garcia BE, et al. Double-blind comparative study of cluster and conventional immunotherapy schedules with Dermatophagoides pteronyssinus. *J Allergy Clin Immunol* 2005;116:109–18.
- [12] Zhang L, Wang C, Han D, Wang X, Zhao Y, Liu J. Comparative study of cluster and conventional immunotherapy schedules with Dermatophagoides pteronyssinus in the treatment of persistent allergic rhinitis. *Int Arch Allergy Immunol* 2009;148:161–9.
- [13] Wang CS, Zhang W, Wang XD, et al. Comparative study on cluster and conventional immunotherapy with Dermatophagoides pteronyssinus in patients with allergic rhinitis. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi [Chinese Journal of Otorhinolaryngology Head and Neck Surgery]* 2011;46:981–5.
- [14] Luo Z, Li B, Wan L, Peng C. Comparative study on cluster and conventional immunotherapy in patients with allergic rhinitis. [Chinese]. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi [Chinese Journal of Otorhinolaryngology Head and Neck Surgery]* 2015;50:105–9.
- [15] Sandilos C, Kontogiorgaki I, Aggelides X, Makris M. Prospective randomized study of cluster immunotherapy in inhalant allergy: safety in comparison with conventional protocol. *Allergy, Asthma & Immunophysiology: Innovative Technologies* 2016:121–30.
- [16] Fan Q, Liu X, Gao J, Huang S, Ni L. Comparative analysis of cluster versus conventional immunotherapy in patients with allergic rhinitis. *Exp Ther Med* 2017;13:717–22.
- [17] Garde J, Ferrer A, Jover V, et al. Tolerance of a Salsola kali extract standardized in biological units administered by subcutaneous route. Multicenter study. *Allergol Immunopathol* 2005;33:100–4.
- [18] Martínez-Canavate A, Eserverri JL, Rodenas R, et al. Evaluation of paediatric tolerance to an extract of Alternaria alternata under two treatment regimes. A multicentre study. *Allergol Immunopathol* 2005;33:138–41.
- [19] Barth C, Anerio F, Pfaar O, Klimek L, Hormann K, Stuck BA. Safety aspects of subcutaneous immunotherapy with multiple allergens—a retrospective analysis on polysensitized patients. *Eur Arch Otorhinolaryngol* 2010;267:1873–9.
- [20] Quiralte J, Justicia JL, Cardona V, et al. Is faster safer? Cluster versus short conventional subcutaneous allergen immunotherapy. *Immunotherapy* 2013;5:1295–303.
- [21] Rodríguez Del Río P, Vidal C, Just J, et al. The European Survey on Adverse Systemic Reactions in Allergen Immunotherapy (EASSI): a paediatric assessment. *Pediatr Allergy Immunol* 2017;28:60–70. <https://doi.org/10.1111/pai.12660>.

- [22] Roger A, Depreux N, Jurgens Y, Heath MD, Garcia G, Skinner MA. A novel and well tolerated mite allergoid subcutaneous immunotherapy: evidence of clinical and immunologic efficacy. *Immunity, Inflammation and Disease* 2014;2(2):92–8. [Aug 2014].
- [23] Gandarias B, Alonso Ma D, Fernandez Rivas M, et al. Retrospective study of tolerance to short initiation schedules in subcutaneous immunotherapy. *Journal of Investigational Allergology and Clinical Immunology* 2005;15:242–8.
- [24] Roger A, Depreux N, Jurgens Y, et al. A prospective observational postauthorisation study of the safety, tolerability and effectiveness of Acarovac Plus: a modified, mite allergoid subcutaneous immunotherapy powered by MCT technology. *Allergy: European Journal of Allergy and Clinical Immunology* 2015;101:260.
- [25] Penas A, Garcia-Gonzalez M, Cruz MJ, et al. Observational study of the safety of a cluster schedule for subcutaneous immunotherapy in a pediatric population. *Journal of Investigational Allergology and Clinical Immunology* 2013;23:63–5.
- [26] Pfaar O, Van Twuijver E, Hecker H, Boot JD, Van Ree R, Klimek L. Accelerated up-dosing of subcutaneous immunotherapy with a registered allergoid grass pollen preparation. *Int Arch Allergy Immunol* 2014;160:420–4.
- [27] Guardia P, Moreno C, Justicia JL, et al. Tolerance and short-term effect of a cluster schedule with pollen-extracts quantified in mass-units. *Allergol Immunopathol* 2004;32:271–7.
- [28] Epstein TG, Liss GM, Murphy-Berendts K, Bernstein DI. AAAAI and ACAAI surveillance study of subcutaneous immunotherapy, year 3: what practices modify the risk of systemic reactions? *Ann Allergy Asthma Immunol* 2013;110:274–8.
- [29] Calderon MA, Vidal C, Rodriguez del Rio P, et al. European Survey on Adverse Systemic Reactions in Allergen Immunotherapy (EASSI): a real-life clinical assessment. *Allergy: European Journal of Allergy and Clinical Immunology* 2017;72:462–72.
- [30] Wang CS, Zhang L, Wang XD, Zhao Y, Liu JH, Zhang W, et al. 变应性鼻炎集群免疫治疗的疗效和安全性临床分析. *Chin J Otorhinolaryngol Head Neck Surg* 2008;43:187–91.
- [31] Serrano P, Justicia JL, Sanchez C, et al. Systemic tolerability of specific subcutaneous immunotherapy with index-of-reactivity-standardized allergen extracts administered using clustered regimens: a retrospective, observational, multicenter study. *Ann Allergy Asthma Immunol* 2009;102:247–52. [https://doi.org/10.1016/s1081-1206\(10\)60088-9](https://doi.org/10.1016/s1081-1206(10)60088-9).
- [32] Higgins JPT GSe. Cochrane handbook for systematic reviews of interventions version 5.1.0 [updated march 2011]. The Cochrane collaboration Available from www.handbook.cochrane.org; 2011.
- [33] Malling HJ, Weeke B. Position paper - immunotherapy. *Allergy* 1993;48. [9-8].
- [34] Wohrl S, Gamper S, Hemmer W, Heinze G, Stingl G, Kinaciyan T. Premedication with montelukast reduces local reactions of allergen immunotherapy. *Int Arch Allergy Immunol* 2007;144:137–42. <https://doi.org/10.1159/000103225>.
- [35] Yoshihiro O, Yoshinori N, Kiyotaka M. Effect of pretreatment with fexofenadine on the safety of immunotherapy in patients with allergic rhinitis. *Ann Allergy Asthma Immunol* 2006;96:600–5. [https://doi.org/10.1016/s1081-1206\(10\)63556-9](https://doi.org/10.1016/s1081-1206(10)63556-9).
- [36] Cook KA, Ford CM, Leyvas EA, Waalen J, White AA. Half of systemic reactions to allergen immunotherapy are delayed, majority require treatment with epinephrine. *Journal of Allergy and Clinical Immunology-In Practice* 2017;5:1415–7. <https://doi.org/10.1016/j.jaip.2017.03.025>.