



Angela K. Shen, MPH, Walter W. Williams, MD, MPH, Alissa C. O'Halloran, MSPH, Amy V. Groom, MPH, Peng-Jun Lu, MD, PhD, Alice Y. Tsai, MPH, Megan C. Lindley, MPH. Promoting Adult Immunization Using Population-Based Data for a Composite Measure. *Am J Prev Med.* 2018;55(4):517–523.

The editorial office has been informed of the following corrections.

In Table 3, the entry showing a % (95% CI) of 8.2 (7.2, 9.3) among those aged  $\geq 65$  years in the “Method 2: Td or Tdap (past 10 years)”, “Does not include influenza” composite measure vaccination coverage group was incorrect. The % (95% CI) for this group should have read 21.3 (20.0–22.7).

Likewise, the Results section incorrectly stated that 8.2% of adults in this age group fell into the Method 2 coverage group excluding influenza vaccination and including receipt of Td or Tdap in the past 10 years. It should have stated that 21.3% fell into this group.

Lastly, there were 9 instances in the Results section where a greater than (>) symbol appeared incorrectly, instead of the correct greater than or equal to ( $\geq$ ) symbol for each age group.

Interpretation of the data in the article was not affected by these errors. The corrected paragraphs in the Results section are reproduced below:

Table 3 presents a composite estimate of overall vaccination coverage among adults who have received the selected vaccines that are recommended for their age groups based on receipt of Tdap in the past 10 years (Method 1) or Td or Tdap in the past 10 years (Method 2), including or not including influenza vaccination. Vaccination coverage for the AIS measure including receipt of Tdap in the past 10 years (Method 1) and influenza vaccine was 11.9% for adults aged  $\geq 19$  years, 13.7% for adults aged 19–59 years, 6.3% for adults aged 60–64 years, and 7.4% for adults aged  $\geq 65$  years (Table 3). The AIS measure including receipt of Td or Tdap in the past 10 years (Method 2) and influenza vaccination was 23.4% for adults aged  $\geq 19$  years, 25.7% for adults aged 19–59 years, 12.6% for adults aged 60–64 years, and 19.2% for adults aged  $\geq 65$  years.

Excluding influenza vaccination, the AIS measure including receipt of Tdap in the past 10 years (Method 1) was 20.7% for adults aged  $\geq 19$  years, 25.2% for adults aged 19–59 years, 8.1% for adults aged 60–64 years, and 8.2% for adults aged  $\geq 65$  years. Vaccination coverage for the AIS measure excluding influenza vaccination and including receipt of Td or Tdap in the past 10 years (Method 2) was 51.4% for adults aged  $\geq 19$  years, 63.0% for adults aged 19–59 years, 15.8% for adults aged 60–64 years, and 21.3% for adults aged  $\geq 65$  years (Table 3).

Coverage with individual component vaccines is also shown in Table 3. Overall and for adults aged  $\geq 19$  years, component coverage was 43.3% for influenza received in the last 12 months, 61.6% for Td or Tdap received in the last 10 years, 23.1% for Tdap received in the past 10 years, 30.6% for herpes zoster ever received, and 63.6% for pneumococcal ever received.