

**Corrigendum**

**Corrigendum to Palliative Care Clinician Overestimation of Survival in Advanced Cancer: Disparities and Association With End-of-Life Care [Journal of Pain and Symptom Management 57 (2019) 233–240]**



The authors regret that incorrect figures were included in Table 3. A corrected table appears below. The errors do not change the study findings and all analyses have been verified. The authors would like to apologize for any inconvenience caused.

Corrected:

	Died	Hospice Ever		Hospice More Than 72 hours	
		<i>n</i>	OR (95% CI)	<i>n</i>	OR (95% CI)
Overestimate	116	75	<b>0.42 (0.21–0.85)</b>	52	<b>0.45 (0.24–0.83)</b>
No Overestimate	70	57	ref	45	ref
Adjusted, EOL preference					
Exposure Variable (Overestimate)			<b>0.38 (0.18–0.79)</b>		<b>0.41 (0.22–0.76)</b>
Covariates					
Strongly Agree			0.93 (0.22–3.86)		1.23 (0.34–4.41)
Somewhat Agree/Unsure/Somewhat Disagree			0.86 (0.20–3.68)		1.61 (0.43–5.97)
Adjusted, full model					
Exposure Variable (Overestimate)			<b>0.33 (0.14–0.78)</b>		<b>0.31 (0.14–0.65)</b>
Covariates					
Lung Cancer					
GI Cancer (non-CRC)					
Breast/CRC/Prostate Cancer					
Age (years, continuous)			1.03 (1.00–1.06)		
Black race			0.56 (0.21–1.52)		0.49 (0.19–1.26)
Latino ethnicity					2.48 (0.71–8.72)
Women					0.88 (0.48–1.61)
Palliative Performance Score (0-100, continuous)					1.15 (0.93–1.44)
Quality of Life (0-10, continuous)			1.14 (0.99–1.30)		
White Blood Cell count (continuous)			0.95 (0.91–1.02)		
Prognosticator: Physician			0.62 (0.29–1.33)		
Participation by Proxy			0.71 (0.54–0.92)		0.76 (0.56–1.02)
Financial strain			0.76 (0.51–1.14)		

## Published:

	Died	<i>n</i>	Hospice Ever	Hospice More Than 72 hours
			OR (95% CI)	<i>n</i> OR (95% CI)
Overestimate	116	75	<b>0.42 (0.21–0.85)</b>	52 <b>0.45 (0.24–0.83)</b>
No Overestimate	70	57	Ref	45 Ref
Adjusted, EOL preference				
Exposure Variable (Overestimate)			<b>0.38 (0.18–0.79)</b>	<b>0.41 (0.22–0.76)</b>
Covariates				
Strongly Agree			0.79 (0.21–2.95)	0.84 (0.22–3.15)
Somewhat Agree/Unsure/Somewhat Disagree			1.16 (0.30–4.50)	1.33 (0.34–5.19)
Adjusted, full model				
Exposure Variable (Overestimate)			<b>0.40 (0.16–0.99)</b>	<b>0.33 (0.15–0.74)</b>
Covariates				
Lung Cancer			1.92 (0.70–5.29)	1.91 (0.68–5.36)
GI Cancer (non-CRC)			0.37 (0.13–1.04)	0.35 (0.12–1.00)
Breast/CRC/Prostate Cancer			1.28 (0.45–3.62)	1.27 (0.44–3.64)
Age (years, continuous)			0.98 (0.95–1.01)	0.98 (0.95–1.01)
Black race			2.38 (0.79–7.21)	2.16 (0.68–6.79)
Latino ethnicity			12.14 (1.24–119.2)	11.36 (1.10–116.7)
Women			0.34 (0.15–0.75)	0.31 (0.14–0.69)
Palliative Performance Score (0-100, continuous)			1.72 (1.32–2.26)	1.77 (1.35–2.33)
Quality of Life (0-10, continuous)			1.11 (0.97–1.26)	1.11 (0.97–1.27)
White Blood Cell count (continuous)			1.05 (0.99–1.12)	1.05 (0.99–1.12)
Prognosticator: Physician			0.65 (0.31–1.39)	0.64 (0.30–1.38)
Participation by Proxy			1.19 (0.86–1.66)	1.24 (0.89–1.73)