



CORRIGENDUM

Corrigendum to “Polyvascular subclinical atherosclerosis in familial hypercholesterolemia: The role of cholesterol burden and gender” [Nutr Metab Cardiovasc Dis 29 (2019) 1068–1076]



Alessandro Mattina^{a,b,*}, Antonina Giammanco^a, Philippe Giral^{c,d}, David Rosenbaum^{c,d,g}, Alain Carrié^{d,e}, Philippe Cluzel^{d,f,g}, Alban Redheuil^{d,f,g}, Randa Bittar^h, Sophie Béliard^{i,j}, Davide Noto^a, Angelo Quartarone^b, Maurizio Averna^a, Éric Bruckert^{c,d}, Antonio Gallo^{c,d,g}

^a Department of Biomedicine, Internal Medicine and Medical Specialties, Division of Internal Medicine and Metabolic diseases, University of Palermo, Via del Vespro, 129, 90127, Palermo, Italy

^b IRCCS Centro Neurolesi “Bonibo-Pulejo”, Via Provinciale Palermo, Contrada Casazza, 95124, Messina, Italy

^c Cardiovascular Prevention Unit, Endocrinology and Metabolism Service, Assistance Publique Hôpitaux de Paris, Pitié-Salpêtrière Hospital, Paris, France

^d Sorbonne Université, Inserm, Institute of Cardiometabolism and Nutrition (ICAN), UMR_S1166, Hôpital de la Pitié, Paris, France

^e Assistance publique Hôpitaux de Paris (APHP), Hôpitaux Universitaires Pitié-Salpêtrière/Charles-Foix, Department of Biochemistry for Endocrinology and Oncology, Obesity and Dyslipidemia Genetics Unit, Paris, France

^f Département d'imagerie cardiovasculaire et de radiologie interventionnelle, Pôle Imagerie e Groupe Hospitalier Pitié-Salpêtrière, Assistance Publique - Hôpitaux de Paris, Paris, France

^g Imaging Core Lab, Institute of Cardiometabolism and Nutrition, ICAN, Paris, France

^h Biochemistry functional Unit for Metabolic Disease Assistance Publique/Hôpitaux de Paris, Groupe Hospitalier Pitié-Salpêtrière, Paris, France

ⁱ C2VN, INRA, INSERM, Aix-Marseille Univ, Marseille, France

^j Endocrinology Department, APHM, Marseille, France

The authors regret that in Table 1, among the lipid parameters, the total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C) and low-density lipoprotein cholesterol (LDL-C) values were reported inverted between men and women. As a consequence, the following sentence in “Results” (“Men showed higher TC and HDL-C levels [...]”) and the following paragraph in “Discussion” (“Interestingly in our cohort,

HDL-C levels were higher in men than in women; despite a role of hypolipidemic treatment cannot be excluded, this might in part explain the higher prevalence of carotid plaque in women than in men.”) is not to be considered. This error, however, does not change the scientific conclusion of the article.

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.numecd.2019.06.015>.

* Corresponding author. Department of Biomedicine, Internal Medicine and Medical Specialties, Division of Internal Medicine and Metabolic diseases, University of Palermo, Via del Vespro, 129, 90127, Palermo, Italy.

E-mail address: alessandromattina@gmail.com (A. Mattina).

<https://doi.org/10.1016/j.numecd.2019.09.014>

0939-4753/© 2019 The Italian Society of Diabetology, the Italian Society for the Study of Atherosclerosis, the Italian Society of Human Nutrition, and the Department of Clinical Medicine and Surgery, Federico II University. Published by Elsevier B.V. All rights reserved.