



Corrigendum to ‘Prevalence of cervical HPV infection in women with systemic lupus erythematosus: a systematic review and meta-analysis’. [AUTREV 18-2 (2019) 184–191]



Mario García-Carrasco^{a,b}, Claudia Mendoza-Pinto^{a,b}, Adriana Rojas-Villarraga^c,
Nicolás Molano-González^d, Verónica Vallejo-Ruiz^e, Pamela Munguía-Realpozo^{a,b},
Aurelio López Colombo^f, Ricard Cervera^{g,*}

^a Systemic Autoimmune Diseases Research Unit, Hospital de Especialidades, UMAE CMNMAC - CIBIOR, Instituto Mexicano del Seguro Social, Puebla, Puebla, Mexico

^b Department of Immunology and Rheumatology, Medicine School, Benemérita Universidad Autónoma de Puebla, Puebla, Puebla, Mexico

^c Fundación Universitaria de Ciencias de la Salud –FUCS, Bogotá, Colombia

^d Center for Autoimmune Diseases Research (CREA), School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia

^e Centro de Investigación Biomédica de Oriente, Instituto Mexicano del Seguro Social, Km. 4.5 Carretera Federal Atlixco-Metepec, Atlixco, C.P. 74360 Puebla, Mexico

^f Puebla Research Coordination, Highly Specialized Medical Unit, Speciality Hospital of Puebla, Instituto Mexicano del Seguro Social, Puebla, Mexico

^g Department of Autoimmune Diseases, Hospital Clinic, Barcelona, Catalonia, Spain

The authors would like to inform the readers of needed clarification regarding errors inserted in Table 3 of this article. (See Table 3.)

In the information on the group analysis for multiple sexual partners, the row corresponding to the information on the estimate of the parameter Σ^2 actually corresponds to the information on the interaction term between control-case status and multiple sexual partners

(Cases:MSP). The information reported for I^2 is actually the information for Σ^2 and the information for I^2 was not presented. In addition, we have changed the name of the column “Mean structure” to parameter, since parameters like Σ^2 and I^2 are not related to the mean structure of the models.

The authors wish to apologise for any inconvenience caused.

Table 3

Results of the most parsimonious models for the groups of studies.

Group	Parameter	Estimate	ci.lb	ci.ub	pval
Case-controls	intrept	-1.7127	-2.4708	-0.9546	< 0.0001
	Cases	1.056	0.7872	1.3247	< 0.0001
	Σ^2	1.2107	0.4373	4.3989	< 0.0001
	I^2				95.42743
Multiple sexual partners	intrept	-1.7222	-2.6954	-0.749	0.0005
	Cases	0.986	0.524	1.4481	< 0.0001
	MSP	0.0117	-0.023	0.0463	0.5091
	Cases: MSP	-0.0042	-0.0305	0.022	0.7513
	Σ^2	0.8324	0.268	41,482	< 0.0001
	I^2				93.96373
Immunosuppressive therapy	intrept	1.2869	0.3323	2.2414	0.0082
	HPV-	-0.0453	-0.5163	0.4257	0.8504
	Σ^2	0.9365	0.1798	6.7102	< 0.0001
	I^2				87.4806

Heterogeneity by means of Higgins's tests.

Abbreviation: ci.lb.: 95% confidence interval lower bound; ci.ub: 95% confidence interval upper bound; intrept: intercept; MSP: multiple sexual partners.

DOI of original article: <https://doi.org/10.1016/j.autrev.2018.09.001>

* Corresponding author at: Servei de Malalties Autoimmunes, Hospital Clínic, Villarroel 170, 08036 Barcelona, Catalonia, Spain.

E-mail address: rcervera@clinic.ub.es (R. Cervera).

<https://doi.org/10.1016/j.autrev.2019.02.001>

Available online 12 February 2019

1568-9972/ © 2019 Elsevier B.V. All rights reserved.