



## Review article

## Decoding the enigma of antiviral crisis: Does one target molecule regulate all?

Abdullah Mahmud-Al-Rafat<sup>a,b,\*</sup>, Apurba Majumder<sup>c</sup>, K.M. Taufiqur Rahman<sup>d</sup>,  
A.M. Mahedi Hasan<sup>e</sup>, K.M. Didarul Islam<sup>b</sup>, Andrew W. Taylor-Robinson<sup>f</sup>, Md Morsaline Billah<sup>b</sup>

<sup>a</sup> Research & Development Division, Incepta Vaccine Ltd., Zirabo, Savar, Dhaka 1341, Bangladesh

<sup>b</sup> Biotechnology and Genetic Engineering Discipline, Khulna University, Khulna 9208, Bangladesh

<sup>c</sup> Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School, Hannover 30625, Germany

<sup>d</sup> Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Ontario K1N6N5, Canada

<sup>e</sup> Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh EH9 3FF, UK

<sup>f</sup> School of Health, Medical & Applied Sciences, Central Queensland University, Brisbane, QLD 4000, Australia

## ARTICLE INFO

## Keywords:

Virus  
Ebola  
SARS-CoV  
Dengue  
Cytokine storm  
Antiviral  
Pro-inflammatory response  
IL-6  
ADAM-17

## ABSTRACT

Disease fatality associated with Ebola, SARS-CoV and dengue infections in humans is attributed to a cytokine storm that is triggered by excessive pro-inflammatory responses. Interleukin (IL)-6 acts as a mediator between pro- and anti-inflammatory reactivity by initiating trans- and classical-signaling, respectively. Hence, IL-6 is assumed to provide a target for a broad range of antiviral agents. Available immunosuppressive antivirals are directed to control an often exaggerated pro-inflammatory response that gives rise to complex clinical conditions such as lymphocytopenia. It is known that IL-6, via its soluble receptor (sIL-6R), initiates a pro-inflammatory response while an anti-inflammatory response is triggered by the membrane-bound IL-6 receptor (IL-6R). Future antivirals should thus aim to target the mechanism that regulates switching between IL-6 trans- and classical-signaling. In this review, we propose that the tumour necrosis factor- $\alpha$  converting enzyme ADAM-17 could be the master molecule involved in regulating IL-6 class switching and through this in controlling pro- and anti-inflammatory responses to viral antigenic stimuli. Therefore, ADAM-17 should be considered as a potential target molecule for novel antiviral drug discovery that would regulate host reactivity to infection and thereby limit or prevent fatal outcomes.

## 1. Introduction

Physiology or Medicine, 1937)

“Research is to see what everybody else has seen, and to think what nobody else has thought.”

Albert Szent-Györgyi, 1957; (Winner of the Nobel Prize in

In recent years broad spectrum antiviral drugs have drawn much attention because of the prospect of their providing humans with greater resistance to major viral diseases. Such drugs should not only be capable of protecting against multiple viruses but should also reduce

**Abbreviations:** ACTH, Adrenocorticotrophic hormone; ADAM-17, a disintegrin and metalloproteinase-17; ADE, antibody-dependent enhancement; ARDS, acute respiratory distress syndrome; BEBOV, Bundibugyo Ebolavirus; CCL5, chemokine (C-C motif) ligand; CTLA-4, cytotoxic T-lymphocyte-associated protein 4; CXCL, chemokine (C-X-C motif) ligand; DC, dendritic cell; DENV, dengue virus; DF, dengue fever; DHF, dengue hemorrhagic fever; DIC, disseminated intravascular coagulation; DSS, dengue shock syndrome; EBOV, Ebolavirus; FasL/FasLG, Fas ligand; ICEBOV, Ivory Coast Ebolavirus; IFN, interferon; IFNAR, interferon- $\alpha/\beta$  receptor; IL-1RA, interleukin-1 receptor antagonist; IL, interleukin; IP-10, interferon gamma-induced protein 10; ISG, interferon-stimulated gene; GRO- $\alpha$ , growth-regulated protein alpha; M-CSF, macrophage colony-stimulating factor; MCP-1, monocyte chemoattractant protein 1; MIF, macrophage migration inhibitory factor; MIP, macrophage inflammatory protein; MMP, matrix metalloproteinases; NF- $\kappa$ B, nuclear factor kappa B; NHP, non-human primate; PD-1, programmed cell death protein 1; RANTES, regulated on activation normal T cell expressed and secreted; REBOV, Reston Ebolavirus; rVSV-ZEBOV, recombinant vesicular stomatitis virus–Zaire Ebola virus; SARS, severe acute respiratory syndrome; SARS-CoV, severe acute respiratory syndrome coronavirus; SEBOV, Sudan Ebolavirus; TGF- $\beta$ 1, transforming growth factor beta 1; TNF, tumour necrosis factor; TRAIL, TNF- $\alpha$  related apoptosis-inducing ligand; T<sub>reg</sub>, T-regulatory cell; VHF, viral hemorrhagic fever; ZEBOV, Zaire Ebolavirus

\* Corresponding author at: Research & Development Division, Incepta Vaccine Ltd., Zirabo, Savar, Dhaka 1341, Bangladesh.

E-mail addresses: [mahmud0907@gmail.com](mailto:mahmud0907@gmail.com), [rafat@inceptavaccine.com](mailto:rafat@inceptavaccine.com) (A. Mahmud-Al-Rafat), [krahm066@uottawa.ca](mailto:krahm066@uottawa.ca) (K.M. Taufiqur Rahman), [A.M.M.Hasan@ed.ac.uk](mailto:A.M.M.Hasan@ed.ac.uk) (A.M. Mahedi Hasan), [a.taylor-robinson@cqu.edu.au](mailto:a.taylor-robinson@cqu.edu.au) (A.W. Taylor-Robinson), [morsaline@yahoo.com](mailto:morsaline@yahoo.com) (M.M. Billah).

<https://doi.org/10.1016/j.cyto.2018.12.008>

Received 10 June 2018; Received in revised form 2 December 2018; Accepted 3 December 2018

Available online 04 January 2019

1043-4666/ © 2018 Elsevier Ltd. All rights reserved.

the cost associated with drug/vaccine development against those specific pathogens. There is a paucity of commercially available, effective preventive therapeutics and vaccines to resist the emergence and re-emergence of virus challenges. In order to establish broad spectrum antiviral therapeutics an innovative approach by which scattered research outcomes could be brought together under one umbrella is essential.

In this article, we postulate a novel hypothesis that aims to provide a guiding principle for future therapeutic antiviral research. This is underpinned by the examination of three infectious diseases of global public health importance; Ebola, severe acute respiratory syndrome (SARS) and dengue. These viral diseases share many features in common and hence an insight into their pathogenic features could reveal a common target molecule for future antiviral development. We propose the role of a master molecule, ADAM-17, as a potential therapeutic target against these three diseases. The cascades of events in the pro-inflammatory response are discussed in search of a common therapeutic target molecule that is evaluated computationally. This leads to the suggestion of a possible novel antiviral remedy. To our knowledge, this is the first paper to consider multiple emerging infectious diseases, to evaluate their common pathway of pathogenesis and to propose a prospective antiviral target molecule.

## 2. General features of Ebola, SARS-coronavirus (SARS-CoV) and Dengue

### 2.1. Ebola

Ebola has a typical incubation period of 4–10 days, following which an infected person develops a fever that rises rapidly to  $> 100^{\circ}\text{F}$  ( $> 37.8^{\circ}\text{C}$ ). Clinical symptoms may include myalgia, nausea, vomiting and/or diarrhoea, impaired coagulation, subconjunctival haemorrhage, bruising and defective clotting at venipuncture sites; blood in urine or faeces may even be visible. In severe cases, patients may suffer visceral organ necrosis along with haemorrhaging into the skin and mucous membranes [1].

### 2.2. SARS-CoV

A typical incubation period of severe acute respiratory syndrome coronavirus (SARS-CoV) is 2–10 days, while the average duration between first clinical symptoms and hospital admission is 3–5 days [2]. The patient usually shows tenacious fever, myalgia, malaise, dry cough, headache and dyspnoea as primary clinical symptoms [3–5]. However, sputum production, sore throat, rhinorrhoea, nausea, vomiting and diarrhoea may also be noticeable as minor signs. Lymphocytopenia along with intravascular coagulation, an elevated level of lactate dehydrogenase and thrombocytopenia, bilateral viral pneumonia, acute respiratory distress syndrome (ARDS) are the most common adverse clinical features of SARS [3–6].

### 2.3. Dengue

Clinical manifestations of dengue range from dengue fever (DF) to more severe forms like dengue haemorrhagic fever (DHF) and, sometimes, dengue shock syndrome (DSS) [7]. Arthralgia, ostealgia, myalgia, headache and skin rash are the most common clinical symptoms of DF [8,9], which persists for around one week, hence the common names of ‘breakbone fever’ and ‘seven-day fever’ [10]. There are four known dengue virus (DENV) serotypes, each of which can cause infection in humans. Secondary dengue infection with a heterotypic serotype to that responsible for primary infection can initiate the immunopathological phenomenon of antibody-dependent enhancement (ADE) of infection, which may result in DHF and DSS. Clinical manifestations of DHF include haemorrhagic tendencies, plasma leakages and thrombocytopenia, each in addition to existing symptoms of DF. Patients who

develop DSS exhibit all the symptoms of DF and DHF along with circulatory failure, hypotension and signs of systemic shock [7,11].

## 3. Pro-inflammatory responses follow a similar pattern of elevated cytokine levels

### 3.1. Ebola

The pathophysiology of Ebola is characterized by the initiation of an uncontrolled cytokine storm that includes overexpression of pro- and anti-inflammatory cytokines, chemokines, free radicals and coagulation factors [12]. All Ebola-related fatalities are notable for triggering the pro-inflammatory cytokines IL-1 $\beta$ , IL-1RA, IL-6, IL-8, TGF-1 $\beta$  and TNF- $\alpha$  and the chemokines MIP-1 $\alpha$ , MIP-1 $\beta$  and MCP-1. [13,14]. The inability of the human immune system to curb this cytokine storm through means of a negative feedback loop results in significant tissue and organ damage that frequently leads to case fatality [15]. At the onset of Ebolavirus (EBOV) infection the RNA polymerase cofactors VP35 and VP24 can inhibit  $\alpha/\beta$  interferon responses, the first line of antiviral defence. VP35 antagonizes the type 1 interferon response while VP24 is believed to disrupt both type 1 and type 2 interferon responses. VP35 and VP24 are also capable of inhibiting DC maturation and of initiating subsequent pro-inflammatory responses by immature DCs [16]. Activated macrophages produce TNF- $\alpha$ , which plays a major role in inflammatory responses. TNF- $\alpha$  promotes migration of immune cells and activates apoptosis caspases. Raised levels of TNF- $\alpha$  are observed in fatal cases of Ebola. TNF- $\alpha$  also disrupts the endothelial barrier, causing cell leakage [17,18].

Clinical laboratory testing performed by several research groups on samples isolated from Ebola-infected patients during the remarkable West African epidemic of 2014 showed substantial immune activation to correlate with excessive inflammatory responses. Increased numbers of activated plasmablasts, CD4<sup>+</sup> T cells and CD8<sup>+</sup> T cells were observed in all infected cases when compared to healthy control individuals. Significantly higher levels of pro-inflammatory cytokines and chemokines were detected in all fatal cases compared to non-fatal cases. Analysis of animal models revealed profound up-regulation of interferon-stimulated genes (ISGs) from the early stage of the infection [14,19]. Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) expressed by CD8<sup>+</sup> T cells was detected at a markedly higher level in non-surviving patients. Both programmed cell death protein 1 (PD-1) and CTLA-4 concentrations were found much higher in fatal cases compared to survivors. Each of these proteins plays a role in regulating T cell homeostasis. Elevated expression of PD-1 and CTLA-4 is correlated with T cell functional exhaustion as well as with poor viral clearance [20,21].

### 3.2. SARS-CoV

SARS-CoV infection is remarkable for the overexpression of cytokines IL-6, IL-2, IL-5, TNF- $\alpha$  and IFN- $\gamma$ , and of chemokines CXCL10, CCL2, CCL3 and CCL5. Disease severity is correlated with the production of a massive cytokine storm, lung infiltration by monocytes and macrophages, and a depletion of the lymphocyte population [22,23]. Studies have revealed that host fatality is in fact not related with SARS-CoV virus replication but with virus-induced unregulated immune responses [24]. Histological examination of a mouse model of SARS two days post-infection showed a high viral titre with an increased presence of macrophages and some other T-lymphocytes in the focal perivascular infiltrate within the lungs. Along with heightened immune cell activation, bronchiolar epithelial necrosis and luminal necrotic debris were also visible. Migration of plasmacytoid DC and inflammatory cells (such as NK T cells, NK cells, macrophages and CD4<sup>+</sup> T cells) to lungs was observed during early stage infection. A significant elevation of inflammatory cytokine and chemokine levels was also detected in early infection compared to the healthy control group. CD8<sup>+</sup> T cells and

neutrophils were found to migrate to the lungs during the late stage of infection [25,26].

### 3.3. Dengue

Infection with DENV activates the innate immune system of the host. *In vitro* investigation has revealed an increase in pro-inflammatory cytokines upon infection. DENV NS5 and NS4B proteins induce infected macrophages and endothelial cells to produce a range of pro-inflammatory cytokines including IL-8, IL-6, MIF, MCP, CXCL10, CXCL11 and RANTES [27–29]. These mediators increase permeability and play role in inflammation and plasma leakage *in vivo* [28,30,31]. Infected DCs produce MMP-2 and MMP-9 which facilitate DC migration to lymph nodes [32]. An increased level of type 1 IFN is positively correlated with viral NS1 protein expression [33]. One of the most studied cytokines in dengue infection is TNF- $\alpha$  which is expressed by both innate and adaptive immune cells. Raised levels of TNF- $\alpha$  are reported particularly in DHF cases [34,35]. Plasma leakage and coagulopathy are two ways by which TNF- $\alpha$  plays a role in disease severity. Elevated IL-10 is observed in late infection, indicative of the activation of Treg cells in response to the preceding cytokine storm [30,36].

Histopathological examination of samples from infected patients has shown pronounced immune activation in specific organs. Monocytes/macrophages and DCs are the major site of DENV infection and replication [7,37,38]. A significant increase of T lymphocytes may be detected in fatal DENV cases compared to non-fatal groups. Further investigation revealed a notable rise in cells producing inflammatory cytokines in the liver of infected patients. The presence of nuclear vacuolar degeneration and swollen mitochondria in infected cases also indicates a mechanism involving cell death. Lung tissue samples from fatal cases show severe damage characterized by increased cellularity, the presence of mononuclear inflammatory infiltrates and hyperplasia of alveolar macrophages [39]. Histopathological and ultrastructural analysis of heart, kidney and spleen detected damage to target organs in DENV-infected patients but no damage in healthy samples. Quantification analysis revealed the presence of lymphocytes, inflammatory cytokines and chemokines at concentrations many fold higher than normal. Significant involvement of apoptosis is also detected in target organ biopsies from fatal cases [39,40].

### 4. Features of a cytokine storm common to viral infections

The three viral infections described above share some common features that establish a cytokine storm, as well as leading to pathogenic outcomes [41]. Following primary exposure to antigens, innate immunity acts as the first line of defence. Studies have revealed that monocyte/macrophage and DCs infiltrate the early sites of infection and virus replication [13,42,43], from where these immune cells containing virus particles later migrate to other organ and tissues [20,39,44]. In order to evade the innate immune response, the virus must inhibit the type-1 interferon (IFN- $\alpha$  and IFN- $\beta$ ) response. Analysis of both *in vitro* and *in vivo* models strongly suggests that EBOV, SARS-CoV and DENV are each able to evade type-1 IFNs responses in order to establish infection [37,45–48]. Viral entry and infection are characterized by the expression of extremely high levels of pro-inflammatory cytokines and chemokines by T lymphocytes [14,19,21,25,39]. Their excessive provokes a cytokine storm that is responsible for plasma leakage, vascular permeability and disseminated intravascular coagulation (DIC) as well as defective adaptive immunity [37,49,50]. These pathophysiological events enable efficient dissemination of virus, providing a metabolic milieu in which a fatal outcome may result [7,51]. Apoptosis of lymphocytes can further facilitate such disease-related mortality [52]. Multiple *in vivo* studies have reported a massive level of lymphocyte apoptosis in these three infections [24,53]. Lymphocytopenia is commonly observed among fatalities but is absent in survivors [13,54].

Focusing on the key mediators involved in this pro-inflammatory response should help to identify any common target molecule(s) of potential therapeutic interest. Among all the different pro-inflammatory cytokines that are caught up in a cytokine storm, IL-6 is of primary interest because of its role as transition cytokine. IL-6 may be considered as the master cytokine that can change its nature to serve in both pro- and anti-inflammatory [55]. Besides this, IL-6 is responsible for bridging the gap between innate and adaptive arms of the immune response [56,57]. In view of this, it is a logical step to review the characteristics of IL-6 and its dual role as well as to determine how its performance may be triggered under infection conditions. Thus, we focus on the function of IL-6 in the following section and discuss how this keystone cytokine contributes to infection immunity.

On the flip side, IL-10 also merits attention because it is the only anti-inflammatory cytokine that is excessively expressed. In almost all fatal cases of viral infections, the level of IL-10 becomes many folds higher than that detected in survivors [13,58,59]. It has been speculated that, as an act of homeostasis, IL-10 is involved in counter-regulating the pro-inflammatory immune response before the cytokine environment becomes extreme [60]. However, the role of IL-10 remains unclear and extensive investigation is required to unravel the complex mechanisms involved. Researchers are particularly interested in dissecting the pathway that triggers the cytokine storm phenomenon and in understanding subsequent cellular events that relate to disease fatality. Below, based on our hypothesis we aim to explain why IL-10 concentrations are elevated in viral infections that are fatal.


### 5. IL-6: Beauty and the beast

IL-6 is sustained at elevated levels in all fatal cases of viral infection, which earmarks it as an important molecule to investigate. In recent years, IL-6 has been considered to be a keystone cytokine in health and disease due to its pleiotropic characteristics [55]. It has a unique ability to function in both pro- and anti-inflammatory responses depending on specific conditions [61]. In light of its decisive role during cytokine storm-mediated immunopathogenesis, IL-6 provides a principal focus of this review – notably, in regard to how IL-6 regulates immune reactivity and the way in which this relates to disease fatality.

IL-6 signaling starts with its binding to the receptor IL-6R, after which the IL-6/IL-6R complex associates with gp130. IL-6R exists in two different forms: membrane-bound receptor (IL-6R) and soluble receptor (sIL-6R). Membrane-bound IL-6 receptors can become soluble through shedding or differential splicing, while the former one is the more frequent event [56,62–64]. Depending on receptor states, IL-6 signaling may be either of two types: classic signaling and trans-signaling. Classic signaling requires binding of IL-6 with membrane-bound IL-6R whereas trans-signaling starts when IL-6 binds with sIL-6R [65–67]. IL-6 trans-signaling is pro-inflammatory while classic signaling is anti-inflammatory in nature. Details of these mechanisms are beyond the scope of this review. Instead, we focus here on the involvement of these two different forms of IL-6 in immunopathogenesis.

IL-6 is the pioneer cytokine to initiate pro-inflammatory responses during infections. IL-6 trans-signaling correlates positively with increased production of pro-inflammatory cytokines and chemokines including IL-10, IL-1 $\beta$ , IL-1RA, TNF, M-CSF, ACTH and cortisol [56,62,67–70]. IL-6 trans-signaling is also shown to initiate monocyte to macrophage differentiation [56,68], to attract other immune cells while also suppressing T<sub>reg</sub> cells [71,72]. In addition, IL-6 trans-signaling is involved in the progression and manifestation of autoimmune diseases [73].

Switching between two different forms of IL-6 is dependent on the concentrations of IL-6 and sIL-6R. Under normal physiological conditions, the plasma level of IL-6 is minute, ranging between 2 and 6 pg/mL [74–76], while sIL-6R is detected at about 75 ng/mL [77,78]. Therefore, in a steady state, the concentration of sIL-6R is more than 10,000 times higher than that of IL-6. During inflammation, however,



**Fig. 1.** General overview of disease progression. A: Virus detected by innate immune cells (macrophage, dendritic cell). Innate immune cells show immune response by secretion of major pro-inflammatory cytokine IL-6 and TNF. GC and IL-10 are also activated by negative feedback system. B: Apoptotic proteins activated by GC, IL-10 and TNF. Necrosis/apoptosis results in lymphocytopenia. ADAM-17 is activated. C: Two different signaling pathway of IL-6. Trans signaling (marked in red arrow) continues the pro-inflammatory response. Classical signaling (marked in blue arrow) facilitates anti-inflammatory response. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

IL-6 levels may increase up to 1 million fold [73,76] whereas sIL-6R rises typically only 2–5 fold [79,80]. Once challenged by an antigen, upon secretion IL-6 will first bind to sIL-6R and initiate pro-inflammatory responses. However, only when IL-6 levels exceed that of sIL-6R does it bind to IL-6R and thereby triggers an anti-inflammatory response. Thus, in order to maintain a sustained pro-inflammatory response and thereby block the anti-inflammatory response, the systemic concentration of sIL-6R must be higher than that of IL-6 [73]. It has been observed that selective blocking of IL-6 trans-signaling rather than the global IL-6 blockade is a more effective means to rescue mouse models from possible death [81]. Hence, IL-6 may be considered as an immunoregulatory master molecule the control of which governs pathogenic outcomes. Yet, there are still many questions surrounding the mechanism of action of IL-6 that need to be answered. Even if IL-6 is held responsible for escalating rates of fatality, theoretically at least, elevated IL-6 concentrations should eventually suppress levels of sIL-6R and thereby facilitate a switch back to anti-inflammatory responses during viral infection. However, quite the opposite is observed in fatal cases, suggestive of a complex mechanism behind this phenomenon. sIL-6R concentrations somehow remain high during the entire period of infection, thereby establishes a persistent pro-inflammatory response via trans-signaling. Thus, it can be reasonably presumed that there are other biomolecules interacting in this process which influence the sustained trans-signaling and result in cytokine storm-mediated fatality.

## 6. ADAM-17: where all light tends to go

Since its discovery in 1997, ADAM-17, commonly known as tumour necrosis factor (TNF)- $\alpha$  converting enzyme, has drawn enormous interest as a potential therapeutic target in multiple disease pathways. ADAM-17 is closely associated with TNF- $\alpha$  and other inflammatory cytokines [82,83]. It has been found that TNF- $\alpha$ , IL-1 $\beta$  and IFN- $\gamma$  up-regulate the transcription and expression of ADAM-17 [84]. Researchers have proposed that ADAM-17 can release the membrane-bound TNF- $\alpha$  precursor to a soluble form which facilitates TNF- $\alpha$ -mediated inflammatory processes [85]. Of most relevance here, ADAM-17 is believed to be the master molecule that may explain uncontrolled IL-6 trans-signaling and increasing pro-inflammatory responses during infection. This is because ADAM-17 is the key protein responsible for membrane-bound IL-6R shedding and thus the production of sIL-6R [56,86]. Apoptosis has been shown to act as a natural stimulus in ADAM-17-mediated IL-6R shedding from the surface of neutrophils and thereby contributes to pro-inflammatory trans-signaling responses [87]. Uncontrolled IL-6 trans-signaling could also be explained by this mechanism, from which a picture of how infection worsens can be envisaged.

During infection, in order to maintain homeostasis production of inflammatory cytokines is followed by a counter release of anti-inflammatory cytokines. IL-6 increases the level of plasma glucocorticoid (cortisol) and IL-10 while also reducing the absolute number of

lymphocytes [69]. The lymphocyte loss observed in pathogenesis is not mediated solely by apoptotic machinery, but also involves necrosis, a non-apoptotic form of cell death. This event is mediated by elevated levels of TNF- $\alpha$  and IL-10 [13,88]. This phenomenon can explain why anti-inflammatory cortisol and IL-10 levels remain high alongside lymphocytopenia in fatal cases. Detailed mechanism of lymphocyte loss is not yet understood, though it is believed that, multiple mediators (Fas/FasL, TNF- $\alpha$  related apoptosis-inducing ligand (TRAIL), pro-apoptotic gene Bim and Bid) triggered by TNF- $\alpha$ , play a pivotal role in this type of cell death [13,89,90]. Interaction of ADAM-17 with these mediators is reported in several studies which show how ADAM-17 is activated during neutrophil loss and then proceeds to generate more sIL-6R by cleaving IL-6R [87,91,92]. We propose this as the mechanism by which the level of sIL-6R remains higher than that of IL-6R throughout infection and explains how IL-6 skips the cis-signaling pathway. This event ultimately disrupts the balance between pro- and anti-inflammatory responses and establishes IL-6 trans-signaling mediated persistent pro-inflammatory responses. There is direct evidence that during apoptosis the presence of a pharmacological inhibitor of ADAM-17 restricts IL-6R shedding [56].

Hence, we postulate that after detection of viral infection IL-6 initiates pro-inflammatory responses via the sIL-6R-mediated trans-signaling pathway. As a consequence, pro-inflammatory cytokine levels rise. The concentrations of cortisol and IL-10 also increase gradually in order to maintain homeostasis and to activate a TNF- $\alpha$ -mediated apoptotic mechanism as part of a negative feedback loop. A combination of cell death and lymphocytopenia further stimulates ADAM-17 to shed more IL-6R by proteolysis. Subsequently, the level of sIL-6R also increases and bypasses IL-6-mediated classic-signaling. As a result, an uncontrolled trans-signaling-mediated inflammatory response prevails (Fig. 1). In this case, therefore, the infection proceeds to a fatal outcome.

The ability of ADAM-17 to control and influence other molecules of the immune system is now the subject of extensive investigation. It is now well established that ADAM-17 mediated shedding of cell surface molecules are involved in progression of multiple pathological condition [82,93–96]. ADAM-17-induced IL-6 trans-signaling-mediated pathogenesis is now known to be a common feature of inflammatory bowel disease as well as of many respiratory diseases [97–101]. Hence, developing a selective inhibitor of ADAM-17 has been a challenge. Several ADAM-17 inhibitors have been tested as therapeutic candidates [83,100]. For several years pharmaceutical companies like British Biotech, Chemicon, Glaxo SmithKline (GSK), Pfizer and Wyeth have also been conducting research on ADAM-17 inhibitors with the aim of developing therapeutics against rheumatoid arthritis [102–106]. Inhibitors like BB-94 developed by British Biotech, GW-3333 by GSK, TMI-005 by Wyeth, and INCB3619 by Incyte Corp have now been discontinued due to not achieving their targeted promise [102,103,107,108]. Inhibitors failed to meet potential because of low efficacy, toxicity, undesired side effects and mostly due to inhibition of multiple proteases by ADAM-17, while targeted for only a specific shedding event [82]. The knowledge gap between *in vivo* and *in vitro* studies also contributed to these failings. Inhibitors designed *in vitro* do not always show the anticipated effect or desired efficacy *in vivo* since when the whole organism is considered multiple pharmacokinetic factors come into play *in vivo* which are not detectable in *in vitro* studies. In addition, our understanding of ADAM-17 biology is incomplete [100].

New research is therefore aiming to selectively block ADAM-17 with targeted inhibitors of higher specificity in order to increase efficacy and to diminish side effects [93]. This type of selective inhibitor should be designed to affect only a specified target subset through inhibiting a specific shedding event driven by ADAM-17 [93]. From the above discussion it also appears that development of a selective ADAM-17 inhibitor can arrest production of sIL-6R by inhibiting shedding of IL-6R. This inhibition may result in suppression of trans-signaling-mediated pro-inflammatory responses and initiation of classical-signaling-

mediated anti-inflammatory responses, thereby balancing elevated cytokine levels and thus reducing disease pathogenicity. Such a selective inhibitor should prevent shedding of only IL-6R and not block all shedding events induced by ADAM-17 [93]. According to our hypothesis if generation of new sIL-6 can be inhibited it may be possible to stop unregulated IL-6 trans-signaling-mediated pro-inflammatory responses. Studies in murine models also suggest that inhibiting IL-6 trans-signaling can increase survival up to 100% [81]. Though failures of previous inhibitors may discourage drug companies to invest more in developing selective inhibitor against ADAM-17 but targeting ADAM-17 with highly selective inhibitor is still an attractive and promising approach to the researchers. If the same or similar mechanisms were to work in our selected viral infections, ADAM-17 inhibitors will not only show potential in future trials but will also bring great relief to infected patients as well. It might then be possible to develop a single therapeutic molecule that will be effective against multiple viral diseases. Although there is a long way to go to achieve commercial realization, the proposition of such a novel hypothesis provides a new dimension to future antiviral research in which a broad range antiviral will not only cure but also be easily affordable globally.


## 7. Key pathogenic biomolecules: interaction in a computational mirror

Bioinformatics and computational biology tools are able to reveal complex biomolecular interactions, thereby systematically integrating the knowledge of traditional reductionist approaches to biomolecular conundrums [109]. This approach has recently been used for decoding the enigma of disease-associated pathways. In this review, based on our hypothesis described above and by pursuing a computational biology approach, we have targeted several key pro- and anti-inflammatory molecules (IL-6, IL-1 $\beta$ , IL-8, IL-6R, IL-10, NR3C1, TNF, ADAM-17, MCP-1/CCL2, Fas, FasL/FasLG & TRAIL/TNFSF10) with an aim to develop an interaction network among these molecules. We have used two interaction databases, STRING [110] and GeneMANIA [111], to build and visualize their interconnectedness.


In undertaking this analysis, we have considered IL-6 as the most important pro-inflammatory cytokine. IL-1 $\beta$ , IL-8 and MCP-1/CCL2 are commonly found in fatal cases and to be closely associated with IL-6, and therefore are also taken into consideration. IL-6R is additionally included due to its instrumental function in determining the pathogenic fate via differential IL-6 signaling. IL-10 and NR3C1 (glucocorticoid receptor) represent the anti-inflammatory regulators. TNF, Fas, FasL, and TRAIL are selected because of their roles in anti-inflammatory regulation and the process of apoptosis. Finally, ADAM-17 was chosen by virtue of its pivotal role in IL-6 signaling regulation and immunopathogenesis, as described earlier.

For the construction of interactome (PPI network) using STRING database, we have set the minimum required interaction score at medium confidence level (0.400) and the maximum number of interactors to be reported for both shells at “not more than 50”, leaving other parameters to their default values. This results in a huge complex network (“hairball”), which has further been remodeled by Cytoscape [112] (Fig. 2). An insightful observation of this complex network shows three most crucial interactors NF- $\kappa$ B, P53 and Notch1. Among all the molecules in this network, these three exert magisterial regulation of our previously selected molecules, where ADAM-17 acts directly as a whistleblower. Several research groups have previously reported the regulatory role of NF- $\kappa$ B, P53 and Notch1 on our selected target molecules. NF- $\kappa$ B is believed to be a key regulator in IL-6 expression [113–117]. Upon infection, when stimulated by TLR ligands, macrophages activate Notch signaling to upregulate IL-6 production as well as to initiate inflammatory responses. Notch mediated upregulation of IL-6 is controlled by NF- $\kappa$ B as well as P53 [118–120]. Interestingly, Notch signaling is further activated by ADAM-17 [121,122].

We further aimed to reconstruct the previous network by



**Fig. 2.** An interactome for the interacting partners of 12 key pro- and anti-inflammatory molecules (IL-6, IL-1b, IL-8, IL-6R, IL-10, NR3C1, TNF, ADAM-17, MCP-1/CCL2, FAS, FASL/FASLG and TRAIL/TNFSF10) was obtained from the STRING database, which was further remodeled using Cytoscape. Along with above mentioned key pro- and anti-inflammatory molecules, NOTCH1, TP53 and KFKB1 (total 15) have been highlighted in “cadet-blue” nodes while the rest are depicted in “orange” nodes. Among those 15 key molecules, nine (IL-1b, IL-8, IL-10, NR3C1, TNF, MCP-1/CCL2, FAS, FASL/FASLG and TRAIL/TNFSF10) are connected with “red” edges and the rest (IL-6, IL-6R, ADAM-17, NOTCH1, NFKB1 and TP53) are connected with “blue” edges for the ease of the readers understanding. The confidence score was set at medium (0.400) and the maximum number of interactors to report was set at “not more than 50” for both the 1st and the 2nd shells during the generation of this interactome. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 3.** Network construction by adding NF-kB, P53 and Notch1 to the 12 key pro- and anti-inflammatory molecules, considering default STRING (a) and GeneMANIA (b) parameters.

integrating these three molecules with our key target set using the default parameters of STRING and GeneMANIA. The newly constructed network, as depicted in Fig. 3(a) and (b), shows the interaction between our target molecules in a comprehensive way to enable a better understanding. The influence of ADAM-17 can be understood easily from this network, as described in our postulation. In addition, we constructed another network (Fig. 4) in GeneMANIA to show the direct interaction between ADAM-17 and our key regulatory molecules (TNF, NOTCH1, IL6R, TP53, TNFSF10). From this network influence of ADAM-17 over other regulatory molecules can be easily understood. Some recent studies on therapeutic development demonstrate that ADAM-17 inhibition downregulates Notch signaling [123,124]. Hence, a rapid screening of molecular interactions to form a target-specific cluster has resolved the stochastic nature of a large number of

molecules in that complex network. Here, as uncovered by this study ADAM-17 may be considered as a potential drug target to form the basis of a broad range therapeutic approach.

### 8. Corticosteroid based therapeutic approaches and their limitations

Corticosteroids are potent suppressors of inflammatory responses. Therefore they are used in antiviral treatment against a diversity of infectious diseases as well as our target viral infections. As this study is focused on proposing a potent antiviral target for multiple diseases, the current status and limitations of corticosteroids is a major consideration. This will help to establish why corticosteroids are not considered suitable as broad range antiviral therapeutics. By addressing existing




Fig. 4. The direct interaction of ADAM-17 with key regulatory molecules based on the GeneMANIA database.

drawbacks we can also gain ideas as to why selective blocking of ADAM-17 might form the basis of an effective treatment, as proposed above. Although immunosuppressive corticosteroid drugs are commonly used in an attempt to reduce fatality rates but the results of clinical trials are not sufficiently satisfactory to approve as an effective class of therapeutic [125,126]. Corticosteroids are believed to reduce concentrations of pro-inflammatory cytokines and to upregulate anti-inflammatory mediators by inhibiting the transcriptional effect of NF- $\kappa$ B [127]. In recent years this type of inhibition has become less popular as a therapeutic intervention due to a combination of the side-effects of glucocorticoid treatment and global inhibition of NF- $\kappa$ B [128,129]. One of the most frequently used corticosteroid drugs is dexamethasone, which has been stated in some case reports as less effective [130]. Glucocorticoid-mediated stimulation of the hypothalamic-pituitary-adrenal axis can also drive lymphocytopenia [24], or it may promote exaggerated pro-inflammatory responses that eventually cause a worsening of the pathogenic condition [126]. Due to concerns regarding these side-effects, the World Health Organization does not at present recommend glucocorticoids for treatment [131,132]. Furthermore, it is now known that the effectiveness of corticosteroid treatment greatly depends on the timing of drug administration. In addition, the genetic make-up of individuals contributes to therapeutic efficacy. Therefore, it has become extremely challenging to determine an appropriate course of drug administration for each patient [126]. Despite the widespread availability of corticosteroid-based common therapeutic strategies many more molecules have been proposed as antiviral candidates against each of Ebola, SARS-CoV and dengue. Some of these have proved effective in experimental models but not in clinical trials.

### 8.1. Ebola

A recent *in silico* study suggested ibuprofen, a non-steroid compound, as a promising antiviral target against ebola infection [133]. Favipiravir, a pyrazinecarboxamide derivative, is effective against aerosol Ebola virus E718 infection in immunodeficient A129 alpha/beta interferon (IFN- $\alpha/\beta$ ) receptor knockout (IFNAR $^{-/-}$ ) mice [134,135]. A phase II trial of favipiravir against EBOV was also completed in Guinea in November 2016 [136]. However, preliminary findings do not suggest any significant relationship between favipiravir and EBOV viral kinetics and failed to achieve the result predicted at the outset of the trial [137]. Brincidofovir, a lipid-conjugated analogue of cidofovir, is given to EBOV patients as an emergency treatment [138]. The effectiveness of brincidofovir against EBOV is still not discernible as clinical, animal, and *in vitro* data are currently withheld [139].

### 8.2. SARS-CoV

Ribavirin is the most frequently used drug to combat SARS-CoV, and is administered together with corticosteroids [3,140–142]. In combination with immunoglobulins or thymosins, IFN- $\alpha$  is also used to stimulate T cells in SARS-infected individuals [143,144]. Sometimes, IFN- $\alpha$  is used as a combined regimen with ribavirin, although the beneficial effect of these therapies is not fully understood [145,146]. The protease inhibitors lopinavir and ritonavir may lead to improved outcomes for SARS patients when used in combination with ribavirin [147].

### 8.3. Dengue

Therapeutic trials against dengue have also proved disappointing. These have been conducted on two different antiviral candidates, balapiravir and celgosivir, in Vietnam and Singapore, respectively [148]. Both of them failed to provide a beneficial outcome against DENV infection [149,150]. Chloroquine showed promising antiviral potential in *in vitro* experiments but in subsequent clinical trials failed to reduce viraemia of adult dengue patients [151,152]. Lovastatin has been shown to increase survival in animal models by interrupting the DENV assembly pathway [153,154]; however, it has not been proved to be effective in humans [155].

## 9. Concluding remarks and future perspectives

This review provides a plausible hypothesis for understanding the molecular basis of selected major infectious diseases – Ebola, SARS and dengue, and furthermore, draws upon empirical data for the proposed design of putatively appropriate therapeutics. In so doing, we have built a framework that might help to overcome the present knowledge gap surrounding the phenomenon of immune hyperstimulation-mediated fatality. We envisage that in the near future a ground-breaking transformation in research direction may possibly be based on our hypothesis. By utilizing computational modelling to analyse a vast volume of publicly accessible data, a systematic approach to understanding various infectious disease episodes can be achieved. This information may be of considerable value in enabling researchers to develop experimental models to test novel antiviral therapeutic designs. However, it is also acknowledged that due to the unavailability of a sizable amount of interactome data, the computational study in this review highlights several intrinsic limitations.

Since most of the animal reservoir hosts do not mimic the pathogenesis in humans of specific viruses, it is very difficult to develop any suitable *in vivo* model that can be exploited in order to dissect pathogenic events and thereby to achieve the hypothesized outcome [9,156]. Furthermore, while animal reservoirs may become infected with the discussed viruses they do not develop any clinical symptoms [157]. Reservoir hosts have some unique immune mechanism to maintain a disease free state against those viruses. This poses another crucial question, how a disease-free state is maintained? Could a cure be found by investigating reservoir animal hosts? Understanding the mechanism how reservoirs do not show any clinical symptoms like humans against these viruses may help us to find out what unique natural mechanism is contributing [9,158,159]. This type knowledge could lead us to shape the *in vivo* studies as well as improve the knowledge gap between *in vitro* and *in vivo*. But there are lots of challenges in studying the reservoir hosts under their natural condition and mimicking their natural habitat not only need huge logistic support but also involve great costs and thus there is a long way to achieve answers from reservoir hosts [158–161]. Our work is restricted to understanding pathogenic events in humans. It is therefore not intended to explain the disease-free state of reservoir animal hosts.

In spite of such acknowledged limitations, this type of computational modeling analysis could shape future antiviral research by narrowing down the targets through logical explanation and rational

hypothesis proposition. The outcomes of numerous research studies performed around the world are known but have not been analysed collectively. In this regard, it is now possible to explore new possibilities by bringing together a considerable body of existing data in a single analytical space. Under such a condition, our work has considered pathogenic events during multiple diseases in order to establish a sequential framework to find a possible nexus for metabolic regulation. Here, we have explained how pathogenic events progress to a fatal outcome through IL-6-mediated trans-signaling, a pathway that is profoundly influenced by ADAM-17.

In summary, we propose ADAM-17 to be a possible antiviral target. If successful regulation between ADAM-17 and IL-6 trans-signaling junction is achieved, it may be possible to keep in check the otherwise uncontrolled pro-inflammatory response that is a common clinical feature of viral infection. We consider that this type of immunopathological regulation may prove to be successful in preventing deaths from Ebola, SARS and dengue infections and may similarly play a critical role in combating other human infectious diseases.

### Conflict of interest

The authors declare no actual or potential conflicts of interest in relation to this article. This article does not contain any studies with human or animal subjects performed by the authors.

### Acknowledgements

The authors wish to thank Dr. Rogier van Doorn, Director, Oxford University Clinical Research Unit, Hanoi, Vietnam, for his inspiration and guidance from the very beginning of this work. We also thank Dr. Linfa Wang, Professor, Duke-NUS Medical School, Singapore, for his support and inspiration during this project. We are also grateful to Mr. Ahmad Ullah, Scientific Officer, Research and Development Division, Incepta Vaccine Ltd., Dhaka, Bangladesh, for the preparation of artwork.

### References

- [1] L. Baseler, D.S. Chertow, K.M. Johnson, H. Feldmann, D.M. Morens, The pathogenesis of ebola virus disease, *Annu. Rev. Pathol. Mech. Dis.* 12 (2017) 387–418, <https://doi.org/10.1146/annurev-pathol-052016-100506>.
- [2] C.A. Donnelly, A.C. Ghani, G.M. Leung, A.J. Hedley, C. Fraser, S. Riley, L.J. Abu-Raddad, L.M. Ho, T.Q. Thach, P. Chau, K.P. Chan, T.H. Lam, L.Y. Tse, T. Tsang, S.H. Liu, J.H.B. Kong, E.M.C. Lau, N.M. Ferguson, R.M. Anderson, Epidemiological determinants of spread of causal agent of severe acute respiratory syndrome in Hong Kong, *Lancet* 361 (2003) 1761–1766, [https://doi.org/10.1016/S0140-6736\(03\)13410-1](https://doi.org/10.1016/S0140-6736(03)13410-1).
- [3] C.M. Booth, L.M. Matukas, G.A. Tomlinson, A.R. Rachlis, D.B. Rose, H.A. Dwosh, S.L. Walmsley, T. Mazzulli, M. Avendano, P. Derkach, I.E. Ephantimos, I. Kitai, B.D. Mederski, S.B. Shadowitz, W.L. Gold, L.A. Hawryluck, E. Rea, J.S. Chenkin, D.W. Cescon, S.M. Poutanen, A.S. Detsky, Clinical features and short-term outcomes of 144 patients with SARS in the greater Toronto area, *JAMA* 289 (2003) 2801, <https://doi.org/10.1001/jama.289.21.JOC30885>.
- [4] L.-Y. Hsu, C.-C. Lee, J.A. Green, B. Ang, N.I. Paton, L. Lee, J.S. Villacian, P.-L. Lim, A. Earnest, Y.-S. Leo, Severe acute respiratory syndrome (SARS) in Singapore: clinical features of index patient and initial contacts, *Emerg. Infect. Dis.* 9 (2003) 713–717, <https://doi.org/10.3201/eid0906.030264>.
- [5] N. Lee, D. Hui, A. Wu, P. Chan, P. Cameron, G.M. Joynt, A. Ahuja, Y. Yung, C.B. Leung, K.F. To, S.F. Lui, C.C. Szeto, S. Chung, J.J.Y. Sung, A major outbreak of severe acute respiratory syndrome in Hong Kong, *N Engl J Med Emerg Med (P. Cameron)* *N Engl J Med* 348 (2003) 1986–1994 (accessed August 26, 2017).
- [6] J.S.M. Peiris, C.M. Chu, V.C.C. Cheng, K.S. Chan, I.F.N. Hung, L.L.M. Poon, K.I. Law, B.S.F. Tang, T.Y.W. Hon, C.S. Chan, K.H. Chan, J.S.C. Ng, B.J. Zheng, W.L. Ng, R.W.M. Lai, Y. Guan, K.Y. Yuen, HKU/UCH SARS Study Group, Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study, *Lancet (London, England)* 361 (2003) 1767–1772 (accessed August 28, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/12781535>.
- [7] R. Islam, M. Salahuddin, M.S.M.S. Ayubi, T. Hossain, A. Majumder, A.W.A.W. Taylor-Robinson, A. Mahmud-Al-Rafat, Dengue epidemiology and pathogenesis: images of the future viewed through a mirror of the past, *Virol. Sin.* 30 (2015) 326–343, <https://doi.org/10.1007/s12250-015-3624-1>.
- [8] D. Guha-Sapir, B. Schimmer, Dengue fever: new paradigms for a changing epidemiology, *Emerg. Themes Epidemiol.* 1 (2005), <https://doi.org/10.1186/1742-7622-2-1>.
- [9] S. Zompi, E. Harris, Animal models of dengue virus infection, *Viruses* 4 (2012) 62–82, <https://doi.org/10.3390/v4010062>.
- [10] A.T. Bäck, A. Lundkvist, Dengue viruses – an overview, *Infect. Ecol. Epidemiol.* 3 (2013) 19839, <https://doi.org/10.3402/iee.v3i0.19839>.
- [11] J. Barnioli, R. Gaczkowski, E.V. Barbato, R.V. da Cunha, D. Salgado, E. Martínez, C.S. Segarra, E.B. Pleites Sandoval, A. Mishra, I.S. Laksono, L.C. Lum, J.G. Martínez, A. Núñez, A. Balsameda, I. Allende, G. Ramírez, E. Dimaaño, K. Thomacheck, N.A. Akbar, E.E. Ooi, E. Villegas, T.T. Hien, J. Farrar, O. Horstick, A. Kroeger, T. Jaenisch, Usefulness and applicability of the revised dengue case classification by disease: multi-centre study in 18 countries, *BMC Infect. Dis.* 11 (2011) 106, <https://doi.org/10.1186/1471-2334-11-106>.
- [12] B.G. Chousterman, F.K. Swirski, G.F. Weber, Cytokine storm and sepsis disease pathogenesis, *Semin. Immunopathol.* 39 (2017) 517–528, <https://doi.org/10.1007/s00281-017-0639-8>.
- [13] L. Falasca, C. Agrati, N. Petrosillo, A. Di Caro, M. Capobianchi, G. Ippolito, M. Piacentini, Molecular mechanisms of Ebola virus pathogenesis: focus on cell death, *Cell Death Differ.* 22 (2015) 1250–1259, <https://doi.org/10.1038/cdd.2015.67>.
- [14] A.K. McElroy, R.S. Akondy, C.W. Davis, A.H. Ellebedy, A.K. Mehta, C.S. Kraft, G.M. Lyon, B.S. Ribner, J. Varkey, J. Sidney, A. Sette, S. Campbell, U. Ströher, I. Damon, S.T. Nichol, C.F. Spiropoulou, R. Ahmed, Human Ebola virus infection results in substantial immune activation, *Proc. Natl. Acad. Sci.* 112 (2015) 4719–4724, <https://doi.org/10.1073/pnas.1502619112>.
- [15] P. Younan, M. Iampietro, A. Bukreyev, Disabling of lymphocyte immune response by Ebola virus, *PLOS Pathog.* 14 (2018) e1006932, <https://doi.org/10.1371/journal.ppat.1006932>.
- [16] N.M. Lubaki, P. Ilinykh, C. Pietzsch, B. Tigabu, A.N. Freiberg, R.A. Koup, A. Bukreyev, The lack of maturation of ebola virus-infected dendritic cells results from the cooperative effect of at least two viral domains, *J. Virol.* 87 (2013) 7471–7485, <https://doi.org/10.1128/JVI.03316-12>.
- [17] S. Bixler, A. Goff, The role of cytokines and chemokines in filovirus infection, *Viruses* 7 (2015) 5489–5507, <https://doi.org/10.3390/v7102892>.
- [18] J. Bradley, TNF-mediated inflammatory disease, *J. Pathol.* 214 (2008) 149–160, <https://doi.org/10.1002/path.2287>.
- [19] I.S. Caballero, A.N. Honko, S.K. Gire, S.M. Winnicki, M. Melé, C. Gerhardinger, A.E. Lin, J.L. Rinn, P.C. Sabeti, L.E. Hensley, J.H. Connor, In vivo Ebola virus infection leads to a strong innate response in circulating immune cells, *BMC Genomics* 17 (2016) 707, <https://doi.org/10.1186/s12864-016-3060-0>.
- [20] P. Ruibal, L. Oestereich, A. Lüdtke, B. Becker-Ziaja, D.M. Wozniak, R. Kerber, M. Korva, M. Cabeza-Cabrerizo, J.A. Bore, F.R. Koundouno, S. Duraffour, R. Weller, A. Thorenz, E. Cimini, D. Viola, C. Agrati, J. Repits, B. Afrough, L.A. Cowley, D. Ngabo, J. Hinzmann, M. Mertens, I. Vitoriano, C.H. Logue, J.P. Boettcher, E. Pallasch, A. Sachse, A. Bah, K. Nitzsche, E. Kuisma, J. Michel, T. Holm, E.-G. Zekeng, I. García-Dorival, R. Wölfel, K. Stoecker, E. Fleischmann, T. Strecker, A. Di Caro, T. Avšič-Županc, A. Kurth, S. Meschi, S. Mély, E. Newman, A. Brockhin, Z. Kis, A. Kelterbaum, P. Molkenhain, F. Carletti, J. Portmann, S. Wolff, C. Castilletti, G. Schudt, A. Fizez, L.J. Ottowell, E. Herker, T. Jacobs, B. Kretschmer, E. Severi, N. Ouedraogo, M. Lago, A. Negro, L. Franco, P. Andra, S. Schmiedel, B. Kreuels, D. Wichmann, M.M. Addo, A.W. Lohse, H. De Clerck, C. Nanclares, S. Jonckheere, M. Van Herp, A. Sprecher, G. Xiaojiang, M. Carrington, O. Miranda, C.M. Castro, M. Gabriel, P. Drury, P. Formenty, B. Diallo, L. Koivogui, N. Magassouba, M.W. Carroll, S. Günther, C. Muñoz-Fontela, Unique human immune signature of Ebola virus disease in Guinea, *Nature* 533 (2016) 100–104, <https://doi.org/10.1038/nature17949>.
- [21] C. Dahlke, S. Lunemann, R. Kasonta, B. Kreuels, S. Schmiedel, M.L. Ly, S.K. Fehling, T. Strecker, S. Becker, M. Altfeld, A. Sow, A.W. Lohse, C. Muñoz-Fontela, M.M. Addo, Comprehensive characterization of cellular immune responses following Ebola virus infection, *J. Infect. Dis.* 215 (2017) 287–292, <https://doi.org/10.1093/infdis/jiw508>.
- [22] H.H. Yiu, A.L. Graham, R.F. Stengel, Dynamics of a Cytokine Storm, *PLoS One* 7 (2012) e45027, <https://doi.org/10.1371/journal.pone.0045027>.
- [23] K.J. Huang, L.J. Su, M. Theron, Y.C. Wu, S.K. Lai, C.C. Liu, H.Y. Lei, An interferon-γ-related cytokine storm in SARS patients, *J. Med. Virol.* 75 (2005) 185–194, <https://doi.org/10.1002/jmv.20255>.
- [24] N.S. Panesar, A. Fauci, G. Tomlinson, et al., Lymphopenia in SARS, *Lancet (London, England)* 10.1016/S0140-6736(03)13557-X 361 (2003) 1985.
- [25] J. Chen, Y.F. Lau, E.W. Lamirande, C.D. Paddock, J.H. Bartlett, S.R. Zaki, K. Subbarao, Cellular immune responses to severe acute respiratory syndrome coronavirus (SARS-CoV) infection in senescent BALB/c mice: CD4+ T cells are important in control of SARS-CoV infection, *J. Virol.* 84 (2010) 1289–1301, <https://doi.org/10.1128/JVI.01281-09>.
- [26] L.E. Gralinski, A. Bankhead, S. Jeng, V.D. Menachery, S. Proll, S.E. Belisle, M. Matzke, B.-J.M. Webb-Robertson, M.L. Luna, A.K. Shukla, M.T. Ferris, M. Bolles, J. Chang, L. Aicher, K.M. Waters, R.D. Smith, T.O. Metz, G.L. Law, M.G. Katze, S. McWeeney, R.S. Baric, Mechanisms of severe acute respiratory syndrome coronavirus-induced acute lung injury, *MBio* 4 (2013), <https://doi.org/10.1128/mBio.00271-13>.
- [27] C.L. Medin, K.A. Fitzgerald, A.L. Rothman, Dengue virus nonstructural protein NS5 induces interleukin-8 transcription and secretion, *J. Virol.* 79 (2005) 11053–11061, <https://doi.org/10.1128/JVI.79.17.11053-11061.2005>.
- [28] J.F. Kelley, P.H. Kafusi, V.R. Nerurkar, Dengue hemorrhagic fever-associated immunomodulators induced via maturation of dengue virus nonstructural 4B protein in monocytes modulate endothelial cell adhesion molecules and human microvascular endothelial cells permeability, *Virology* 422 (2012) 326–337, <https://doi.org/10.1016/j.virol.2011.10.030>.
- [29] V.V. Costa, C.T. Fagundes, D.G. Souza, M.M. Teixeira, Inflammatory and innate immune responses in dengue infection, *Am. J. Pathol.* 182 (2013) 1950–1961, <https://doi.org/10.1016/j.ajpath.2013.02.027>.
- [30] A. Srikiatkachorn, A. Mathew, A.L. Rothman, Immune-mediated cytokine storm and its role in severe dengue, *Semin. Immunopathol.* 39 (2017) 563–574, <https://doi.org/10.1007/s00281-017-0625-1>.

- [31] G.N. Malavige, G.S. Ogg, Pathogenesis of vascular leak in dengue virus infection, *Immunology* 151 (2017) 261–269, <https://doi.org/10.1111/imm.12748>.
- [32] N. Luplertlop, N. Luplertlop, D. Missé, D. Bray, V. Deleuze, J.-P. Gonzalez, V. Leardkamolkarn, H. Yssel, F. Veas, Dengue-virus-infected dendritic cells trigger vascular leakage through metalloproteinase overproduction, *EMBO Rep.* 7 (2006) 1176–1181, <https://doi.org/10.1038/sj.embor.7400814>.
- [33] D.H. Libraty, P.R. Young, D. Pickering, T.P. Endy, S. Kalayanaraj, S. Green, D.W. Vaughn, A. Nisalak, F.A. Ennis, A.L. Rothman, High circulating levels of the dengue virus nonstructural protein NS1 early in dengue illness correlate with the development of dengue hemorrhagic fever, *J. Infect. Dis.* 186 (2002) 1165–1168, <https://doi.org/10.1086/343813>.
- [34] E.L. Azeredo, L.M. De Oliveira-Pinto, S.M. Zagne, D.I.S. Cerqueira, R.M.R. Nogueira, C.F. Kubelka, NK cells, displaying early activation, cytotoxicity and adhesion molecules, are associated with mild dengue disease, *Clin. Exp. Immunol.* 143 (2006) 345–356, <https://doi.org/10.1111/j.1365-2249.2006.02996.x>.
- [35] M.A. Lopez-Ramirez, R. Fischer, C.C. Torres-Badillo, H.A. Davies, K. Logan, K. Pfizenmaier, D.K. Male, B. Sharrack, I.A. Romero, Role of caspases in cytokine-induced barrier breakdown in human brain endothelial cells, *J. Immunol.* 189 (2012) 3130–3139, <https://doi.org/10.4049/jimmunol.1103460>.
- [36] J. Huang, W. Liang, S. Chen, Y. Zhu, H. Chen, C.K.P. Mok, Y. Zhou, Serum cytokine profiles in patients with dengue fever at the acute infection phase, *Dis. Markers* 2018 (2018) 1–8, <https://doi.org/10.1155/2018/8403937>.
- [37] B.E.E. Martina, P. Koraka, A.D.M.E. Osterhaus, Dengue virus pathogenesis: an integrated view, *Clin. Microbiol. Rev.* 22 (2009) 564–581, <https://doi.org/10.1128/CMR.00035-09>.
- [38] Seema, S.K. Jain, Molecular mechanism of pathogenesis of dengue virus: entry and fusion with target cell, *Indian J. Clin. Biochem.* 20 (2005) 92–103, <https://doi.org/10.1007/BF02867407>.
- [39] T.F. Póvoa, E.R.A. Oliveira, C.A. Basílio-de-Oliveira, G.J. Nuovo, V.L.A. Chagas, N.G. Salomão, E.M. Mota, M.V. Paes, Peripheral organs of dengue fatal cases present strong pro-inflammatory response with participation of IFN- $\gamma$ , TNF- $\alpha$  and RANTES-producing cells, *PLoS One* 11 (2016) e0168973, <https://doi.org/10.1371/journal.pone.0168973>.
- [40] T.F. Póvoa, A.M.B. Alves, C.A.B. Oliveira, G.J. Nuovo, V.L.A. Chagas, M.V. Paes, The pathology of severe dengue in multiple organs of human fatal cases: histopathology, ultrastructure and virus replication, *PLoS One* 9 (2014) e83386, <https://doi.org/10.1371/JOURNAL.PONE.0083386>.
- [41] J.R. Tisoncik, M.J. Korth, C.P. Simmons, J. Farrar, T.R. Martin, M.G. Katze, Into the eye of the cytokine storm, *Microbiol. Mol. Biol. Rev.* 76 (2012) 16–32, <https://doi.org/10.1128/MMBR.05015-11>.
- [42] E. Kindler, V. Thiel, SARS-CoV and IFN: too little, too late, *Cell Host Microbe* 19 (2016) 139–141, <https://doi.org/10.1016/j.chom.2016.01.012>.
- [43] M.A. Schmid, E. Harris, Monocyte recruitment to the dermis and differentiation to dendritic cells increases the targets for dengue virus replication, *PLoS Pathog.* 10 (2014) e1004541, <https://doi.org/10.1371/journal.ppat.1004541>.
- [44] J. Gu, E. Gong, B. Zhang, J. Zhong, Z. Gao, Y. Zhong, W. Zou, J. Zhan, S. Wang, Z. Xie, H. Zhuang, B. Wu, H. Zhong, H. Shao, W. Fang, D. Gao, F. Pei, X. Li, Z. He, D. Xu, X. Shi, V.M. Anderson, A.S.-Y. Leong, Multiple organ infection and the pathogenesis of SARS, *J. Exp. Med.* 202 (2005) 415–424, <https://doi.org/10.1084/jem.20050828>.
- [45] M. Frieman, M. Heise, R. Baric, SARS coronavirus and innate immunity, *Virus Res.* 133 (2008) 101–112, <https://doi.org/10.1016/j.virusres.2007.03.015>.
- [46] C.F. Basler, G.K. Amarasinghe, Evasion of interferon responses by ebola and marburg viruses, *J. Interf. Cytokine Res.* 29 (2009) 511–520, <https://doi.org/10.1089/jir.2009.0076>.
- [47] A.A. Ansari, Clinical features and pathobiology of Ebolavirus infection, *J. Autoimmun.* 55 (2014) 1–9, <https://doi.org/10.1016/j.jaut.2014.09.001>.
- [48] J. Misasi, N.J. Sullivan, Camouflage and misdirection: the full-on assault of ebola virus disease, *Cell* 159 (2014) 477–486, <https://doi.org/10.1016/j.cell.2014.10.006>.
- [49] C.W.K. Lam, M.H.M. Chan, C.K. Wong, Severe acute respiratory syndrome: clinical and laboratory manifestations, *Clin. Biochem. Rev.* 25 (2004) 121–132 (accessed August 24, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/18458712>.
- [50] H. Feldmann, T.W. Geisbert, Ebola haemorrhagic fever, *Lancet* 377 (2011) 849–862, [https://doi.org/10.1016/S0140-6736\(10\)60667-8](https://doi.org/10.1016/S0140-6736(10)60667-8).
- [51] J.C. Kash, E. Mühlberger, V. Carter, M. Grosch, O. Perwitasari, S.C. Proll, M.J. Thomas, F. Weber, H.-D. Klenk, M.G. Katze, Global suppression of the host antiviral response by Ebola- and Marburgviruses: increased antagonism of the type I interferon response is associated with enhanced virulence, *J. Virol.* 80 (2006) 3009–3020, <https://doi.org/10.1128/JVI.80.6.3009-3020.2006>.
- [52] N. Wauquier, P. Becquart, C. Padilla, S. Baize, E.M. Leroy, Human fatal zaire ebola virus infection is associated with an aberrant innate immunity and with massive lymphocyte apoptosis, *PLoS Negl. Trop. Dis.* 4 (2010) e837, <https://doi.org/10.1371/journal.pntd.0000837>.
- [53] S. De, T. Martins, G.F. Silveira, L.R. Alves, C. Nunes, D. Dos Santos, J. Bordignon, Dendritic cell apoptosis and the pathogenesis of dengue, *Viruses* 4 (2012) 2736–2753, <https://doi.org/10.3390/v4112736>.
- [54] R. O'Donnell, R.C. Tasker, M.F.E. Roe, SARS: understanding the coronavirus: apoptosis may explain lymphopenia of SARS, *BMJ* 327 (2003) 620, <https://doi.org/10.1136/bmj.327.7415.620-b>.
- [55] C.A. Hunter, S.A. Jones, IL-6 as a keystone cytokine in health and disease, *Nat. Immunol.* 16 (2015) 448–457, <https://doi.org/10.1038/ni.3153>.
- [56] J. Scheller, A. Chalaris, D. Schmidt-Arras, S. Rose-John, The pro- and anti-inflammatory properties of the cytokine interleukin-6, *BBA – Mol. Cell Res.* 2011 (2011) 878–888, <https://doi.org/10.1016/j.bbamcr.2011.01.034>.
- [57] S.P. Commins, L. Borish, J.W. Steinke, Immunologic messenger molecules: Cytokines, interferons, and chemokines, *J. Allergy Clin. Immunol.* 125 (2010) S53–S72, <https://doi.org/10.1016/j.jaci.2009.07.008>.
- [58] G.N. Malavige, L. Gomes, L. Alles, T. Chang, M. Salimi, S. Fernando, K.D. Nanayakkara, S. Jayaratne, G.S. Ogg, Serum IL-10 as a marker of severe dengue infection, *BMC Infect. Dis.* 13 (2013) 341, <https://doi.org/10.1186/1471-2334-13-341>.
- [59] C.K. Li, H. Wu, H. Yan, S. Ma, L. Wang, M. Zhang, X. Tang, N.J. Temperton, R.A. Weiss, J.M. Brechley, D.C. Douek, J. Mongkolsapaya, B.-H. Tran, C.S. Lin, G.R. Screaton, J. Hou, A.J. McMichael, X.-N. Xu, T cell responses to whole SARS coronavirus in humans, *J. Immunol.* 181 (2008) 5490–5500 (accessed April 6, 2018), <http://www.ncbi.nlm.nih.gov/pubmed/18832706>.
- [60] K.L. Hutchinson, P.E. Rollin, Cytokine and chemokine expression in humans infected with Sudan Ebola virus, *J. Infect. Dis.* 196 (2007) S357–S363, <https://doi.org/10.1086/520611>.
- [61] W. Wu, K.K. Dietze, K. Gibbert, K.S. Lang, M. Trilling, H. Yan, J. Wu, D. Yang, M. Lu, M. Roggendorf, U. Dittmer, J. Liu, TLR ligand induced IL-6 counter-regulates the anti-viral CD8 T cell response during an acute retrovirus infection, *Nat. Publ. Gr.* (2015), <https://doi.org/10.1038/srep10501>.
- [62] P. Stenvinkel, M. Ketteler, R.J. Johnson, B. Lindholm, R. Pecoits-Filho, M. Riella, H. Unger, T. Cederholm, M. Girndt, IL-10, IL-6, and TNF- $\alpha$ : Central factors in the altered cytokine network of uremia—The good, the bad, and the ugly, *Kidney Int.* 67 (2005) 1216–1233, <https://doi.org/10.1111/j.1523-1755.2005.00200.x>.
- [63] S. Dimitrov, T. Lange, C. Benedict, M.A. Nowell, S.A. Jones, J. Scheller, S. Rose-John, J. Born, Sleep enhances IL-6 trans-signaling in humans, *FASEB J.* 20 (2006) 2174–2176, <https://doi.org/10.1096/fj.06-5754fe>.
- [64] S.A. Jones, S. Horiuchi, N. Topley, N. Yamamoto, G.M. Fuller, The soluble interleukin 6 receptor: mechanisms of production and implications in disease, *FASEB J.* 15 (2001) 43–58, <https://doi.org/10.1096/fj.99-1003rev>.
- [65] R.J. Simpson, A. Hammacher, D.K. Smith, J.M. Matthews, L.D. Ward, Interleukin-6: structure-function relationships, *Protein Sci.* 6 (1997) 929–955, <https://doi.org/10.1002/pro.5560060501>.
- [66] P.C. Heinrich, J. Behrmann, S. Haan, H.M. Hermans, G. Uller-Newen, F. Schaper, Principles of interleukin (IL)-6-type cytokine signalling and its regulation 1, *Biochem. J.* 374 (2003) 1–20 (accessed August 28, 2017), <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1223585/pdf/12773095.pdf>.
- [67] M. Lacroix, F. Rousseau, F. Guilhot, P. Malinge, G. Magistrelli, S. Herren, S.A. Jones, G.W. Jones, J. Scheller, R. Lissilaa, M. Kosco-Vilbois, Z. Johnson, V. Buatois, W. Ferlin, Novel Insights into Interleukin 6 (IL-6) Cis- and Trans- signaling Pathways by Differentially Manipulating the Assembly of the IL-6 Signaling Complex \*, 2015, <https://doi.org/10.1074/jbc.M115.682138>.
- [68] P. Chomarat, J. Banchereau, J. Davoust, A. Karolina, Palucka, IL-6 switches the differentiation of monocytes from dendritic cells to macrophages, *Nat. Immunol.* 1 (2000) 510–514, <https://doi.org/10.1038/10338763>.
- [69] A. Steensberg, C.P. Fischer, C. Keller, K. Møller, B.K. Pedersen, IL-6 enhances plasma IL-1ra, IL-10, and cortisol in humans, *Am. J. Physiol. – Endocrinol. Metab.* 285 (2003) E433–E437, <https://doi.org/10.1152/ajpendo.00074.2003>.
- [70] H. Urbaniśka-Ryś, A. Wiersbowska, H. Stepien, T. Robak, Relationship between circulating interleukin-10 (IL-10) with interleukin-6 (IL-6) type cytokines (IL-6, interleukin-11 (IL-11), oncostatin M (OSM)) and soluble interleukin-6 (IL-6) receptor (sIL-6R) in patients with multiple myeloma, *Eur. Cytokine Netw.* 11 (2000) 443–451 (accessed August 24, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/11022130>.
- [71] S. Dominitzki, M.C. Fantini, C. Neufert, A. Nikolaev, P.R. Galle, J. Scheller, G. Monteleone, S. Rose-John, M.F. Neurath, C. Becker, Cutting edge: trans-signaling via the soluble IL-6R abrogates the induction of FoxP3 in naive CD4+ CD25– T cells, *J. Immunol.* 179 (2007) (accessed August 28, 2017), <http://www.jimmunol.org/content/179/4/2041>.
- [72] M.H. Sofi, W. Li, M.H. Kaplan, C.-H. Chang, Elevated IL-6 expression in CD4 T cells via PKC $\theta$  and NF- $\kappa$ B induces Th2 cytokine production, *Mol. Immunol.* 46 (2009) 1443–1450, <https://doi.org/10.1016/j.molimm.2008.12.014>.
- [73] S. Rose-John, IL-6 trans-signaling via the soluble IL-6 receptor: importance for the pro-inflammatory activities of IL-6, *Int. J. Biol. Sci.* 8 (2012) 1237–1247, <https://doi.org/10.7150/ijbs.4989>.
- [74] G. Scambia, U. Testa, P.B. Panici, R. Martucci, E. Foti, M. Petrini, M. Amoroso, V. Masciullo, C. Peschle, S. Mancuso, Interleukin-6 serum levels in patients with gynecological tumors, *Int. J. Cancer.* 57 (1994) 318–323, <https://doi.org/10.1002/ijc.2910570305>.
- [75] M.A. Nowell, P.J. Richards, S. Horiuchi, N. Yamamoto, S. Rose-John, N. Topley, A.S. Williams, S.A. Jones, Soluble IL-6 receptor governs IL-6 activity in experimental arthritis: blockade of arthritis severity by soluble glycoprotein 130, *J. Immunol.* 171 (2003) 3202–3209 (accessed December 9, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/12960349>.
- [76] A. Waage, P. Brandtzaeg, A. Halstensen, P. Kierulf, T. Espevik, The complex pattern of cytokines in serum from patients with meningococcal septic shock. Association between interleukin 6, interleukin 1, and fatal outcome, *J. Exp. Med.* 169 (1989) 333–338 (accessed December 9, 2017).
- [77] Y. Saito, T. Osugi, T. Tokunaga, M. Kishimoto, S. Honda, M. Yamamoto, K. Cheng, H. Yasukawa, T. Suzuki, M. Honda, S. Yamamoto, M. Cheng, K. Yasukawa, H. Suzuki, T. Saito, Y. Osugi, T. Tokunaga, T. Kishimoto, Human soluble IL-6 receptor: its detection human soluble il-6 receptor: its detection and enhanced release by HIV infection, *J. Immunol.* 148 (1992) 2175–2180 (accessed August 26, 2017).
- [78] F. Padberg, W. Feneberg, S. Schmidt, M.J. Schwarz, D. Körschenhausen, B.D. Greenberg, T. Nolde, N. Müller, H. Trapmann, N. König, H.J. Möller, H. Hampel, N. Müller, T. Sunderland, S. Rapoport, H.-J. Möller, CSF and serum levels of soluble interleukin-6 receptors (sIL-6R and sgp130), but not of interleukin-6 are altered in multiple sclerosis, *J. Neuroimmunol.* 99 (1999) 218–223, [https://doi.org/10.1016/S0165-5728\(99\)00120-4](https://doi.org/10.1016/S0165-5728(99)00120-4).
- [79] A. Doganci, T. Eigenbrod, N. Krug, G.T. De Sanctis, M. Hausding, V.J. Erpenbeck, E.-B. Haddad, E. Schmitt, T. Bopp, K.-J. Kallen, U. Herz, S. Schmitt, C. Luft, O. Hecht, J.M. Hohlfield, H. Ito, N. Nishimoto, K. Yoshizaki, T. Kishimoto, S. Rose-John, H. Renz, M.F. Neurath, P.R. Galle, S. Finotto, The IL-6 $\alpha$  chain controls lung CD4 + CD25 + Treg development and function during allergic airway

- inflammation in vivo, *J. Clin. Invest.* 115 (2005), <https://doi.org/10.1172/JCI200522433>.
- [80] H. Schuett, R. Oestreich, G.H. Waetzig, W. Annema, M. Luchtfeld, A. Hillmer, U. Bavendiek, J. von Felden, D. Divchev, T. Kempf, K.C. Wollert, D. Seeger, S. Rose-John, U.J.F. Tietge, B. Schieffer, K. Grote, Transsignaling of interleukin-6 crucially contributes to atherosclerosis in mice, *Arterioscler. Thromb. Vasc. Biol.* 32 (2012) (accessed August 28, 2017), <http://atvb.ahajournals.org/content/32/2/281.long>.
- [81] T. Barkhausen, T. Tschernig, P. Rosenstiel, M. van Griensven, R.-P. Vonberg, M. Dorsch, A. Mueller-Heine, A. Chalaris, J. Scheller, S. Rose-John, D. Seeger, C. Krettek, G.H. Waetzig, Selective blockade of interleukin-6 trans-signaling improves survival in a murine polymicrobial sepsis model, *Crit. Care Med.* 39 (2011) 1407–1413, <https://doi.org/10.1097/CCM.0b013e318211ff56>.
- [82] D.C. Blaydon, P. Biancheri, W.-L. Di, V. Plagnol, R.M. Cabral, M.A. Brooke, D.A. van Heel, F. Ruschendorf, M. Toynbee, A. Walne, E.A. O'Toole, J.E. Martin, K. Lindley, T. Vulliamy, D.J. Abrams, T.T. MacDonald, J.I. Harper, D.P. Kelsell, Inflammatory skin and bowel disease linked to ADAM17 deletion, *N. Engl. J. Med.* 365 (2011) 1502–1508, <https://doi.org/10.1056/NEJMoa1100721>.
- [83] M. Goetz, ADAM-17: the enzyme that does it all, *Crit. Rev. Biochem. Mol. Biol.* 45 (2010) 146–169, <https://doi.org/10.3109/10409231003628015>.
- [84] M. Bzowska, N. Jura, A. Lassak, R.A. Black, J. Bereta, Tumour necrosis factor- $\alpha$  stimulates expression of TNF- $\alpha$  converting enzyme in endothelial cells, *Eur. J. Biochem.* 271 (2004) 2808–2820, <https://doi.org/10.1111/j.1432-1033.2004.04215.x>.
- [85] R.S. McMahan, K.J. Riehle, N. Fausto, J.S. Campbell, A disintegrin and metalloproteinase 17 regulates TNF and TNFR1 levels in inflammation and liver regeneration in mice, *Am. J. Physiol. Gastrointest. Liver Physiol.* 305 (2013) G25–G34, <https://doi.org/10.1152/ajpgi.00326.2012>.
- [86] J. Rgen Scheller, A. Chalaris, C. Garbers, S. Rose-John, ADAM17: a molecular switch to control inflammation and tissue regeneration, *Trends Immunol.* 32 (2011) 380–387, <https://doi.org/10.1016/j.it.2011.05.005>.
- [87] A. Chalaris, B. Rn Rabe, K. Paliga, H. Lange, T. Laskay, C.A. Fielding, S.A. Jones, S. Rose-John, J. Rgen Scheller, Apoptosis is a natural stimulus of IL6R shedding and contributes to the proinflammatory trans-signaling function of neutrophils, *Blood* 110 (2007) 1748–1755, <https://doi.org/10.1182/blood-2007-01-067918>.
- [88] A. Torrentes-Carvalho, E.L. Azeredo, S.R. Reis, A.S. Miranda, M. Gandini, L.S. Barbosa, C.F. Kubelka, Dengue-2 infection and the induction of apoptosis in human primary monocytes, *Mem Inst Oswaldo Cruz Rio Janeiro* 104 (2009) 1091–1099 (accessed August 24, 2017), <http://www.scielo.br/pdf/mioc/v104n8/05.pdf>.
- [89] M. Bray, T.W. Geisbert, Ebola virus: the role of macrophages and dendritic cells in the pathogenesis of Ebola hemorrhagic fever, *Int. J. Biochem. Cell Biol.* 37 (2005) 1560–1566, <https://doi.org/10.1016/j.biocel.2005.02.018>.
- [90] S.B. Bradfute, P.E. Swanson, M.A. Smith, E. Watanabe, J.E. McDunn, R.S. Hothkiss, S. Bavari, Mechanisms and consequences of ebolavirus-induced lymphocyte apoptosis, *J. Immunol.* 184 (2009) (accessed August 28, 2017), <http://www.jimmunol.org/content/184/1/327.long>.
- [91] B. Walcheck, A.H. Herrera, C. St, P.E. Hill, A.R. Mattila, F.R. DeLeo Whitney, ADAM17 activity during human neutrophil activation and apoptosis, *Eur. J. Immunol.* 36 (2006) 968–976, <https://doi.org/10.1002/eji.200535257>.
- [92] Y. Wang, J.D. Robertson, B. Walcheck, J.D. Robertson, B. Walcheck, Different signaling pathways stimulate a disintegrin and metalloproteinase-17 (ADAM17) in neutrophils during apoptosis and activation, *J. Biol. Chem.* 286 (2011) 38980–38988, <https://doi.org/10.1074/jbc.M111.277087>.
- [93] M.L. Moss, D. Minond, Recent Advances in ADAM17 Research: A Promising Target for Cancer and Inflammation, *Mediators Inflamm.* 2017 (2017) 1–21, <https://doi.org/10.1155/2017/9673537>.
- [94] A. Nishimi, T. Isozaki, S. Nishimi, S. Ishii, T. Tokunaga, H. Furuya, K. Wakabayashi, T. Kasama, SAT0322 ADAM-17 is expressed in the inflammatory myopathy, and is involved with interstitial lung disease (ILD), in: Poster Present., BMJ Publishing Group Ltd and European League Against Rheumatism, 2017, p. 894.1–894, <http://doi.org/10.1136/annrheumdis-2017-eular.3402>.
- [95] A. Bertram, S. Lovric, A. Engel, M. Beese, K. Wyss, B. Hertel, J.-K. Park, J.U. Becker, J. Kegel, H. Haller, M. Haubitz, T. Kirsch, Circulating ADAM17 level reflects disease activity in proteinase-3 ANCA-associated vasculitis, *J. Am. Soc. Nephrol.* 26 (2015) 2860–2870, <https://doi.org/10.1681/ASN.2014050477>.
- [96] R. Menghini, L. Fiorentino, V. Casagrande, R. Lauro, M. Federici, The role of ADAM17 in metabolic inflammation, *Atherosclerosis* 228 (2013) 12–17, <https://doi.org/10.1016/j.atherosclerosis.2013.01.024>.
- [97] C. Greene, G. Lowe, C. Taggart, P. Gallagher, N. McElvaney, S. O'Neill, Tumor necrosis factor- $\alpha$ -converting enzyme: its role in community-acquired pneumonia, *J. Infect. Dis.* 186 (2002) 1790–1796, <https://doi.org/10.2307/30077980>.
- [98] V. Marin, F. Montero-Julian, S. Grès, P. Bongrand, C. Farnarier, G. Kaplanski, Chemotactic agents induce IL-6R $\alpha$  shedding from polymorphonuclear cells: involvement of a metalloproteinase of the TNF- $\alpha$ -converting enzyme (TACE) type, *Eur. J. Immunol.* 32 (2002) 2965–2970, [https://doi.org/10.1002/1521-4141\(200210\)32:10<2965::AID-IMMU2965>3.0.CO;2-V](https://doi.org/10.1002/1521-4141(200210)32:10<2965::AID-IMMU2965>3.0.CO;2-V).
- [99] M.A. Berry, B. Hargadon, M. Shelley, D. Parker, D.E. Shaw, R.H. Green, P. Bradding, C.E. Brightling, A.J. Wardlaw, I.D. Pavord, Evidence of a role of tumor necrosis factor  $\alpha$  in refractory asthma, *N Engl J Med.* 354 (2006) 697–708 (accessed August 24, 2017), <http://www.nejm.org/doi/pdf/10.1056/NEJMoa050580>.
- [100] J. Arribas, C. Esseles, ADAM17 as a therapeutic target in multiple diseases, *Curr. Pharm. Des.* 15 (2009) 2319–2335, <https://doi.org/10.2174/138161209788682398>.
- [101] G. Kolios, V. Valatas, S.G. Ward, Nitric oxide in inflammatory bowel disease: a universal messenger in an unsolved puzzle, *Immunology* 113 (2004) 427–437, <https://doi.org/10.1111/j.1365-2567.2004.01984.x>.
- [102] P.D. Brown, Matrix metalloproteinase inhibitors: a novel class of anticancer agents, *Adv. Enzyme Regul.* 35 (1995) 293–301 (accessed December 10, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/7572350>.
- [103] J.I. Levin, Y. Gu, F.C. Nelson, A. Zask, J.F. DiJoseph, M.A. Sharr, A. Sung, G. Jin, R. Cowling, P. Chanda, S. Cosmi, C.L. Hsiao, W. Edris, J. Wilhelm, L.M. Killar, J.S. Skotnicki, Heteroaryl and cycloalkyl sulfonamide hydroxamic acid inhibitors of matrix metalloproteinases, *Bioorg. Med. Chem. Lett.* 11 (2001) 239–242 (accessed December 10, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/11206468>.
- [104] M. Blacker, M.C. Noe, T.J. Carty, C.G. Goodyer, A.C. LeBlanc, Effect of tumor necrosis factor- $\alpha$  converting enzyme (TACE) and metalloproteinase inhibitor on amyloid precursor protein metabolism in human neurons, *J. Neurochem.* 83 (2002) 1349–1357, <https://doi.org/10.1046/j.1471-4159.2002.01228.x>.
- [105] M.A. Letavic, M.Z. Axt, J.T. Barberia, T.J. Carty, D.E. Danley, K.F. Geoghegan, N.S. Halim, L.R. Hoth, A.V. Kamath, E.R. Laird, L.L. Lopresti-Morrow, K.F. McClure, P.G. Mitchell, V. Natarajan, M.C. Noe, J. Pandit, L. Reeves, G.K. Schulte, S.L. Snow, F.J. Sweeney, D.H. Tan, C.H. Yu, Synthesis and biological activity of selective peipicol acid-based TNF- $\alpha$  converting enzyme (TACE) inhibitors, *Bioorg. Med. Chem. Lett.* 12 (2002) 1387–1390 (accessed December 10, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/11992783>.
- [106] C. Hundhausen, D. Misztela, T.A. Berkhout, N. Broadway, P. Saftig, K. Reiss, D. Hartmann, F. Fahrenholz, R. Postina, V. Matthews, K.-J. Kallen, S. Rose-John, A. Ludwig, The disintegrin-like metalloproteinase ADAM10 is involved in constitutive cleavage of CX3CL1 (fractalkine) and regulates CX3CL1-mediated cell-cell adhesion, *Blood* 102 (2003) 1186–1195, <https://doi.org/10.1182/blood-2002-12-3775>.
- [107] M.M. Thabet, T.W.J. Huizinga, Drug evaluation: apratastat, a novel TACE/MMP inhibitor for rheumatoid arthritis, *Curr. Opin. Investig. Drugs.* 7 (2006) 1014–1019 (accessed August 31, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/17117591>.
- [108] R.C. Newton, E.C. Bradley, R.S. Levy, D. Doval, S. Bondarde, T.P. Sahoo, D. Lokanatha, P.K. Julka, R. Nagarkar, S.M. Friedman, Clinical benefit of INCB7839, a potent and selective ADAM inhibitor, in combination with trastuzumab in patients with metastatic HER2+ breast cancer, 3025–3025, *J. Clin. Oncol.* 28 (2010), <https://doi.org/10.1200/jco.2010.28.15.suppl.3025>.
- [109] H. Kitano, Computational systems biology, *Nature* 420 (2002) 206–210, <https://doi.org/10.1038/nature01254>.
- [110] D. Szklarczyk, A. Franceschini, S. Wyder, K. Forslund, D. Heller, J. Huerta-Cepas, M. Simonovic, A. Roth, A. Santos, K.P. Tsafou, M. Kuhn, L.J. Jensen, C. Von Mering, STRING v10: protein-protein interaction networks, integrated over the tree of life, *Nucleic Acids Res.* 43 (2015), <https://doi.org/10.1093/nar/gku1003>.
- [111] D. Warde-Farley, S.L. Donaldson, O. Comes, K. Zuberi, R. Badrawi, P. Chao, M. Franz, C. Grouios, F. Kazi, C.T. Lopes, A. Maitland, S. Mostafavi, J. Montojo, Q. Shao, G. Wright, G.D. Bader, Q. Morris, The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function, *Nucleic Acids Res.* 38 (2010) W214–W220, <https://doi.org/10.1093/nar/gkq537>.
- [112] P. Shannon, A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, N. Amin, B. Schwikowski, T. Ideker, Cytoscape: a software environment for integrated models of biomolecular interaction networks, *Genome Res.* 13 (2003) 2498–2504, <https://doi.org/10.1101/gr.1239303>.
- [113] T.A. Libermann, D. Baltimore, Activation of interleukin-6 gene expression through the NF- $\kappa$ B transcription factor, *Mol. Cell. Biol.* (1990) 2327–2334 (accessed August 26, 2017).
- [114] L. Tornatore, A.K. Thotakura, J. Bennett, M. Moretti, G. Franzoso, The nuclear factor kappa B signaling pathway: integrating metabolism with inflammation, *Trends Cell Biol.* 22 (2012) 557–566, <https://doi.org/10.1016/j.tcb.2012.08.001>.
- [115] W. Wongchana, T. Palaga, Direct regulation of interleukin-6 expression by Notch signaling in macrophages, *Cell Mol. Immunol.* 9 (2012) 155–162, <https://doi.org/10.1038/cmi.2011.36>.
- [116] K. Miyazawa, A. Mori, K. Yamamoto, H. Okudaira, Transcriptional roles of CCAAT/enhancer binding protein-beta, nuclear factor-kappaB, and C-promoter binding factor 1 in interleukin (IL)-1beta-induced IL-6 synthesis by human rheumatoid fibroblast-like synoviocytes, *J. Biol. Chem.* 273 (1998) 7620–7627 (accessed December 10, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/9516466>.
- [117] A.R. Brasier, The nuclear factor- $\kappa$ B-interleukin-6 signalling pathway mediating vascular inflammation, *Cardiovasc. Res.* 86 (2010) 211–218, <https://doi.org/10.1093/cvr/cvq076>.
- [118] X. Hu, A.Y. Chung, I. Wu, J. Foldi, J. Chen, J.D. Ji, T. Tateya, Y.J. Kang, J. Han, M. Gessler, R. Kageyama, L.B. Ivashkiv, Integrated regulation of toll-like receptor responses by notch and interferon- $\gamma$  pathways, *Immunity* 29 (2008) 691–703, <https://doi.org/10.1016/j.immuni.2008.08.016>.
- [119] T. Palaga, C. Buranaruk, S. Rengpipat, A.H. Fauq, T.E. Golde, S.H.E. Kaufmann, B.A. Osborne, Notch signaling is activated by TLR stimulation and regulates macrophage functions, *Eur. J. Immunol.* 38 (2008) 174–183, <https://doi.org/10.1002/eji.200636999>.
- [120] S. Jin, A.P. Mutvei, I.V. Chivukula, E.R. Andersson, D. Ramsköld, R. Sandberg, K.L. Lee, P. Kronqvist, V. Mamaeva, P. Östling, J.-P. Mpindi, O. Kallioniemi, I. Screpanti, L. Poellinger, C. Sahlgren, U. Lendahl, Non-canonical Notch signaling activates IL-6/JAK/STAT signaling in breast tumor cells and is controlled by p53 and IKK $\alpha$ /IKK $\beta$ , *Oncogene* 32 (2013) 4892–4902, <https://doi.org/10.1038/onc.2012.517>.
- [121] A. Murthy, Y. Washington Shao, S.R. Narala, S.D. Molyneux, J. Carlos Zúñiga-Pflicker, R. Khokha, Notch activation by the metalloproteinase ADAM17 regulates myeloproliferation and atopic barrier immunity by suppressing epithelial cytokine synthesis, *Immunity* 36 (2012) 105–119, <https://doi.org/10.1016/j.immuni.2012.01.005>.
- [122] E.C. Bozkulak, G. Weinmaster, Selective use of ADAM10 and ADAM17 in activation of Notch1 signaling, *Mol. Cell. Biol.* 29 (2009) 5679–5695, <https://doi.org/10.1128/MCB.00406-09>.
- [123] N. Kaviani, A. Servettaz, C. Mongaret, A. Wang, C. Nicco, C. Chéreau, P. Grange, V. Vuiblet, P. Birembaut, M.-D. Diebold, B. Weill, N. Dupin, F. Bateau, Targeting

- ADAM-17/notch signaling abrogates the development of systemic sclerosis in a murine model, *Arthritis Rheum.* 62 (2010) 3477–3487, <https://doi.org/10.1002/art.27626>.
- [124] Z. Guo, X. Jin, H. Jia, Inhibition of ADAM-17 more effectively down-regulates the Notch pathway than that of  $\gamma$ -secretase in renal carcinoma, *J. Exp. Clin. Cancer Res.* 32 (2013) 26, <https://doi.org/10.1186/1756-9966-32-26>.
- [125] G.-T. Ho, P. Chiam, H. Drummond, J. Loane, I.D.R. Arnott, J. Satsangi, The efficacy of corticosteroid therapy in inflammatory bowel disease: analysis of a 5-year UK inception cohort, *Aliment. Pharmacol. Ther.* 24 (2006) 319–330, <https://doi.org/10.1111/j.1365-2036.2006.02974.x>.
- [126] X. Jun, T. Dingyu, F. Yangyang, W. Joseph, Y. Xuezhong, Science China do corticosteroids have a role in treating Ebola virus disease? *Sci China Life Sci.* 58 (2015) 111–113, <https://doi.org/10.1007/s11427-014-4790-3>.
- [127] I.M. Adcock, Corticosteroids: limitations and future prospects for treatment of severe inflammatory disease, *Drug Discov. Today Ther. Strateg.* 1 (2004) 321–328, <https://doi.org/10.1016/j.ddstr.2004.11.015>.
- [128] P.J. Rutgeerts, The limitations of corticosteroid therapy in Crohn's disease, *Aliment. Pharmacol. Ther.* 15 (2001) 1515–1525, <https://doi.org/10.1046/j.1365-2036.2001.01060.x>.
- [129] M. Karin, Y. Yamamoto, Q.M. Wang, The IKK NF- $\kappa$ B system: a treasure trove for drug development, *Nat. Rev. Drug Discov.* 3 (2004) 17–26, <https://doi.org/10.1038/nrd1279>.
- [130] A.C. Sayce, J.L. Miller, N. Zitzmann, Glucocorticosteroids as dengue therapeutics: resolving clinical observations with a primary human macrophage model, *Clin. Infect. Dis.* 56 (2013) 901–903, <https://doi.org/10.1093/cid/cis1048>.
- [131] S. Rajapakse, C. Rodrigo, S. Maduranga, A.C. Rajapakse, Corticosteroids in the treatment of dengue shock syndrome, *Infect. Drug Resist.* 7 (2014) 137, <https://doi.org/10.2147/IDR.S55380>.
- [132] WHO, Dengue: guidelines for diagnosis, treatment, prevention and control, 2009. [http://apps.who.int/iris/bitstream/10665/44188/1/9789241547871\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44188/1/9789241547871_eng.pdf) (accessed December 7, 2017).
- [133] S. Paessler, C. Huang, M. Sencanski, N. Veljkovic, V. Perovic, S. Glisic, V. Veljkovic, Ibuprofen as a template molecule for drug design against Ebola virus, *Front. Biosci. (Landmark Ed.)* 23 (2018) 947–953 (accessed October 26, 2018).
- [134] Y. Furuta, K. Takahashi, M. Kuno-Maekawa, H. Sangawa, S. Uehara, K. Kozaki, N. Nomura, H. Egawa, K. Shiraki, Mechanism of Action of T-705 against Influenza Virus, *Antimicrob. Agents Chemother.* 49 (2005) 981–986, <https://doi.org/10.1128/AAC.49.3.981-986.2005>.
- [135] S.J. Smither, L.S. Eastaugh, J.A. Steward, M. Nelson, R.P. Lenk, M.S. Lever, Post-exposure efficacy of Oral T-705 (Favipiravir) against inhalational Ebola virus infection in a mouse model, *Antiviral Res.* 104 (2014) 153–155, <https://doi.org/10.1016/j.antiviral.2014.01.012>.
- [136] ClinicalTrials.gov, Efficacy of Favipiravir Against Ebola (JIKI) (JIKI), 2014. <https://clinicaltrials.gov/ct2/show/NCT02329054>.
- [137] T.H.T. Nguyen, J. Guedj, X. Anglaret, C. Laouénan, V. Madelain, A.-M. Taburet, S. Baize, D. Sissoko, B. Pastorino, A. Rodallec, G. Piorowski, S. Carazo, M.N. Conde, J.-L. Gala, J.A. Bore, C. Carboneffe, F. Jacquot, H. Raouil, D. Malvy, X. De, Lamballerie, F. Mentre, on behalf of the J. study group, Favipiravir pharmacokinetics in Ebola-infected patients of the JIKI trial reveals concentrations lower than targeted, *PLoS Negl. Trop. Dis.* 11 (2017) e0005389, <https://doi.org/10.1371/journal.pntd.0005389>.
- [138] K.Y. Hostetler, Alkoxyalkyl prodrugs of acyclic nucleoside phosphonates enhance oral antiviral activity and reduce toxicity: current state of the art, *Antivir. Res.* 82 (2009) 84–98, <https://doi.org/10.1016/j.antiviral.2009.01.005>.
- [139] K. Kupferschmidt, J. Cohen, Ebola drug trials lurch ahead, *Science* (80-) 347 (2015) (accessed August 26, 2017).
- [140] S.M. Poutanen, D.E. Low, B. Henry, S. Finkelstein, D. Rose, K. Green, R. Tellier, R. Draker, D. Adachi, M. Ayers, A.K. Chan, D.M. Skowronski, I. Salit, A.E. Simor, A.S. Slutsky, P.W. Doyle, M. Kraiden, M. Petric, R.C. Brunham, A.J. McGeer, National Microbiology Laboratory, Canada, Canadian Severe Acute Respiratory Syndrome Study Team, Identification of Severe Acute Respiratory Syndrome in Canada, *N. Engl. J. Med.* 348 (2003) 1995–2005, <https://doi.org/10.1056/NEJMoã34>.
- [141] L.K.-Y. So, A.C.W. Lau, L.Y.C. Yam, T.M.T. Cheung, E. Poon, R.W.H. Yung, K.Y. Yuen, Development of a standard treatment protocol for severe acute respiratory syndrome, *Lancet* (London, England). 361 (2003) 1615–1617 (accessed December 19, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/12747883>.
- [142] K.W. Tsang, P.L. Ho, G.C. Ooi, W.K. Yee, T. Wang, M. Chan-Yeung, W.K. Lam, W.H. Seto, L.Y. Yam, T.M. Cheung, P.C. Wong, B. Lam, M.S. Ip, J. Chan, K.Y. Yuen, K.N. Lai, A cluster of cases of severe acute respiratory syndrome in Hong Kong, *N. Engl. J. Med.* 348 (2003) 1977–1985, <https://doi.org/10.1056/NEJMoã66>.
- [143] M.R. Loutfy, L.M. Blatt, K.A. Siminovitich, S. Ward, B. Wolff, H. Lho, D.H. Pham, H. Deif, E.A. LaMere, M. Chang, K.C. Kain, G.A. Farcas, P. Ferguson, M. Latchford, G. Levy, J.W. Dennis, E.K.Y. Lai, E.N. Fish, Interferon alfacon-1 plus corticosteroids in severe acute respiratory syndrome, *JAMA* 290 (2003) 3222, <https://doi.org/10.1001/jama.290.24.3222>.
- [144] Z. Zhao, F. Zhang, M. Xu, K. Huang, W. Zhong, W. Cai, Z. Yin, S. Huang, Z. Deng, M. Wei, J. Xiong, P.M. Hawkey, Description and clinical treatment of an early outbreak of severe acute respiratory syndrome (SARS) in Guangzhou, PR China, *J. Med. Microbiol.* 52 (2003) 715–720, <https://doi.org/10.1099/jmm.0.05320-0>.
- [145] K.S. Chan, S.T. Lai, C.M. Chu, E. Tsui, C.Y. Tam, M.M.L. Wong, M.W. Tse, T.L. Que, J.S.M. Peiris, J. Sung, V.C.W. Wong, K.Y. Yuen, Treatment of severe acute respiratory syndrome with lopinavir/ritonavir: a multicentre retrospective matched cohort study, *Hong Kong Med. J.* 9 (2003) 399–406 (accessed December 19, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/14660806>.
- [146] C.M. Chu, V.C.C. Cheng, I.F.N. Hung, M.M.L. Wong, K.H. Chan, K.S. Chan, R.Y.T. Kao, L.L.M. Poon, C.L.P. Wong, Y. Guan, J.S.M. Peiris, K.Y. Yuen, HKU/UCH SARS Study Group, Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings, *Thorax* 59 (2004) 252–256 (accessed December 19, 2017), <http://www.ncbi.nlm.nih.gov/pubmed/14985565>.
- [147] N. Spanakis, S. Tsiodras, B.L. Haagmans, V.S. Raj, K. Pontikis, A. Koutsoukou, N.G. Koulouris, A.D.M.E. Osterhaus, M.P.G. Koopmans, A. Tsakris, Virological and serological analysis of a recent Middle East respiratory syndrome coronavirus infection case on a triple combination antiviral regimen, *Int. J. Antimicrob. Agents.* 44 (2014) 528–532, <https://doi.org/10.1016/j.ijantimicag.2014.07.026>.
- [148] S. Yacoub, J. Mongkolsapaya, G. Screaton, Recent advances in understanding dengue, *PLoS One* 5 (2010) e12688, <https://doi.org/10.1371/journal.pone.0126888>.
- [149] N.M. Nguyen, C.N.B. Tran, L.K. Phung, K.T.H. Duong, H. Le, A. Huynh, J. Farrar, Q.T.H. Nguyen, H.T. Tran, C.V.V. Nguyen, L. Merson, L.T. Hoang, M.L. Hibberd, P.P.K. Aw, A. Wilm, N. Nagarajan, D.T. Nguyen, M.P. Pham, T.T. Nguyen, H. Javanbakht, K. Klumpp, J. Hammond, R. Petric, M. Wolbers, C.T. Nguyen, C.P. Simmons, A randomized, double-blind placebo controlled trial of Balapiravir, a polymerase inhibitor, in adult dengue patients, *J. Infect. Dis.* 207 (2013) 1442–1450, <https://doi.org/10.1093/infdis/jis470>.
- [150] J.G. Low, C. Sung, L. Wijaya, Y. Wei, A.P.S. Rathore, S. Watanabe, B. Hian Tan, L. Toh, L. Tee Chua, an Hou, A. Chow, S. Howe, W. Ki Chan, K. Hin Tan, J.S. Chung, B.P. Cherng, D.C. Lye, P.A. Tambayah, L. Ching Ng, J. Connolly, M.L. Hibberd, Y. Sin Leo, Y. Bun Cheung, E. Eong Ooi, S.G. Vasudevan, Efficacy and safety of celgosivir in patients with dengue fever (CELADEN): a phase 1b, randomised, double-blind, placebo-controlled, proof-of-concept trial, *Lancet Infect. Dis.* 14 (2014) 706–715, [https://doi.org/10.1016/S1473-3099\(14\)70730-3](https://doi.org/10.1016/S1473-3099(14)70730-3).
- [151] V. Tricou, N.N. Minh, T.P. Van, S.J. Lee, J. Farrar, B. Wills, H.T. Tran, C.P. Simmons, A randomized controlled trial of chloroquine for the treatment of dengue in vietnamese adults, *PLoS Negl. Trop. Dis.* 4 (2010) e785, <https://doi.org/10.1371/journal.pntd.0000785>.
- [152] K.J.S. Farias, P.R.L. Machado, R.F. de Almeida Junior, A.A. de Aquino, B.A.L. da Fonseca, Chloroquine interferes with dengue-2 virus replication in U937 cells, *Microbiol. Immunol.* 58 (2014) 318–326, <https://doi.org/10.1111/1348-0421.12154>.
- [153] M. Martínez-Gutiérrez, J.E. Castellanos, J.C. Gallego-Gómez, Statins reduce dengue virus production via decreased virion assembly, *Intervirology* 54 (2011) 202–216, <https://doi.org/10.1159/000321892>.
- [154] M. Martínez-Gutiérrez, L.A. Correa-Londoño, J.E. Castellanos, J.C. Gallego-Gómez, J.E. Osorio, Lovastatin delays infection and increases survival rates in AG129 mice infected with dengue virus serotype 2, *PLoS One* 9 (2014) e87412, <https://doi.org/10.1371/journal.pone.0087412>.
- [155] J. Whitehorn, C.V.V. Nguyen, L.P. Khanh, D.T.H. Kien, N.T.H. Quyen, N.T.T. Tran, N.T. Hang, N.T. Truong, L.T. Hue Tai, N.T. Cam Huong, V.T. Nhon, T. Van Tram, J. Farrar, M. Wolbers, C.P. Simmons, B. Wills, Lovastatin for the treatment of adult patients with dengue: a randomized, double-blind, placebo-controlled trial, *Clin. Infect. Dis.* 62 (2015) 468–476, <https://doi.org/10.1093/cid/civ949>.
- [156] J.A. Pallister, D.J. Middleton, Animal models of recently emerged bat-borne viruses, in: L.-F. Wang, C. Cowled (Eds.), *Bats Viruses*, John Wiley & Sons, Inc, Hoboken, NJ, 2015, p. 295 <http://doi.org/10.1002/9781118818824.ch12>.
- [157] D. Bente, J. Gren, J.E. Strong, H. Feldmann, Disease modeling for Ebola and Marburg viruses, *Dis. Model. Mech.* 2 (2009) 12–17, <https://doi.org/10.1242/dmm.000471>.
- [158] K.J. Olival, C.C. Weekley, P. Daszak, Are Bats Really “Special” as Viral Reservoirs? What We Know and Need to Know, in: *Bats Viruses*, John Wiley & Sons Inc, Hoboken, NJ, 2015, pp. 281–294, <https://doi.org/10.1002/9781118818824.ch11>.
- [159] M.L. Baker, T. Schountz, L.-F. Wang, Antiviral immune responses of bats: a review, *Zoonoses Publ. Health* 60 (2013) 104–116, <https://doi.org/10.1111/j.1863-2378.2012.01528.x>.
- [160] K.B. Clark, N. Onlamoon, H.-M. Hsiao, G.C. Perng, F. Villinger, Can non-human primates serve as models for investigating dengue disease pathogenesis? *Front. Microbiol.* 4 (2013) 305, <https://doi.org/10.3389/fmicb.2013.00305>.
- [161] T. Schountz, Immunology of bats and their viruses: challenges and opportunities, *Viruses* 6 (2014) 4880–4901, <https://doi.org/10.3390/v6124880>.