

---

# Dermatologic care for lesbian, gay, bisexual, and transgender persons



## Epidemiology, screening, and disease prevention

Howa Yeung, MD,<sup>a</sup> Kevin M. Luk, BS,<sup>a</sup> Suephy C. Chen, MD, MS,<sup>a,b</sup> Brian A. Ginsberg, MD,<sup>c</sup> and Kenneth A. Katz, MD, MSc, MSCE<sup>d</sup>

*Atlanta and Decatur, Georgia, New York, New York, and San Francisco, California*

### Learning objectives

After completing this learning activity, participants should be able to recognize skin diseases for which men who have sex with men (MSM), women who have sex with women (WSW), and transgender patients are disproportionately at risk and to provide appropriate treatment or referral; to deliver appropriate preventative health screening and counseling for MSM, WSW, and transgender patients for infectious and noninfectious skin diseases; and to identify cutaneous effects of cross-sex hormones for transgender patients and the roles dermatologists can play in gender affirmation therapy.

### Disclosures

#### Editors

The editors involved with this CME activity and all content validation/peer reviewers of the journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

#### Authors

The authors involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

#### Planners

The planners involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s). The editorial and education staff involved with this journal-based CME activity have reported no relevant financial relationships with commercial interest(s).

Lesbian, gay, bisexual, and transgender (LGBT) persons face important health issues relevant to dermatologists. Men who have sex with men (MSM) are at higher risk of certain infectious diseases, including HIV, syphilis and other sexually transmitted diseases (STDs), methicillin-resistant *Staphylococcus aureus* infections, and invasive meningococcal disease, and might be at higher risk of non-infectious conditions, including skin cancer. Recommendations for preventive health care, including screening for HIV and other STDs, sexual health-related vaccinations, and HIV pre-exposure prophylaxis, differ for MSM compared with non-MSM. Women who have sex with women experience disparities in STDs, including chlamydia and HPV. Transgender patients have unique, and often unmet, dermatologic needs during gender transition (also called gender affirmation), related to hormonal therapy and gender-affirming surgery. Familiarity with LGBT health issues and disease-prevention guidelines can enable dermatologists to provide medically appropriate and culturally competent care to LGBT persons. (J Am Acad Dermatol 2019;80:591-602.)

**Key words:** bisexual; cross-sex hormone; dermal fillers; dermatology; gay; gender affirmation; HIV; indoor tanning; lesbian; LGBT; sexually transmitted diseases; sexual minority; skin cancers; transgender.

---

From the Department of Dermatology,<sup>a</sup> Emory University School of Medicine, Atlanta, and the Division of Dermatology,<sup>b</sup> Atlanta Veterans Affairs Medical Center, Decatur; and the Departments of Dermatology at Mount Sinai Hospital,<sup>c</sup> New York, and Kaiser Permanente,<sup>d</sup> San Francisco.

Supported in part by the Dermatology Foundation and the National Center for Advancing Translational Sciences of the National Institutes of Health under grant numbers UL1TR002378 and KL2TR002381 (Dr Yeung). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Dr Yeung received honorarium from InVentiv Health. Dr Katz was a stockholder in Prevention Health Labs and Arrowhead

---

Pharmaceuticals. All other authors had no potential conflicts of interest to disclose.

Correspondence to: Howa Yeung, MD, Emory University School of Medicine, Department of Dermatology, 1525 Clifton Rd NE, Ste 100, Atlanta, GA 30322. E-mail: [howa.yeung@emory.edu](mailto:howa.yeung@emory.edu). 0190-9622/\$36.00

© 2018 by the American Academy of Dermatology, Inc. Published by Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.jaad.2018.02.045>

**Date of release: March 2019**

**Expiration date: March 2022**

*Abbreviations used:*

CDC:	Centers for Disease Control and Prevention
HHV-8:	human herpesvirus-8
HPV:	human papillomavirus
HSV-2:	herpes simplex virus type 2
LGBT:	lesbian, gay, bisexual, and transgender
MRSA:	methicillin-resistant <i>Staphylococcus aureus</i>
MSM:	men who have sex with men
nPEP:	nonoccupational postexposure prophylaxis
PrEP:	preexposure prophylaxis
STD:	sexually transmitted disease
USPSTF:	US Preventive Services Task Force
WSW:	women who have sex with women

Lesbian, gay, bisexual, and transgender (LGBT) persons face important health disparities that include conditions relevant to dermatology. Knowing about those disparities and about public health approaches to mitigating them can enable dermatologists to provide medically appropriate and culturally competent care to LGBT persons, including referrals to other providers if needed. Dermatologic care for LGBT patients has not been widely discussed in the literature.<sup>1</sup> Because many LGBT health issues relevant to dermatology relate to sexual behavior, rather than sexual orientation, this review largely focuses on behavioral categories, such as men who have sex with men (MSM) and women who have sex with women (WSW). This review describes dermatology-related health concerns among MSM, WSW, and transgender persons.

**MSM****Key points**

- **MSM are at higher risk than men who have sex with women for certain infectious conditions, including HIV and other sexually transmitted diseases, Kaposi sarcoma, viral hepatitis, methicillin-resistant *Staphylococcus aureus* skin infections, and invasive meningococcal disease and might be at higher risk for certain noninfectious conditions, including skin cancer**
- **Recommendations regarding HIV and sexually transmitted disease screening, sexual health-related vaccinations, and HIV preexposure prophylaxis differ for MSM compared with men who have sex with women**
- **Recommendations for HIV nonoccupational postexposure prophylaxis apply to MSM**

Compared with men who have sex with women (MSW), MSM are at higher risk for certain

dermatology-related infectious diseases (Table I).<sup>2-18</sup> As discussed in the first article in this continuing medical education series, eliciting a sexual history, including gender(s) of sex partners, can help physicians stratify patients' risk for those conditions. Eliciting a sexual history can also help physicians determine which, if any, preventive health services their patients might need, because recommendations are in many cases different for MSM than MSW. Those preventive health services could be provided by dermatologists, but if needed dermatologists can and should refer patients to other providers to obtain those services.

**Infectious diseases**

**HIV.** HIV disproportionately affects MSM, who accounted for 615,400 (56%) of 1.1 million persons living with HIV and 26,200 (70%) of 39,513 newly diagnosed HIV infections in the United States in 2014.<sup>2,3</sup> Risk factors for HIV infection include multiple sexual partners, unprotected intercourse (particularly receptive anal sex), coinfection with other sexually transmitted diseases (STDs), and substance use during sex.<sup>19</sup> Acute HIV infection can present with morbilliform eruption, lymphadenopathy, and influenza-like or other signs or symptoms,<sup>20</sup> and cutaneous manifestations of chronic HIV infection include seborrheic dermatitis, folliculitis, psoriasis, prurigo nodularis, and other conditions.<sup>21</sup> Testing for acute HIV infection should include an HIV antigen/antibody immunoassay (also known as a fourth-generation HIV test) and an HIV RNA test (either concurrently or if the immunoassay was negative).<sup>3</sup>

**Syphilis.** The incidence of primary and secondary syphilis from *Treponema pallidum* infection increased from 2.1 cases per 100,000 persons in 2000 to 8.7 cases in 2016, with MSM accounting for 81% of all male cases in 2016.<sup>14</sup> Ocular syphilis and neurosyphilis can develop during any stage of syphilis infection, presenting with cognitive dysfunction, motor or sensory deficits, ophthalmic or auditory symptoms, cranial nerve palsies, symptoms or signs of meningitis or stroke, and uveitis or other ocular manifestations (eg, neuroretinitis and optic neuritis).<sup>3</sup> More than 200 cases of ocular syphilis were reported between 2014 and 2015, mostly affecting HIV-positive MSM.<sup>22</sup> Symptomatic neurosyphilis developed in 2.1% and 0.3%, respectively, of HIV-positive and -negative MSM diagnosed with syphilis in Los Angeles between 2001 and 2004.<sup>23</sup> All patients diagnosed with syphilis, regardless of stage or HIV status, should be asked about neurologic and ocular symptoms and should undergo a neurologic examination, including a cranial nerve examination.<sup>3</sup> Patients with

**Table I.** Selected infectious diseases disproportionately affecting MSM

Viral infections	
HIV	<ul style="list-style-type: none"> <li>• MSM accounted for 70% of 39,513 new HIV cases in the United States in 2014<sup>2,3</sup></li> <li>• The prevalence of HIV infection among MSM in the United States was estimated at 15% in 2012<sup>4</sup></li> <li>• HIV diagnosis from 2005-2014 declined by 18% among white MSM but increased by 22% among black MSM and by 24% among Latino MSM<sup>5</sup></li> </ul>
Genital herpes simplex virus	<ul style="list-style-type: none"> <li>• HSV-2 seroprevalence was 18.4% in MSM and 12.5% in non-MSM<sup>6</sup></li> </ul>
HPV	<ul style="list-style-type: none"> <li>• Anal cancer incidence (cases per 100,000 person-years) was 45.9 among HIV-positive MSM, 5.1 among HIV-negative MSM, and 1.5 among US men overall<sup>7,8</sup></li> <li>• 74% of HIV-positive and 37% of HIV-negative MSM had high-risk anal infection with HPV types 16 or 18<sup>7</sup></li> <li>• 18.2% and 12.7% of men reporting ever having had a same-sex sexual partner had oral HPV infection and high-risk oral HPV infection in 2011-2014, compared with 10.8% and 6.8% of men reporting never having had a same-sex sexual partner<sup>9</sup></li> </ul>
Kaposi sarcoma	<ul style="list-style-type: none"> <li>• Observed in MSM with advanced HIV/AIDS, in MSM with well-controlled HIV infection, and in HIV-negative MSM<sup>10,11</sup></li> </ul>
Viral hepatitis	<ul style="list-style-type: none"> <li>• MSM accounted for 10% and 20% of hepatitis A virus and hepatitis B virus infections in US adults<sup>12</sup></li> <li>• Hepatitis C incidence in HIV-positive MSM tripled between 1991-2012<sup>13</sup></li> </ul>
Bacterial infections	
Syphilis	<ul style="list-style-type: none"> <li>• Cases of primary or secondary syphilis increased to 27,814 in 2016 from 5979 in 2000<sup>3,14</sup></li> <li>• 81% of primary and secondary syphilis cases in 2016 occurred in MSM, 47% of whom were coinfecting with HIV<sup>14</sup></li> <li>• Black MSM had 3.5 times higher rates of syphilis than white MSM in 2014<sup>15</sup></li> </ul>
Gonorrhea and chlamydia	<ul style="list-style-type: none"> <li>• Higher rates of gonorrhea and chlamydia compared with heterosexual men, based on sentinel surveillance at STD clinics<sup>14</sup></li> <li>• Infections can be in the pharynx, rectum, or urethra, underscoring the importance of screening or testing at all 3 sites, if exposed<sup>16</sup></li> </ul>
MRSA	<ul style="list-style-type: none"> <li>• Clusters of multidrug-resistant, community-associated MRSA skin and soft tissue infections have been reported<sup>17</sup></li> </ul>
Meningococcal disease	<ul style="list-style-type: none"> <li>• Meningitis and invasive meningococcal disease outbreaks with high fatality rate have occurred among MSM in North American and European cities<sup>18</sup></li> </ul>

HPV, Human papillomavirus; HSV-2, herpes simplex virus type 2; MRSA, methicillin-resistant *Staphylococcus aureus*; MSM, men who have sex with men; STD, sexually transmitted disease.

ocular or neurologic complaints should also receive ophthalmologic or neurologic evaluation and lumbar puncture with cerebrospinal fluid examination.<sup>22</sup>

**Gonorrhea and chlamydia.** Infections from *Neisseria gonorrhoea* and *Chlamydia trachomatis* may affect the pharynx, rectum, or urethra.<sup>24,25</sup> In 1 study, approximately 70% and 85% of gonorrhea and chlamydia infections in asymptomatic MSM were detected in the pharynx or the rectum rather than the urethra, underscoring the importance of screening at all three sites, if exposed, rather than only the urethra.<sup>16</sup> Urethral gonorrhea typically presents with discharge and dysuria, and can disseminate hematogenously to infect the skin and joints.<sup>26</sup> Lymphogranuloma venereum outbreaks, caused by *C trachomatis* serovars L1 to L3, have been reported among MSM in North America and Europe.<sup>27-29</sup> Unlike the classic presentation of urogenital ulcer and suppurative

inguinal lymphadenopathy, most lymphogranuloma venereum cases presented as proctocolitis without lymphadenopathy.<sup>28,30,31</sup>

**MRSA.** Clusters of multidrug-resistant, community-associated, methicillin-resistant *Staphylococcus aureus* (MRSA) infections have occurred among MSM, mostly affecting skin and soft tissue in the buttocks, genitals, or perineum.<sup>17,32-34</sup> Most MRSA infections are caused by the USA300 clone, which confers resistance to quinolones, mupirocin, tetracycline, and clindamycin.<sup>17</sup> Transmission can occur by direct skin-to-skin contact during sex; risk factors include nasal and perianal colonization, illicit drug use, multiple sexual partners, and previous MRSA infection.<sup>32-34</sup>

**Genital herpes.** Genital herpes, mostly caused by herpes simplex virus type 2 (HSV-2), is characterized by recurrent, painful ulcers. HSV-2 seroprevalence is higher among MSM than heterosexual men.<sup>6</sup> HSV-2 infection can increase HIV acquisition

risk among MSM by 70%.<sup>35</sup> However, suppressive treatment of HSV-seropositive, HIV-negative MSM with acyclovir did not decrease risk of HIV acquisition, although it decreased genital ulcer incidence by 47%.<sup>36</sup>

**Human papillomavirus.** Human papillomavirus (HPV) infection, which disproportionately affect MSM (Table I), can cause anogenital warts, premalignant and malignant anal and penile lesions, and oropharyngeal cancer.<sup>37</sup> Compared with heterosexual men, MSM have a higher prevalence of anal,<sup>38</sup> penile,<sup>38</sup> and oral<sup>9</sup> HPV infection, including infection with high-risk HPV types. According to the Centers for Disease Control and Prevention (CDC), patients with perianal warts might benefit from digital anal examination or referral for standard or high-resolution anoscopy to detect intraanal warts.<sup>3</sup> However, improvements in morbidity or mortality associated with the detection and treatment of intraanal warts have not been shown. Anal cancer incidence among MSM is comparable—and among MSM living with HIV, greater than—the incidence of cervical cancer among women before widespread screening with cervical Papanicolaou smears.<sup>7</sup> According to the CDC, there is insufficient evidence to recommend routine anal cancer screening among MSM, including HIV-positive MSM, although some institutions do perform that screening, particularly for HIV-positive MSM, using anal Papanicolaou smears and high-resolution anoscopy.<sup>3</sup>

**Meningococcal disease.** Outbreaks of invasive *Neisseria meningitidis* infections occurred in MSM, disproportionately affecting HIV-positive MSM, in New York, Los Angeles, Chicago, Toronto, and Europe between 2001 and 2015.<sup>18,39-41</sup> Early symptoms include nonspecific influenza-like illness followed by retiform purpura, hemorrhagic bulla, and disseminated intravascular coagulation. Rapid diagnosis and treatment are critical, given the high case fatality rate of 42%.<sup>18</sup>

**Kaposi sarcoma.** Kaposi sarcoma, caused by human herpesvirus-8, disproportionately affects MSM outside endemic regions, and human herpesvirus-8 seroprevalence among MSM ranges from 8% to 38%.<sup>42-46</sup> In addition to its association with advanced HIV infection, Kaposi sarcoma has also been reported in HIV-positive MSM with high CD4 counts and low HIV viral loads and in HIV-negative MSM.<sup>10,11</sup>

**Viral hepatitis.** MSM accounted for 10% and 20% of incident hepatitis A virus and hepatitis B virus infections, respectively, in US adults.<sup>12</sup> Hepatitis A and B virus infections can present as jaundice and urticaria.<sup>47</sup> Hepatitis A virus is transmitted by the

fecal–oral route, which can include oral–anal intercourse.<sup>48,49</sup> Cutaneous manifestations of hepatitis C virus infection can include lichen planus, mixed cryoglobulinemia, leukocytoclastic vasculitis, and porphyria cutanea tarda.<sup>47</sup> Hepatitis C virus is most commonly transmitted by shared injection drug use, but can also be transmitted by sexual intercourse; risk factors among HIV-positive MSM include mucosally traumatic sex and having sex while using methamphetamine.<sup>13</sup>

## Noninfectious diseases

**Skin cancer and indoor tanning.** Skin cancer and indoor tanning might be more common among MSM.<sup>50-54</sup> In national and California surveys from 2001 to 2013, gay and bisexual men were twice as likely as heterosexual men to report having had nonmelanoma skin cancers.<sup>51</sup> In those surveys, indoor tanning and frequent indoor tanning were 2 to 6 times more common among gay and bisexual men than heterosexual men.<sup>50-52</sup> MSM may also have lower rates of sun protective clothing use despite higher rates of skin cancer screening examinations and sunscreen use.<sup>50</sup> The higher prevalence of body image dissatisfaction and psychological distress among MSM have been proposed as possible mediators for increased indoor tanning.<sup>55</sup> Targeted behavioral counseling and public health interventions may be warranted to reduce indoor tanning and skin cancer risks among MSM.<sup>55</sup>

**“Poppers” dermatitis.** Volatile alkyl nitrites, commonly called poppers, are used by some MSM as recreational inhalants to induce a brief rush with vasodilation, euphoria, smooth muscle relaxation, and sexual arousal.<sup>56</sup> Poppers are often sold as “video head cleaners” and “room odorizers.”<sup>56</sup> Poppers dermatitis presents as an eczematous eruptions around the nasal orifices, perioral region, cheeks, or upper aspect of the chest from either irritant or allergic contact dermatitis caused by amyl nitrite or admixed fragrances.<sup>57</sup> A yellowish crust is commonly present because of the xanthoproteic reaction from nitric acid.<sup>57</sup> Identification of dermatitis in specific areas (eg, in areas corresponding to jean pockets or socks where small glass vials of poppers can be carried, or on the penis onto which poppers were spilled) requires careful history taking and high clinical suspicion.<sup>57,58</sup> Chemical leukoderma in the perinasal region, acrocyanosis, and methemoglobinemia have been reported with chronic use.<sup>59-61</sup> Life-threatening hypotension can occur if poppers are used concurrently with phosphodiesterase-5 inhibitors, such as sildenafil.<sup>62</sup>

**Table II.** Recommendations from the CDC and other public health agencies for HIV and STD screening in MSM\*

Screening	Recommendations	Comment
HIV <sup>64,66</sup>	Unknown or negative HIV status, who have had (or whose sexual partner[s] have had) >1 sexual partner since the last test	Screen at least annually; consider screening every 3-6 months if risk behavior persists or if the man or his sex partner(s) have multiple partners <sup>‡§</sup>
Syphilis <sup>65</sup>	Sexually active in the past year or since the last test	Screen at least annually; consider screening every 3-6 months if risk behavior persists or if the man or his sex partner(s) have multiple partners <sup>‡§</sup>
Gonorrhea and chlamydia*		Screen at least annually; consider screening every 3-6 months if risk behavior persists or if the man or his sex partner(s) have multiple partners <sup>‡§</sup>
Urethra	Any insertive oral or anal intercourse in the past year, regardless of reported condom use	
Rectum	Any receptive anal intercourse in the past year, regardless of reported condom use	
Pharynx <sup>†</sup>	Any receptive oral intercourse in the past year, regardless of reported condom use	
Hepatitis B*	No known previous or current infection, or previous vaccination	One-time screening
Hepatitis C*	HIV-positive persons, including HIV-positive MSM	One-time screening; periodic screening can be considered

CDC, Centers for Disease Control and Prevention; MSM, men who have sex with men; STD, sexually transmitted disease.

\*Data from Workowski et al.<sup>3</sup>

<sup>†</sup>The CDC recommends pharyngeal screening for gonorrhea only, but many available tests screen for both gonorrhea and chlamydia.

<sup>‡</sup>MSM receiving preexposure prophylaxis; should receive HIV testing every 3 months and immediate testing if acute HIV infection is suspected, as well as syphilis, gonorrhea, and chlamydia testing at least every 6 months.<sup>63,64</sup>

<sup>§</sup>Nucleic acid amplification tests rather than cultures for gonorrhea and chlamydia are preferred for all anatomic sites.<sup>3</sup>

### Preventive health services

MSM-specific recommendations from the CDC or the US Preventive Services Task Force (USPSTF) for HIV and STD screening, sexual health-related vaccinations, and HIV preexposure prophylaxis (PrEP) are as follows (Table II).<sup>3,63,64</sup>

**HIV and STD screening.** Screening means testing for a disease in persons without signs or symptoms of that disease. Screening for HIV and other STDs, including syphilis, urethral and rectal gonorrhea and chlamydia, and pharyngeal gonorrhea, should be done annually for nonmonogamous, sexually active MSM, with more frequent screening (every 3-6 months) recommended for MSM with persistent risk behaviors or if they or their sex partners have multiple sex partners.<sup>3</sup> One-time screening for hepatitis B virus infection is recommended for all MSM not known to be or to have been infected, or to have been vaccinated. Screening for hepatitis C virus infection is recommended for HIV-positive MSM.<sup>3</sup> The CDC does not recommend routine screening for HSV-2 infection for MSM. The USPSTF recommends screening for MSM for HIV (at least annually) and syphilis (uncertain optimal frequency; every 3 months appears more effective than annually).<sup>65,66</sup> Both of those USPSTF recommendations have an “A” grade, meaning that there is high certainty that the net benefit is substantial and that they must be covered by US

insurance companies without patients incurring out-of-pocket expenses.<sup>67,68</sup>

**Vaccinations.** MSM should receive the 4- or 9-valent HPV vaccine through 26 years of age (which is the same age cutoff as for women; heterosexual men should be vaccinated through 21 years of age and may be vaccinated through 26 years of age)<sup>69</sup>; hepatitis A and B vaccine if not previously infected<sup>3</sup>; and meningococcal vaccine if HIV-positive or if living in or traveling to cities with outbreaks (Table III).<sup>3,41,63,69-72</sup>

**PrEP.** PrEP refers to chemoprevention for persons at high risk of HIV acquisition, including subgroups of MSM, heterosexually active men and women, and intravenous drug users. Notably, survey data from 2007 to 2012 showed that 24.7% of sexually active, HIV-negative MSM 18 to 59 years of age met the CDC criteria for consideration of PrEP (Table I).<sup>73</sup> The antiretroviral medicine emtricitabine-tenofovir disoproxil fumarate, taken once daily, is approved by the US Food and Drug Administration for PrEP and can reduce HIV incidence by up to 92%.<sup>74</sup> Concurrent safer sex practices, including condom use, are recommended while taking PrEP.<sup>3,63</sup> MSM taking PrEP should be screened for HIV every 3 months and for STDs every 3 to 6 months, depending on risk.<sup>63,64</sup> Although 1 study has shown high STD rates among MSM taking PrEP,<sup>75</sup> another study predicted

**Table III.** Recommendations from the CDC and other public health agencies for sexual health–related vaccinations and HIV pre- and postexposure prophylaxis in MSM\*

Vaccination	Indications	Comment
HPV vaccination <sup>69</sup>	Through 26 years of age, regardless of previous or current HPV infection status	3 doses of 4- or 9-valent vaccine; 2 doses if 11-12 years of age
Hepatitis A virus vaccination <sup>3</sup>	No known previous or current infection, or previous vaccination	2-3 dose series depending on vaccine
Hepatitis B virus vaccination <sup>3</sup>	No known previous or current infection, or previous vaccination	3-dose series
Meningococcal vaccination <sup>41,70</sup>	HIV-positive persons above 2 years of age, including HIV-positive MSM	2-dose series of conjugated vaccine for serotypes A/C/W/Y
HIV prophylaxis PrEP <sup>63</sup>	<ul style="list-style-type: none"> <li>• Adult man</li> <li>• Without acute or established HIV infection</li> <li>• Any male sex partners in past 6 months</li> <li>• Not in a monogamous partnership with a recently tested, HIV-negative man AND at least one of the following</li> <li>• Any anal sex without condoms (receptive or insertive) in past 6 months</li> <li>• A bacterial STD (syphilis, gonorrhea, or chlamydia) diagnosed or reported in past 6 months</li> </ul>	Discuss and refer to PrEP providers to consider PrEP initiation, in addition to safer sex counseling to decrease risk of HIV acquisition <sup>†</sup>
nPEP <sup>72</sup>	Exposure with substantial risk for HIV acquisition within 72 hours from a source known to be HIV-positive; for other exposures with risk, consider on a case-by-case basis	Refer immediately for evaluation and treatment; must initiate nPEP within 72 hours of exposure

CDC, Centers for Disease Control and Prevention; HPV, human papillomavirus; MSM, men who have sex with men; nPEP, nonoccupational postexposure prophylaxis; PrEP, preexposure prophylaxis; STD, sexually transmitted disease.

\*Data from Workowski et al<sup>3</sup> and MacNeil et al.<sup>70</sup>

†A national directory of PrEP providers is available at [prelocator.org](http://prelocator.org).<sup>71</sup>

that frequent STD screening and treatment associated with receiving PrEP could result in a decline in STD incidence.<sup>76</sup> A national directory of PrEP providers can be found at [www.prelocator.org](http://www.prelocator.org).<sup>71</sup>

#### Nonoccupational postexposure prophylaxis.

Nonoccupational postexposure prophylaxis is indicated for HIV-negative persons following exposures that carry substantial risks of HIV acquisition. For MSM, relevant exposures include unprotected receptive anal intercourse with an untreated HIV-positive person. The 28-day course of non-occupational postexposure prophylaxis, consisting of antiretroviral medications, must be initiated within 72 hours after exposure, making prompt referral to urgent care or an emergency department critical.<sup>72</sup>

## WSW

### Key points

- WSW are at risk for HIV and other STDs

- **Guidelines for HIV, STD, and cervical cancer screening and HPV vaccination apply to both WSW and heterosexual women**

Dermatologic concerns among WSW are understudied in the literature. Previously thought to be at low-risk for HIV and other STDs, WSW are at risk for acquiring HIV and STDs from current and previous female or male partners.<sup>77</sup> Few WSW use safer sex practices because many do not face pregnancy risks and perceive themselves to be at low risk for STD acquisition.<sup>78-80</sup> However, transmission of HIV, genital warts, HSV, trichomoniasis, syphilis, hepatitis A virus, and bacterial vaginosis among WSW have been reported.<sup>81,82</sup> Notably, women who have sex with both men and women have higher self-reported STD rates than women who have sex with men only.<sup>77</sup> In some settings, chlamydia prevalence (7.1% vs 5.3%),<sup>83</sup> HSV-2 seropositivity (30-36% vs 24%),<sup>84</sup> bacterial vaginosis (45% vs 29%),<sup>85,86</sup> oral HPV

infection (6.6% vs 2.9%),<sup>9</sup> and oral high-risk HPV infections (3.6% vs 1.2%)<sup>9</sup> were higher among WSW than among women who have sex with men. However, many WSW believe they have less need for screening and report lower Papanicolaou smear use rates.<sup>87</sup> HPV vaccination initiation and completion rates may also be lower among lesbians than heterosexual women.<sup>88-90</sup> Guidelines for HIV, syphilis, chlamydia, gonorrhea, and cervical cancer screening and HPV vaccination do not differ between WSW and women who have sex with men.<sup>3</sup> Safer sex counseling for WSW should highlight the possibility of HIV and STD transmission and acquisition among WSW and encourage safer sex practices.<sup>78-80</sup>

## TRANSGENDER INDIVIDUALS

### Key points

- **Transgender individuals, particularly those receiving gender-affirming hormone and surgical treatments, have unique skin health needs**
- **Dermatologists should be aware of potential complications of gender-affirming treatments and offer appropriate counseling for patients seeking those treatments**

Transgender individuals, particularly those who undergo gender-affirmation treatments (see the first article in this continuing medical education series), experience unmet and unique dermatologic needs. Dermatologists can play important roles in transgender health by helping manage the cutaneous adverse effects of hormonal and surgical treatments, performing safe and effective procedures that contribute to gender affirmation, and facilitating screening and preventive care in a welcoming environment.<sup>91</sup>

### Transgender men

Transgender men who receive cross-sex hormone therapy may receive testosterone through intramuscular or subcutaneous injections or transdermal patch, gel, or cream.<sup>92</sup> Desired effects may include increases in facial and body hair, redistribution of subcutaneous fat, changes in sweat and odor pattern, deepening of voice, cessation of menses, decrease in breast size, clitoral enlargement, and reduction of gender dysphoria.<sup>92,93</sup>

Common cutaneous adverse effects of testosterone treatment include acne vulgaris and androgenetic alopecia. Testosterone significantly increases sebum production, and acne on the face, back, or chest develops in 88% to 94% of patients within 4 to 6 months of initiation of testosterone therapy.<sup>94-96</sup> Most cases decrease in severity after

12 months of testosterone therapy and respond to topical retinoids and topical or oral antibiotics.<sup>95,96</sup> Severe acne in transgender men has been successfully treated with isotretinoin.<sup>97</sup> The US Food and Drug Administration requires patients initiating isotretinoin to register with iPLEDGE, a risk evaluation and management strategy that aims to minimize fetal exposure to isotretinoin, which is a teratogen. iPLEDGE mandates classification of patients according to sex assigned at birth, which is not acceptable to some transgender patients; for that reason, some have advocated that iPLEDGE classify persons by pregnancy potential rather than sex or gender.<sup>93,98</sup> Notably, pregnancies have been reported in transgender men who receive testosterone and are amenorrheic.<sup>99</sup> When considering isotretinoin for transgender patients, clinicians should discuss contraception and pregnancy testing based on a patient's anatomy, pregnancy potential, sexual behavior, and iPLEDGE requirements.<sup>100</sup>

Androgenetic alopecia may be desirable for some transgender men who consider it as a masculine feature and undesirable for others. Severity correlates with duration of testosterone treatment.<sup>93</sup> Thirty-three percent of transgender men develop mild alopecia and 31% develop moderate to severe alopecia after an average of 10 years of testosterone treatment.<sup>95</sup> In a case series, 10 transgender men with grade IV alopecia on the Norwood–Hamilton scale were treated with finasteride 1 mg daily, showed 1 grade improvement after a mean of 5.5 months, and reported no significant side effects after a mean of 16.2 months.<sup>101</sup> Optimal timing and use of selective 5 $\alpha$ -reductase inhibitors or minoxidil to treat testosterone-induced alopecia in transgender men, without blocking desired secondary sex characteristics development, is not established. The Endocrine Society Clinical Practice Guidelines recommends similar alopecia treatments for transgender and cisgender men.<sup>102</sup>

Some transgender men undergo individualized combinations of gender-affirming “top” and “bottom” surgeries. “Top” surgery includes chest reconstruction (mastectomy, nipple–areola complex reduction and reposition, or chest contouring).<sup>103-105</sup> Before such surgery, many transgender men bind their chests, which flattens its appearance but can also commonly result in cutaneous side effects, including pain, swelling, itch, skin breakdown, acne, miliaria, fungal infections, contact dermatitis, and scarring.<sup>106</sup> “Bottom” surgery can include metoidioplasty (which alters clitoral appearance), phalloplasty, urethroplasty, hysterectomy, oophorectomy, and vaginectomy.<sup>103-105</sup> Surgical

scars, particularly keloids from chest reconstruction, may be prominent and distressing and can be treated like other postsurgical keloids with intralesional corticosteroids, radiofrequency devices, or lasers.<sup>103</sup>

### Transgender women

Transgender women who receive cross-sex hormone therapy may receive estrogen through intramuscular injections, transdermal patch, orally, or sublingually.<sup>92</sup> Ethinyl estradiol, common in oral contraceptives, is not recommended because of the high risk of venous thromboembolism.<sup>92</sup> Antiandrogens, such as intramuscular medroxyprogesterone acetate, oral progesterone, spironolactone, finasteride, or dutasteride, may be used with or without estrogen.<sup>92</sup> Desired effects of hormone therapy include breast development, reduction of body and facial hair, redistribution of subcutaneous fat, changes in sweat and odor pattern, arrest or reversal of hair loss, decrease in sebum production, improvement in acne, and reduction in gender dysphoria.<sup>92</sup> Melasma from exogenous estrogens can occur.<sup>107</sup> Facial hair growth is often resistant to hormonal therapy, and hair removal procedures represent the most performed facial procedure among transgender women.<sup>91</sup> Topical eflornithine, electrolysis, photoepilation, or laser hair removal may reduce the need for shaving or depilatory use.<sup>92</sup>

Some transgender women may also undergo gender-affirming surgeries. “Top surgery” may include breast augmentation.<sup>105</sup> Speech therapy or vocal cord surgery may increase the vocal pitch reduce thyroid cartilage prominence.<sup>105</sup> “Bottom surgery” may include vaginoplasty (neovagina reconstruction) with or without penectomy and orchiectomy.<sup>103,105</sup> Transgender men or women undergoing some types of “bottom surgery” may require preoperative laser hair removal at the donor skin site; in transgender women, intravaginal hair growth can cause irritation, infection, formation of hairballs and calculi, and poorer satisfaction with surgical outcomes.<sup>108</sup> However, preoperative donor-site electrolysis did not reduce postoperative intravaginal hair complications in 1 study.<sup>108</sup> Lack of hair regrowth should be confirmed 3 months after the most recent laser session before proceeding with surgery.<sup>109</sup> Condyloma acuminata<sup>110-113</sup> and carcinomas<sup>114-119</sup> of the neovagina have been reported; referral for internal examination can be considered, especially for patients who have previously had external genital warts.

Transgender women often seek feminizing facial and body contouring procedures, which can improve quality of life.<sup>120</sup> Many transgender women have received illicit “silicone” or “filler” injections in

the buttocks, hips, breasts, face, or calves from unlicensed, low-cost “pumpers.”<sup>92,121-123</sup> Injected substances have included food- or industrial-grade silicone, paraffin, petroleum jelly, lanolin, beeswax, various oils, tire sealant, cement glue, and automobile transmission fluid; volumes have ranged from 2 oz to 8 L.<sup>121</sup> Serious complications, including foreign-body granulomas, bacterial or atypical mycobacterial infections, bleeding, pain, scarring, ulceration, fistula formation, gross disfigurement, lymphedema, silicone migration or embolism, sepsis, hypersensitivity pneumonitis, and death have occurred hours to decades later.<sup>92,121</sup> Treatments for filler-induced nodules include intralesional corticosteroids, topical tacrolimus, oral doxycycline, minocycline, or isotretinoin, etanercept, carbon dioxide ablative laser, or surgical excision.<sup>124</sup> Infection should be excluded before immunosuppressant use. Persons injected with “fillers” should be screened for hepatitis C virus infection.<sup>92</sup>

Dermatologists can provide gender-affirming injectable treatments.<sup>91</sup> Botulinum toxin injections can lift, shape, or flatten the forehead and eyebrows, reduce the appearance of periorbital rhytides, or reduce masseter hypertrophy for lower face contouring.<sup>125</sup> Soft tissue augmentation of the cheeks, lips, or chin can be considered.<sup>125</sup> High costs and lack of access to culturally competent providers represent major barriers to care.<sup>91</sup>

Transgender women are at risk for HIV and STDs and should receive screening and prevention based on anatomy-specific sexual behaviors.<sup>92</sup> Notably, 28% of transgender women overall, and 56% of black transgender women, tested HIV-positive in the United States between 1990 and 2003.<sup>126</sup> PrEP referrals can be considered for transgender women who have sex with men, following the same criteria as for MSM (Table II).<sup>92</sup> PrEP appears to be effective when taken appropriately by transgender women,<sup>127</sup> although substantial social and structural barriers to PrEP uptake and adherence need to be addressed.<sup>127</sup>

In conclusion, many LGBT health concerns are relevant to dermatologists. By familiarizing themselves with those health concerns, adhering to disease prevention guidelines, and providing appropriate counseling, treatment, and preventive health care (including referrals when necessary), dermatologists can provide medically appropriate and culturally competent care to LGBT persons.

### REFERENCES

1. Katz KA, Furnish TJ. Dermatology-related epidemiologic and clinical concerns of men who have sex with men, women

- who have sex with women, and transgender individuals. *Arch Dermatol*. 2005;141:1303-1310.
- Centers for Disease Control and Prevention. CDC Fact Sheet: HIV among Gay and Bisexual Men. Available at: <https://www.cdc.gov/nchhstp/newsroom/docs/factsheets/cdc-msm-508.pdf>; 2017. Accessed September 20, 2017.
  - Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Morb Mortal Wkly Rep*. 2015;64(3). Available at: <https://www.cdc.gov/std/tg2015/tg-2015-print.pdf>. Accessed July 7, 2017.
  - Rosenberg ES, Grey JA, Sanchez TH, Sullivan PS. Rates of prevalent HIV infection, prevalent diagnoses, and new diagnoses among men who have sex with men in US states, metropolitan statistical areas, and counties, 2012-2013. *JMIR Public Health Surveill*. 2016;2:e22.
  - Centers for Disease Control and Prevention. CDC Fact Sheet: Trends in U.S. HIV Diagnoses, 2005-2014. Available at: <https://www.cdc.gov/nchhstp/newsroom/docs/factsheets/hiv-data-trends-fact-sheet-508.pdf>; 2016. Accessed September 24, 2017.
  - Xu F, Sternberg MR, Markowitz LE. Men who have sex with men in the United States: demographic and behavioral characteristics and prevalence of HIV and HSV-2 infection: results from National Health and Nutrition Examination Survey 2001-2006. *Sex Transm Dis*. 2010;37:399-405.
  - Machalek DA, Poynten M, Jin F, et al. Anal human papillomavirus infection and associated neoplastic lesions in men who have sex with men: a systematic review and meta-analysis. *Lancet Oncol*. 2012;13:487-500.
  - SEER Cancer Stat Facts: Anal Cancer. Available at: <https://seer.cancer.gov/statfacts/html/anus.html>. Accessed August 9, 2017.
  - Sonawane K, Suk R, Chiao EY, et al. Oral human papillomavirus infection: differences in prevalence between sexes and concordance with genital human papillomavirus infection, NHANES 2011 to 2014. *Ann Intern Med*. 2017;167(10):714-724.
  - Lanternier F, Lebbe C, Scharzt N, et al. Kaposi's sarcoma in HIV-negative men having sex with men. *AIDS*. 2008;22:1163-1168.
  - Maurer T, Ponte M, Leslie K. HIV-associated Kaposi's sarcoma with a high CD4 count and a low viral load. *N Engl J Med*. 2007;357:1352-1353.
  - Centers for Disease Control and Prevention. Hepatitis Fact Sheet for Gay and Bisexual Men. Available at: <http://www.cdc.gov/hepatitis/populations/pdfs/hepgay-factsheet.pdf>; 2013. Accessed July 19, 2016.
  - Hagan H, Jordan AE, Neurer J, Cleland CM. Incidence of sexually transmitted hepatitis C virus infection in HIV-positive men who have sex with men. *AIDS*. 2015;29:2335-2345.
  - Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2016. Atlanta, GA: U.S. Department of Health and Human Services; 2017. Available at: [https://www.cdc.gov/std/stats16/CDC\\_2016\\_STDS\\_Report-for508WebSep21\\_2017\\_1644.pdf](https://www.cdc.gov/std/stats16/CDC_2016_STDS_Report-for508WebSep21_2017_1644.pdf). Accessed September 27, 2017.
  - Grey JA, Bernstein KT, Sullivan PS, et al. Rates of primary and secondary syphilis among white and black non-hispanic men who have sex with men, US states, 2014. *J Acquir Immune Defic Syndr*. 2017;76(3):e65-e73.
  - Patton ME, Kidd S, Llata E, et al. Extragenital gonorrhea and chlamydia testing and infection among men who have sex with men—STD Surveillance Network, United States, 2010-2012. *Clin Infect Dis*. 2014;58:1564-1570.
  - David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev*. 2010;23:616-687.
  - Simon MS, Weiss D, Gulick RM. Invasive meningococcal disease in men who have sex with men. *Ann Intern Med*. 2013;159:300-301.
  - Ackers ML, Greenberg AE, Lin CY, et al. High and persistent HIV seroincidence in men who have sex with men across 47 U.S. cities. *PLoS One*. 2012;7:e34972.
  - Hecht FM, Busch MP, Rawal B, et al. Use of laboratory tests and clinical symptoms for identification of primary HIV infection. *AIDS*. 2002;16:1119-1129.
  - Rigopoulos D, Paparizos V, Katsambas A. Cutaneous markers of HIV infection. *Clin Dermatol*. 2004;22:487-498.
  - Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2016. Atlanta, GA: U.S. Department of Health and Human Services; 2017. Available at: [https://www.cdc.gov/std/stats16/CDC\\_2016\\_STDS\\_Report-for508WebSep21\\_2017\\_1644.pdf](https://www.cdc.gov/std/stats16/CDC_2016_STDS_Report-for508WebSep21_2017_1644.pdf). Accessed September 27, 2017.
  - Taylor MM, Aynalem G, Olea LM, He P, Smith LV, Kerndt PR. A consequence of the syphilis epidemic among men who have sex with men (MSM): neurosyphilis in Los Angeles, 2001-2004. *Sex Transm Dis*. 2008;35:430-434.
  - Centers for Disease Control and Prevention. Clinic-based testing for rectal and pharyngeal *Neisseria gonorrhoeae* and *Chlamydia trachomatis* infections by community-based organizations—five cities, United States, 2007. *MMWR Morb Mortal Wkly Rep*. 2009;58:716-719.
  - Park J, Marcus JL, Pandori M, Snell A, Philip SS, Bernstein KT. Sentinel surveillance for pharyngeal chlamydia and gonorrhea among men who have sex with men—San Francisco, 2010. *Sex Transm Dis*. 2012;39:482-484.
  - Brown TJ, Yen-Moore A, Tyring SK. An overview of sexually transmitted diseases. Part I. *J Am Acad Dermatol*. 1999;41:511-532.
  - Nieuwenhuis RF, Ossewaarde JM, Gotz HM, et al. Resurgence of lymphogranuloma venereum in Western Europe: an outbreak of *Chlamydia trachomatis* serovar I2 proctitis in The Netherlands among men who have sex with men. *Clin Infect Dis*. 2004;39:996-1003.
  - de Vries HJ, Zingoni A, Kreuter A, Moi H, White JA. European Branch of the International Union against Sexually Transmitted Infections et al. 2013 European guideline on the management of lymphogranuloma venereum. *J Eur Acad Dermatol Venereol*. 2015;29:1-6.
  - Centers for Disease Control and Prevention. Lymphogranuloma venereum among men who have sex with men—Netherlands, 2003-2004. *MMWR Morb Mortal Wkly Rep*. 2004;53:985-988.
  - Pathela P, Blank S, Schillinger JA. Lymphogranuloma venereum: old pathogen, new story. *Curr Infect Dis Rep*. 2007;9:143-150.
  - Saxon C, Hughes G, Ison C, UK LGV Case-Finding Group. Asymptomatic lymphogranuloma venereum in men who have sex with men, United Kingdom. *Emerg Infect Dis*. 2016;22:112-116.
  - Galindo GR, Casey AJ, Yeung A, Weiss D, Marx MA. Community associated methicillin resistant *Staphylococcus aureus* among New York City men who have sex with men: qualitative research findings and implications for public health practice. *J Community Health*. 2012;37:458-467.
  - Centers for Disease Control and Prevention. Outbreaks of community-associated methicillin-resistant *Staphylococcus aureus* skin infections—Los Angeles County, California, 2002-2003. *MMWR Morb Mortal Wkly Rep*. 2003;52:88.

34. Szumowski JD, Wener KM, Gold HS, et al. Methicillin-resistant *Staphylococcus aureus* colonization, behavioral risk factors, and skin and soft-tissue infection at an ambulatory clinic serving a large population of HIV-infected men who have sex with men. *Clin Infect Dis*. 2009;49:118-121.
35. Freeman EE, Weiss HA, Glynn JR, Cross PL, Whitworth JA, Hayes RJ. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *AIDS*. 2006;20:73-83.
36. Celum C, Wald A, Hughes J, et al. Effect of aciclovir on HIV-1 acquisition in herpes simplex virus 2 seropositive women and men who have sex with men: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2008;371:2109-2119.
37. Viens LJ, Henley SJ, Watson M, et al. Human papillomavirus-associated cancers - United States, 2008-2012. *MMWR Morb Mortal Wkly Rep*. 2016;65:661-666.
38. Goldstone S, Palefsky JM, Giuliano AR, et al. Prevalence of and risk factors for human papillomavirus (HPV) infection among HIV-seronegative men who have sex with men. *J Infect Dis*. 2011;203:66-74.
39. Folaranmi TA, Kretz CB, Kamiya H, et al. Increased risk for meningococcal disease among men who have sex with men in the United States, 2012-2015. *Clin Infect Dis*. 2017;65(5):756-763.
40. Tsang RS, Kiefer L, Law DK, et al. Outbreak of serogroup C meningococcal disease caused by a variant of *Neisseria meningitidis* serotype 2a ET-15 in a community of men who have sex with men. *J Clin Microbiol*. 2003;41:4411-4414.
41. Kamiya H, MacNeil J, Blain A, et al. Meningococcal disease among men who have sex with men - United States, January 2012-June 2015. *MMWR Morb Mortal Wkly Rep*. 2015;64:1256-1257.
42. Martin JN, Ganem DE, Osmond DH, Page-Shafer KA, Macrae D, Kedes DH. Sexual transmission and the natural history of human herpesvirus 8 infection. *N Engl J Med*. 1998;338:948-954.
43. Engels EA, Atkinson JO, Graubard BI, et al. Risk factors for human herpesvirus 8 infection among adults in the United States and evidence for sexual transmission. *J Infect Dis*. 2007;196:199-207.
44. Gogineni E, Marshall V, Miley W, et al. Quantitative determinations of anti-Kaposi sarcoma-associated herpesvirus antibody levels in men who have sex with men. *Diagn Microbiol Infect Dis*. 2013;76:56-60.
45. Regamey N, Cathomas G, Schwager M, Wernli M, Harr T, Erb P. High human herpesvirus 8 seroprevalence in the homosexual population in Switzerland. *J Clin Microbiol*. 1998;36:1784-1786.
46. Rohner E, Wyss N, Trelle S, et al. HHV-8 seroprevalence: a global view. *Syst Rev*. 2014;3:11.
47. Dogra S, Jindal R. Cutaneous manifestations of common liver diseases. *J Clin Exp Hepatol*. 2011;1:177-184.
48. Urbanus AT, van Houdt R, van de Laar TJ, Coutinho RA. Viral hepatitis among men who have sex with men, epidemiology and public health consequences. *Euro Surveill*. 2009;14.
49. Corey L, Holmes KK. Sexual transmission of hepatitis A in homosexual men: incidence and mechanism. *N Engl J Med*. 1980;302:435-438.
50. Yeung H, Chen SC. Disparities in sunburns, photoprotection, indoor tanning, and skin cancer screening among U.S. men and women in same- and opposite-sex relationships [abstract]. *The Journal of investigative dermatology*. 2015;135(Suppl 1):S49.
51. Mansh M, Katz KA, Linos E, Chren MM, Arron S. Association of skin cancer and indoor tanning in sexual minority men and women. *JAMA Dermatol*. 2015;151:1308-1316.
52. Yeung H, Chen SC. Sexual orientation and indoor tanning device use: a population-based study. *JAMA Dermatol*. 2016;152:99-101.
53. Mansh M, Arron ST. Indoor tanning and melanoma: are gay and bisexual men more at risk? *Melanoma Management*. 2016;3:89-92.
54. Blashill AJ. Indoor tanning and skin cancer risk among diverse US youth: results from a National sample. *JAMA Dermatol*. 2017;153:344-345.
55. Blashill AJ, Pagoto S. Skin cancer risk in gay and bisexual men: a call to action. *JAMA Dermatol*. 2015;151:1293-1294.
56. Colfax G, Coates TJ, Husnik MJ, et al. Longitudinal patterns of methamphetamine, popper (amyl nitrite), and cocaine use and high-risk sexual behavior among a cohort of san francisco men who have sex with men. *J Urban Health*. 2005;82:i62-i70.
57. Schaubert J, Herzinger T. 'Poppers' dermatitis. *Clin Exp Dermatol*. 2012;37:587-588.
58. Latini A, Lora V, Zaccarelli M, Cristaudo A, Cota C. Unusual Presentation of Poppers Dermatitis. *JAMA Dermatol*. 2017;153(2):233-234.
59. Vine K, Meulener M, Shieh S, Silverberg NB. Vitiliginous lesions induced by amyl nitrite exposure. *Cutis*. 2013;91:129-136.
60. Hoegl L, Thoma-Greber E, Poppinger J, Rocken M. Butyl nitrite-induced acrocyanosis in an HIV-infected patient. *Arch Dermatol*. 1999;135:90-91.
61. Hunter L, Gordge L, Dargan PI, Wood DM. Methaemoglobinemia associated with the use of cocaine and volatile nitrites as recreational drugs: a review. *Br J Clin Pharmacol*. 2011;72:18-26.
62. Schwartz BG, Kloner RA. Drug interactions with phosphodiesterase-5 inhibitors used for the treatment of erectile dysfunction or pulmonary hypertension. *Circulation*. 2010;122:88-95.
63. Centers for Disease Control and Prevention. US Public Health Service: Preexposure prophylaxis for the prevention of HIV infection in the United States—2017 Update: a clinical practice guideline. Available at: <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2017.pdf>; 2018. Accessed May 15, 2018.
64. DiNenno EA, Prejean J, Irwin K, et al. Recommendations for HIV screening of gay, bisexual, and other men who have sex with men — United States, 2017. *MMWR Morb Mortal Wkly Rep*. 2017;66:830-832.
65. U.S. Preventive Services Task Force, Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for Syphilis Infection in Nonpregnant Adults and Adolescents: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2016;315:2321-2327.
66. Moyer VA, U.S. Preventive Services Task Force. Screening for HIV: U.S. preventive services task force recommendation statement. *Ann Intern Med*. 2013;159:51-60.
67. Agency for Healthcare Research and Quality and U.S. Preventive Services Task Force. The Guide to Clinical Preventive Services 2014. Preface. Available at: <https://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/guide/preface.html>. Accessed September 25, 2017.
68. U.S. Preventive Services Task Force. Grade Definitions. Available at: <https://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>; 2016. Accessed September 27, 2017.

69. Markowitz LE, Dunne EF, Saraiya M, Centers for Disease Control and Prevention. Human papillomavirus vaccination: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2014; 63:1-30.
70. MacNeil JR, Rubin LG, Patton M, Ortega-Sanchez IR, Martin SW. Recommendations for use of meningococcal conjugate vaccines in HIV-infected persons - Advisory Committee on Immunization Practices, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:1189-1194.
71. Siegler AJ, Wirtz SS, Weber S, Sullivan PS. Developing a geolocated directory of PrEP-providing clinics: the PrEP Locator Protocol and Operating Procedures. *JMIR Public Health Surveill*. 2017;3(3):e58.
72. Centers for Disease Control and Prevention. Updated guidelines for antiretroviral postexposure prophylaxis after sexual, injection-drug use, or other nonoccupational exposure to HIV - United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:458.
73. Smith DK, Van Handel M, Wolitski RJ, et al. Vital signs: estimated percentages and numbers of adults with indications for preexposure prophylaxis to prevent HIV acquisition--United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2015;64:1291-1295.
74. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med*. 2010;363:2587-2599.
75. Volk JE, Marcus JL, Phengrasamy T, et al. No new HIV infections with increasing use of HIV preexposure prophylaxis in a clinical practice setting. *Clin Infect Dis*. 2015;61:1601-1603.
76. Jenness SM, Weiss KM, Goodreau SM, et al. Incidence of gonorrhea and chlamydia following HIV preexposure prophylaxis among men who have sex with men: a modeling study. *Clin Infect Dis*. 2017;65(5):712-718.
77. Everett BG. Sexual orientation disparities in sexually transmitted infections: examining the intersection between sexual identity and sexual behavior. *Arch Sex Behav*. 2013;42: 225-236.
78. Cox P, McNair R. Risk reduction as an accepted framework for safer-sex promotion among women who have sex with women. *Sex Health*. 2009;6:15-18.
79. Marrazzo JM, Coffey P, Bingham A. Sexual practices, risk perception and knowledge of sexually transmitted disease risk among lesbian and bisexual women. *Perspect Sex Reprod Health*. 2005;37:6-12.
80. Dolan KA, Davis PW. Nuances and shifts in lesbian women's constructions of STI and HIV vulnerability. *Soc Sci Med*. 2003; 57:25-38.
81. Gorgos LM, Marrazzo JM. Sexually transmitted infections among women who have sex with women. *Clin Infect Dis*. 2011;53(Suppl 3):S84-S91.
82. Koh AS, Gomez CA, Shade S, Rowley E. Sexual risk factors among self-identified lesbians, bisexual women, and heterosexual women accessing primary care settings. *Sex Transm Dis*. 2005;32:563-569.
83. Singh D, Fine DN, Marrazzo JM. Chlamydia trachomatis infection among women reporting sexual activity with women screened in family planning clinics in the Pacific Northwest, 1997 to 2005. *Am J Public Health*. 2011;101:1284-1290.
84. Xu F, Sternberg MR, Markowitz LE. Women who have sex with women in the United States: prevalence, sexual behavior and prevalence of herpes simplex virus type 2 infection--results from national health and nutrition examination survey 2001-2006. *Sex Transm Dis*. 2010;37:407-413.
85. Berger BJ, Kolton S, Zenilman JM, Cummings MC, Feldman J, McCormack WM. Bacterial vaginosis in lesbians: a sexually transmitted disease. *Clin Infect Dis*. 1995;21:1402-1405.
86. Bailey JV, Farquhar C, Owen C. Bacterial vaginosis in lesbians and bisexual women. *Sex Transm Dis*. 2004;31:691-694.
87. Valanis BG, Bowen DJ, Bassford T, Whitlock E, Charney P, Carter RA. Sexual orientation and health: comparisons in the women's health initiative sample. *Archives of family medicine*. 2000;9:843-853.
88. Agenor M, Peitzmeier S, Gordon AR, Haneuse S, Potter JE, Austin SB. Sexual orientation identity disparities in awareness and initiation of the human papillomavirus vaccine among U.S. women and girls: a National survey. *Ann Intern Med*. 2015;163:99-106.
89. Charlton BM, Reisner SL, Agenor M, Gordon AR, Sarda V, Austin SB. Sexual orientation disparities in human papillomavirus vaccination in a longitudinal cohort of U.S. males and females. *LGBT Health*. 2017;4:202-209.
90. Agenor M, Peitzmeier SM, Gordon AR, et al. Sexual orientation identity disparities in human papillomavirus vaccination initiation and completion among young adult US women and men. *Cancer Causes Control*. 2016;27: 1187-1196.
91. Ginsberg BA, Calderon M, Seminara NM, Day D. A potential role for the dermatologist in the physical transformation of transgender people: A survey of attitudes and practices within the transgender community. *J Am Acad Dermatol*. 2016;74:303-308.
92. Center of Excellence for Transgender Health, Department of Family and Community Medicine, University of California San Francisco. *Guidelines for the Primary and Gender-Affirming Care of Transgender and Gender Nonbinary People*. 2nd edition. Deutsch MB; 2016.
93. Irwig MS. Testosterone therapy for transgender men. *Lancet Diabetes Endocrinol*. 2017;5(4):301-311.
94. Giltay EJ, Gooren LJ. Effects of sex steroid deprivation/ administration on hair growth and skin sebum production in transsexual males and females. *J Clin Endocrinol Metab*. 2000;85:2913-2921.
95. Wierckx K, Van de Peer F, Verhaeghe E, et al. Short- and long-term clinical skin effects of testosterone treatment in trans men. *J Sex Med*. 2014;11:222-229.
96. Nakamura A, Watanabe M, Sugimoto M, et al. Dose-response analysis of testosterone replacement therapy in patients with female to male gender identity disorder. *Endocr J*. 2013;60: 275-281.
97. Turrion-Merino L, Urech-Garcia-de-la-Vega M, Miguel-Gomez L, Harto-Castano A, Jaen-Olasolo P. Severe Acne in Female-to-Male Transgender Patients. *JAMA Dermatol*. 2015;151(11): 1260-1261.
98. Katz KA. Transgender patients, isotretinoin, and US Food and Drug Administration-mandated risk evaluation and mitigation strategies: a prescription for inclusion. *JAMA dermatology*. 2016;152:513-514.
99. Light AD, Obedin-Maliver J, Sevelius JM, Kerns JL. Transgender men who experienced pregnancy after female-to-male gender transitioning. *Obstet Gynecol*. 2014; 124:1120-1127.
100. Yeung H, Chen SC, Katz KA, Stoff BK. Prescribing isotretinoin in the United States for transgender individuals: ethical considerations. *J Am Acad Dermatol*. 2016;75:648-651.
101. Moreno-Arrones OM, Becerra A, Vano-Galvan S. Therapeutic experience with oral finasteride for androgenetic alopecia in female-to-male transgender patients. *Clin Exp Dermatol*. 2017;42(7):743-748.

102. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: an Endocrine Society clinical practice guideline. *Endocr Pract*. 2017;23(12):1437.
103. Ginsberg BA. Dermatologic care of the transgender patient. *Int J Womens Dermatol*. 2017;3:65-67.
104. Frey JD, Poudrier G, Chiodo MV, Hazen A. An update on genital reconstruction options for the female-to-male transgender patient: a review of the literature. *Plast Reconstr Surg*. 2017;139:728-737.
105. Berli JU, Knudson G, Fraser L, et al. What surgeons need to know about gender confirmation surgery when providing care for transgender individuals: a review. *JAMA Surg*. 2017;152:394-400.
106. Peitzmeier S, Gardner I, Weinand J, Corbet A, Acevedo K. Health impact of chest binding among transgender adults: a community-engaged, cross-sectional study. *Cult Health Sex*. 2017;19:64-75.
107. Erickson-Schroth L. *Trans Bodies, Trans Selves: A Resource for the Transgender Community*. Oxford University Press; 2014.
108. Lawrence AA. Patient-reported complications and functional outcomes of male-to-female sex reassignment surgery. *Arch Sex Behav*. 2006;35:717-727.
109. Zhang WR, Garrett GL, Arron ST, Garcia MM. Laser hair removal for genital gender affirming surgery. *Transl Androl Urol*. 2016;5:381-387.
110. Liguori G, Trombetta C, Bucci S, et al. Condylomata acuminata of the neovagina in a HIV-seropositive male-to-female transsexual. *Urol Int*. 2004;73:87-88.
111. Buscema J, Rosenshein NB, Shah K. Condylomata acuminata arising in a neovagina. *Obstet Gynecol*. 1987;69:528-530.
112. Wasef W, Sugunendran H, Alawattegama A. Genital warts in a transsexual. *Int J STD AIDS*. 2005;16:388-389.
113. Yang C, Liu S, Xu K, Xiang Q, Yang S, Zhang X. Condylomata gigantea in a male transsexual. *Int J STD AIDS*. 2009;20:211-212.
114. Fernandes HM, Manolitsas TP, Jobling TW. Carcinoma of the neovagina after male-to-female reassignment. *J Low Genit Tract Dis*. 2014;18:E43-E45.
115. Harder Y, Erni D, Banic A. Squamous cell carcinoma of the penile skin in a neovagina 20 years after male-to-female reassignment. *Br J Plast Surg*. 2002;55:449-451.
116. Rotmensch J, Rosenshein N, Dillon M, Murphy A, Woodruff JD. Carcinoma arising in the neovagina: case report and review of the literature. *Obstet Gynecol*. 1983;61:534-536.
117. Steiner E, Woernle F, Kuhn W, et al. Carcinoma of the neovagina: case report and review of the literature. *Gynecol Oncol*. 2002;84:171-175.
118. Bollo J, Balla A, Rodriguez Luppi C, Martinez C, Quaresima S, Targarona EM. HPV-related squamous cell carcinoma in a neovagina after male-to-female gender confirmation surgery. *Int J STD AIDS*. 2017, 956462417728856.
119. Schober JM. Cancer of the neovagina. *J Pediatr Urol*. 2007;3:167-170.
120. Ainsworth TA, Spiegel JH. Quality of life of individuals with and without facial feminization surgery or gender reassignment surgery. *Qual Life Res*. 2010;19:1019-1024.
121. Wilson E, Rapues J, Jin H, Raymond HF. The use and correlates of illicit silicone or "fillers" in a population-based sample of transwomen, San Francisco, 2013. *J Sex Med*. 2014;11:1717-1724.
122. Benotsch EG, Zimmerman RS, Cathers L, et al. Non-medical use of prescription drugs and HIV risk behaviour in transgender women in the Mid-Atlantic region of the United States. *Int J STD AIDS*. 2016;27:776-782.
123. Kobrak P, White B. Transgender Women and HIV Prevention in New York City: A Needs Assessment. Available at: <https://www1.nyc.gov/assets/doh/downloads/pdf/ah/transgender-women.pdf>; 2010. Accessed August 16, 2017.
124. Paul S, Goyal A, Duncan LM, Smith GP. Granulomatous reaction to liquid injectable silicone for gluteal enhancement: review of management options and success of doxycycline. *Dermatol Ther*. 2015;28:98-101.
125. Ginsberg BA. *Dermatology*: Eckstrand KL, Ehrenfeld JM, eds. *Lesbian, Gay, Bisexual, and Transgender Healthcare: A Clinical Guide to Preventive, Primary, and Specialist Care*. Springer; 2016.
126. Herbst JH, Jacobs ED, Finlayson TJ, et al. Estimating HIV prevalence and risk behaviors of transgender persons in the United States: a systematic review. *AIDS Behav*. 2008;12:1-17.
127. Deutsch MB, Glidden DV, Sevelius J, et al. HIV pre-exposure prophylaxis in transgender women: a subgroup analysis of the iPrEx trial. *Lancet HIV*. 2015;2:e512-e519.