

Seminar Article

Development and translation of novel therapeutics targeting tumor-associated macrophages

David Kosoff, M.D.^{a,b,*}, Joshua M. Lang, M.D., M.S.^{a,b}

^a Department of Medicine, University of Wisconsin-Madison, Madison, WI

^b Carbone Cancer Center, University of Wisconsin-Madison, Madison, WI

Received 6 July 2018; received in revised form 25 September 2018; accepted 3 October 2018

Abstract

Tumor-associated macrophages (TAMs) regulate an array of tumor functions and have critical roles in both the progression and the eradication of cancer. Numerous therapies targeting TAMs are under development in cancer and many have demonstrated success at the preclinical and clinical levels. Most of these therapies fall within 3 main categories: systemic depletion of TAMs, inhibition of TAM recruitment and polarization, and promoting the antitumor functions of TAMs. In this article, the rationale behind these various therapies and approaches is reviewed along with supporting preclinical and clinical data. © 2018 Elsevier Inc. All rights reserved.

Keywords: Tumor-associated macrophages; Therapy development; Cancer; Tumor microenvironment

1. Introduction

Macrophages are myeloid immune cells with integral roles in a wide range of physiologic processes. Within these roles, macrophages can perform a diverse range of functions, including matrix remodeling, growth factor secretion, angiogenesis, pathogen eradication, and immune regulation [1–7]. In cancer, this broad functional capacity enables critical roles in both tumor progression and tumor destruction. These roles are dependent on their phenotype, which are grouped into 2 main categories, termed M1 and M2. M1 macrophages support tumor eradication through stimulation of innate and adaptive immune responses. M2 macrophages promote tumor growth, invasion, and therapeutic resistance through their roles in growth factor secretion, angiogenesis, and immune suppression. Within the tumor microenvironment (TME), the phenotype of these tumor-associated macrophages (TAMs) is regulated in large part by an array of factors, which are secreted by tumor, immune, and stromal cells as well as physical interactions with the extracellular matrix.

Disclosures: The authors have no relevant financial disclosures.

*Corresponding author. Tel: +1-847-207-0383.

E-mail address: dkosoff83@gmail.com (D. Kosoff).

In both solid and hematologic malignancies, the extent of TAM infiltration within the TME as well as the predominant TAM phenotype has direct correlation with patient outcomes [8]. In cancers where TAMs are predominantly of the M2 phenotype, increased TAM infiltration is associated with more advanced disease at diagnosis as well as worse progression-free survival (PFS) and overall survival (OS) [9]. However, when TAMs are skewed more toward an M1 phenotype, their infiltration is associated with better cancer-related outcomes, including a decreased risk of death.

Given their numerous tumor supportive functions, their potential antitumor properties, and their strong correlation with patient outcomes, TAMs represent an attractive therapeutic target with vast potential. A number of approaches for targeting TAMs in cancer are under investigation and have demonstrated success at both the preclinical and clinical levels.

2. Systemic depletion of TAMs

Several TAM-directed strategies have focused on the systemic depletion of monocyte/macrophage populations through the administration of compounds that have macrophage-selective toxicity. Among these is trabectedin, a

DNA binding agent that promotes apoptosis of macrophages through tumor necrosis factor (TNF)-related apoptosis-inducing ligand receptors 1 and 2 [10]. In both Europe and the United States, trabectedin has been approved for treatment of sarcomas based on improvements in PFS and clinical benefit rates. This therapy was overall well tolerated in phase III studies, with hematologic toxicity and transient transaminitis as the most frequent complications. Rates of alopecia, cardiotoxicity, and neurotoxicity were notably low. The macrophage-directed toxicity has been found to be a key component of the antitumor effect of trabectedin [10,11] and multiple clinical trials are now investigating trabectedin as an antineoplastic agent with a focus on its macrophage-mediated effects (Table 1).

Bisphosphonates are another class of drugs that induce macrophage apoptosis [12]. Their effect is mediated through multiple pathways, which include impairing mitochondrial function and inhibiting protein prenylation, both of which lead to induction of cellular apoptosis. While these agents have roles in the prevention of skeletal-related events as antiresorptive agents in multiple cancers including breast and prostate cancer, their impact on PFS and OS is limited. In an effort to maximize the antitumor activity of these agents, numerous approaches have been

developed that enhance their macrophage toxicity. This includes the loading of bisphosphonates into liposomes, which are preferentially endocytosed by macrophages following systemic administration. In preclinical studies, which include murine teratocarcinoma models, this strategy has resulted in depletion of systemic and tumor macrophage populations with associated slowed tumor growth, decreased vascular density [13], and delayed tumor recovery from radiation [14].

Immunotherapy has also been utilized to target macrophages in cancer and has demonstrated success at the preclinical level. Legumain is an endopeptidase that is up-regulated in M2 macrophages with little to no expression in M1 macrophages. Vaccination against legumain in murine cancer models has resulted in CD8-mediated depletion of M2 macrophages with resultant decreased metastatic burden and increased survival [1,15]. Other surface molecules have also been targeted. Alemtuzumab, a monoclonal antibody directed against CD52, a surface glycoprotein expressed by monocyte/macrophage, has been found to induce complement-mediated lysis of monocytes in vitro. In a murine ovarian tumor model, administration of alemtuzumab resulted in decreased angiogenesis as well as slowed tumor growth [16].

Table 1
Ongoing clinic trials for TAM-targeted agents

Target	TAM-directed therapy	Cancer	Combination therapy	Phase	Trial		
TRAIL 1,2	Trabectedin	Sarcoma	–	2	NCT03397186		
		Mesothelioma	–	2	NCT02194231		
CCR2	PF-04136309	Pancreatic	Nab-paclitaxel/gemcitabine	1b/2	NCT02732938		
CCR2/5	BMS 813160	Pancreatic	Nivolumab/gemcitabine/Nab-paclitaxel	1/2	NCT03496662		
CSF-1R	Pexidartinib	Solid tumors	PLX9486	1/2	NCT02401815		
		Solid tumors	–	1	NCT01004861		
		Solid/hematologic	–	1/2	NCT02390752		
		Melanoma	–	1/2	NCT02975700		
		Glioblastoma	Radiation/temozolamide	1/2	NCT01790503		
		Pancreatic/colon	Durvalumab	1	NCT02777710		
		Breast	Eribulin	1/2	NCT01596751		
		Prostate	Antiandrogen/hormone	1	NCT02472275		
		Peripheral nerve sheath tumors	Sirolimus	1/2	NCT02584647		
		Melanoma	Pembrolizumab	1/2	NCT02452424		
		Melanoma	–	2	NCT02071940		
		CD40	RG7155	Solid tumors	Atezolizumab	1	NCT02323191
				Female genital	Paclitaxel/bevacizumab	2	NCT02923739
				Female genital	Paclitaxel/bevacizumab	2	NCT02923739
		CD47	CP-870,893	Melanoma	Tremelimumab	1	NCT01103635
CD47	CC-90002	Hematologic	–	1	NCT02641002		
		Solid/hematologic	Rituximab	1	NCT02367196		
	Hu5F9-G4	TTI-621	Solid/hematologic	–	1	NCT02678338	
			Hematologic	Rituximab/PD-1 inhibitor/proteasome Inhibitor	1	NCT03530683	
	SRF231	ALX148	Solid/hematologic	Rituximab/nivolumab	1	NCT02663518	
			Solid/hematologic	–	1	NCT03512340	
	Hu5F9-G4	Hu5F9-G4	Solid/hematologic	Rituximab/trastuzumab/pembrolizumab	1	NCT03013218	
			Hematologic	Azacitidine	1	NCT03248479	
			Solid tumors	Cetuximab	1/2	NCT02953782	

3. Targeting recruitment and reprogramming

Aiming to limit or even alter the role of macrophages within the TME, multiple TAM-directed strategies have focused on targeting the pathways involved in TAM recruitment and polarization. One of the most extensively studied targets of macrophage recruitment is the CCL2. CCL2 is a chemokine that is secreted by cells within the TME that recruits TAMs into tumors through interaction with the macrophage CCR2 receptor. Targeting CCL2 has shown considerable promise at the preclinical level, resulting in enhanced tumor responses and improved survival through a reduction in TAM infiltration [17–20]. Other strategies targeting this pathway include inhibition of the CCR2 receptor on macrophages. This strategy offers theoretical benefit over CCL2-directed therapies as CCR2 is the receptor for multiple macrophage chemoattractant proteins and recruitment pathways. PF-04136309 is a potent antagonist of the CCR2 receptor that is currently under investigation as a cancer therapy. After successful preclinical evaluation, administration of PF-04136309 in combination with FOLFIRINOX chemotherapy resulted in objective response rates approaching 50% in a phase 1b study of patients with locally advanced pancreatic adenocarcinoma [21]. Additional trials targeting CCR2 in cancer patients are ongoing (Table 1).

The CSF-1 pathway is another target for TAM-directed therapies that has shown considerable promise in preclinical and clinical investigations. In addition to involvement in TAM recruitment, CSF-1 also promotes M2 polarization of TAMs within the TME. This pathway is of particular value as CSF-1R is expressed exclusively on myeloid cells and may potentially be targeted with lower toxicity. Numerous agents are under development targeting this pathway and preclinical studies have demonstrated efficacy in a range of cancers, including prostate, brain, breast, pancreas, ovaries, sarcomas, and hematologic malignancies [22–25]. A number of these therapies have been further evaluated at the clinical level and have already demonstrated antitumor efficacy. In early-phase trials, emactuzumab (RG7155), a monoclonal antibody against the CSF-1R, elicited objective responses in up to 93% of patient with diffuse-type giant cell tumors [22]. Pexidartinib (PLX3397), a CSF-1R TKI, was also evaluated in patients with diffuse-type giant cell tumors in a phase I/II dose expansion study and objected responses were demonstrated in 52% of patients [26]. Based on these data, pexidartinib was granted breakthrough status by the Food and Drug Administration (FDA) for treatment of patients with tenosynovial giant cell tumors.

CSF-1R targeted therapies such as GW2580 and pexidartinib have been combined with chemotherapy in preclinical cancer models, including pancreatic and breast cancers. This combination resulted in an increase in cytotoxic immune responses, decreased metastases, and improved survival compared with chemotherapy alone [27–29]. Administration

of pexidartinib has also improved the efficacy of immunotherapies, including adoptive cell-transfer immunotherapy in melanoma and PDL-1 therapy in pancreatic cancer [30,31]. These exciting preclinical data have led to numerous clinical trials, which are currently investigating combinations of CSF-1 directed therapies with hormone therapies, chemotherapies, and immunotherapy (Table 1). Specific examples include NCT02452424, which is evaluating the combination pexidartinib and the immune checkpoint inhibitor, pembrolizumab, in a phase 1/2 trial of melanoma and other solid tumors. Checkpoint inhibitors, which promote antitumor immune responses, have gained considerable momentum recently due to their ability to induce durable responses and improve survival in a range of cancers. This trial and others, such as NCT02777710, evaluating pexidartinib in combination with durvalumab in pancreas and colon cancers, and NCT02323191, evaluating RG7155 in combination with atezolizumab in solid tumors, are seeking to enhance the effectiveness of these checkpoint inhibitors through the combinatorial targeting of macrophages.

In addition to systemic therapies, radiation therapy is also augmented by the administration of CSF-1 directed therapies. In murine cancer models, tumor responses were significantly improved when CSF-1 therapies were combined with radiation therapy [32,33]. In a glioblastoma model, this combination even resulted in improved survival compared with radiation therapy alone. The combination of pexidartinib and radiation therapy is now also being evaluated in NCT01790503, which is combining pexidartinib with standard of care radiation and temozolamide (chemotherapy) in patients with newly diagnosed glioblastoma (Table 1).

Rather than blocking M2 polarization, another approach to reprogramming macrophages is to utilize therapeutic agents that promote M1 polarization. In clinical trials, administration of interferon (IFN)- γ , which has been shown to stimulate transformation from M2 to M1 *in vitro*, resulted in objective responses in minimal residual ovarian cancer [34–36]. Therapies that activate CD40 have up-regulated MHC class II and CD86 on macrophages in a mouse model of pancreatic cancer along with increased systemic levels of M1 cytokines [37]. Furthermore, when a fully human CD40 agonist antibody CP-870,893 was administered in combination with gemcitabine chemotherapy in a phase I trial for patients with pancreatic cancer, partial responses were identified in approximately 20% of patients [38].

In genitourinary cancers, numerous studies have also demonstrated a promising role for targeting macrophage recruitment/reprogramming and this has led to further research at both the preclinical and clinical levels. Combining GW2580, a CSF-1R kinase inhibitor, with hormonal therapy (enzalutamide) resulted in slowed tumor growth and delayed development of therapeutic resistance in a murine prostate cancer model [23]. The combination of hormone therapy with CSF-1 directed therapy is now being investigated at the clinical level in NCT02472275. This

trial combines pexidartinib with antiandrogen therapy in patients with prostate cancer without distant metastases. CSF-1 directed therapies (pexidartinib) have also been shown to improve the efficacy of radiation therapy in prostate cancer. In these mice, the addition of pexidartinib decreased macrophage infiltration following radiotherapy and slowed the growth of tumors [32].

Other investigated approaches that have demonstrated success in genitourinary cancers include the targeting of macrophage recruitment. The CCL2 inhibitor, carlumab, demonstrated efficacy in numerous preclinical studies, leading to the evaluation of this therapy at the clinical level [17–19]. Phase I and II trials have investigated carlumab both as a monotherapy in prostate cancer and in combination with chemotherapy in a range of malignancies [20]. Although these trials failed to demonstrate antitumor responses, CCL2 levels were noted not to be effectively suppressed following carlumab administration. Thus, the limited efficacy of carlumab in the clinical setting may have been primarily driven by suboptimal pharmacokinetics or ineffective dosing.

4. Supporting the antitumor functions of TAMs

Direct phagocytosis of malignant cells is an important mechanism of TAM-mediated tumor eradication [25]. Unfortunately, tumor cells are able to avoid this method of destruction through up-regulation of surface receptors, such as CD47, which inhibit macrophage phagocytic activity [39]. Multiple therapies are under development, which target this defense mechanism by tumors. This includes monoclonal antibodies, which can block surface CD47. At the preclinical level, administration of these agents has resulted in increased macrophage phagocytosis as well as impressive antitumor effect in both solid and hematologic malignancies. Included among these is bladder cancer, which has been found to express high levels of CD47. In *in vitro* models using human bladder cancer cells, blocking CD47 increased the phagocytosis of these cells by macrophages [40]. Clinical investigation of these agents in a range of cancers is now under way (Table 1).

5. Discussion

While tumor-directed therapies such as hormone therapies, chemotherapies, and radiation therapies remain the foundation of cancer treatment, therapies that target the TME are rapidly becoming an invaluable component of oncologic care. This includes genitourinary malignancies, where a number of these therapies are now standard of care and even frontline treatments. Notable examples include checkpoint inhibitors in renal and bladder cancers, antiangiogenic agents in renal cell carcinoma, and sipuleucel-T in prostate cancer. The continued development of novel, TME-directed therapies holds great potential for the further advancement of cancer treatment and TAMs, which are an

integral component of the TME, are a promising target. Numerous therapies targeting TAMs have already demonstrated success at the preclinical level in cancers of renal, bladder, prostate, and testicular origin. The effective clinical translation of these therapies will rely on identifying the contexts of use for which these therapies are able to improve patient outcomes, such as disease morbidity, PFS, and OS, as well as optimizing dosing to achieve maximum biologic effect.

In evaluating for the most effective contexts of use for TAM-directed therapies, key questions that must be addressed by current and future trials include the following: in which cancers will TAM-directed therapies improve patient outcomes, in which stage of disease (i.e., localized, locally advanced, metastatic) are they most effective, where do TAM-directed therapies fit best among the current sequencing of therapies, and should these therapies be administered as a monotherapy or can they be combined with other therapeutic modalities for maximum efficacy? Many early-phase clinical trials are currently investigating TAM-directed therapies through cohort designs, which include patients with a wide range of solid and hematologic malignancies. This strategy provides a convenient and cost-effective way to screen these drugs in an array of cancers and may potentially identify useful roles for TAM-directed therapies. Caution in the interpretation of these results, however, must be exercised as often times in these types of trials, very limited numbers of patients in each cancer type are exposed to a drug that may not be optimally dosed. In this setting, it can be easy to miss a potentially valuable role for these agents if there are not enough patients or if the signal is not strong enough in the small number of patients who were treated. Disease-focused phase II and III trials are therefore also needed, particularly in genitourinary malignancies where immunotherapies have already demonstrated efficacy at the clinical level.

Successful clinical translation of TAM-directed therapies will also include identifying whether each agent should be utilized as a monotherapy or if they can be more effective as part of a combination approach with hormonal therapies, chemotherapies, radiation therapies, or immune therapies. The majority of clinical trials now under way are investigating TAM-directed therapies alone and in combination with other modalities. The key for these trials moving forward will need to be a continued focus on rational therapeutic combinations. Important questions should focus on where mechanisms of resistance for these therapies lie. Many clinical trials are addressing this question from the standpoint of traditional therapies, which are known to have resistance mechanisms that are mediated by TAMs. However, a focus on the resistance pathways for TAM-directed therapies is also important, such as how PDL-1 expression by macrophages may limit immune-mediated depletion of TAMs. Future trials will need to focus on these questions at the preclinical level as well as in clinical trials.

Furthermore, it is critical to address where TAM-directed therapies fit within the sequencing of therapies for each cancer. Rational arguments can certainly be made that TAM-directed therapies have the potential to be effective throughout the stages of disease treatment. However, in genitourinary cancers, there is compelling data that these therapies may be able to play a particularly influential role when administered early on in cancer treatment. In prostate cancer, the development of castrate-resistant disease is a critical phase of disease progression, which signifies the development of more aggressive, treatment resistant disease and poorer prognosis for patients. Escamilla et al. demonstrated that in murine models of prostate cancer, the administration of TAM-directed delayed the development of castrate resistance [23]. Studies such as this one provide strong rationale for administering TAM-directed therapies up front in the treatment of patients with prostate cancer as a strategy to prevent or delay the development of castrate-resistant disease and improve both morbidity and mortality. In other cancers, such as renal cell carcinoma and urothelial bladder cancer, immune checkpoint inhibitors are gaining approval for use in earlier lines of treatment, including in the first line. Zhu et al. demonstrated that TAM-directed therapies can improve the efficacy of these checkpoint inhibitors [31]. Given the potential of checkpoint inhibition to induce complete and durable responses in patients, combining TAM-directed therapies early in the course of the disease may ultimately be able to increase the number of patients who experience these impressive responses and dramatically delay the need for subsequent therapies.

While the inclusion of a broad range of malignancies can be an effective strategy for identifying contexts of use, most of these trials are still relying on traditional dosing strategies such as dosing to the maximum tolerated dose. Unfortunately, in the age of biologic therapies, this type of trial design frequently fails to identify the most effective therapeutic doses. In the case of carlumab, the phase II trial evaluating this medication in men with metastatic prostate cancer failed to demonstrate a significant antitumor effect. The dosing of carlumab in this trial, however, was derived from the highest doses investigated in phase I trials despite the fact that this dose was noted to ineffectively neutralize CCL2 in patients [41]. In the phase I study, this predefined maximal dose of 15 mg/kg every 2 weeks was based on a number of factors, including pharmacokinetics and preclinical data. This dose, however, was ultimately not found to be the maximum tolerated dose based on safety and tolerability analysis in the trial. While the authors did consider evaluating higher doses of carlumab, due to preclinical data in nonhuman primates demonstrating ineffective suppression of CCL2 even at considerably higher doses, this investigation was ultimately not pursued. It is therefore not known whether higher doses of carlumab could be tolerable as well as effectively suppress CCL2 and/or improve patient outcomes. Carlumab still holds considerable promise for the treatment of prostate cancer and future clinical trials

must identify doses that are biologically relevant and able to improve patient outcomes. Biomarkers, in particular those that are predictive or pharmacodynamic, can be an invaluable asset in this pursuit. For carlumab, titrating the dose to demonstrated suppression of systemic CCL2 will be an important next step in the clinical investigation of this drug and one that may ultimately identify a role for this therapy in cancer treatment. The development and validation of biomarker assays will certainly be a critical step in this process and the focus of such research should include the biologic relevance of the assays as well as reliability and reproducibility [42].

Bisphosphonates are another example of the importance of dosing. In clinical trials, these medications have had little to no antitumor activity despite demonstrated efficacy at the preclinical level. However, the dosing for bisphosphonates in these trials was based on approved dosing for prevention of bone resorption and not for antitumor activity. As a consequence, this strategy resulted in far lower and less frequent exposure to active drug than has been shown to be effective at the preclinical level. Thus, the limited antitumor effect in patients may also be attributable to how this medication is administered and not a lack of biologic efficacy. Therefore, tailoring the dosing to the context of use (i.e., treatment of cancer) with higher or more frequent dosing may lead to a more effective role for these agents in cancer treatment.

Continued development of novel TAM-directed therapeutic strategies is critical as well. A number of alternative targets such as IL-4, CXCL12/CXCR4, Toll-like receptor, STAT3, STAT6, antiphosphatidylserine antibodies, and Nrpl [28,43–47] are under investigation. The identification of high-value therapeutic targets with robust likelihood of clinical translation will be supported by improved TME modeling in vitro. The TME is a complex, multicellular environment governed by coordinated signaling pathways. Investigation and analysis of such a multifaceted environment is limited by traditional preclinical platforms, such as mouse models and commercial cell culture devices, which have a restricted ability to model and analyze such a multifaceted environment. A number of new tools such as microfluidic cell culture platforms are under development, which have the capacity for multicellular tumor modeling in both 2D and 3D using primary, patient-derived cells. These platforms, as well as others in development, hold great potential for future drug development, which will ultimately be realized through high-quality translational research.

References

- [1] Schmidt T, Carmeliet P. Blood-vessel formation: bridges that guide and unite. *Nature* 2010;465:697–9.
- [2] Murdoch C, Muthana M, Coffelt SB, Lewis CE. The role of myeloid cells in the promotion of tumour angiogenesis. *Nat Rev Cancer* 2008;8:618–31.

- [3] Bottazzi B, Doni A, Garlanda C, Mantovani A. An integrated view of humoral innate immunity: pentraxins as a paradigm. *Annu Rev Immunol* 2010;28:157–83.
- [4] Biswas SK, Mantovani A. Macrophage plasticity and interaction with lymphocyte subsets: cancer as a paradigm. *Nat Immunol* 2010;11:889–96.
- [5] Chanmee T, Ontong P, Konno K, Itano N. Tumor-associated macrophages as major players in the tumor microenvironment. *Cancers* 2014;6:1670–90.
- [6] Giannoni E, Bianchini F, Masieri L, et al. Reciprocal activation of prostate cancer cells and cancer-associated fibroblasts stimulates epithelial-mesenchymal transition and cancer stemness. *Cancer Res* 2010;70:6945–56.
- [7] Park KY, Li G, Platt MO. Monocyte-derived macrophage assisted breast cancer cell invasion as a personalized, predictive metric to score metastatic risk. *Sci Rep* 2015;5:13855.
- [8] Zhang QW, Liu L, Gong CY, et al. Prognostic significance of tumor-associated macrophages in solid tumor: a meta-analysis of the literature. *PLoS One* 2012;7:e50946.
- [9] Lundholm M, Hägglöf C, Wikberg ML, et al. Secreted factors from colorectal and prostate cancer cells skew the immune response in opposite directions. *Sci Rep* 2015;5:15651.
- [10] Germano G, Frapolli R, Belgiovine C, et al. Role of macrophage targeting in the antitumor activity of trabectedin. *Cancer Cell* 2013;23:249–62.
- [11] Germano G, Frapolli R, Simone M, et al. Antitumor and anti-inflammatory effects of trabectedin on human myxoid liposarcoma cells. *Cancer Res* 2010;70:2235–44.
- [12] Seiler P, Aichele P, Odermatt B, Hengartner H, Zinkernagel RM, Schwendener RA. Crucial role of marginal zone macrophages and marginal zone metallophilic cells in the clearance of lymphocytic choriomeningitis virus infection. *Eur J Immunol* 1997;27:2626–33.
- [13] Zeisberger SM, Odermatt B, Marty C, Zehnder-Fjällman AH, Ballmer-Hofer K, Schwendener RA. Clodronate-liposome-mediated depletion of tumour-associated macrophages: a new and highly effective antiangiogenic therapy approach. *Br J Cancer* 2006;95:272–81.
- [14] Meng Y, Beckett MA, Liang H, et al. Blockade of tumor necrosis factor alpha signaling in tumor-associated macrophages as a radiosensitizing strategy. *Cancer Res* 2010;70:1534–43.
- [15] Smahel M, Duskova M, Polakova I, Musil J. Enhancement of DNA vaccine potency against legumain. *J Immunother* 2014;37:293–303.
- [16] Pulaski HL, Spahlinger G, Silva IA, et al. Identifying alemtuzumab as an anti-myeloid cell antiangiogenic therapy for the treatment of ovarian cancer. *J Transl Med* 2009;7:49.
- [17] Mizutani K, Sud S, McGregor NA, et al. The chemokine CCL2 increases prostate tumor growth and bone metastasis through macrophage and osteoclast recruitment. *Neoplasia* 2009;11:1235–42.
- [18] Loberg RD, Ying C, Craig M, Yan L, Snyder LA, Pienta KJ. CCL2 as an important mediator of prostate cancer growth in vivo through the regulation of macrophage infiltration. *Neoplasia* 2007;9:556–62.
- [19] Rozel S, Galban CJ, Nicolay K, et al. Synergy between anti-CCL2 and docetaxel as determined by DW-MRI in a metastatic bone cancer model. *J Cell Biochem* 2009;107:58–64.
- [20] Moisan F, Francisco EB, Brozovic A, et al. Enhancement of paclitaxel and carboplatin therapies by CCL2 blockade in ovarian cancers. *Mol Oncol* 2014;8:1231–9.
- [21] Nywening TM, Wang-Gillam A, Sanford DE, et al. Targeting tumour-associated macrophages with CCR2 inhibition in combination with FOLFIRINOX in patients with borderline resectable and locally advanced pancreatic cancer: a single-centre, open-label, dose-finding, non-randomised, phase 1b trial. *Lancet Oncol* 2016;17:651–62.
- [22] Ries CH, Cannarile MA, Hoves S, et al. Targeting tumor-associated macrophages with anti-CSF-1R antibody reveals a strategy for cancer therapy. *Cancer Cell* 2014;25:846–59.
- [23] Escamilla J, Schokrpur S, Liu C, et al. CSF1 receptor targeting in prostate cancer reverses macrophage-mediated resistance to androgen blockade therapy. *Cancer Res* 2015;75:950–62.
- [24] Pyonteck SM, Akkari L, Schuhmacher AJ, et al. CSF-1R inhibition alters macrophage polarization and blocks glioma progression. *Nat Med* 2013;19:1264–72.
- [25] Mantovani A, Marchesi F, Malesci A, Laghi L, Allavena P. Tumour-associated macrophages as treatment targets in oncology. *Nat Rev Clin Oncol* 2017;14:399–416.
- [26] Tap WD, Wainberg ZA, Anthony SP, et al. Structure-guided blockade of CSF1R kinase in tenosynovial giant-cell tumor. *N Engl J Med* 2015;373:428–37.
- [27] Weizman N, Krelin Y, Shabtay-Orbach A, et al. Macrophages mediate gemcitabine resistance of pancreatic adenocarcinoma by upregulating cytidine deaminase. *Oncogene* 2014;33:3812–9.
- [28] DeNardo DG, Barreto JB, Andreu P, et al. CD4(+) T cells regulate pulmonary metastasis of mammary carcinomas by enhancing protumor properties of macrophages. *Cancer Cell* 2009;16:91–102.
- [29] DeNardo DG, Brennan DJ, Rexhepaj E, et al. Leukocyte complexity predicts breast cancer survival and functionally regulates response to chemotherapy. *Cancer Discov* 2011;1:54–67.
- [30] Mok S, Koya RC, Tsui C, et al. Inhibition of CSF-1 receptor improves the antitumor efficacy of adoptive cell transfer immunotherapy. *Cancer Res* 2014;74:153–61.
- [31] Zhu Y, Knolhoff BL, Meyer MA, et al. CSF1/CSF1R blockade reprograms tumor-infiltrating macrophages and improves response to T-cell checkpoint immunotherapy in pancreatic cancer models. *Cancer Res* 2014;74:5057–69.
- [32] Xu J, Escamilla J, Mok S, et al. CSF1R signaling blockade stanches tumor-infiltrating myeloid cells and improves the efficacy of radiotherapy in prostate cancer. *Cancer Res* 2013;73:2782–94.
- [33] Stafford JH, Hirai T, Deng L, et al. Colony stimulating factor 1 receptor inhibition delays recurrence of glioblastoma after radiation by altering myeloid cell recruitment and polarization. *Neuro Oncol* 2016;18:797–806.
- [34] Duluc D, Corvaisier M, Blanchard S, et al. Interferon-gamma reverses the immunosuppressive and protumoral properties and prevents the generation of human tumor-associated macrophages. *Int J Cancer* 2009;125:367–73.
- [35] Colombo N, Peccatori F, Paganin C, et al. Anti-tumor and immunomodulatory activity of intraperitoneal IFN-gamma in ovarian carcinoma patients with minimal residual tumor after chemotherapy. *Int J Cancer* 1992;51:42–6.
- [36] Pujade-Lauraine E, Guastalla JP, Colombo N, et al. Intraperitoneal recombinant interferon gamma in ovarian cancer patients with residual disease at second-look laparotomy. *J Clin Oncol* 1996;14:343–50.
- [37] Beatty GL, Chiorean EG, Fishman MP, et al. CD40 agonists alter tumor stroma and show efficacy against pancreatic carcinoma in mice and humans. *Science* 2011;331:1612–6.
- [38] Beatty GL, Torigian DA, Chiorean EG, et al. A phase I study of an agonist CD40 monoclonal antibody (CP-870,893) in combination with gemcitabine in patients with advanced pancreatic ductal adenocarcinoma. *Clin Cancer Res* 2013;19:6286–95.
- [39] Rodriguez PL, Harada T, Christian DA, Pantano DA, Tsai RK, Discher DE. Minimal "self" peptides that inhibit phagocytic clearance and enhance delivery of nanoparticles. *Science* 2013;339:971–5.
- [40] Chan KS, Espinosa I, Chao M, et al. Identification, molecular characterization, clinical prognosis, and therapeutic targeting of human bladder tumor-initiating cells. *Proc Natl Acad Sci USA* 2009;106:14016–21.
- [41] Sandhu SK, Papadopoulos K, Fong PC, et al. A first-in-human, first-in-class, phase I study of carlumab (CNTO 888), a human monoclonal antibody against CC-chemokine ligand 2 in patients with solid tumors. *Cancer Chemother Pharmacol* 2013;71:1041–50.
- [42] King JD, Casavant BP, Lang JM. Rapid translation of circulating tumor cell biomarkers into clinical practice: technology

- development, clinical needs and regulatory requirements. *Lab Chip* 2014;14:24–31.
- [43] Shime H, Matsumoto M, Oshiumi H, et al. Toll-like receptor 3 signaling converts tumor-supporting myeloid cells to tumoricidal effectors. *Proc Natl Acad Sci USA* 2012;109:2066–71.
- [44] Robinson SC, Scott KA, Wilson JL, Thompson RG, Proudfoot AE, Balkwill FR. A chemokine receptor antagonist inhibits experimental breast tumor growth. *Cancer Res* 2003;63:8360–5.
- [45] Yin Y, Huang X, Lynn KD, Thorpe PE. Phosphatidylserine-targeting antibody induces M1 macrophage polarization and promotes myeloid-derived suppressor cell differentiation. *Cancer Immunol Res* 2013;1:256–68.
- [46] Casazza A, Laoui D, Wenes M, et al. Impeding macrophage entry into hypoxic tumor areas by Sema3A/Nrp1 signaling blockade inhibits angiogenesis and restores antitumor immunity. *Cancer Cell* 2013;24:695–709.
- [47] Jia X, Yu F, Wang J, et al. Emodin suppresses pulmonary metastasis of breast cancer accompanied with decreased macrophage recruitment and M2 polarization in the lungs. *Breast Cancer Res Treat* 2014;148:291–302.