



# Developmental Outcomes of Extremely Preterm Infants with a Need for Child Protective Services Supervision

Elisabeth C. McGowan, MD<sup>1</sup>, Abbot. R. Luptook, MD<sup>1</sup>, Jean Lowe, PhD<sup>2</sup>, Myriam Peralta-Carcelen, MD, MPD<sup>3</sup>, Dhuly Chowdhury, MS<sup>4</sup>, Rosemary D. Higgins, MD<sup>5</sup>, Susan R. Hintz, MD, MS Epi<sup>6</sup>, Betty R. Vohr, MD<sup>1</sup>, for the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Neonatal Research Network\*

**Objective** To evaluate neurodevelopmental outcomes of preterm infants with need for Child Protective Services (CPS) supervision at hospital discharge compared with those discharged without CPS supervision.

**Study design** For infants born at <27 weeks of gestation between 2006 and 2013, prospectively collected maternal and neonatal characteristics and 18- to 26-month corrected age follow-up data were analyzed. Bayley-III cognitive and language scores of infants with discharge CPS supervision were compared with infants without CPS supervision using regression analysis while adjusting for potentially confounding variables, including entering CPS after discharge from the hospital.

**Results** Of the 4517 preterm infants discharged between 2006 and 2013, 255 (5.6%) were discharged with a need for CPS supervision. Mothers of infants with CPS supervision were significantly more likely to be younger, single, and gravida  $\geq 3$ ; to have less than a high school education; and to have a singleton pregnancy and less likely to have received prenatal care or antenatal steroids. Despite similar birth weight and medical morbidities, the CPS group had longer hospital stays compared with the non-CPS group. In adjusted analysis, cognitive scores were points lower ( $B = -1.94$ ; 95% CI,  $-3.88$  to  $-0.08$ ;  $P = .04$ ) in the CPS at discharge group compared with the non-CPS group. In children who entered CPS supervision after hospital discharge (an additional 106 infants), cognitive scores were 4 points lower ( $\beta = -4.76$ ; 95% CI,  $-7.47$  to  $-2.05$ ;  $P < .001$ ) and language scores were 5 points lower ( $\beta = -4.93$ ; 95% CI,  $-8.00$  to  $-1.86$ ;  $P = .002$ ).

**Conclusion** Extremely preterm infants discharged from the hospital with CPS supervision or entering CPS post-discharge are at increased risk for cognitive delay at 2 years of age. Opportunities exist to intervene and potentially improve outcomes in this vulnerable group of children. (*J Pediatr* 2019;215:41-9).

Child Protective Services (CPS) investigations increased by approximately 12% from 2013 to 2017.<sup>1</sup> Although rates vary, developmental delays have been reported in up to 40% of infants and 90% of toddlers involved in CPS.<sup>2</sup> Significant cognitive delays have been reported in nearly one-third of toddlers in the child welfare system.<sup>3</sup> Increased behavioral problems, including externalizing and dysregulation, have also been reported.<sup>4</sup>

Preterm infants are at risk for poor developmental and behavioral outcomes,<sup>5-9</sup> but most CPS cohorts do not distinguish between term and preterm status. A 1995 cross-sectional analysis of a California cohort reported that 36% of children entering foster care before 1 year of age were born preterm.<sup>10</sup> In a *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) Neonatal Research Network (NRN) study of extremely preterm (EPT) infants born to adolescent mothers, CPS supervision was a predictor of low cognitive and language scores, although the subset of CPS infants was small.<sup>11</sup>

Given this scarcity of data, the additional impact of the need for CPS involvement in preterm infants already at high risk for developmental delay is unknown. The main objective of this study was to evaluate differences in

|            |   |       |   |
|------------|---|-------|---|
| Bayley III | Bayley Scales of Infant and Toddler Development III | EPT   | Extremely preterm   |
| BITSEA     | Brief Infant Social and Emotional Assessment        | IVH   | Intraventricular hemorrhage   |
| BPD        | Bronchopulmonary dysplasia                          | NEC   | Necrotizing enterocolitis   |
| CP         | Cerebral palsy                                      | NICHD | <i>Eunice Kennedy Shriver</i> National Institute of Health Child Health and Human Development |
| CPS        | Child Protective Services                           | NICU  | Neonatal intensive care unit  |
| cPVL       | Cystic periventricular leukomalacia                 | NRN   | Neonatal Research Network   |
|            |   | ROP   | Retinopathy of prematurity  |

From the <sup>1</sup>Division of Neonatal Medicine, Department of Pediatrics, Brown University, Providence, RI; <sup>2</sup>Department of Pediatrics, University of New Mexico, Albuquerque, NM; <sup>3</sup>Department of Pediatrics, University of Alabama at Birmingham, Birmingham, AL; <sup>4</sup>Social, Statistical, and Environmental Sciences Unit, RTI International, Rockville, MD; <sup>5</sup>*Eunice Kennedy Shriver* National Institute of Child Health and Human Development, Rockville, MD; and <sup>6</sup>Division of Neonatal and Developmental Medicine, School of Medicine, Stanford University and Lucile Packard Children's Hospital, Palo Alto, CA

\*A list of additional members of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Neonatal Research Network is available at [www.jpeds.com](http://www.jpeds.com) (Appendix).

Portions of this study were presented at the American Academy of Pediatrics National Conference, September 17, 2017, Chicago, IL.

0022-3476/\$ - see front matter. © 2019 Published by Elsevier Inc. <https://doi.org/10.1016/j.jpeds.2019.07.063>

cognitive, language, and behavior outcomes between children discharged from the hospital with the need for CPS supervision and those without the need for CPS supervision. Our primary hypothesis was that EPT infants with CPS supervision at discharge would have lower 2-year Bayley Scales of Infant and Toddler Development III (Bayley-III) cognitive and language scores compared with EPT infants without CPS supervision at discharge. A secondary hypothesis was that EPT infants with CPS supervision would have increased behavior/social problems, decreased competence on the Brief Infant Social and Emotional Assessment (BITSEA), and decreased growth outcomes at age 2 years. A post hoc analysis explored further relationships between changes in CPS status from discharge to 2 years of age and developmental delay measured by Bayley-III scores.

## Methods

This was a retrospective cohort analysis of prospectively collected data (in the NICHD NRN generic database and follow-up database) by 21 NRN centers. Local Institutional Review Boards approved data collection. Infants born at <27 weeks of gestation between January 2006 and December 2013 at a participating NRN hospital and with a follow-up assessment completed at 2 years (defined as 18-26 months corrected age) were included. Infants with a major congenital anomaly or a syndrome associated with adverse developmental outcomes were excluded.

Two cohorts were evaluated: EPT infants who were discharged from the hospital with CPS supervision and those who were discharged without CPS supervision. CPS status was defined as any infant supervised as a ward of the court, in temporary custody by a state agency/court system and in the home of a relative (or other person), or mother and child together but supervised by state agency/court system. CPS status was recorded at hospital discharge and at a 2-year follow-up assessment.

Research personnel prospectively collected maternal data, including age, gravida status, race/ethnicity, education, prenatal care, and antenatal steroid use. Neonatal data included gestational age, growth measures (including *z*-scores),<sup>12</sup> sex, and morbidities, including but not limited to small for gestational age,<sup>13</sup> intraventricular hemorrhage (IVH) grade III-IV, cystic periventricular leukomalacia (cPVL), necrotizing enterocolitis (NEC), early-onset and late-onset sepsis, bronchopulmonary dysplasia (BPD), and severe retinopathy of prematurity (ROP).

At discharge and at 2-year follow-up assessment, living arrangement data collected included primary caretaker, household composition, education level, primary language, insurance status, and the number of places the child had lived between discharge and the 2-year assessment.

Certified examiners performed neurodevelopmental assessments at corrected age 18-22 months (for infants born between January 1, 2006 and June 30, 2012) or corrected

age 22-26 months (for infants born between July 1, 2012, and December 31, 2013) secondary to a protocol change for assessment age.<sup>14</sup> The Bayley-III includes cognitive and language composite scores (mean,  $100 \pm 15$ ); a score <85 represents 1 SD below the mean.<sup>15</sup> Bayley-III motor scores were not collected by the NICHD NRN until January 1, 2010, and are not included in the present analysis. The BITSEA, which captures socioemotional and behavioral information, was completed by the primary caregiver (for infants born between January 1, 2006, and May 30, 2012). This standardized, normative value-referenced behavioral screening instrument includes 2 scales: a Problem Scale (31 items), with higher scores reflecting increased behavior/social problems, and a Competence Scale (11 items), with lower scores reflecting less competence.<sup>16</sup>

The primary study outcomes were Bayley-III cognitive and language composite scores. Secondary outcomes were BITSEA Problem Scale score  $\geq 75$ th percentile; BITSEA Competence score  $\leq 15$ th percentile; moderate to severe cerebral palsy (CP), as demonstrated by an abnormal neurologic examination and Palisano Gross Motor Function Classification System level  $\geq 2$ <sup>17</sup>; and growth measures (weight, length, and head circumference).

## Statistical Analyses

Maternal, infant, and social characteristics and 2-year outcomes were compared between infants with CPS supervision (CPS group) and those without CPS supervision (non-CPS group) documented at initial hospital discharge. The  $\chi^2$  test was used for categorical variables and the *t* test or Wilcoxon rank-sum test was used for continuous variables, as appropriate. Bayley-III composite scores were calculated for corrected age. Unadjusted group comparisons were performed for the primary and secondary outcomes. Two separate generalized linear mixed models were used to evaluate the associations between CPS at discharge and cognitive and language scores. The unadjusted comparisons of outcomes for each factor accounted only for NRN center as a random effect (reflecting differences in population, clinical, and CPS practices).

Adjusted analysis controlled for center and other potentially confounding variables of the mother (ie, prenatal care, antenatal steroids, age, and nonwhite race), the infant (ie, gestational age, sex, BPD, NEC, early or late sepsis, IVH or cPVL, and severe ROP), the postdischarge home environment (ie, 2-year caregiver education of less than high school and non-English speaking, infant living in more than 1 home), and age at testing. Because some of the children entered CPS between discharge and 2 years of age, the final models included "CPS after discharge," because this potential confounder remained a significant factor in the adjusted model. The available sample provided at least 80% power for detecting a small difference (Cohen *d* = 0.20) between the CPS and non-CPS groups at discharge at a *P* value of .05 for the primary outcome. All *P* values were 2-tailed, with a value  $\leq .05$  considered to indicate statistical

significance. Analyses were conducted with SAS 9.4 (SAS Institute, Cary, North Carolina).

To explore the movement of children entering and exiting CPS over the 2-year period in more detail, an unadjusted post hoc analysis was performed for the primary outcomes (Bayley-III cognitive and language scores) and BITSEA scores for 4 subgroups of CPS exposure: (1) CPS at discharge only (child under CPS supervision at hospital discharge but no longer at 2 years), (2) CPS at 2 years only (child entered CPS supervision any time after initial hospital discharge), (3) child in CPS supervision at both discharge and until 2 years, and (4) child never in CPS. The overall *P* values for comparing all 4 groups were calculated, and pairwise comparisons between groups were performed only for outcomes with a significant overall *P* value.

## Results

The study population included a total of 8995 EPT infants born at <27 weeks of gestation between January 1, 2006, and December 31, 2013, and admitted to a participating NRN center at ≤72 hours of age. Study cohort derivation is depicted in the [Figure](#). There were no significant differences in rates of death, incomplete follow-up, or loss to follow-up between the CPS and non-CPS groups. The study cohort comprised 4517 EPT infants with available 2-year Bayley-III cognitive composite scores, of whom 255 (5.6%) were discharged from the neonatal intensive care unit (NICU) under CPS supervision.

[Table I](#) presents maternal and neonatal characteristics in the CPS and non-CPS groups at hospital discharge. Compared with mothers of infants in the non-CPS group, mothers of infants in the CPS group were more likely to be younger, single, of higher gravida, and less educated and to have a singleton pregnancy and less likely to have received prenatal care or antenatal steroids. Despite similar rates of sepsis, BPD, NEC, IVH, cPVL, and severe ROP in the 2 groups, the CPS group had longer NICU stays than the non-CPS group. At 36 weeks postmenstrual age, CPS infants were slightly heavier and had a larger head circumference.

[Table II](#) (available at [www.jpeds.com](http://www.jpeds.com)) presents living arrangements at discharge and at 2 years. More than one-half (58%) of infants with CPS supervision were discharged to family, 36% were placed with a foster parent, and 2% were placed with an adoptive parent. At 2 years, the majority (62%) of infants with CPS supervision continued to have a family member as the primary caretaker, and the primary caretaker was more likely to be an English speaker compared with the non-CPS group. At both hospital discharge and 2 years, infants with CPS supervision were more likely to reside in larger households, to have lived in more than 1 home, and to have public insurance. Early intervention participation was similar in the 2 groups.

In unadjusted comparisons at 2 years ([Table III](#)), the CPS group had lower Bayley-III cognitive composite scores (mean,  $85.3 \pm 15.8$  vs  $88.6 \pm 15.3$ ;  $P < .001$ ) and a greater

proportion of scores <85 (39% vs 30%;  $P = .003$ ). There were no significant between-group differences in Bayley-III language scores, BITSEA scores, or weight. The CPS group had a smaller head circumference, shorter length, and greater likelihood of moderate to severe CP and were more likely to have been rehospitalized.


Linear regression analysis adjusted for biological and social risks ([Table IV](#)) showed that CPS at discharge was independently associated with an almost 2-point lower Bayley-III cognitive score ( $-1.94$ ; 95% CI,  $-3.80$  to  $-0.08$ ;  $P = .04$ ), and that entering CPS after discharge was associated with almost 5-point lower cognitive and language scores ( $-4.76$ ; 95% CI,  $-7.47$  to  $-2.05$ ;  $P < .001$  and  $-4.93$ ; 95% CI,  $-8.0$  to  $-1.86$ ;  $P = .002$ , respectively). Additional characteristics associated with lower Bayley-III scores included single mother, nonwhite race, and less than high school education of the 2-year primary caretaker. Male sex and major neonatal morbidities were identified as independent risk factors for lower cognitive and language scores.

Exploratory comparisons among CPS subgroups and infants never exposed to CPS are presented in [Table V](#). Cognitive and language composite scores were lowest for infants who at 2 years either remained in CPS since discharge or entered CPS sometime after discharge. Compared with infants under CPS supervision at discharge only, the subgroup of infants under CPS supervision at both discharge and 2 years was more likely to have cognitive scores <85 (51% vs 32%;  $P = .004$ ) and language scores <70 (29% vs 18%;  $P = .04$ ). Infants who had never been under CPS supervision had the highest mean cognitive and language scores. In addition, mean BITSEA competency scores were lower for infants with CPS supervision at discharge and 2 years compared with those who had never been under CPS supervision ( $15.5 \pm 4.6$  vs  $16.6 \pm 3.6$ ;  $P = .017$ ), although mean behavior problem scores were similar in the 2 groups (data not shown).

## Discussion

This study investigated differences in cognition and language and secondary outcomes of behavior and growth at 2 years of age among EPT infants with and without CPS involvement. Our findings support our primary hypothesis that infants discharged from the hospital with CPS supervision are more likely to have lower 2-year cognitive scores compared with infants discharged without CPS supervision. Language scores were similar in the 2 primary study groups, however.

The CPS group was characterized by a greater number of maternal social and environmental risk factors, including multigravida, single mother, public insurance, and less education, prenatal care, and antenatal steroids. It did not appear that more medically complex infants were placed into CPS care at NICU discharge, given that there was no difference in neonatal morbidities between the 2 groups. For those infants leaving the hospital under CPS supervision, the



**Figure.** Study flow chart. The cohort of infants prospectively planned for analyses of primary and secondary outcomes. The cohort of infants for which post hoc analyses were performed.

majority were living with biological parent(s) or family members at discharge and at 2 years of age. This preference toward kinship care aligns with the goals of stability and family preservation advocated by the Adoptions and Safe Family Act.<sup>18</sup>

Early cognitive assessments of children with CPS involvement, consisting of data analyzed from the 1999-2000 National Survey of Child and Adolescent Well-Being cohort, were published by Stahmer et al.<sup>19</sup> These authors reported

that 30% of infants aged 0-2 years had cognitive scores >2 SD below the mean. In a California child welfare cohort, 35% of toddlers scored >1 SD below the mean on the Bayley-II Mental Developmental Index, and 30% scored >2 SD below the mean.<sup>3</sup> Although not specified, it is likely that both cohorts included children born prematurely. In our EPT cohort, 39% and 13% had Bayley-III cognitive scores >1 SD and >2 SD below the mean, respectively.

**Table I. Maternal and neonatal characteristics**

| Characteristics                                    | CPS group      | Non-CPS group     | P value |
|--|----------------|-------------------|---------|
| Maternal   | N = 244 (5.9%) | N = 3877 (94.1 %) |         |
| Age, y, mean $\pm$ SD                              | 26.0 $\pm$ 6.7 | 27.5 $\pm$ 6.3    | <.001   |
| Gravidity, n (%)                                   |                |                   |         |
| 1  | 44 (18.0)      | 1237 (31.9)       | <.001   |
| 2  | 49 (20.1)      | 950 (24.5)        | .12     |
| 3+   | 151 (61.9)     | 1689 (43.6)       | <.001   |
| Singleton pregnancy, n (%)                         | 217 (88.9)     | 3219 (83.0)       | .02     |
| Prenatal care ( $\geq$ 1 visit), n (%)             | 186 (76.2)     | 3730 (96.2)       | <.001   |
| Antenatal steroids, n (%)                          | 190 (77.9)     | 3447 (88.9)       | <.001   |
| Highest grade completed, n (%)                     |                |                   |         |
| Less than high school graduate                     | 81 (33.2)      | 674 (17.4)        | <.001   |
| High school graduate                               | 44 (18.0)      | 868 (22.4)        | .11     |
| Some college                                       | 23 (9.4)       | 718 (18.5)        | <.001   |
| College graduate                                   | 10 (4.1)       | 667 (17.2)        | <.001   |
| Unknown  | 86 (35.2)      | 950 (24.5)        | <.001   |
| Race/ethnicity, n (%)                              |                |                   |         |
| White  | 97 (39.8)      | 1454 (37.5)       | .48     |
| African American                                   | 105 (43.0)     | 1597 (41.2)       | .57     |
| Hispanic/Latino                                    | 31 (12.7)      | 643 (16.6)        | .11     |
| Other  | 9 (3.6)        | 173 (4.4)         | .57     |
| Unknown  | 2 (0.8)        | 10 (0.3)          | .11     |
| Single, n (%)                                      | 203 (83.2)     | 2130 (54.9)       | <.001   |
| Neonatal   | N = 255 (5.6%) | N = 4262 (94.4%)  |         |
| Male sex, n (%)                                    | 125 (49.0)     | 2115 (49.6)       | .84     |
| Birth weight, g, mean $\pm$ SD                     | 774 $\pm$ 157  | 756 $\pm$ 154     | .07     |
| Birth length, cm, mean $\pm$ SD                    | 32.5 $\pm$ 2.5 | 32.4 $\pm$ 2.5    | .71     |
| Birth head circumference, cm, mean $\pm$ SD        | 22.9 $\pm$ 1.7 | 22.8 $\pm$ 1.6    | .30     |
| Gestational age, wk, mean $\pm$ SD                 | 25 $\pm$ 1     | 25 $\pm$ 1        | .12     |
| Postnatal steroids, n (%)                          | 49 (19.2)      | 808 (18.8)        | .96     |
| Small for gestational age, n (%) <sup>*</sup>      | 12 (4.7)       | 219 (5.1)         | .76     |
| Early-onset sepsis, n (%) <sup>†</sup>             | 9 (3.5)        | 88 (2.1)          | .12     |
| Late-onset sepsis, n (%) <sup>‡</sup>              | 75 (29.4)      | 1430 (33.6)       | .17     |
| NEC, n (%) <sup>§</sup>                            | 26 (10.2)      | 403 (9.5)         | .69     |
| IVH <sup>¶</sup>                                   | 43 (16.9)      | 655 (15.4)        | .52     |
| cPVL   | 16 (6.3)       | 235 (5.5)         | .59     |
| Severe ROP <sup>**</sup>                           | 64 (25.1)      | 998 (23.4)        | .54     |
| BPD <sup>††</sup>                                  | 142 (55.7)     | 2449 (57.5)       | .59     |
| Days of ventilation, mean $\pm$ SD                 | 29 $\pm$ 29    | 30 $\pm$ 26       | .71     |
| Days in hospital, mean $\pm$ SD                    | 127 $\pm$ 61   | 116 $\pm$ 45      | <.001   |
| Weight at 36 weeks PMA, g, mean $\pm$ SD           | 2116 $\pm$ 389 | 2064 $\pm$ 399    | .06     |
| z-score, mean $\pm$ SD                             | -1.2 $\pm$ 0.8 | -1.3 $\pm$ 0.8    | .04     |
| Length at 36 weeks PMA, cm, mean $\pm$ SD          | 42.3 $\pm$ 3.0 | 42.0 $\pm$ 2.9    | .22     |
| z-score, mean $\pm$ SD                             | -1.8 $\pm$ 1.0 | -1.9 $\pm$ 0.9    | .16     |
| Head circumference 36 weeks PMA, cm, mean $\pm$ SD | 30.9 $\pm$ 2.0 | 30.5 $\pm$ 1.8    | .01     |
| z-score, mean $\pm$ SD                             | -1.1 $\pm$ 1.1 | -1.3 $\pm$ 1.0    | .01     |

PMA, postmenstrual age.

<sup>\*</sup>Small for gestational age defined as <10% at birth as determined by the Alexander growth curve.<sup>13</sup>

<sup>†</sup>Early-onset sepsis defined as blood culture positive within first 3 days of life.

<sup>‡</sup>Late-onset sepsis defined as blood culture positive after day of life 3.

<sup>§</sup>NEC greater than or equal to Bell classification IIA.

<sup>¶</sup>IVH defined as grade III-IV.

<sup>\*\*</sup>Severe ROP defined as greater than or equal to stage 3 and/or intervention.

<sup>††</sup>BPD defined as oxygen requirement at 36 weeks PMA.

Mean scores for Bayley-III have been reported to be higher than expected, and thus comparing Bayley-III and Bayley-II results is challenging; nonetheless, this suggests that our results may be underestimates of delays.<sup>20,21</sup>

Mean cognitive scores were 3 points lower for infants in the CPS group compared with the non-CPS group. Although many of the social risk factors necessitating CPS supervision are also risk factors for developmental delay, cognitive findings remained significant after statistical adjustment. Regression analysis also revealed potentially greater negative impacts on cognitive and language scores when entering CPS supervision after discharge. This could be an effect of

timing and/or duration of exposure to environments that necessitate supervision. The additional lowering of cognitive and language scores for those infants with CPS involvement, even by 2-5 points, can be thought of as increasing the risk of special education needs for EPT infants, as reflected in their higher rate of cognitive composite scores <85.

There were no between-group differences in mean Bayley-III language scores; however, 22% of the CPS group had language scores <70, in contrast to data reported by Stahmer et al, in which approximately 10% of children with CPS supervision at 0-2 years of age had language assessment scores >2 SD below the mean.<sup>19</sup> It is unclear whether Stahmer et al included

**Table III. Two-year outcomes of EPT infants with or without CPS supervision at NICU discharge**

| Outcomes                                 | CPS (N = 255; 5.6%) | Non-CPS (N = 4262; 94.4%) | P value |
|--|---------------------|---------------------------|---------|
| Age at follow up, mo, mean ± SD          | 21.3 ± 3.1          | 21.3 ± 3.1                | .79     |
| Bayley-III                               |                     |                           |         |
| Cognitive composite score, mean ± SD     | 85.3 ± 15.8         | 88.6 ± 15.3               | <.001   |
| Language composite score, mean ± SD      | 82.4 ± 16.5         | 83.9 ± 17.1               | .19     |
| Expressive language score, mean ± SD     | 7.2 ± 2.8           | 7.5 ± 3.0                 | .13     |
| Receptive language score, mean ± SD      | 7.2 ± 2.7           | 7.3 ± 2.9                 | .57     |
| Cognitive composite <85, n/N (%)         | 97/252 (38.5)       | 1250/4224 (30.5)          | .003    |
| Language composite <85, n/N (%)          | 134/246 (54.5)      | 2070/4153 (50.0)          | .16     |
| Cognitive composite <70, n/N (%)         | 32/252 (12.7)       | 425/4224 (10.1)           | .18     |
| Language composite <70, n/N (%)          | 54/246 (22.0)       | 784/4153 (18.9)           | .23     |
| BITSEA                                   |                     |                           |         |
| Behavior problems total score, mean ± SD | 12.7 ± 6.9          | 12.3 ± 7.2                | .38     |
| Competencies total score, mean ± SD      | 16.2 ± 4.1          | 16.6 ± 3.6                | .13     |
| Behavior problems ≥75%, n/N (%)          | 72/193 (37.3)       | 1076/3176 (33.9)          | .33     |
| Competency problems <15% n/N (%)         | 59/193 (30.6)       | 826/3113 (26.5)           | .22     |
| CP moderate-severe, n/N (%)              | 26/255 (10.2)       | 283/4259 (6.6)            | .03     |
| Weight, kg, mean ± SD                    | 10.8 ± 1.4          | 10.9 ± 1.6                | .24     |
| Length, cm, mean ± SD                    | 81.6 ± 5.0          | 82.2 ± 4.9                | .04     |
| Head circumference, cm, mean ± SD        | 46.3 ± 2.3          | 46.9 ± 2.2                | <.001   |
| Rehospitalization                        |                     |                           |         |
| Hospitalized since discharge, n/N (%)    | 144/253 (56.9)      | 2052/4239 (48.4)          | .007    |
| Number of times, mean ± SD               | 2.5 ± 2.3           | 2.2 ± 2.0                 | .05     |

preterm infants in their analysis. Many infants in our cohort were tested at 18-22 months of corrected age, which may be too early to detect language delays in EPT children. The time spent in disruptive environments, particularly during critical periods of language development, may adversely affect skills. Because language delays may increase over time in children under CPS supervision,<sup>19</sup> close surveillance is warranted.

BITSEA scores were similar in our CPS and non-CPS groups; approximately one-third of all children had high problem or low competency scores, similar to rates of behavioral problems reported in other preterm cohorts.<sup>9,22</sup> This could be due to parental underreporting of problems due to perception biases or socially desirable responding,<sup>23,24</sup> or because prematurity is a stronger determinant of behavioral problems than environments prompting CPS oversight. However, behavioral difficulties among children in foster care are well documented, with a 3- to 4-fold increased odds of anxiety, depression, and attention/hyperactivity disorders compared with the general population.<sup>25,26</sup> Because preterm infants are at increased risk for similar behavioral problems over time, longitudinal mental health evaluation of the EPT children in CPS is warranted.<sup>27,28</sup>

Poor growth is common among EPT infants.<sup>29</sup> The CPS group had a larger head circumference at 36 weeks postmenstrual age, but at 2 years demonstrated a greater lag in head circumference, and length, compared with the non-CPS group. Whether this was due to environmental influence is unclear, given that multiple variables, including poverty, stress, inflammation, and nutritional deficiencies, may contribute to poor growth.<sup>30,31</sup> The CPS group had a higher percentage of infants with moderate to severe CP, which has been associated with delayed head growth,<sup>32-34</sup> which may partially explain some of the growth findings. Although there

were no differences in IVH or cPVL between the 2 groups, other markers of brain injury, such as cerebellar and white matter injury,<sup>35,36</sup> which were not captured in the NRN database, may have been higher in the CPS group, leading to higher rates of motor impairment. In addition, the infants with CPS supervision were less likely to receive antenatal steroids, which has a protective effect against CP.<sup>37</sup>

Health care utilization for medically and socially vulnerable populations is an important consideration. For this EPT cohort, infants discharged under CPS supervision were in the hospital an average of 11 days longer than infants not under CPS supervision. Because neonatal morbidities were similar in the 2 groups, prolonged hospitalization may have been due to social factors, including decreased parent/caregiver visitation, or the complexities of caregiver identification/placement. Rehospitalization rates were also higher for infants under CPS supervision. Although public insurance has been associated with an increased rate of rehospitalization,<sup>38</sup> CPS-supervised families may encounter challenges when navigating the healthcare environment, resulting in lapses in preventative care or frequent provider changes that lead to escalation of health care needs or services. It is encouraging that early intervention, which provides support to both infant and caregiver, was comparable in the 2 groups.

In post hoc analysis, the proportions of children scoring >1 SD and >2 SD below the mean for cognitive and language scores were highest for those in under CPS supervision from discharge to 2 years. One-third of the EPT infants discharged under CPS supervision remained under supervision at 2 years, and in this group, one-half had a mean cognitive score <85. The effect of duration of exposure to environmental stressors on neurodevelopment remains

**Table IV. Unadjusted and adjusted linear regression of Bayley-III cognitive and language composite scores**

| Variables                                       | Bayley-III cognitive composite  |         |                                 |         | Bayley-III language composite   |         |                                 |         |
|---|---------------------------------|---------|---------------------------------|---------|---------------------------------|---------|---------------------------------|---------|
|   | Unadjusted                      |         | Adjusted                        |         | Unadjusted                      |         | Adjusted                        |         |
|   | $\beta$ coefficient<br>(95% CI) | P value |
| Age at testing                                  | -0.37 (-0.52 to -0.23)          | <.001   | -0.39 (-0.52 to -0.25)          | <.001   | -0.16 (-0.32 to 0.01)           | .06     | -0.16 (-0.32 to -0.01)          | .05     |
| CPS at discharge                                | -2.97 (-4.90 to -1.04)          | .003    | -1.94 (-3.80 to -0.08)          | .04     | -1.12 (-3.29 to 1.06)           | .31     | -0.90 (-3.02 to 1.22)           | .41     |
| CPS entered after discharge                     | -5.02 (-7.97 to -2.07)          | <.001   | -4.76 (-7.47 to -2.05)          | <.001   | -4.71 (-8.01 to -1.41)          | .005    | -4.93 (-8.0 to -1.86)           | .002    |
| No prenatal care                                | -2.62 (-4.73 to -0.51)          | .01     | -1.67 (-3.71 to 0.37)           | .11     | 0.05 (-2.33 to 2.44)            | .96     | 1.05 (-1.28 to 3.39)            | .38     |
| No antenatal steroids                           | -2.1 (-3.53 to -0.68)           | .003    | 0.11 (-1.27 to 1.49)            | .88     | -2.32 (-3.92 to -0.73)          | .004    | -0.05 (-1.62 to 1.52)           | .95     |
| Single mother                                   | -2.58 (-3.48 to -1.68)          | <.001   | -1.19 (-2.18 to -0.20)          | .02     | -3.01 (-4.02 to -2.0)           | <.001   | -1.72 (-2.85 to -0.60)          | .003    |
| Maternal age                                    | 0.37 (0.02 to 0.72)             | .04     | -0.07 (-0.43 to 0.30)           | .72     | 0.38 (-0.01 to 0.78)            | .06     | -0.11 (-0.52 to 0.31)           | .62     |
| Nonwhite  | -3.69 (-4.61 to -2.77)          | <.001   | -3.17 (-4.17 to -2.18)          | <.001   | -5.47 (-6.50 to -4.40)          | <.001   | -4.44 (-5.57 to -3.31)          | <.001   |
| Less than high school education (2-y caregiver) | -3.31 (-4.48 to -2.14)          | <.001   | -1.91 (-3.11 to -0.71)          | .002    | -4.42 (-5.73 to -3.10)          | <.001   | -2.72 (-4.09 to -1.35)          | <.001   |
| Lived in more than 1 place since discharge      | -0.73 (-1.64 to 0.17)           | .11     | -0.33 (-1.22 to 0.55)           | .46     | -0.36 (-1.38 to 0.66)           | .48     | 0.0 (-1.00 to 1.01)             | .99     |
| Non-English speaker (2-y caregiver)             | -1.96 (-3.27 to -0.66)          | .003    | -0.02 (-1.44 to 1.39)           | .97     | -5.16 (-6.63 to -3.70)          | <.001   | -2.78 (-4.40 to -1.16)          | <.001   |
| Gestational age                                 | 2.86 (2.43 to 3.30)             | <.001   | 1.33 (0.88 to 1.77)             | <.001   | 2.82 (2.33 to 3.31)             | <.001   | 1.45 (0.94 to 1.95)             | <.001   |
| Male sex  | -3.38 (-4.27 to -2.50)          | <.001   | -3.05 (-3.87 to -2.22)          | <.001   | -5.72 (-6.7 to -4.73)           | <.001   | -5.45 (-6.39 to -4.51)          | <.001   |
| BPD   | -5.21 (-6.1 to -4.31)           | <.001   | -3.54 (-4.46 to -2.62)          | <.001   | -4.74 (-5.74 to -3.73)          | <.001   | -3.38 (-4.42 to -2.34)          | <.001   |
| NEC   | -5.25 (-6.77 to -3.73)          | <.001   | -3.51 (-4.95 to -2.08)          | <.001   | -4.32 (-6.03 to -2.62)          | <.001   | -2.21 (-3.85 to -0.57)          | .008    |
| Early-onset sepsis                              | -2.93 (-6.0 to 0.15)            | .06     | -1.13 (-3.99 to 1.72)           | .44     | -0.26 (-3.73 to 3.21)           | .88     | 0.63 (-2.65 to 3.91)            | .71     |
| Late-onset sepsis                               | -4.1 (-5.04 to -3.16)           | <.001   | -1.84 (-2.76 to -0.93)          | <.001   | -4.80 (-5.85 to -3.75)          | <.001   | -2.40 (-3.45 to -1.36)          | <.001   |
| cPVL or IVH (grade 3-4)                         | -7.83 (-8.98 to -6.67)          | <.001   | -6.58 (-7.69 to -5.47)          | <.001   | -5.77 (-7.08 to -4.47)          | <.001   | -4.45 (-5.72 to -3.19)          | <.001   |
| Severe ROP                                      | -6.94 (-7.97 to -5.91)          | <.001   | -4.08 (-5.84 to -3.75)          | <.001   | -6.54 (-7.7 to -5.37)           | <.001   | -4.16 (-5.35 to -2.97)          | <.001   |

Each variable included in the regression reflects statistically significant ( $P \leq .05$ ) associations with the outcomes or prior data have shown associations with the outcome.

controversial, owing to the complexity and timing of exposures and variations in living environments and caregiver skills.<sup>39-42</sup> The results of our exploratory analysis should be interpreted cautiously but nonetheless suggest the need for a detailed study of the support available to and adversities faced by EPT infants under CPS supervision.

In our analysis, CPS supervision is likely a proxy for an inadequate home environment. There are state-to-state differences in the details of CPS policy, referrals, and management, as described in the National Study of Child Protective Services Systems and Reform Efforts of State

CPS Policy<sup>43</sup> and the state statutes database.<sup>44</sup> Factors prompting CPS referral in an NICU may include maternal substance abuse, adverse maternal mental health, previous custody loss, or other concerns for potential abuse, neglect, or endangerment. In EPT infants, a long NICU stay allows for identification of psychosocial risks and implementation of parental support as needed. After discharge, it may be more difficult to identify risk factors and provide support, thus increasing the infants' exposure to adverse environments. Risk factors necessitating CPS supervision may impact the quality of parenting,<sup>45-47</sup> a critical factor for

**Table V. Two-year outcomes of EPT infants in CPS supervision subgroups**

| Outcomes mean $\pm$ SD; n (%)            | CPS supervision                       |                                 |  | Never in CPS supervision (N = 4129; 92%) | Overall P value* |
|--|---------------------------------------|---------------------------------|--|--|------------------|
|  | CPS at discharge only (N = 171; 3.8%) | CPS at 2 y only (N = 106; 2.4%) | CPS at both discharge and 2 y (N = 80; 1.8%) |  |                  |
| Age at follow-up, mo, mean $\pm$ SD      | 21.6 $\pm$ 3.3                        | 21.7 $\pm$ 3.3                  | 20.5 $\pm$ 2.8                               | 21.3 $\pm$ 3.1                           | .072             |
| Bayley-III                               |                                       |                                 |  |  |                  |
| Cognitive composite score, mean $\pm$ SD | 85.9 $\pm$ 14.5 <sup>a</sup>          | 83.9 $\pm$ 15.3 <sup>b</sup>    | 83.9 $\pm$ 18.4 <sup>c</sup>                 | 88.7 $\pm$ 15.3                          | <.001            |
| Language composite score, mean $\pm$ SD  | 83.6 $\pm$ 16.0 <sup>d</sup>          | 79.6 $\pm$ 18.2                 | 80.2 $\pm$ 17.7                              | 84.0 $\pm$ 17.0                          | .02              |
| Cognitive composite <85, n/N (%)         | 55/170 (32) <sup>e</sup>              | 44/104 (42) <sup>f</sup>        | 40/78 (51) <sup>g</sup>                      | 1198/4094 (29)                           | <.001            |
| Language composite <85, n/N (%)          | 86/167 (53)                           | 59/103 (57)                     | 45/75 (60)                                   | 2000/4025 (50)                           | .14              |
| Cognitive composite <70, n/N (%)         | 19/170 (11)                           | 15/104 (14)                     | 13/78 (17)                                   | 408/4094 (10)                            | .11              |
| Language composite <70, n/N (%)          | 30/167 (18) <sup>h</sup>              | 28/103 (27) <sup>i</sup>        | 22/75 (29) <sup>j</sup>                      | 751/4025 (19)                            | .02              |

\*Overall P values indicate where there is an overall difference between variables. Localization of differences between subgroups is indicated by superscript letters.<sup>a</sup>Discharge only vs never CPS ( $P = .02$ ).<sup>b</sup>2 y only vs never CPS ( $P = .001$ ).<sup>c</sup>Discharge and 2 y vs never CPS ( $P = .006$ ).<sup>d</sup>2 y only vs never CPS ( $P = .01$ ).<sup>e</sup>Discharge only vs discharge and 2 y ( $P = .004$ ).<sup>f</sup>2 y only vs never CPS ( $P = .004$ ).<sup>g</sup>Discharge and 2 y vs never CPS ( $P < .001$ ).<sup>h</sup>Discharge only vs discharge and 2 y ( $P = .04$ ).<sup>i</sup>2 y only vs never CPS ( $P = .004$ ).<sup>j</sup>Discharge and 2 y vs never CPS ( $P = .02$ ).

optimizing outcomes. Responsive, structured, and independence-promoting parenting styles are associated with improved child cognition and behavior,<sup>48,49</sup> and CPS goals include child placement with caregivers who are able to provide such nurturing interactions.

Our study has several strengths and limitations. We evaluated prospectively collected neonatal and 2-year outcome data in a relatively large cohort of EPT newborns with CPS supervision. The NICHD NRN database, which includes a large and diverse population represented by 21 tertiary care centers throughout the US, allowed exploration of outcomes at a critical time point of 0-2 years of age. Limitations include a retrospective analysis and the absence of data capturing reasons for CPS supervision and maternal risk factors, such as drug/substance use, domestic violence, or mental illness. In addition, the study was not powered to evaluate differences among CPS subgroups. Generalizations about foster homes and kinship care cannot be made, because placement strategies vary and case workers face the challenge of securing a safe environment while minimizing the disruptions of removal.

EPT infants discharged to CPS are a highly vulnerable population and are at increased risk for cognitive delays. Environments necessitating either the continued need for CPS supervision at 2 years of age or the need to enter CPS after discharge appear to convey even greater risk for low cognitive and language skills. The combination of being born EPT, an identified need for CPS supervision, and exposure to multiple social and environmental risk factors should trigger close surveillance. More research is needed to identify potential interventions for caregivers and infants. ■

*We are indebted to our medical and nursing colleagues (Appendix) and the infants and their parents who participated in this study.*

Submitted for publication Apr 5, 2019; last revision received Jun 11, 2019; accepted Jul 24, 2019.

Reprint requests: Elisabeth C. McGowan, MD, Division of Neonatal Medicine, Alpert Medical School of Brown University, Women & Infants Hospital, 101 Plain Street, Providence, RI 02905. E-mail: emcgowan@wihri.org

## Data statement

Data sharing statement available at [www.jpeds.com](http://www.jpeds.com).

## References

- US Department of Health and Human Services. Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau. Child maltreatment. 2017. Available at: <https://www.acf.hhs.gov/cb/research-data-technology/statistics-research/child-maltreatment>. Accessed June 4, 2019.
- Jee SH, Szilagyi M, Ovenshire C, Norton A, Conn AM, Blumkin A, et al. Improved detection of developmental delays among young children in foster care. *Pediatrics* 2010;125:282-9.
- Leslie LK, Gordon JN, Meneken L, Premji K, Michelmore KL, Ganger W. The physical, developmental, and mental health needs of young children in child welfare by initial placement type. *J Dev Behav Pediatr* 2005;26:177-85.
- Jacobsen H, Moe V, Ivarsson T, Wentzel-Larsen T, Smith L. Cognitive development and social-emotional functioning in young foster children: a follow-up study from 2 to 3 years of age. *Child Psychiatry Hum Dev* 2013;44:666-77.
- Hintz SR, Kendrick DE, Wilson-Costello DE, Das A, Bell EF, Vohr BR, et al. Early-childhood neurodevelopmental outcomes are not improving for infants born at <25 weeks' gestational age. *Pediatrics* 2011;127:62-70.
- Vohr BR, Wright LL, Poole WK, McDonald SA. Neurodevelopmental outcomes of extremely low birth weight infants <32 weeks' gestation between 1993 and 1998. *Pediatrics* 2005;116:635-43.
- Adams-Chapman I, Bann C, Carter SL, Stoll BJ. Language outcomes among ELBW infants in early childhood. *Early Hum Dev* 2015;91:373-9.
- Arpi E, Ferrari F. Preterm birth and behaviour problems in infants and preschool-age children: a review of the recent literature. *Dev Med Child Neurol* 2013;55:788-96.
- Peralta-Carcelen M, Bailey K, Rector R, Gantz M. Behavioral and socio-emotional competence problems of extremely low birth weight children. *J Perinatol* 2013;33:887-92.
- Halfon N, Mendonca A, Berkowitz G. Health status of children in foster care. The experience of the Center for the Vulnerable Child. *Arch Pediatr Adolesc Med* 1995;149:386-92.
- Hoffman L, Bann C, Higgins R, Vohr B. Developmental outcomes of extremely preterm infants born to adolescent mothers. *Pediatrics* 2015;135:1082-92.
- Olsen IE, Groveman SA, Lawson ML, Clark RH, Zemel BS. New intra-uterine growth curves based on United States data. *Pediatrics* 2010;125:e214-24.
- Alexander GR, Himes JH, Kaufman RB, Mor J, Kogan M. A United States national reference for fetal growth. *Obstet Gynecol* 1996;87:163-8.
- Newman JE, Bann CM, Vohr BR, Dusick AM, Higgins RD. Improving the Neonatal Research Network annual certification for neurologic examination of the 18-22 month child. *J Pediatr* 2012;161:1041-6.
- Bayley N. Bayley scales of infant and toddler development (Bayley-III). 3rd ed. San Antonio (TX): Harcourt; 2006.
- Briggs-Gowan MJ, Carter AS. Brief infant-toddler social and emotional assessment (BITSEA) manual. Version 2.0. New Haven (CT): Yale University; 2002.
- Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol* 1997;39:214-23.
- Adoption and Safe Families Act of 1997 (Public Law 105-89).
- Stahmer AC, Leslie LK, Hurlburt M, Barth RP, Webb MB, Landsverk J, et al. Developmental and behavioral needs and service use for young children in child welfare. *Pediatrics* 2005;116:891-900.
- Anderson PJ, Burnett A. Assessing developmental delay in early childhood - concerns with the Bayley-III scales. *Clin Neuropsychol* 2017;31:371-81.
- Anderson PJ, De Luca CR, Hutchinson E, Roberts G, Doyle LW. Underestimation of developmental delay by the new Bayley-III Scale. *Arch Pediatr Adolesc Med* 2010;164:352-6.
- Peralta-Carcelen M, Carlo WA, Pappas A, Vaucher YE, Yeates KO, Phillips VA, et al. Behavioral problems and socioemotional competence at 18 to 22 months of extremely premature children. *Pediatrics* 2017;139:e20161043.
- Berg-Nielsen TS, Solheim E, Belsky J, Wichstrom L. Preschoolers' psychosocial problems: in the eyes of the beholder? Adding teacher characteristics as determinants of discrepant parent-teacher reports. *Child Psychiatry Hum Dev* 2012;43:393-413.
- Bou Malham P, Saucier G. The conceptual link between social desirability and cultural normativity. *Int J Psychol* 2016;51:474-80.
- Turney K, Wildeman C. Mental and physical health of children in foster care. *Pediatrics* 2016;138:e20161118.
- Forkey H, Szilagyi M. Foster care and healing from complex childhood trauma. *Pediatr Clin North Am* 2014;61:1059-72.
- Mathewson KJ, Chow CH, Dobson KG, Pope EI, Schmidt LA, Van Lieshout RJ. Mental health of extremely low birth weight survivors: a systematic review and meta-analysis. *Psychol Bull* 2017;143:347-83.
- Linsell L, Johnson S, Wolke D, Morris J, Kurinczuk JJ, Marlow N. Trajectories of behavior, attention, social and emotional problems from childhood to early adulthood following extremely preterm birth: a prospective cohort study. *Eur Child Adolesc Psychiatry* 2019;28:531-42.

29. Dusick AM, Poindexter BB, Ehrenkranz RA, Lemons JA. Growth failure in the preterm infant: can we catch up? *Semin Perinatol* 2003;27:302-10.
30. Suchdev PS, Boivin MJ, Forsyth BW, Georgieff MK, Guerrant RL, Nelson CA III. Assessment of neurodevelopment, nutrition, and inflammation from fetal life to adolescence in low-resource settings. *Pediatrics* 2017;139(Suppl 1):S23-37.
31. Cook JT. Clinical implications of household food security: definitions, monitoring, and policy. *Nutr Clin Care* 2002;5:152-67.
32. Raghuram K, Yang J, Church PT, Cieslak Z, Synnes A, Mukerji A, et al. Head growth trajectory and neurodevelopmental outcomes in preterm neonates. *Pediatrics* 2017;140:e20170216.
33. Hack M, Breslau N, Weissman B, Aram D, Klein N, Borawski E. Effect of very low birth weight and subnormal head size on cognitive abilities at school age. *N Engl J Med* 1991;325:231-7.
34. Strand KM, Dahlseng MO, Lydersen S, Ro TB, Finbraten AK, Jahnsen RB, et al. Growth during infancy and early childhood in children with cerebral palsy: a population-based study. *Dev Med Child Neurol* 2016;58:924-30.
35. Hintz SR, Barnes PD, Bulas D, Slovis TL, Finer NN, Wrage LA, et al. Neuroimaging and neurodevelopmental outcome in extremely preterm infants. *Pediatrics* 2015;135:e32-42.
36. Woodward LJ, Anderson PJ, Austin NC, Howard K, Inder TE. Neonatal MRI to predict neurodevelopmental outcomes in preterm infants. *N Engl J Med* 2006;355:685-94.
37. Linsell L, Malouf R, Morris J, Kurinczuk JJ, Marlow N. Prognostic factors for cerebral palsy and motor impairment in children born very preterm or very low birthweight: a systematic review. *Dev Med Child Neurol* 2016;58:554-69.
38. Vohr B, McGowan E, Keszler L, Alksninis B, O'Donnell M, Hawes K, et al. Impact of a transition home program on rehospitalization rates of preterm infants. *J Pediatr* 2017;181:86-92.e1.
39. Judge S. Developmental recovery and deficit in children adopted from Eastern European orphanages. *Child Psychiatry Hum Dev* 2003;34:49-62.
40. Rutter M. Developmental catch-up, and deficit, following adoption after severe global early privation. English and Romanian Adoptees (ERA) Study Team. *J Child Psychol Psychiatry* 1998;39:465-76.
41. Beckett C, Maughan B, Rutter M, Castle J, Colvert E, Groothues C, et al. Do the effects of early severe deprivation on cognition persist into early adolescence? Findings from the English and Romanian adoptees study. *Child Dev* 2006;77:696-711.
42. Goemans A, van Geel M, Vedder P. Over three decades of longitudinal research on the development of foster children: a meta-analysis. *Child Abuse Negl* 2015;42:121-34.
43. U.S. Department of Health and Human Services. National study of child protective services systems and reform efforts. Review of state CPS policy. 2013. Available at: <https://aspe.hhs.gov/national-study-child-protective-services-systems-and-reform-efforts>. Accessed June 4, 2019.
44. Child Welfare Information Gateway. State statutes search. Available at: <https://www.childwelfare.gov/topics/systemwide/laws-policies/state/>. Accessed June 4, 2019.
45. Leschied AW, Chiodo D, Whitehead PC, Hurley D. The relationship between maternal depression and child outcomes in a child welfare sample: implications for treatment and policy. *Child Fam Social Work* 2005;10:281-91.
46. Casanueva C, Cross TP, Ringeisen H, Christ S. Prevalence, trajectories, and risk factors for depression among caregivers of young children involved in maltreatment investigations. *J Emot Behav Disord* 2011;19:98-116.
47. Dorsey S, Mustillo SA, Farmer EM, Elbogen E. Caseworker assessments of risk for recurrent maltreatment: association with case-specific risk factors and re-reports. *Child Abuse Negl* 2008;32:377-91.
48. Neel MLM, Stark AR, Maitre NL. Parenting style impacts cognitive and behavioural outcomes of former preterm infants: a systematic review. *Child Care Health Dev* 2018;44:507-15.
49. Hoffenkamp HN, Braeken J, Hall RA, Tooten A, Vingerhoets AJ, van Bakel HJ. Parenting in complex conditions: does preterm birth provide a context for the development of less optimal parental behavior? *J Pediatr Psychol* 2015;40:559-71.

## Appendix

**Additional Members of the Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network**

NRN Steering Committee Chair: Richard A. Polin, MD, Division of Neonatology, College of Physicians and Surgeons, Columbia University

Alpert Medical School of Brown University and Women & Infants Hospital of Rhode Island (U10 HD27904): Abbott R. Laptook, MD; Martin Keszler, MD; Angelita M. Hensman, MS, RNC-NIC, Barbara Alksninis, PNP, Kristin M. Basso, BSN, MaT, Robert Burke, MD, Melinda Caskey, MD, Katharine Johnson, MD, Mary Lenore Keszler, MD, Andrea M. Knoll, Theresa M. Leach, MEd, CAES, Emilee Little, RN, BSN, Elisabeth C. McGowan, MD, Elisa Vieira, RN, BSN, Victoria E. Watson, MS, CAS, Suzy Ventura.

Case Western Reserve University, Rainbow Babies & Children's Hospital (U10 HD21364, M01 RR80): Michele C. Walsh, MD, MS, Avroy A. Fanaroff, MD, Anna Marie Hibbs, MD, Deanne E. Wilson-Costello, MD, Nancy S. Newman, BA, RN, Allison H. Payne, MD, MS, Bonnie S. Siner, RN, Monika Bhola, MD, Gulgun Yalcinkaya, MD, Harriet G. Friedman, MA.

Children's Mercy Hospital, University of Missouri Kansas City School of Medicine (U10 HD68284): William E. Truog, MD, Eugenia K. Pallotto, MD, MSCE, Howard W. Kilbride, MD, Cheri Gauldin, RN, BS, CCRC, Anne Holmes, RN, MSN, MBA-HCM, CCRC, Kathy Johnson RN, CCRC, Allison Knutson, BSN, RNC-NIC.

Cincinnati Children's Hospital Medical Center, University Hospital, and Good Samaritan Hospital (U10 HD27853, M01 RR8084): Kurt Schibler, MD, Edward F. Donovan, MD, Cathy Grisby, BSN, CCRC, Kate Bridges, MD, Barbara Alexander, RN, Estelle E. Fischer, MHSA, MBA, Holly L. Mincey, RN, BSN, Jody Hessling, RN, Teresa L. Gratton, PA, Lenora Jackson, CRC, Kristin Kirker, CRC, Greg Muthig, BS, Jean J. Steichen, MD, Stacey Tepe, BS, Kimberly Yolton, PhD.

Duke University School of Medicine, University Hospital, University of North Carolina, and Duke Regional Hospital (U10 HD40492, UL1 TR1117, M01 RR30, UL1 TR1111): Ronald N. Goldberg, MD, C. Michael Cotten, MD, MHS, Ricki F. Goldstein, MD, Patricia L. Ashley, MD, PhD, William F. Malcolm, MD, Kathy J. Auten, MSHS, Kimberley A. Fisher, PhD, FNP-BC, IBCLC, Sandra Grimes, RN, BSN, Kathryn E. Gustafson, PhD, Melody B. Lohmeyer, RN, MSN, Joanne Finkle, RN, JD, Matthew M. Laughon, MD, MPH, Carl L. Bose, MD, Janice Bernhardt, MS, RN, Gennie Bose, RN, Cindy Clark, RN, Linda Manor, RPh, Diane Warner, MD, MPH, Janice Wereszczak, NNP.

Emory University, Children's Healthcare of Atlanta, Grady Memorial Hospital, and Emory University Hospital Midtown (U10 HD27851, M01 RR39): David P. Carlton, MD, Barbara J. Stoll, MD, Ira Adams-Chapman, MD, Ellen C. Hale, RN, BS, CCRC, Yvonne Loggins, RN, BSN.

*Eunice Kennedy Shriver* National Institute of Child Health and Human Development: Stephanie Wilson Archer, MA

Indiana University, University Hospital, Methodist Hospital, Riley Hospital for Children, and Wishard Health Services (U10 HD27856, M01 RR750): Gregory M. Sokol, MD, Brenda B. Poindexter, MD, MS, Anna M. Dusick, MD (deceased), LuAnn Papile, MD, Susan Gunn, NNP, CCRC, Faithe Hamer, BS, Dianne E. Herron, RN, CCRC, Abbey C. Hines, PsyD, Carolyn Lytle, MD, MPH, Heike M. Minnich, PsyD, HSPP, Lucy Smiley, CCRC, Leslie Dawn Wilson, BSN, CCRC.

Nationwide Children's Hospital and the Ohio State University Medical Center (U10 HD68278): Pablo J. Sanchez, MD, Leif D. Nelin, MD, Sudarshan R. Jadcherla, MD, Patricia Luzader, RN, Christine A. Fortney, PhD, RN, Gail E. Besner, MD, Nehal A. Parikh, MD.

RTI International (U10 HD36790): Abhik Das, PhD, Dennis Wallace, PhD, Marie G. Gantz, PhD, W. Kenneth Poole, PhD (deceased), Jamie E. Newman, PhD, MPH, Jeanette O'Donnell Auman, BS, Margaret M. Crawford, BS, CCRP, Carolyn M. Petrie Huitema, MS, CCRP, Kristin M. Zaterka-Baxter, RN, BSN, CCRP.

Stanford University, Dominican Hospital, El Camino Hospital, and Lucile Packard Children's Hospital (U10 HD27880, M01 RR70): Krisa P. Van Meurs, MD, David K. Stevenson, MD, Susan R. Hintz, MD, MS, Epi, M. Bethany Ball, BS, CCRC, Alexis S. Davis, MD, MS, Epi, Andrew W. Palmquist, RN, Melinda S. Proud, RCP, Barbara Bentley, PsychD, MEd, Elizabeth Bruno, PhD, Maria Elena DeAnda, PhD, Anne M. DeBattista, RN, PNP, Beth Earhart, PhD, Lynne C. Huffman, MD, Jean G. Kohn, MD, MPH, Casey Krueger, PhD, Hali E. Weiss, MD.

Tufts Medical Center, Floating Hospital for Children (U10 HD53119, M01 RR54): Ivan D. Frantz III, MD, John M. Fiascone, MD, Brenda L. MacKinnon, RNC, Anne Furey, MPH, Ellen Nysten, RN, BSN, Elisabeth C. McGowan, MD.

University of Alabama at Birmingham Health System and Children's Hospital of Alabama (U10 HD34216, M01 RR32): Waldemar A. Carlo, MD, Namasivayam Ambalavanan, MD, Myriam Peralta-Carcelen, MD, MPH, Monica V. Collins, RN, BSN, MaEd, Shirley S. Cosby, RN, BSN, Fred J. Biasini, PhD, Kristen C. Johnston, MSN, CRNP, Kathleen G. Nelson, MD, Cryshelle S. Patterson, PhD, Vivien A. Phillips, RN, BSN, Sally Whitley, MA, OTR-L, FAOTA.

University of California Los Angeles, Mattel Children's Hospital, Santa Monica Hospital, Los Robles Hospital and Medical Center, and Olive View Medical Center (U10 HD68270): Uday Devaskar, MD, Meena Garg, MD, Isabell B. Purdy, PhD, CPNP, Teresa Chanlaw, MPH, Rachel Geller, RN, BSN.

University of California San Diego Medical Center and Sharp Mary Birch Hospital for Women and Newborns (U10 HD40461): Neil N. Finer, MD, Yvonne E. Vaucher, MD, MPH, David Kaegi, MD, Maynard R. Rasmussen, MD, Kathy Arnell, RNC, Clarence Demetrio, RN, Martha G. Fuller, RN, MSN, Wade Rich, BSHS, RRT.

University of Iowa and Mercy Medical Center (U10 HD53109, M01 RR59): Edward F. Bell, MD, Tarah T.

Colaizy, MD, MPH, Michael J. Acarregui, MD, Dan L. Ellsbury, MD, John A. Widness, MD, Karen J. Johnson, RN, BSN, Donia B. Campbell, RNC-NC, Diane L. Eastman, RN, CPNP, MA, Jacky R. Walker, RN, Jane E. Brumbaugh, MD.

University of Miami, Holtz Children's Hospital (U10 HD21397, M01 RR16587): Shahnaz Duara, MD, Charles R. Bauer, MD, Ruth Everett-Thomas, RN, MSN, Sylvia Fajardo-Hiriart, MD, Arielle Rigaud, MD, Maria Calejo, MS, Silvia M. Frade Eguaras, MA, Michelle Harwood Berkwits, PhD, Andrea Garcia, MS, Helina Pierre, BA, Alexandra Stoerger, BA.

University of New Mexico Health Sciences Center (U10 HD53089, M01 RR997): Kristi L. Watterberg, MD, Jean R. Lowe, PhD, Janell F. Fuller, MD, Robin K. Ohls, MD, Conra Backstrom Lacy, RN, Andrea F. Duncan, MD, MSc, Rebecca Montman, BSN.

University of Pennsylvania, Hospital of the University of Pennsylvania, Pennsylvania Hospital, and Children's Hospital of Philadelphia (U10 HD68244): Barbara Schmidt, MD, MSc, Hareesh Kirpalani, MB, MSc, Sara B. DeMauro, MD, MSCE, Aasma S. Chaudhary, BS, RRT, Soraya Abbasi, MD, Toni Mancini, RN, BSN, CCRC, Dara M. Cucinotta, RN, Judy C. Bernbaum, MD, Marsha Gerdes, PhD, Hallam Hurt, MD.

University of Rochester Medical Center, Golisano Children's Hospital, and the University of Buffalo Women's and Children's Hospital of Buffalo (U10 HD68263, U10 HD40521, M01 RR44, UL1 TR42): Carl T. D'Angio, MD, Dale L. Phelps, MD, Ronnie Guillet, MD, PhD, Satyan Lakshminrusimha, MD, Julie Babish Johnson, MSW, Linda J. Reubens, RN, CCRC, Cassandra A. Horihan, MS, Diane Hust, MS, RN, CS, Rosemary L. Jensen, Emily Kushner, MA, Joan Merzbach, LMSW, Gary J. Myers, MD, Mary Rowan, RN, Holly I. M. Wadkins, MA, Melissa Bowman, MSN, Julianne Hunn, BS, Stephanie Guilford, BS, Deanna Maffett, RN, Farooq Osman, MD, Diane Prinzing, Anne Marie Reynolds, MD, MPH, Mary Rowan, RN, Michael G. Sacilowski, BS, Ashley Williams, MEd, Karen Wynn, RN, Kelley Yost, PhD, William Zorn, MD, Lauren Zwetsch, RN, MS, PNP.

University of Texas Health Science Center at Houston Medical School, Children's Memorial Hermann Hospital, and Lyndon Baines Johnson General Hospital/Harris County Hospital District (U10 HD21373): Kathleen A. Kennedy, MD, MPH, Jon E. Tyson, MD, MPH, Georgia E. McDavid, RN, Nora I. Alaniz, BS, Julie Arldt-McAlister, RN, BSN, Katrina Burson, RN, BSN, Patricia W. Evans, MD, Andrea F. Duncan, MD, Carmen Garcia, RN, CCRP, Charles Green, PhD, Beverly Foley Harris, RN, BSN, Margarita Jiminez, MD, MPH, Janice John, CPNP, Patrick M. Jones, MD, Layne M. Lillie, RN, BSN, Anna E. Lis, RN, BSN, Karen Martin, RN, Sara C. Martin, RN, BSN, Brenda H. Morris, MD, M. Layne Poundstone, RN, BSN, Peggy Robichaux, RN, BSN, Shawna Rodgers, RN, Saba Siddiki, MD, Maegan C. Simmons, RN, Daniel Sperry, RN, Patti L. Pierce Tate, RCP, Sharon L. Wright, MT(ASCP).

University of Texas Southwestern Medical Center at Dallas, Parkland Health & Hospital System, and Children's Med-

ical Center Dallas (U10 HD40689, M01 RR633): Myra H. Wyckoff, MD, Luc P. Brion, MD, Roy J. Heyne, MD, Walid A. Salhab, MD, Charles R. Rosenfeld, MD, Diana M. Vasil, RNC-NIC, Lijun Chen, PhD, RN, Alicia Guzman, Gaynelle Hensley, RN, Melissa H. Leps, RN, Nancy A. Miller, RN, Janet S. Morgan, RN, Sally S. Adams, MS, RN, CPNP, Catherine Twell Boatman, MS, CIMI, Elizabeth T. Heyne, MS, MA, PA-C, PsyD, Linda A. Madden, RN, CPNP, Lizette E. Torres, RN.

University of Utah Medical Center, Intermountain Medical Center, LDS Hospital, and Primary Children's Medical Center (U10 HD53124, M01 RR64, UL1 RR25764): Roger G. Faix, MD, Bradley A. Yoder, MD, Karen A. Osborne, RN, BSN, CCRC, Cynthia Spencer, RNC, BSN, Kimberlee Weaver-Lewis, RN, MS, Shawna Baker, RN, Karie Bird, RN, BSN, Jill Burnett, RNC, BSN, Michael Steffen, MS, CPM, Jennifer J. Jensen, RN, BSN, Sarah Winter, MD, Karen Zanetti, RN.

Wake Forest University, Baptist Medical Center, Forsyth Medical Center, and Brenner Children's Hospital (U10 HD40498, M01 RR7122): T. Michael O'Shea, MD, MPH, Robert G. Dillard, MD, Lisa K. Washburn, MD, Barbara G. Jackson, RN, BSN, Nancy Peters, RN, Korinne Chiu, MA, Deborah Evans Allred, MA, LPA, Donald J. Goldstein, PhD, Raquel Halfond, MA, Carroll Peterson, MA, Ellen L. Waldrep, MS, Cherrie D. Welch, MD, MPH, Melissa Whalen Morris, MA, Gail Wiley Hounshell, PhD.

Wayne State University, Hutzel Women's Hospital and Children's Hospital of Michigan (U10 HD21385): Seetha Shankaran, MD, Athina Pappas, MD, John Barks, MD, Rebecca Bara, RN, BSN, Laura A. Goldston, MA, Girija Natarajan, MD, Mary Christensen, RT, Stephanie A. Wiggins, MS, Diane White, RT.

Yale University, Yale-New Haven Children's Hospital, and Bridgeport Hospital (U10 HD27871, ULTR142, M01 RR125): Richard A. Ehrenkranz, MD, Harris Jacobs, MD, Christine G. Butler, MD, Patricia Cervone, RN, Sheila Greisman, RN, Monica Konstantino, RN, BSN, JoAnn Poulsen, RN, Janet Taft, RN, BSN, Joanne Williams, RN, BSN, Elaine Romano, MSN.

Supported by the National Institutes of Health, Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) (U10 HD21364, U10 HD27871, U10 HD21373, U10 HD21385, U10 HD21397, U10 HD27851, U10 HD27853, U10 HD27856, U10 HD27880, U10 HD27904, U10 HD34216, U10 HD36790, U10 HD40461, U10 HD40498, U10 HD40689, U10 HD40492, U10 HD40521, U10 HD53089, U10 HD53109, U10 HD53119, U10 HD53124, U10 HD68244, U10 HD68263, U10 HD68270, U10 HD68278, U10 HD68284) and the National Center for Advancing Translational Sciences (M01 RR30, M01 RR32, M01 RR39, M01 RR44, M01 RR54, M01 RR59, M01 RR64, M01 RR70, M01 RR80, M01 RR125, M01 RR633, M01 RR750, M01 RR997, M01 RR7122, M01 RR8084, M01 RR16587, UL1 TR42, ULTR142, UL1 TR1111, UL1 TR1117, UL1 RR25764). The NIH, NICHD, National Center for Research Resources, and National Center for Advancing Translational Sciences

provided grant support for the Neonatal Research Network's (NRN) generic database and follow-up studies through cooperative agreements. Recruitment was between January 1, 2006 through December 31, 2013, and follow-up was between approximately June 1, 2008, and October 31, 2015. Although the NICHD staff did have input into the study design, conduct, analysis, and manuscript drafting, the content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Data

collected at participating sites of the NICHD NRN were transmitted to RTI International, the data coordinating center (DCC) for the network, which stored, managed, and analyzed the data for this study. On behalf of the NRN, Abhik Das (DCC Principal Investigator) and Douglas Kendrick (DCC statistician) had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. The authors declare no conflicts of interest.

**Table II. Social and environmental characteristics at NICU discharge and at 2 years in EPT infants with or without CPS supervision at discharge**

| Characteristics   | At NICU discharge      |                              |         | At 2 years          |                           |         |
|---|------------------------|------------------------------|---------|---------------------|---------------------------|---------|
|   | CPS<br>(N = 255; 5.6%) | Non-CPS<br>(N = 4262; 94.4%) | P value | CPS (N = 255; 5.6%) | Non-CPS (N = 4262; 94.4%) | P value |
| Primary caretaker, n (%)                                |                        |                              |         |                     |                           |         |
| Kinship care  | 149 (58.4)             | 4226 (99.2)                  | <.001   | 159/251 (63.3)      | 4171/4236 (98.5)          | <.001   |
| Biological mother                                       | 107 (42.0)             | 4184 (98.2)                  | <.001   | 111 (44.2)          | 4041 (95.4)               | <.001   |
| Biological father                                       | 5 (2.0)                | 24 (0.6)                     | .006    | 13 (5.2)            | 63 (1.5)                  | <.001   |
| Grandparent(s)  | 24 (9.4)               | 16 (0.3)                     | <.001   | 26 (10.4)           | 54 (1.3)                  | <.001   |
| Other relative  | 13 (5.1)               | 2 (0.05)                     | <.001   | 9 (3.6)             | 13 (0.3)                  | <.001   |
| Foster parent (nonkinship)                              | 91 (35.7)              | 6 (0.1)                      | <.001   | 49 (19.5)           | 34 (0.8)                  | <.001   |
| Adoptive parent   | 5 (2.0)                | 8 (0.2)                      | <.001   | 40 (15.9)           | 18 (0.4)                  | <.001   |
| Other nonrelative/congregate care                       | 7 (2.7)                | 3 (0.07)                     | <.001   | 2 (0.8)             | 10 (0.2)                  | .09     |
| Still hospitalized                                      | 0                      | 3 (0.07)                     | .67     | 1 (0.4)             | 3 (0.1)                   | .09     |
| Primary caretaker married, n (%)                        | 127 (49.8)             | 2071 (48.6)                  | .40     | 128/248 (51.6)      | 2168/4224 (51.3)          | .90     |
| Living/household arrangements, n (%)                    |                        |                              |         |                     |                           |         |
| Single biological parent                                | 54 (21.2)              | 1161 (27.2)                  | .04     | 61/250 (24.4)       | 1303 (30.8)               | .03     |
| 2-parent biological                                     | 63 (24.7)              | 3048 (71.5)                  | <.001   | 70 (28.0)           | 2806 (66.3)               | <.001   |
| Grandparent(s)  | 16 (6.3)               | 19 (0.4)                     | <.001   | 21 (8.4)            | 43 (1.0)                  | <.001   |
| Adoptive parent (relative)                              | 2 (.8)                 | 1 (0.5)                      | <.001   | 14 (5.6)            | 10 (0.2)                  | <.001   |
| Adoptive parent (nonrelative)                           | 1 (0.4)                | 4 (0.1)                      | .16     | 27 (10.8)           | 9 (0.2)                   | <.001   |
| Preadoptive home  | 5 (2.0)                | 6 (0.1)                      | <.001   | 6 (2.4)             | 6 (0.1)                   | <.001   |
| Foster family (relative)                                | 12 (4.7)               | 0                            | <.001   | 2 (0.8)             | 6 (0.1)                   | .02     |
| Foster family (nonrelative)                             | 88 (34.5)              | 3 (0.1)                      | <.001   | 41 (16.4)           | 29 (0.7)                  | <.001   |
| Other   | 11 (4.3)               | 12 (0.3)                     | <.001   | 8 (3.2)             | 18 (0.4)                  | <.001   |
| Number of people in home, median (IQR)                  | 4 (3-6)                | 4 (3-5)                      | <.001   | 5 (4-6)             | 4 (3-5)                   | <.001   |
| Medical insurance, n (%)                                |                        |                              |         |                     |                           |         |
| Public  | 214 (84.0)             | 2669 (62.6)                  | <.001   | 214/251 (85.3)      | 2669 (63.1)               | <.001   |
| Private   | 20 (7.8)               | 1063 (24.9)                  | <.001   | 20 (8.0)            | 1063 (25.1)               | <.001   |
| Uninsured   | 2 (0.8)                | 56 (1.3)                     | .47     | 2 (0.8)             | 56 (1.3)                  | .47     |
| Both public and private                                 | 15 (5.9)               | 442 (10.4)                   | .02     | 15 (6.0)            | 442 (10.4)                | .02     |
| Highest grade completed of 2-y primary caretaker, n (%) |                        |                              |         |                     |                           |         |
| Less than high school                                   | n/a                    | n/a                          |         | 52/251 (20.7)       | 682 (16.1)                | .05     |
| High school graduate                                    | n/a                    | n/a                          |         | 71 (28.2)           | 1180 (28.0)               | .88     |
| Some college  | n/a                    | n/a                          |         | 74 (29.5)           | 1179 (28.0)               | .57     |
| College graduate  | n/a                    | n/a                          |         | 49 (19.5)           | 1138 (26.9)               | .01     |
| Unknown   | n/a                    | n/a                          |         | 5 (2.0)             | 57 (1.3)                  | .39     |
| Primary language in 2-y home, n (%)                     |                        |                              |         |                     |                           |         |
| English   | n/a                    | n/a                          |         | 234/251 (93.2)      | 3649/4236 (86.1)          | .001    |
| Spanish   | n/a                    | n/a                          |         | 14 (5.6)            | 445 (10.5)                | .01     |
| Other   | n/a                    | n/a                          |         | 3 (1.2)             | 141 (3.3)                 | .06     |
| Bilingual home in 2-y household, n (%)                  | n/a                    | n/a                          |         | 48/251 (19.1)       | 1043/4233 (24.6)          | .04     |
| Places lived from discharge to 2 y, mean $\pm$ SD       | n/a                    | n/a                          |         | 1.8 $\pm$ 0.9       | 1.6 $\pm$ 0.8             | .003    |
| 1, n (%)  | n/a                    | n/a                          |         | 114/248 (46.0)      | 2432/4192 (58.0)          | .002    |
| 2, n (%)  | n/a                    | n/a                          |         | 97 (39.1)           | 1311 (31.2)               | .009    |
| 3+, n (%)   | n/a                    | n/a                          |         | 37 (14.9)           | 449 (10.7)                | .04     |
| Receipt of early intervention at 2 y, n (%)             | n/a                    | n/a                          |         | 181/251 (72.1)      | 2848/4235 (67.2)          | .11     |