



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Original article

Diabetes and dyslipidaemia are associated with oxidative stress independently of inflammation in long-term antiretroviral-treated HIV-infected patients

J.-P. Bastard^a, C. Couffignal^b, S. Fellahi^a, J.-M. Bard^c, F. Mentre^b, D. Salmon^d, C. Katlama^e, F. Raffi^{f,1}, C. Leport^{b,1}, J. Capeau^{a,*}, ANRS CO, APROCO-COPILOTE Cohort study group

^a Faculty of medicine, Sorbonne université, inserm UMR_S938, ICAN, AP-HP, hôpital Tenon, 27, rue Chaligny, 75571 Paris cedex 12, Paris, France

^b Université Paris Diderot, Sorbonne Paris Cité, inserm UMR_S1137, COREB APHP, 16, rue Henri-Huchard, 75890 Paris cedex 18, France

^c UFR des sciences pharmaceutiques et biologiques, MMS - EA 2160, IUML FR3473 CNRS, Nantes and institut de cancérologie de l'Ouest, 4, rue Bras France, BP61112, 44035 Nantes cedex 1, France

^d Service des maladies infectieuses et tropicales, hôpital Cochin, AP-HP, 27, rue du Faubourg Saint-Jacques, 75014 Paris, France

^e Service de maladies infectieuses et tropicales hôpital Pitié-Salpêtrière, AP-HP, Sorbonne Universités, UPMC Université Paris-6, inserm UMR_S1136 IPLESP, 47–83, boulevard de l'Hôpital, 75013 Paris, France

^f Service des maladies infectieuses et tropicales, inserm CIC 1413, CHU de Nantes, 1, place Alexis-Ricordeau, 44000 Nantes, France

ARTICLE INFO

Article history:

Received 4 October 2018

Received in revised form 15 February 2019

Accepted 24 February 2019

Available online 9 March 2019

Keywords:

Diabetes
 Dyslipidaemia
 HIV-infected patients
 Oxidative stress
 Inflammation

ABSTRACT

Aim. – Ageing HIV-infected patients controlled by antiretroviral therapy (ART) frequently present age-related comorbidities, such as cardiovascular (CV) events, diabetes, dyslipidaemia, hypertension and chronic kidney disease (CKD). The prevalence of these comorbidities was evaluated in a cohort of long-term-monitored ART-controlled HIV-infected patients, then followed by a search into whether oxidative stress, like inflammation, might be associated with metabolic parameters and/or comorbidities.

Methods. – Included were 352 long-term ART patients who started with protease inhibitors (PIs) in 1997–1999. They were evaluated at their final visit, 11 years later, for previous CV events, prevalence of diabetes, LDL-related and atherogenic (high TG/HDL) dyslipidaemias, hypertension and CKD. Also measured were circulating biomarkers to explore oxidative stress (Lp-PLA2, oxLDL, oxLDL/LDL ratio, paraoxonase and arylesterase activities), inflammation/immune activation (hsCRP, hslL-6, D dimer, soluble CD14, β 2 microglobulin, cystatin C), adipokines and insulin resistance. Levels were compared in patients with and without each comorbidity or condition using non-parametric correlation tests and multivariate adjusted analyses.

Results. – At the final visit, 81.5% of patients were male and were aged (median, IQR) 49 years (45–56); BMI was 23.0 kg/m² (21.1–25.4), CD4+ lymphocytes were 620 cells/mm³ (453–790) and 91.5% had undetectable HIV-1 viral loads. The prevalence of diabetes was 11%, and LDL-related dyslipidaemia 28%, atherogenic dyslipidaemia 9%, hypertension 28%, CKD 9% and previous CV events 9%. Diabetes and atherogenic dyslipidaemia were associated with increased oxidative stress and independently with inflammation. LDL-related dyslipidaemia and impaired fasting glucose were associated with increased oxidative stress. No association of these biomarkers was detected with hypertension, CKD and previous CV events.

Conclusion. – In long-term-treated HIV-infected patients with frequent comorbid conditions, oxidative stress could be contributing to diabetes and LDL-related and atherogenic dyslipidaemias independently of inflammation.

© 2019 Elsevier Masson SAS. All rights reserved.

Introduction

Ageing human immunodeficiency virus (HIV)-infected patients present with a high prevalence of age-related comorbidities,

including cardiovascular disease (CVD) and those associated with increased CV risk, such as diabetes, hypertension, dyslipidaemia and chronic kidney disease (CKD) [1–3]. This prevalence is higher than that observed in HIV-negative subjects [2], and could result from impaired levels of metabolic parameters commonly observed in such patients as well as from other classic factors, such as smoking, age, and familial histories of diabetes and CVD. Atherogenic dyslipidaemia [low levels of high-density lipoprotein (HDL) and high levels of triglycerides (TG)] can also contribute to

* Corresponding author at: Faculty of medicine Saint-Antoine site, 27, rue Chaligny 75571 Paris cedex 12, France.

E-mail address: jacqueline.capeau@inserm.fr (J. Capeau).

¹ These authors contributed equally to the study.

CVD risk in these patients, and may be assessed by measuring TG/HDL ratios [4,5].

Impaired levels of metabolic factors could result from HIV infection itself, but also from antiretroviral treatment (ART). In particular, increased risks of diabetes, dyslipidaemia and/or CVD have been reported in patients receiving a protease inhibitor (PI) boosted by ritonavir (r) [6–9]. In addition, increased risks of diabetes, hypertension, hypercholesterolaemia and low HDL have been associated with previous treatment with first-generation non-nucleoside analogues, such as stavudine and zidovudine, and possibly related to long-lasting alterations in adipose tissue distribution [10–13].

Previously, our team evaluated the risk of diabetes in 1046 HIV-infected patients from the APROCO-COPILOTE cohort who started treatment in 1997–1998 with a first-generation PI-containing regimen, which were regularly followed and ultimately evaluated 10 years after inclusion. A high incidence of diabetes was observed, linked to the use of some ART agents, including indinavir, stavudine, zidovudine and didanosine [11]. Moreover, our team also previously reported that 352 patients from the same cohort, upon close evaluation 11 years after inclusion and despite having good immunovirological status, presented with increased levels of inflammation and immune activation compared with healthy controls [14].

The present report of the same group of 352 patients here presents additional findings on a panel of biomarkers evaluating oxidative stress [lipoprotein-associated phospholipase A2 (Lp-PLA2), oxidized low-density lipoprotein (oxLDL), oxLDL/LDL ratios, paraoxonase and arylesterase activities], insulin resistance and adipokines [adiponectin, leptin, homoeostasis model assessment of insulin resistance (HOMA-IR) and plasminogen activator inhibitor 1 (PAI-1)]. Our main objective was to search for any association between these biomarkers and diabetes, LDL-linked and atherogenic dyslipidaemias, hypertension, CKD and previous CV events in addition to inflammation and immune activation.

Patients and methods

Patients

The study population has previously been described elsewhere [14]. Briefly, the 352 patients included here were from the French APROCO-COPILOTE cohort, which enrolled patients between 1997 and 1999 when they initiated PI-containing ART. All received first-generation ART, stavudine and/or zidovudine in combination with indinavir, nelfinavir, saquinavir or full-dose ritonavir.

The ethics committee of Cochin Hospital in Paris, France, approved the study, and informed consent was obtained from all participants. Patients were prospectively followed using standardized case-report forms, and all underwent complete clinical and biological evaluations after a median duration of 11 years of follow-up [interquartile range (IQR): 10–12]. All patients' characteristics (including clinical evaluation of fat distribution), metabolic and adipose-related parameters, biomarkers of oxidative stress, and inflammatory and immune activation biomarkers were recorded or measured during the same final visit. Thus, the median ART duration was 12.3 years (IQR: 11.4–13.9), CD4 cell count was 620 cells/mm³ (IQR: 453–790) and CD4/CD8 ratio was 0.81 (IQR: 0.56–1.10), with a viral load (VL) < 50 and < 400 copies/mL for 91.5% and 97.2% of patients, respectively.

Diagnosis of comorbidities

Comorbidities were also assessed at the final visit. Diabetes was diagnosed by fasting glycaemia ≥ 7.0 mmol/L and/or treatment,

impaired fasting glucose (IFG) by glycaemia at 6.1–6.9 mmol/L, hypertension by systolic blood pressure (SBP) ≥ 140 mmHg and/or diastolic blood pressure (DBP) ≥ 90 mmHg, LDL-linked dyslipidaemia by LDL ≥ 4.1 mmol/L and/or statin treatment, CKD by Modification of Diet in Renal Disease (MDRD) Study equation with an estimated glomerular filtration rate (eGFR) ≤ 60 mL/min, and atherogenic dyslipidaemia by a TG (mg/dL)/HDL (mg/dL) ratio ≥ 3.5 in men and ≥ 2.5 in women [5]. Insulin resistance was also evaluated by HOMA-IR score: fasting glycaemia (mmol/L) \times fasting insulinaemia (mU/L)/22.5 ≥ 2.5 . The lipid accumulation product (LAP) index was used as a surrogate marker of steatosis [15]: for men, waist circumference (WC) – 65 \times TG (mmol/L); and for women, WC – 58 \times TG (mmol/L). Also addressed were any previous CV events, defined as a diagnosis of cardiac or vascular disease reported in the patient's medical records during the 11-year follow-up.

Biomarkers

Whole blood was recovered at the final visit with/without ethylenediaminetetraacetic acid (EDTA) or citrate, and the plasma/serum was stored at -80° until assayed. The following markers were also analyzed:

- oxidative stress biomarkers as oxLDL by enzyme-linked immunosorbent assay (ELISA) (Mercodia, Uppsala, Sweden); Lp-PLA2 by immunoturbidimetric assay (Le PLAC Test, Eurobio, Courtaboeuf, France) adapted for an automated chemistry analyzer (AU480, Beckman Coulter, Brea, CA, USA); and antioxidant HDL-linked paraoxonase activities using hydrolysis rates of its two substrates, paraoxon (PON) and phenylacetate (arylesterase, ARE), as previously described [16];
- inflammatory and immune-activation biomarkers, high-sensitivity interleukin 6 (hsIL-6) and soluble CD14 (sCD14) by ELISA (Quantikine[®], R&D Systems, Abingdon, OX, UK); high-sensitivity C-reactive protein (hsCRP) by immunonephelometry (IMMAGE[®] Immunochemistry System, Beckman Coulter); D dimer (plasma) and $\beta 2$ microglobulin (B2M) by enzyme-linked fluorescent assay (ELFA) (VIDAS[®], Biomérieux, Marcy-l'Étoile, France); and cystatin C by immunoturbidimetry (Architect, Abbott Laboratories, Abbott Park, IL, USA); and;
- adipose-derived and metabolic biomarkers by ELISA, including leptin (Quantikine, R&D Systems), adiponectin (ALPCO Diagnostics, Salem, NH, USA) and PAI-1 (Diagnostica Stago, Asnières-sur-Seine, France).

Statistics

Continuous variables are presented as medians and 25th–75th percentile values (IQR), and categorical variables as numbers (*n*) and proportions (%). Prevalences of each comorbidity and multimorbidity were presented with 95% confidence intervals (CI).

Correlations between quantitative parameters were assessed by calculation of item-to-item Spearman correlation coefficients, whereas Kruskal–Wallis, Wilcoxon rank-sum and Fisher's exact tests were used to compare continuous and categorical variables between patients with vs without each comorbidity or condition. Given the number of comorbidities evaluated (diabetes, LDL-related and atherogenic dyslipidaemias, hypertension, CKD, previous CV events), differences were considered statistically significant when the two-sided *P*-value was < 0.01 . These variables were also compared after the exclusion of patients taking statins.

In addition, levels of metabolic parameters and biomarkers were evaluated according to ongoing treatment with antiretroviral agents (PIs, efavirenz, nevirapine). For multivariate analyses of

levels of oxLDL and oxLDL/LDL ratios, values for age, gender, body mass index (BMI) and inflammation were adjusted for three biomarkers: hsCRP, hsIL-6 and D dimer.

All analyses were performed by a dedicated statistician using either R version 3.4.2 software (R Foundation for Statistical Computing, Vienna, Austria) or SAS version 9.4 software (SAS Institute Inc., Cary, NC, USA).

Results

Patients' main clinical and HIV-related characteristics at the final visit and levels of metabolic and adipose-derived biomarkers are presented in Table 1. Patients were aged (median) 49 years (IQR: 45–56) with a median BMI of 23 kg/m² (IQR: 21.1–25.4). In these long-term ART patients, clinical features of lipodystrophy, associated mainly with first-generation ART regimens [17], were often evident: 27% (*n* = 96) presented with clinical signs of lipoatrophy; 11% (*n* = 37) showed trunk fat hypertrophy; and 22% (*n* = 76) had both peripheral lipoatrophy and trunk fat hypertrophy. The median SBP level was 123 mmHg (IQR: 118–137) with a median DBP of 80 mmHg (IQR: 70–80). In addition, 97% had received nucleoside analogues, 52% had taken PIs, 47% had been treated with non-nucleoside analogues (21% efavirenz, 26% nevirapine) and 2% with an integrase inhibitor (raltegravir).

The prevalences of metabolic comorbidities were high according to age: previous CV events, 9% (32 out of 352, 95% CI: 7.5–10.5%); diabetes, 11% (37 out of 346, 95% CI: 8–14%); hypertension, 28% (90 out of 322, 95% CI: 23–33%); LDL-related dyslipidaemia, 27.5% (90 out of 328, 95% CI: 23–32%); atherogenic dyslipidaemia, 8% (29 out of 352, 95% CI: 5–11%); and CKD, 9% (30 out of 348, 95% CI: 6–11%). The mean HOMA-IR score was 2.0 (IQR: 1.4–3.2), and 36% (122 out of 339) were insulin-resistant.

Regarding the prevalence of multimorbidity, including previous CV events, diabetes, LDL-related and atherogenic dyslipidaemias, hypertension and CKD, 42.6% (95% CI: 40–45.2%) had no comorbidity, 34.9% (95% CI: 32.4–37.4%) had one, 16.5% (95% CI: 14.5–18.5%) had two, 4.3% (95% CI: 3.2–5.4%) had three and 1.7% (95% CI: 1.0–2.4%) had four.

Biomarker levels according to diabetes

Patients with diabetes had, as expected, higher BMIs (25.7 vs. 22.9 kg/m²; *P* = 0.005) and waist-to-hip ratios (WHRs; 0.99 vs. 0.95; *P* < 0.001). Body fat distribution differed, with 61% of diabetic patients presenting with trunk fat accumulation (alone or associated with peripheral lipoatrophy) vs. 32% of non-diabetic subjects. Only 9% had peripheral lipoatrophy on its own vs. 29% of non-diabetic subjects (*P* < 0.001). Diabetic patients also had increased levels of TG, TG/HDL ratios and HOMA-IR scores (5.6 vs. 1.9; *P* < 0.00001), reduced levels of adiponectin (3.4 vs. 6.1 mg/L; *P* < 0.0001), but higher oxLDL/LDL ratios (17.8 vs. 15.6 U/L-mmol; *P* < 0.01) and CRP levels (3.85 vs. 1.94 mg/L; *P* < 0.005; Table 2). On multivariate analyses adjusted for age, gender, BMI and inflammation, diabetes was associated with increased oxLDL/LDL ratios [adjusted odds ratio (ORa): 1.15, 95% CI: 1.05–1.25]. Thus, diabetes was independently associated with oxidative stress and inflammation (Fig. 1).

Patients with impaired fasting glucose (IFG) were also evaluated and compared with normoglycaemic patients: BMI did not differ and WHR was only slightly increased in IFG subjects (0.96 vs. 0.94; *P* = 0.04). Interestingly, oxLDL levels were 60.2 U/L in IFG patients vs. 49.9 U/L in normoglycaemic subjects (*P* = 0.01). HOMA-IR Scores were also higher (3.3 vs. 1.8; *P* < 0.0001), although CRP levels did not differ. In addition, glycaemia did not correlate with biomarkers of inflammation, immune activation or oxidative stress except for cystatin C (*R* = −0.16; Table 3).

Table 1

Characteristics of patients included from the French APROCO-COPILOTE cohort, including levels of metabolic and adipose-derived parameters and biomarkers at the final follow-up visit.

HIV-infected patients (<i>n</i> = 352)	Median (IQR) or <i>n</i> (%) ^a
Demographic, blood pressure and anthropometric parameters	
Age (years), current	49 (45–56)
Male gender	287 (81.5%)
Body mass index (kg/m ²)	23.0 (21.1–25.4)
Waist circumference (cm)	88 (81–95)
Waist-to-hip ratio	0.95 (0.90–1.00)
No fat redistribution (<i>n</i> = 349)	140 (40%)
Lipoatrophy	96 (27%)
Lipohypertrophy	37 (11%)
Lipoatrophy + lipohypertrophy	76 (22%)
Systolic blood pressure (mmHg; <i>n</i> = 322)	123 (118–137)
Diastolic blood pressure (mmHg; <i>n</i> = 322)	80 (70–80)
Never smokers (<i>n</i> = 342)	112 (32.7%)
Current smokers	138 (40.4%)
Past smokers	92 (26.9%)
Hepatitis C virus co-infection (<i>n</i> = 333)	63 (18.9%)
HIV infection-related characteristics	
Current CD4 cell count (cells/mm ³)	620 (453–790)
Current CD8 cell count (cells/mm ³)	756 (544–1025)
Current CD4/CD8 ratio	0.81 (0.56–1.10)
Current HIV-1 RNA < 50 copies/mL	322 (91.5%)
Total duration of ART (years)	12.3 (11.5–13.8)
Current ART (<i>n</i> = 349)	
Protease inhibitors	183 (52.4%)
Nucleoside analogues	337 (96.6%)
Non-nucleoside analogues	164 (46.7%)
Efavirenz	74 (21.2%)
Nevirapine	90 (25.8%)
Integrase inhibitors	7 (2%)
Metabolic and adipose-related parameters	
Total cholesterol (mmol/L; <i>n</i> = 341)	4.94 (3.96–5.93)
LDL cholesterol (mmol/L; <i>n</i> = 328)	2.91 (2.01–3.74)
HDL cholesterol (mmol/L; <i>n</i> = 334)	1.16 (0.87–1.49)
Triglycerides (mmol/L; <i>n</i> = 341)	1.56 (1.02–2.13)
Glycaemia (mmol/L; <i>n</i> = 347)	5.3 (4.9–5.8)
Insulin (mU/L; <i>n</i> = 344)	8.5 (6.3–12.6)
HOMA-IR (<i>n</i> = 339)	2.02 (1.42–3.16)
PAI-1 (ng/mL; <i>n</i> = 331)	31.9 (19.7–50.9)
Adiponectin (mg/L; <i>n</i> = 347)	5.75 (3.53–9.6)
Leptin (pt; <i>n</i> = 342)	4.14 (2.41–8.04)
eGFR (mL/min/1.73 m ² ; <i>n</i> = 328)	93 (79–109)
Bilirubin (mol/L; <i>n</i> = 329)	6.8 (4.6–11.3)
Oxidative stress biomarkers	
Lp-PLA2 (ng/mL; <i>n</i> = 347)	183 (153–213)
oxLDL (U/L; <i>n</i> = 350)	50.6 (38.3–64.5)
oxLDL/LDL (U/mmol; <i>n</i> = 326)	15.9 (13.2–18.5)
Arylesterase (mmol/L-min; <i>n</i> = 350)	21.9 (17.5–27.3)
Paraoxonase (mol/L-min; <i>n</i> = 350)	133 (67–204)
Inflammatory biomarkers	
hsCRP (mg/L; <i>n</i> = 347)	2.1 (0.9–4.5)
hsIL-6 (pg/mL; <i>n</i> = 347)	1.3 (0.7–2.6)
D dimer (ng/mL; <i>n</i> = 331)	252 (177–374)
Immune activation biomarkers	
sCD14 (ng/mL; <i>n</i> = 347)	1356 (1027–1818)
β2 microglobulin (mg/L; <i>n</i> = 347)	2.4 (2.0–3.1)
Cystatin C (mg/L; <i>n</i> = 347)	0.93 (0.82–1.10)

HIV: human immunodeficiency virus; IQR: interquartile range; RNA: ribonucleic acid; ART: antiretroviral therapy; LDL/HDL: low-density/high-density lipoprotein; HOMA-IR: homeostasis model assessment of insulin resistance; PAI-1: plasminogen activator inhibitor 1; eGFR: estimated glomerular filtration rate; Lp-PLA2: lipoprotein-associated phospholipase A2; oxLDL: oxidized LDL; hsCRP/hsIL-6: high-sensitivity C-reactive protein/interleukin 6; sCD14: soluble CD14.

^a Continuous or categorical variables.

Patients with insulin resistance had higher levels of CRP and oxLDL/LDL ratios compared with those without insulin resistance [2.9 vs. 1.8 mg/L (*P* < 0.005) and 16.4 vs. 15.4 U/L-mmol/L (*P* < 0.01), respectively]. PAI-1, a marker of insulin resistance and increased CVD risk, correlated positively with oxidative stress (oxLDL and oxLDL/LDL ratio; Table 3), whereas adiponectin

Table 2
Median levels of oxidative stress, inflammation and immune-activation biomarkers in patients with/without diabetes, low-density lipoprotein (LDL)-related and atherogenic dyslipidaemias (DYSLIP), hypertension and chronic kidney disease (CKD) and previous CV events.

Oxidative stress biomarkers	Diabetes Yes/No n = 37 n = 310	LDL-related DYSLIP Yes No n = 90 n = 238	Atherogenic DYSLIP Yes/No n = 29 n = 323	Hypertension Yes/No n = 90 n = 232	CKD Yes No n = 30 n = 318	Previous CV events Yes No n = 32 n = 320
Lp-PLA2 (ng/mL)	166 186	181 184	192 183	178 185	181 184	181 184
oxLDL (U/L)	54.0 50.2	61.5 48.3 ^c	65.5 49.8 ^c	52.5 49.8	49.5 50.6	49.5 50.6
oxLDL/LDL (U/mmol)	17.8 15.6 ^b	15.0 16.2	18.7 15.5 ^c	16.1 15.3	15.9 15.9	18.1 15.7
Arylesterase (mmol/L-min)	21.2 22.2	23.6 21.6	22.8 21.7	22.4 21.6	24.8 21.6	21.0 22.2
Paraoxonase (mol/L-min)	92 136	119 141	175 132	158 128	73 145	133 133
Inflammatory biomarkers						
hsCRP (mg/L)	3.85 1.94*	2.25 1.94	3.81 1.93 ^a	2.19 2.11	2.83 1.96	2.95 1.96
hsIL-6 (pg/mL)	1.43 1.28	1.12 1.34	2.11 1.25	1.22 1.27	1.92 1.28	1.91 1.25
D dimer (ng/mL)	269 252	237 259	234 252	289 249	287 252	287 251
Immune activation biomarkers						
sCD14 (ng/mL)	1260 1360	1290 1378	1452 1353	1442 1330	1472 1329	1343 1356
Cystatin C (mg/L)	0.91 0.93	0.93 0.92	0.95 0.92	0.94 0.92	1.31 0.91 ^c	1.13 0.92 ^c
β2 microglobulin(mg/L)	2.40 2.40	2.30 2.40	2.65 2.40	2.35 2.40	3.80 2.40 ^c	3.25 2.40 ^b

PCE: previous cardiovascular events; Lp-PLA2: lipoprotein-associated phospholipase A2; oxLDL: oxidized low-density lipoprotein; hsCRP/hsIL-6: highsensitivity. C-reactive protein/interleukin 6; sCD14: soluble CD14.

^a $P < 0.01$.

^b $P < 0.001$.

^c $P < 0.0001$.

correlated negatively with CRP and oxidative stress (Lp-PLA2, oxLDL and oxLDL/LDL ratio; [Table 3](#)).

The median LAP value was 39 (IQR: 20–60). As expected, LAP was associated with metabolic parameters: HOMA-IR 0.421 ($P < 0.0001$); adiponectin -0.279 ($P < 0.0001$); leptin 0.336 ($P < 0.0001$); and PAI-1 0.319 ($P < 0.0001$). LAP was also associated with oxidative stress [oxLDL 0.439 ($P < 0.0001$) and oxLDL/LDL ratio 0.384 ($P < 0.0001$)] and with inflammation [CRP 0.251 ($P < 0.0001$) and IL-6 0.184 ($P = 0.003$)].

Biomarker levels according to LDL-related or atherogenic dyslipidaemia

Patients with vs without LDL-related dyslipidaemia had increased values of oxLDL (61.5 vs. 48.3 U/L; $P < 0.0001$; [Table 2](#)). On multivariate analyses adjusted for age, gender, BMI and inflammation, LDL-related dyslipidaemia was associated with increased oxLDL (ORa: 1.05, 95% CI: 1.03–1.07) and decreased oxLDL/LDL ratio (ORa: 0.90, 95% CI: 0.82–0.97; [Fig. 1](#)). Because statins alter inflammatory/immune-activation profiles, these parameters were also evaluated in patients not receiving statins. In these patients ($n = 291$), those with LDL-related dyslipidaemia exhibited higher levels of oxLDL (64.4 vs. 48.3 U/L; $P < 0.0001$) and ARE (23.8 vs. 21.6 mmol/L-min; $P < 0.01$), and lower levels of D dimer (201 vs. 259.5 ng/mL; $P < 0.005$) and oxLDL/LDL ratios (14.5 vs. 16.2 U/L-mmol; $P < 0.001$) than those without dyslipidaemia.

Thus, LDL-related dyslipidaemia was not related to treatment with PI and was associated with oxidative stress, but not with inflammation or immune activation. Moreover, LDL levels were related to biomarkers of oxidative stress (Lp-PLA2, oxLDL, oxLDL/LDL ratios, ARE; [Table 3](#)), but not of inflammation and immune activation [except for B2M ($R = -0.14$)].

Patients with atherogenic dyslipidaemia had higher levels of oxidative stress biomarkers: oxLDL (65.5 vs. 49.8 U/L; $P < 0.0001$); oxLDL/LDL ratio (18.7 vs. 15.5 U/L-mmol; $P < 0.0001$); and CRP (3.81 vs. 1.93 mg/L; $P = 0.003$; [Table 2](#)). On multivariate analyses adjusted for age, gender, BMI and inflammation, atherogenic dyslipidaemia was associated with increased oxLDL (ORa: 1.05, 95% CI: 1.02–1.08) and oxLDL/LDL ratio (ORa: 1.25, 95% CI: 1.11–1.40; [Fig. 1](#)), and also with PI use ($P = 0.01$). Thus, atherogenic

dyslipidaemia was independently associated with oxidative stress and inflammation.

HDL was negatively correlated with biomarkers assessing oxidative stress (Lp-PLA2, oxLDL, oxLDL/LDL ratios), inflammation (CRP, IL-6) and immune activation (cystatin C, B2M), and positively with PON and ARE ([Table 3](#)), whereas TG levels were positively associated with most of the biomarkers negatively associated with HDL (oxLDL, oxLDL/LDL ratios, CRP, cystatin C, B2M). TG/HDL ratios correlated positively with anthropometric parameters, WHR ($r = 0.293$, $P < 0.0001$) and HOMA-IR ($R = 0.309$, $P < 0.0001$), and negatively with adiponectin ($R = -0.307$, $P < 0.0001$). In addition, TG/HDL ratios were associated with biomarkers of oxidative stress (Lp-PLA2, oxLDL, oxLDL/LDL ratios) as well as inflammation and immune activation (CRP, IL-6, cystatin C, B2M; [Table 3](#)).

Biomarker levels according to hypertension, CKD or previous cardiovascular events

Patients with hypertension presented with higher values of HOMA, BMI and WHR (data not shown), but had similar levels of oxidative stress and inflammatory/immune-activation biomarkers ([Table 2](#)). In multivariate analyses, oxLDL and oxLDL/LDL ratio were not associated with hypertension (ORa: 1.01, 95% CI: 0.99–1.03 and ORa: 1.02, 95% CI: 0.95–1.08, respectively). SBP and DBP correlated (with low Spearman coefficients) with oxLDL ([Table 3](#)), but not with other biomarkers.

CKD was associated with no biomarkers except for cystatin C and B2M, as expected, as these are markers excreted by the kidney ([Table 2](#)). On multivariate analyses, oxLDL and oxLDL/LDL ratio were not associated with CKD (ORa: 1.00, 95% CI: 0.97–1.02 and ORa: 1.01, 95% CI: 0.91–1.12, respectively). Creatinine levels correlated with IL-6 and oxLDL ($r = 0.15$ and $r = 0.16$, respectively; [Table 3](#)) in addition to cystatin C and B2M.

Patients with a history of CV events during follow-up were of older age (54 vs. 49 years; $P = 0.002$), and eGFR was decreased (75.6 vs. 87.3 mL/min; $P = 0.003$) while, as expected, B2M and cystatin C were increased. However, inflammatory and oxidative stress biomarkers did not differ ([Table 2](#)). On multivariate analyses adjusted for age, BMI, gender and inflammation, oxLDL and oxLDL/LDL ratio were not associated with previous CV events




Fig. 1. Levels of oxidized low-density lipoprotein (oxLDL) and oxLDL/LDL ratios according to presence or absence of LDL-related or atherogenic dyslipidaemias or diabetes in patients from the APROCO-COPILOTE study. Medians and interquartile ranges (IQRs) are shown as boxplots and as individual values. Results were adjusted for age, gender, body mass index and inflammatory markers.

(ORa: 0.99, 95% CI: 0.96–1.02 and ORa: 1.07, 95% CI: 0.99–1.16, respectively).

Biomarker levels according to antiretroviral molecules

Patients treated with PIs had lower levels of HDL (1.23 vs. 1.36 mmol/L; $P = 0.002$), and higher levels of TG (1.77 vs. 1.35 mmol/L; $P < 0.0001$), IL-6 (1.6 vs. 1.1 pg/mL; $P < 0.0001$), cystatin C (0.98 vs. 0.87 mg/L; $P < 0.0001$), B2M (2.6 vs. 2.3 mg/L; $P < 0.0001$) and oxLDL/LDL ratio (16.4 vs. 14.9 U/L-mmol/L; $P = 0.004$) compared with those following a PI-free regimen.

Regarding non-nucleoside reverse transcriptase inhibitors (NNRTIs), 21% were receiving an efavirenz (EFV)-based and 26% a nevirapine-based regimen of combination ART (cART). Levels of eGFR were higher in those receiving vs. not receiving EFV (93 vs. 85 mL/min; $P = 0.005$) whereas levels of D dimer were lower (204 vs. 266 ng/mL, respectively; $P = 0.004$). Treatment with vs without nevirapine was associated with higher HDL (1.45 vs. 1.21 mmol/L, respectively; $P = 0.00001$), but with lower PAI-1 (26.5 vs. 34.5 ng/mL, respectively; $P = 0.006$), cystatin C (0.86 vs. 0.96 mg/L, respectively; $P = 0.002$) and oxLDL/LDL ratio (14.3 vs. 16.3 U/L-mmol/L, respectively; $P = 0.0001$).

Biomarker levels according to smoking, body fat redistribution and HIV infection severity

No oxidative stress, inflammation or immune-activation biomarkers differed between groups of patients according to fat redistribution (absence or presence of lipoatrophy and/or trunk fat hypertrophy). However, oxLDL and oxLDL/LDL ratios were positively associated with WHR (0.319 and 0.266, respectively; both $P < 0.0001$) and oxLDL was associated with BMI (0.21; $P < 0.0001$). In addition, CRP was associated with WHR and BMI [0.224 ($P = 0.0003$) and 0.200 ($P = 0.0002$), respectively] whereas, as expected, adiponectin was negatively related to WHR and BMI [−0.200 ($P = 0.0002$) and −0.324 ($P < 0.0001$), respectively]. Thus, increased trunk fat was associated with oxidative stress and inflammation as well as with dysmetabolic features.

As regards smoking, increased levels of IL-6 in current vs. never and past smokers had previously been reported [14]. Moreover, oxLDL/LDL ratios differed across the three groups ($P = 0.03$) and between past and never smokers (16.76 U/mmol, 95% CI: 13.45–19.49 vs. 15.14 U/mmol, 95% CI: 12.62–17.62, respectively; $P = 0.01$).

Levels of biomarkers did not differ in patients with VL < or > 50 copies/mL, CD4/CD8 ratios < or > 0.4,

Table 3
Correlations between metabolic, anthropometric and blood pressure parameters and biomarkers assessing oxidative stress, inflammation and innate immune activation in 352 human immunodeficiency virus (HIV)-infected patients.

	Oxidative stress biomarkers					Inflammatory biomarkers			Immune-activation biomarkers		
	Lp-PLA2	oxLDL	oxLDL/LDL	PON	ARE	hsCRP	hsIL-6	D dimer	sCD14	Cystatin C	B2M
Glucose metabolism-related parameters											
Glycaemia	-	-	-	-	-	-	-	-	-	-0.16	-
										<i>P</i> < 0.05	
Insulin	-	-	0.16	-	-	0.17	-	-	-	0.16	-
			<i>P</i> < 0.005			<i>P</i> < 0.005				<i>P</i> < 0.005	
HOMA	-	-	0.16	-	-	0.14	-	-	-	-	-
			<i>P</i> < 0.005			<i>P</i> < 0.01					
PAI-1	-	0.18	0.23	-	-	-	-	-	-	-	-
		<i>P</i> < 0.001	<i>P</i> < 0.0001								
Adipo	-0.23	-0.28	-0.35	-	-	-0.14	-	-	-	-	-
	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.0001			<i>P</i> < 0.01					
Leptin	-	-	-	0.16	-	-	-	0.16	-	-	-
				<i>P</i> < 0.005				<i>P</i> < 0.005			
Lipid parameters											
LDL	0.27	0.68	-0.32	-	0.22	-	-	-	-	-	-0.14
	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.0001		<i>P</i> < 0.0001						<i>P</i> < 0.01
HDL	-0.37	-0.29	-0.47	0.17	0.17	-0.16	-0.27	-	-	-0.34	-0.33
	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.005	<i>P</i> < 0.002	<i>P</i> < 0.005	<i>P</i> < 0.0001			<i>P</i> < 0.0001	<i>P</i> < 0.0001
TG/HDL	0.23	0.42	0.44	-	-	0.20	0.20	-	-	0.30	0.28
	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.0001			<i>P</i> < 0.001	<i>P</i> < 0.001			<i>P</i> < 0.0001	<i>P</i> < 0.0001
TG	-	0.41	0.35	-	0.15	0.20	-	-	-	0.20	0.19
		<i>P</i> < 0.0001	<i>P</i> < 0.0001		<i>P</i> < 0.01	<i>P</i> < 0.001				<i>P</i> < 0.001	<i>P</i> < 0.001
Anthropometric parameters											
BMI	-	0.21	-	-	-	0.20	-	-	-	-	-
		<i>P</i> < 0.0001				<i>P</i> < 0.005					
WHR	-	0.32	0.27	-	-	0.22	-	-	-	-	-
		<i>P</i> < 0.0001	<i>P</i> < 0.0001			<i>P</i> < 0.001					
Blood pressure											
Diastolic	-	0.18	-	-	-	-	-	-	-	-	-
		<i>P</i> < 0.005									
Systolic	-	0.15	-	-	-	-	-	-	-	-	-
		<i>P</i> < 0.01									
Other biochemical parameters											
Creatinine	-	0.15	-	-	-	-	0.16	-	-	0.41	0.33
		<i>P</i> < 0.005					<i>P</i> < 0.005			<i>P</i> < 0.0001	<i>P</i> < 0.0001

NB: Only *P*-values < 0.01 are presented (Spearman's correlation coefficients).

Lp-PLA2: lipoprotein-associated phospholipase A2; oxLDL: oxidized LDL; PON: paraoxonase activity; ARE: arylesterase activity; hsCRP/hsIL-6: high-sensitivity C-reactive protein/interleukin 6; sCD14: soluble CD14; B2M: β 2 microglobulin; HOMA: homoeostasis model assessment; PAI-1: plasminogen activator inhibitor 1; Adipo: adiponectin; LDL/HDL: low-density/high-density lipoprotein; TG: triglycerides; BMI: body mass index; WHR: waist-to-hip ratio.

CD4 < or > 350 cells/mm³, < or > 500 copies/mL at the last follow-up visit, or CD4 < or > 200 cells/mm³ on entry into the study, as previously reported for inflammatory/immune-activation biomarkers [14].

Discussion

The present study has found that, in long-term HIV-infected and ART-controlled patients from the *Agence Nationale de Recherches sur le Sida et les Hépatites Virales* (ANRS; French National Agency for Research on AIDS and Viral Hepatitis) CO3 APROCO-COPILOTE cohort, diabetes and atherogenic dyslipidaemia are each independently associated with oxidative stress and inflammation, whereas LDL-related dyslipidaemia and IFG are associated with only oxidative stress.

Our study population is representative of the long-term HIV-infected and treated population with prevalent age-related comorbidities, as observed in our participants and in the total group of 1227 patients [18]. Similar prevalences of comorbidities were reported in 876 patients of the French Chronic Immune Activation and Senescence (CIADIS) study (ANRS CO3 Aquitaine cohort) who were aged 50.5 years [1]. In addition, in a cohort of 524 HIV-infected patients aged 52 years from the AGEHIV study, the prevalence of CVD-linked comorbidities was similar to that observed in our present study, yet was greater overall than that

observed in a well-matched control group of 540 HIV-negative subjects [2].

Regarding diabetes, a prevalence of 11% was found in our cohort of adults (median age: 49 years) with normal BMI (median 23 kg/m²) and receiving long-term ART (median duration: 12.3 years), which is a high proportion compared with the general population. However, this prevalence is comparable to that reported in the ANRS CO3 Aquitaine cohort (12.8%), which also included long-term ART patients (median duration: 13.2 years) [1], albeit higher than that reported in the AGEHIV cohort (5.5%) [2], which had a median ART duration of 10 years. This difference might account for the greater prevalence of diabetes, as those treated for longer durations had probably received first-generation ART, which was associated with a greater risk of diabetes [11,12]. In fact, some ART agents are clearly involved in addition to ageing, family history of diabetes, ethnic origin, overweight/obesity, lipodystrophy and dysmetabolic profiles [19]. Interestingly, our diabetic patients frequently presented with a lipodystrophic phenotype and trunk fat redistribution, which accords with the deleterious role of trunk fat in metabolic disorders and the insulin resistance that often precedes overt diabetes [20]. Indeed, long-lasting alterations in adipose tissue distribution have recently been confirmed in patients with previous exposures to stavudine, zidovudine and didanosine in the Copenhagen Comorbidity in HIV Infection (COCOMO) study [13]. Moreover, a recent report of HIV-

infected ART-controlled subjects from sub-Saharan Africa found an increased prevalence of diabetes, with increased LDL levels, which was attributed to ART [21].

LDL-linked dyslipidaemia, also attributed in part to ART, is frequent in HIV-infected patients [22]. Previous exposures to thymidine analogues and didanosine have been associated with excess risks of hypercholesterolaemia and low HDL levels [13], while PIs have been associated with dyslipidaemia [17]. In addition, our present study has shown that 9% had atherogenic dyslipidaemia, which is predictive of increased CV risk as well as insulin resistance [4,5] and diabetes [23]. Furthermore, in HIV-infected patients, high TG/HDL ratios have been shown to predict risk of new-onset diabetes independently of other traditional risk factors [24], whereas atherogenic dyslipidaemia has recently proved to be more prevalent in HIV-infected vs non-infected subjects and associated with low adiponectin [25]. In our present study population, TG/HDL ratio was associated positively with diabetes and insulin resistance (by HOMA-IR and PAI-1), and negatively with adiponectin. In HIV-negative subjects, this ratio has been associated with CRP and oxLDL [26] and, in our present study, was associated with oxidative stress, immune activation and inflammatory markers, which could reflect causal relationships that contribute, together with insulin resistance, to higher CVD risk.

Regarding oxidative stress, it was observed that oxLDL levels were higher in subjects with trunk fat accumulation, as indicated by WHR, which accords with the deleterious role of this fat depot and its involvement in the insulin-resistant phenotype [27]. Also, an association between markers of oxidative stress and insulin resistance has been reported in the general population, suggesting that oxidative stress is an important trigger of insulin resistance [28], an association that was confirmed in our study with our patients who presented with high HOMA-IR scores according to their BMI. Our study also found that diabetes is associated with oxidative stress independently of inflammation. Interestingly, patients with IFG had increased levels of oxLDL as well as increased HOMA-IR scores, suggesting a causal relationship. In addition, while increased levels of oxLDL have been positively associated with type 2 diabetes (T2D) in the general population [29], no such significant association was found between oxLDL and incident T2D in our study patients. In fact, our analysis indicating increased levels of oxLDL in IFG patients could be suggesting that oxidative stress plays a greater role in the development of diabetes in HIV-infected patients than in the general population.

Our study also found that a surrogate marker of liver steatosis, LAP, is associated with oxidative stress and inflammation. Increased oxidative stress has been proposed as a critical factor in the pathogenesis of non-alcoholic fatty liver disease [30].

In HIV-infected ART patients, PAI-1, a marker reflecting insulin resistance, was also associated with oxidative stress [31]. Otherwise, high levels of PAI-1 were associated with a risk of myocardial infarction independently of CVD risk factors, HIV parameters and ART, suggesting that PAI-1 may be used for risk stratification and prediction of CVD [32].

In our present cohort, Lp-PLA2 was associated with lipid parameters and adiponectin, although its levels did not differ with the presence or absence of the evaluated comorbidities. Increased levels of Lp-PLA2 have been associated with increased risk of coronary heart disease in the general population [33] and in HIV-infected patients [34,35], and Lp-PLA2 has been associated with TG, LDL, HDL and insulin in HIV-infected patients initiated with either darunavir/r or atazanavir/r [36]. However, in a population of ART-controlled patients with normal LDL, Lp-PLA2 was associated with sCD14, but not with lipid parameters or inflammatory biomarkers [34].

OxLDL is thought to play a central role in atherosclerosis development [35] and is clearly associated with atherosclerotic CVD [37]. OxLDL is increased in HIV-infected compared with HIV-negative individuals [38], and is associated with the Framingham coronary heart disease risk score and subclinical atherosclerosis as evaluated by carotid intima-media thickness (cIMT) [39]. These data argue for atherosclerosis being particularly related to an increased oxidative and inflammatory status in HIV patients. In addition, as reported here, oxidative stress is associated with LDL-related and atherogenic dyslipidaemias independently of inflammation. However, when levels of oxidative stress markers were evaluated in patients with a history of CV events during the 11-year follow-up, no increased levels were observed in multivariate analyses, perhaps due to the time lag since the occurrence of such events.

Importantly, statin treatment decreased oxidative stress biomarker levels, and increased antioxidant PON and ARE activities in the general population [40,41]. Also, in HIV-infected patients, statins reduced oxidative stress markers (Lp-PLA2, oxLDL), leading to a reduced number of coronary plaques independently of LDL levels [35,42]. These data further emphasize the deleterious impact of oxidative stress on the CV complications encountered by HIV-infected patients.

Taken as a whole, our study found that oxLDL and oxLDL/LDL ratios were strongly associated with diabetes and atherogenic dyslipidaemia independently of inflammation. In fact, a causal relationship with insulin resistance is suggested. Further studies are now required to validate their involvement in ageing HIV-infected patients. Moreover, these biomarkers were not related to the severity of the current HIV infection, as evaluated by HIV-1 VL, CD4/CD8 ratios or CD4 numbers. However, increased oxidative stress could be related to persistent effects of stavudine and/or zidovudine, drugs that are responsible for mitochondrial toxicity. Indeed, 80.5% of our study patients had received stavudine and 54% were taking zidovudine.

A previous study had shown that biomarkers of inflammation and immune activation were increased in patients in the APROCO-COPILOTE cohort [14]. In addition, as noted in the present study, they were also associated with diabetes and atherogenic dyslipidaemia, but not with the other evaluated comorbidities.

Several studies have suggested that, in the general population, diabetes is associated with increased inflammation [43]. In HIV-infected patients, inflammatory biomarkers were previously associated with the development or presence of diabetes [11,44–46]. In addition, the following markers have previously been associated with CVD risk in HIV-infected patients: increased levels of IL-6, CRP and/or D dimer with myocardial infarction [32,47,48], and sCD14 with progressive or pathological cIMT and calcified coronary plaques [49,50].

However, no consistent association was detected between hypertension and the tested biomarkers. Likewise, in the AGEHIV cohort, hypertension was not associated with markers of either inflammation or immune activation [51]. Regarding CKD, our study found that, except for cystatin C and B2M as expected, there was no difference in biomarker levels between patients with or without CKD. In the CIADIS study, only 4% of patients presented with CKD, and levels of CRP, IL-6 and sCD14 did not differ between patients with or without CKD, although a soluble weighted score of inflammation [including soluble tumour necrosis factor receptor 1 (sTNFR1) and D dimer] was independently associated with CKD [52]. In the ACTIVIH study, some biomarkers were associated with some comorbidities, but not with others, and different comorbid conditions presented with specific patterns of alteration [53]. Nevertheless, the absence of an association between the tested biomarkers and hypertension or CKD in our present study does not

preclude an association with other biomarkers related to oxidative stress or inflammation.

Some ART molecules can modify patients' metabolic profiles and, thus, could be involved in metabolic comorbidities. Indeed, our study found that PI treatment was, as expected, associated with differences in lipid levels, but also with biomarkers of oxidative stress, inflammation and immune activation that might participate in elevating CVD risk. Our study also confirmed the generally favourable effect of nevirapine on metabolic and inflammatory markers.

Nevertheless, it is necessary to acknowledge the limitations of our study. Regarding diabetes, hypertension, LDL-related and atherogenic dyslipidaemias and CKD, it was decided to consider only the situation at the time of evaluation and, therefore, only previous CV events were evaluated. Also, our study was cross-sectional in design and, as such, cannot address causality of the different biomarkers in the development of the various studied comorbidities.

In conclusion, in our present APROCO-COPILOTE cohort of long-term virologically suppressed patients, a high level of metabolic comorbidities was observed in association, to variable extents, with biomarkers of oxidative stress independently of inflammation. These biomarkers are most probably involved in diabetes and LDL-related and atherogenic dyslipidaemias, and less so with CKD and hypertension. Increased oxidative stress was observed in diabetic patients and those with IFG, and was also associated with insulin resistance. Moreover, there was an association with trunk fat accumulation that might be related to previous ART regimens, but also to HIV present in fat reservoirs and responsible for fat dysfunction and fibrosis [54].

Our findings suggest that oxidative stress could play a particular role in patients presenting with lipodystrophic features and insulin resistance and explain, at least in part, the high frequencies of some comorbidities, especially diabetes and dyslipidaemia, in HIV-infected ART-controlled patients. Thus, therapeutic strategies to control these comorbidities should target oxidative stress in addition to inflammation, and should also consider their differential associations with these biomarkers.

Funding

This work was supported by the French Agence Nationale de Recherches sur le Sida et les Hépatites Virales (ANRS) and Institut National de la Santé et de la Recherche Médicale.

Part of this work was presented as a poster at the 18th International Workshop on Comorbidities and Adverse Drug Reactions in HIV, held in September 2016 in New York City, NY, USA.

Disclosure of interest

The authors declare that they have no competing interest.

Acknowledgments

The authors would like to thank all participating patients, nurses and physicians at the study clinical sites. We wish to thank Elodie Heude and Lydie Doufoudji for their excellent technical expertise in testing markers at the Unit of Inflammatory and Metabolic Biomarkers in Tenon Hospital. The APROCO-COPILOTE cohort was supported by the ANRS and Collège des Universitaires de Maladies Infectieuses et Tropicales (CMIT).

The ANRS CO8 APROCO-COPILOTE Study Group included the following:

Scientific Committee

Steering Committee principal investigators: C. Lepout, F. Raffi; Methodology: G. Chêne, R. Salamon; Social sciences: J.P. Moatti, J. Pierret, B. Spire; Virology: F. Brun-Vézinet, H. Fleury, B. Masquelier†; Pharmacology: G. Peytavin, R. Garraffo

Other members: D. Costagliola, P. Dellamonica, C. Katlama, L. Meyer, D. Salmon, A. Sobel

Events validation committee

L. Cuzin, M. Dupon, X. Duval, V. Le Moing, B. Marchou, T. May, P. Morlat, C. Rabaud, A. Waldner-Combernoux

Project coordination

L. Hardel, P. Reboud

ANRS representatives

S. Couffin-Cadiergues, L. Marchand

Methodological, statistical and other contributions

A. Assuied, P. Carrieri, S. Habak, F. Couturier, C. Jadand, A. Perrier, M. Préau, C. Protopopescu

Promotion

Agence Nationale de Recherches sur le Sida et les Hépatites Virales (ANRS Action Coordonnée No. 7).

Other support

CMIT ex APPIT, Sidaction Ensemble contre le Sida and associated pharmaceutical compagnies (Abbott Laboratories, Boehringer Ingelheim, Bristol-Myers Squibb, GlaxoSmithKline, Gilead Sciences, Pfizer and Roche).

Clinical Centres (investigators)

Amiens (Pr J.L. Schmit), Angers (Dr J.M. Chenebault), Belfort (Dr J.P. Faller), Besançon (Pr C. Chirouze, Pr N. Magy-Bertrand, Pr P. Humbert), Bordeaux (Pr M. Dupon, Pr M. Longy-Boursier, Pr S. Morlat, Pr D. Neau), Bourg-en-Bresse (Dr P. Granier), Brest (Pr S. Ansart), Caen (Pr R. Verdon), Compiègne (Dr D. Merrien), Corbeil Essonne (Dr P. Chevojon), Créteil (Pr Y. Levy, Pr A. Sobel), Dijon (Pr L. Piroth), Garches (Pr C. Perronne), Lagny (Dr E. Froguel), Libourne (Dr J. Ceccaldi), Lyon (Pr C. Chidiac), Meaux (Dr V. Grégoire), Montpellier (Pr J. Reynes), Nancy (Pr T. May), Nantes (Pr F. Raffi), Nice (Pr J.G. Fuzibet, Pr P. Dellamonica), Orléans (Dr P. Arsac), Paris (Pr E. Bouvet, Pr F. Bricaire, Dr J. Monsonego, Pr P.M. Girard, Pr S. Herson, Pr C. Lepout, Pr G. Pialoux, Pr O. Sain, Pr D. Salmon, Dr P. Sellier), Poitiers (Pr P. Roblot), Reims (Pr F. Bani-Sadr), Rennes (Pr C. Michelet), Saint-Etienne (Pr F. Lucht), Saint-Mandé (Pr T. Debord), Strasbourg (Pr T. Martin), Toulon (Dr J.P. De Jaureguiberry), Toulouse (Pr B. Marchou), Tours (Pr L. Bernard).

References

- [1] Duffau P, Wittkop L, Lazaro E, le Marec F, Cogne C, Blanco P, et al. Association of immune-activation and senescence markers with non-AIDS-defining comorbidities in HIV-suppressed patients. *AIDS* 2015;29:2099–108.
- [2] Schouten J, Wit FW, Stolte IG, Kootstra NA, van der Valk M, Geerlings SE, et al. Cross-sectional comparison of the prevalence of age-associated comorbidities and their risk factors between HIV-infected and uninfected individuals: the AGEHIV cohort study. *Clin Infect Dis* 2014;59:1787–97.
- [3] Serrano-Villar S, Gutierrez F, Miralles C, Berenguer J, Rivero A, Martinez E, et al. Human immunodeficiency virus as a chronic disease: evaluation and management of nonacquired immune deficiency syndrome-defining conditions. *Open Forum Infect Dis* 2016;3:ofw097.
- [4] Kannel WB, Vasani RS, Keyes MJ, Sullivan LM, Robins SJ. Usefulness of the triglyceride-high-density lipoprotein versus the cholesterol-high-density lipoprotein ratio for predicting insulin resistance and cardiometabolic risk (from the Framingham Offspring Cohort). *Am J Cardiol* 2008;101:497–501.
- [5] Salazar MR, Carbajal HA, Espeche WG, Aizpurua M, Maciel PM, Reaven GM. Identification of cardiometabolic risk: visceral adiposity index versus triglyceride/HDL cholesterol ratio. *Am J Med* 2014;127:152–7.
- [6] DADS Group, Friis-Moller N, Reiss P, Sabin CA, Weber R, Monforte A, et al. Class of antiretroviral drugs and the risk of myocardial infarction. *N Engl J Med* 2007;356:1723–35.
- [7] Lang S, Mary-Krause M, Cotte L, Gilquin J, Partisani M, Simon A, et al. Impact of individual antiretroviral drugs on the risk of myocardial infarction in human immunodeficiency virus-infected patients: a case-control study nested within the French Hospital Database on HIV ANRS cohort CO4. *Arch Intern Med* 2010;170:1228–38.
- [8] Eckard AR, Meissner EG, Singh I, McComsey GA. Cardiovascular Disease, Statins, and HIV. *J Infect Dis* 2016;214(Suppl 2):S83–92.
- [9] Ryom L, Lundgren JD, El-Sadr W, Reiss P, Kirk O, Law M, et al. Cardiovascular disease and use of contemporary protease inhibitors: the D:A:D international prospective multicohort study. *Lancet HIV* 2018;5:e291–300.
- [10] van Zoest RAWF, Kooij KW, van der Valk M, Schouten J, Kootstra NA, Wiersinga WJ, Prins M, van den Born BJ, Reiss P. AGEHIV Cohort Study Group. Higher prevalence of hypertension in HIV-infected patients on combination antire-

- troviral therapy is associated with changes in body composition and prior stavudine exposure. *Clin Infect Dis* 2016;63:205–8.
- [11] Capeau J, Bouteloup V, Katlama C, Bastard JP, Guiyedi V, Salmon-Ceron D, et al. Ten-year diabetes incidence in 1046 HIV-infected patients started on a combination antiretroviral treatment. *AIDS* 2012;26:303–14.
- [12] De Wit S, Sabin CA, Weber R, Worm SW, Reiss P, Cazanave C, et al. Incidence and risk factors for new-onset diabetes in HIV-infected patients: the Data Collection on Adverse Events of Anti-HIV Drugs (D:A:D) study. *Diabetes Care* 2008;31:1224–9.
- [13] Gelpi M, Afzal S, Fuchs A, Lundgren J, Knudsen AD, Drivsholm N, et al. Prior exposure to thymidine analogues and didanosine is associated with long-lasting alterations in adipose tissue distribution and cardiovascular risk factors. *AIDS* 2018 [Epub ahead of print].
- [14] Bastard JP, Fellahi S, Couffignal C, Raffi F, Gras G, Hardel L, et al. Increased systemic immune activation and inflammatory profile of long-term HIV-infected ART-controlled patients is related to personal factors, but not to markers of HIV infection severity. *J Antimicrob Chemother* 2015;70(6):1816–24.
- [15] Bedogni G, Kahn HS, Bellentani S, Tiribelli C. A simple index of lipid overaccumulation is a good marker of liver steatosis. *BMC Gastroenterol* 2010;10:98.
- [16] Bobin-Dubigieon C, Jaffre I, Joalland MP, Classe JM, Campone M, Herve M, et al. Paraoxonase 1 (PON1) as a marker of short term death in breast cancer recurrence. *Clin Biochem* 2012;45:1503–5.
- [17] Caron-Debarle M, Lagathu C, Boccara F, Vigouroux C, Capeau J. HIV-associated lipodystrophy: from fat injury to premature aging. *Trends Mol Med* 2010;16:218–29.
- [18] Hema MN, Ferry T, Dupon N, Cuzin L, Verdon R, Thiebaut R, et al. Low CD4/CD8 ratio is associated with non aids-defining cancers in patients on antiretroviral therapy: ANRS CO8 (Aproco/Copilote) Prospective Cohort Study. *PLoS One* 2016;11:e0161594.
- [19] Nansseu JR, Bigna JJ, Kaze AD, Noubiap JJ. Incidence and risk factors for prediabetes and diabetes mellitus among HIV-infected adults on antiretroviral therapy: a systematic review and meta-analysis. *Epidemiology* 2018;29:431–41.
- [20] Smith U. Abdominal obesity: a marker of ectopic fat accumulation. *J Clin Invest* 2015;125:1790–2.
- [21] Mathabire Rucker SC, Tayea A, Bitilinyu-Bangoh J, Bermudez-Aza EH, Salumu L, Quiles IA, et al. High rates of hypertension, diabetes, elevated low-density lipoprotein cholesterol, and cardiovascular disease risk factors in HIV-infected patients in Malawi. *AIDS* 2018;32:253–60.
- [22] Myerson M, Poltavskiy E, Armstrong EJ, Kim S, Sharp V, Bang H. Prevalence, treatment, and control of dyslipidemia and hypertension in 4278 HIV outpatients. *J Acquir Immune Defic Syndr* 2014;66:370–7.
- [23] Vega GL, Barlow CE, Grundy SM, Leonard D, DeFina LF. Triglyceride-to-high-density-lipoprotein-cholesterol ratio is an index of heart disease mortality and of incidence of type 2 diabetes mellitus in men. *J Investig Med* 2014;62:345–9.
- [24] Squillace N, Lorenzini P, Lapadula G, Bandera A, Cozzi-Lepri A, Rusconi S, et al. Triglyceride/HDL ratio and its impact on the risk of diabetes mellitus development during ART. *J Antimicrob Chemother* 2016;71:2663–9.
- [25] Gelpi M, Afzal S, Roen A, Mocroft A, Lebech AM, Lindegaard B, et al. Increased risk of low adiponectin and atherogenic dyslipidemia in HIV infection. Conference on Retroviruses and Opportunistic Infections 2018;706.
- [26] Pihl E, Zilmer K, Kullisaar T, Kairane C, Pulges A, Zilmer M. High-sensitive C-reactive protein level and oxidative stress-related status in former athletes in relation to traditional cardiovascular risk factors. *Atherosclerosis* 2003;171:321–6.
- [27] Matsuda M, Shimomura I. Roles of adiponectin and oxidative stress in obesity-associated metabolic and cardiovascular diseases. *Rev Endocr Metab Disord* 2014;15:1–10.
- [28] Kopprasch S, Srirangan D, Bergmann S, Graessler J, Schwarz PE, Bornstein SR. Association between systemic oxidative stress and insulin resistance/sensitivity indices - the PREDIAS study. *Clin Endocrinol (Oxf)* 2016;84:48–54.
- [29] Njajou OT, Kanaya AM, Holvoet P, Connelly S, Strotmeyer ES, Harris TB, et al. Association between oxidized LDL, obesity and type 2 diabetes in a population-based cohort, the Health, Aging and Body Composition Study. *Diabetes Metab Res Rev* 2009;25:733–9.
- [30] Spahis S, Delvin E, Borys JM, Levy E. Oxidative stress as a critical factor in nonalcoholic fatty liver disease pathogenesis. *Antioxid Redox Signal* 2017;26(10):519–41.
- [31] Cassar S, Teede HJ, Harrison CL, Joham AE, Moran LJ, Stepto NK. Biomarkers and insulin sensitivity in women with polycystic ovary syndrome: characteristics and predictive capacity. *Clin Endocrinol (Oxf)* 2015;83:50–8.
- [32] Knudsen A, Katzenstein TL, Benfield T, Jorgensen NR, Kronborg G, Gerstoft J, et al. Plasma plasminogen activator inhibitor-1 predicts myocardial infarction in HIV-1-infected individuals. *AIDS* 2014;28:1171–9.
- [33] Lp PLASC, Thompson A, Gao P, Orfei L, Watson S, Di Angelantonio E, et al. Lipoprotein-associated phospholipase A(2) and risk of coronary disease, stroke, and mortality: collaborative analysis of 32 prospective studies. *Lancet* 2010;375:1536–44.
- [34] Ross Eckard A, Longenecker CT, Jiang Y, Debanne SM, Labbato D, Storer N, et al. Lipoprotein-associated phospholipase A2 and cardiovascular disease risk in HIV infection. *HIV Med* 2014;15:537–46.
- [35] Nou E, Lu MT, Looby SE, Fitch KV, Kim EA, Lee H, et al. Serum oxidized low-density lipoprotein decreases in response to statin therapy and relates independently to reductions in coronary plaque in patients with HIV. *AIDS* 2016;30:583–90.
- [36] Saumoy M, Ordonez-Llanos J, Martinez E, Ferrer E, Domingo P, Ribera E, et al. Atherogenic properties of lipoproteins in HIV patients starting atazanavir/ritonavir or darunavir/ritonavir: a substudy of the ATADAR randomized study. *J Antimicrob Chemother* 2015;70:1130–8.
- [37] Gao S, Zhao D, Wang M, Zhao F, Han X, Qi Y, et al. Association Between Circulating Oxidized LDL and Atherosclerotic Cardiovascular Disease: A Meta-analysis of Observational Studies. *Can J Cardiol* 2017;33:1624–32.
- [38] Zidar DA, Juchnowski S, Ferrari B, Claggett B, Pilch-Cooper HA, Rose S, et al. Oxidized LDL Levels Are Increased in HIV Infection and May Drive Monocyte Activation. *J Acquir Immune Defic Syndr* 2015;69:154–60.
- [39] Parra S, Coll B, Aragones G, Marsillach J, Beltran R, Rull A, et al. Nonconcordance between subclinical atherosclerosis and the calculated Framingham risk score in HIV-infected patients: relationships with serum markers of oxidation and inflammation. *HIV Med* 2010;11:225–31.
- [40] Tsimikas S. In vivo markers of oxidative stress and therapeutic interventions. *Am J Cardiol* 2008;101:34D–42D.
- [41] Ferretti G, Bacchetti T, Sahebkar A. Effect of statin therapy on paraoxonase-1 status: A systematic review and meta-analysis of 25 clinical trials. *Prog Lipid Res* 2015;60:50–73.
- [42] Toribio M, Fitch KV, Sanchez L, Burdo TH, Williams KC, Sponseller CA, et al. Effects of pitavastatin and pravastatin on markers of immune activation and arterial inflammation in HIV. *AIDS* 2017;31:797–806.
- [43] Ma X, Chen Z, Wang L, Wang G, Wang Z, Dong X, et al. The pathogenesis of diabetes mellitus by oxidative stress and inflammation: its inhibition by Berberine. *Front Pharmacol* 2018;9:782.
- [44] Brown TT, Tassiopoulos K, Bosch RJ, Shikuma C, McComsey GA. Association between systemic inflammation and incident diabetes in HIV-infected patients after initiation of antiretroviral therapy. *Diabetes Care* 2010;33:2244–9.
- [45] Betene ADC, De Wit S, Neuhaus J, Palfreeman A, Pepe R, Pankow JS, et al. Interleukin-6, high sensitivity C-reactive protein, and the development of type 2 diabetes among HIV-positive patients taking antiretroviral therapy. *J Acquir Immune Defic Syndr* 2014;67:538–46.
- [46] Hove-Skovsgaard M, Gaardbo JC, Kolte L, Winding K, Seljeflot I, Svardal A, et al. HIV-infected persons with type 2 diabetes show evidence of endothelial dysfunction and increased inflammation. *BMC Infect Dis* 2017;17:234.
- [47] Triant VA, Meigs JB, Grinspoon SK. Association of C-reactive protein and HIV infection with acute myocardial infarction. *J Acquir Immune Defic Syndr* 2009;51:268–73.
- [48] Duprez DA, Neuhaus J, Kuller LH, Tracy R, Bellosso W, De Wit S, et al. Inflammation, coagulation and cardiovascular disease in HIV-infected individuals. *PLoS One* 2012;7:e44454.
- [49] Kelesidis T, Kendall MA, Yang OO, Hodis HN, Currier JS. Biomarkers of microbial translocation and macrophage activation: association with progression of subclinical atherosclerosis in HIV-1 infection. *J Infect Dis* 2012;206:1558–67.
- [50] Merlini E, Luzi K, Suardi E, Barassi A, Cerrone M, Martinez JS, et al. T-cell phenotypes, apoptosis and inflammation in HIV+ patients on virologically effective cART with early atherosclerosis. *PLoS One* 2012;7:e46073.
- [51] van Zoest RA, Wit FW, Kooij KW, van der Valk M, Schouten J, Kootstra NA, et al. Higher prevalence of hypertension in HIV-1-infected patients on combination antiretroviral therapy is associated with changes in body composition and prior stavudine exposure. *Clin Infect Dis* 2016;63:205–13.
- [52] Ozanne A, Duffau P, Dauchy FA, Rigother C, Terrien C, Lazaro E, et al. Activation, senescence and inflammation markers in HIV patients: association with renal function. *AIDS* 2017;31:1119–28.
- [53] Younas M, Psomas C, Reynes J, Corbeau P. Immune activation in the course of HIV-1 infection: Causes, phenotypes and persistence under therapy. *HIV Med* 2016;17:89–105.
- [54] Gorwood J, Bourgeois C, Mantecon M, Atlan M, Pourcher V, Pourcher G, et al. Impact of HIV/SIV infection and viral proteins on adipose tissue fibrosis and adipogenesis. *AIDS* 2019. <http://dx.doi.org/10.1097/QAD00000000000012.2168>.