

# Contents

## Articles from the Special Issue on "Diabetes and Ramadan Fasting"; Edited by Mohamed Hassanein and Abdul Jabbar

- A. Chowdhury, H. Khan, S.S. Lasker and T.A. Chowdhury* Fasting outcomes in people with diabetes and chronic kidney disease in East London during Ramadan 2018: The East London diabetes in Ramadan survey 166
- Z. Dabbous, M. Bashir, A.-N. Elzouki, M.S. Ahmed, S. Farouk, M. Hassanien, R.A. Malik, A.B. Abou Samra, T. Elhadd, for the PROFAST Study Group* Differential effects of gender and patient background diversity on the changes in metabolic and biophysical profiles in people with type-2 diabetes from different ethnicities who fast during Ramadan (H1439); a prospective study from Qatar 171
- E.M. Ba-Essa, M. Hassanein, S. Abdulrhman, M. Alkhalifa and Z. Alsafar* Attitude and safety of patients with diabetes observing the Ramadan fast 177

## Reviews

- S. Faal, P. Abedi, S. Jahanfar, J.M. Ndeke, Z. Mohaghegh, F. Sharifipour and M. Zahedian* Sex hormone binding globulin for prediction of gestational diabetes mellitus in pre-conception and pregnancy: A systematic review 39
- M.S. Hamed, M. Samy, H. Mahmoud and N. Yehia* Study of the difficult glycemic control in relation to the presence of diabetes-autoantibodies in a sample of Egyptians with type 1 diabetes 53

## Research Articles

- Y. Matsushita, Y. Takata, R. Kawamura, M. Takakado, T. Hadate and H. Osawa* The fluctuation in sympathetic nerve activity around wake-up time was positively associated with not only morning but also daily glycemic variability in subjects with type 2 diabetes 1
- M. Meloni, V. Izzo, L. Giurato, R. Gandini and L. Uccioli* Below-the-ankle arterial disease severely impairs the outcomes of diabetic patients with ischemic foot ulcers 9
- A.M.J. Saad, D. Hiyasat, H. Jaddou and N. Obeidat* The prevalence of high risk obstructive sleep apnoea among patients with type 2 diabetes in Jordan 16
- M. Yang, C. Ni, B. Chang, Z. Jiang, Y. Zhu, Y. Tang, Z. Li, C. Li and B. Li* Association between serum total bilirubin levels and the risk of type 2 diabetes mellitus 23
- T.P.L. Nguyen, H. Edwards, T.N.D. Do and K. Finlayson* Effectiveness of a theory-based foot care education program (3STEPFUN) in improving foot self-care behaviours and foot risk factors for ulceration in people with type 2 diabetes 29
- M.F. Keller, M. Vestgaard, P. Damm, E.R. Mathiesen and L. Ringholm* Treatment with the long-acting insulin analog degludec during pregnancy in women with type 1 diabetes: An observational study of 22 cases 58
- T. Magan, A. Pouncey, K. Gadhvi, M. Katta, M. Posner and C. Davey* Prevalence and severity of diabetic retinopathy in patients attending the endocrinology diabetes clinic at Mulago Hospital in Uganda 65
- W.-L. Chiu, L. Churilov, C.-H. Lim, A. Tan, R. Nedumanni, L.-H. Lau, J. Lew, M. Hachem, A. Kong, R. Robbins, H. Sutcliffe, Q. Lam, A. Lee, F. Djukiadmodjo, N. Nanayakkara, J.D. Zajac and E.I. Ekinici* Routine HbA1c among hematology and oncology inpatients: Diabetes-status and hospital-outcomes 71
- M. Iwanishi, T. Kusakabe, C. Azuma, Y. Tezuka, Y. Yamamoto, J. Ito-Kobayashi, M. Washiyama, M. Morimoto and K. Ebihara* Clinical characteristics in two patients with partial lipodystrophy and Type A insulin resistance syndrome due to a novel heterozygous missense mutation in the insulin receptor gene 79
- B. Ásbjörnsdóttir, H. Ronneby, M. Vestgaard, L. Ringholm, V.L. Nichum, D.M. Jensen, A. Raben, P. Damm and E.R. Mathiesen* Lower daily carbohydrate consumption than recommended by the Institute of Medicine is common among women with type 2 diabetes in early pregnancy in Denmark 88



<i>B. Wolnik, E. Orłowska-Kunikowska, M. Błaszowska, B. Graff, J. Wolf, L. Czupryniak and K. Narkiewicz</i> The phenomenon of HbA1c stability and the risk of hypoglycemia in long-standing type 1 diabetes	96
<i>H.-K. Huang and J.-I. Yeh</i> Comparison of mortality and cardiovascular event risk associated with various insulin secretagogues: A nationwide real-world analysis	103
<i>G.G. Massa, I. Gys, E. Bevilacqua, A. Wijnands and R. Zeevaert</i> Comparison of flash glucose monitoring with real time continuous glucose monitoring in children and adolescents with type 1 diabetes treated with continuous subcutaneous insulin infusion	111
<i>E.M. Koc, H. Aksoy, D. Ayhan Başer, A. Baydar Artantaş and R. Kahveci</i> Quality assessment of clinical practice guidelines for management of type 2 diabetes mellitus. Assessment of type 2 diabetes mellitus guidelines	119
<i>B.C. Shaddinger, J. Soffer, G. Vlasakakis, M. Shabbout, C. Weston and A. Nino</i> Efficacy and safety of an albiglutide liquid formulation compared with the lyophilized formulation: A 26-week randomized, double-blind, repeat-dose study in patients with type 2 diabetes mellitus	125
<i>M.M. Hashemi, E. Mousavi, Z. Arab-Bafrani, A. Nezhadebrahimi and A. Marjani</i> The most effective polymorphisms of paraoxonase-1 gene on enzyme activity and concentration of paraoxonase-1 protein in type 2 diabetes mellitus patients and non-diabetic individuals: A systematic review and meta-analysis	135
<i>K.K. Vesco, A.J. Sharma, J. Bulkley, T. Kimes, W.M. Callaghan, L.J. England and M.C. Hornbrook</i> Association of glucose levels in pregnancy with use of health care services	146
<i>K. Micielska, A. Gmiat, M. Zychowska, M. Kozłowska, A. Walentukiewicz, A. Lysak-Radomska, J. Jaworska, E. Rodziewicz, B. Duda-Biernacka and E. Ziemann</i> The beneficial effects of 15 units of high-intensity circuit training in women is modified by age, baseline insulin resistance and physical capacity	156
<b>International Diabetes Federation</b>	
Type 1 diabetes camp is much more than blood glucose	186
<b>Letters to the Editor</b>	
<i>M.B. Aykan and I. Tasci</i> Interpretation of blood apelin level across different clinical pictures of diabetes mellitus	183
<i>A. Noori-Zadeh and S. Bakhtiyari</i> Reply to comments on: Elevated blood apelin levels in type 2 diabetes mellitus: A systematic review and meta-analysis	185