

Contents

Articles from the Special Issue on "Diabetes and Ramadan Fasting"; Edited by Mohamed Hassanein and Abdul Jabbar

- E. Al-Ozairi, M.M. AlAwadhi, A. Al-Ozairi, E. Taghadom and K. Ismail* A prospective study of the effect of fasting during the month of Ramadan on depression and diabetes distress in people with type 2 diabetes 145
- I. Saadane, T. Ashraf, T. Ali and N. Lessan* Diabetes and Ramadan: Utility of flash-glucose monitoring derived markers of glycaemic control and comparison with glycosylated haemoglobin 150
- D. Alsaeed, J. Al-Kandari and E. Al-Ozairi* Experiences of people with type 1 diabetes fasting Ramadan following structured education: A qualitative study 157
- "M.A."E. Faris, M.I. Madkour, A.K. Obaideen, E.Z. Dalah, H.A. Hasan, H. Radwan, H.A. Jahrami, O. Hamdy and M.G. Mohammad* Effect of Ramadan diurnal fasting on visceral adiposity and serum adipokines in overweight and obese individuals 166
- M.M. Abushady, M. Samy, M. Bekhet and A. Abdullah* Effect of Ramadan fasting on renal function in patients with type 2 diabetes 176

Review

- D.G. Marrero, M.E. Hilliard, D.M. Maahs, A.H. McAuliffe-Fogarty and C.M. Hunter* Using patient reported outcomes in diabetes research and practice: Recommendations from a national workshop 23

Research Articles

- M. Nowaczewska, A. Kamińska, B. Kukulska-Pawluczuk, R. Junik and K. Pawlak-Osińska* Effect of hyperglycemia on cerebral blood flow in patients with diabetes 1
- S.-M. Chuang, H.-M. Shih, M.-N. Chien, S.-C. Liu, C.-H. Wang and C.-C. Lee* Risk factors in metabolic syndrome predict the progression of diabetic nephropathy in patients with type 2 diabetes 6
- M.H.A. Muskiet, M.C. Bunck, R.J. Heine, A. Cornér, H. Yki-Järvinen, B. Eliasson, J.A. Joles, M. Diamant, L. Tonneijck and D.H. van Raalte* Exenatide twice-daily does not affect renal function or albuminuria compared to titrated insulin glargine in patients with type 2 diabetes mellitus: A post-hoc analysis of a 52-week randomised trial 14
- H. Ito, S. Ando, E. Tsugami, R. Araki, E. Kusano, S. Matsumoto, K. Uemura, S. Nishio, S. Antoku, T. Yamasaki, T. Mori and M. Togane* Changes in medication adherence and unused drugs after switching from daily dipeptidyl peptidase-4 inhibitors to once-weekly trelagliptin in patients with type 2 diabetes 41
- C.I. Elizalde-Barrera, A.F. Rubio-Guerra, J.J. Lozano-Nuevo and J.L. Olvera-Gomez* Triglycerides and waist to height ratio are more accurate than visceral adiposity and body adiposity index to predict impaired fasting glucose 49
- W.M. Kattan and A.A. Abduljawad* Predicting different factors that affect hospital utilization and outcomes Among diabetic patients admitted with hypoglycemia using structural equation modeling 55
- I. Ezquerro-Lázaro, L. Cea-Soriano, C. Giraldez-García, A. Ruiz, J. Franch-Nadal, J. Diez-Espino, P. Nogales, F. Carramiñana, F. Javier Sangros, E. Regidor and for the PREDAPS Study Group* Lifestyle factors do not explain the difference on diabetes progression according to type of prediabetes: Results from a Spanish prospective cohort of prediabetic patients 66
- Y.-J. Kwon, K.-W. Hong, B.J. Park and D.-H. Jung* Serotonin receptor 3B polymorphisms are associated with type 2 diabetes: The Korean Genome and Epidemiology Study 76
- G.J. José, E. Jorge, F. Lujan, Q. Ivanna and C. Jean-Marc* Type 2 diabetes: Prescription patterns and treatment outcomes of IDMPS survey in Argentina 86
- U.P. Gujral, D. Prabhakaran, R. Pradeepa, N.R. Kandula, D. Kondal, M. Deepa, N.A. Zakai, R.M. Anjana, G. Rautela, V. Mohan, K.M.V. Narayan, N. Tandon and A.M. Kanaya* Isolated HbA1c identifies a different subgroup of individuals with type 2 diabetes compared to fasting or post-challenge glucose in Asian Indians: The CARRS and MASALA studies 93



<i>Y.-T. Liu, L.-Y. Lin, C.-W. Tuan, C.-Y. Yang and P.-L. Tang</i> Analyzing the Association HbA1c control by Depression, social participation and Utilizing Self-management Questionnaire	103
<i>S.G. Sosa-Rubi, L. Dainelli, I. Silva-Zolezzi, P. Detzel, S. Espino y Sosa, E. Reyes-Muñoz, C. Chivardi, E. Ortiz-Panozo and R. Lopez-Ridaura</i> Short-term health and economic burden of gestational diabetes mellitus in Mexico: A modeling study	114
<i>T. Xia, L. Turner, J. Enticott, D. Mazza and P. Schattner</i> Glycaemic control of Type 2 diabetes in older patients visiting general practitioners: An examination of electronic medical records to identify risk factors for poor control	125
<i>U. Juttada, T.P. Smina, S. Kumpatla and V. Viswanathan</i> Seroprevalence and risk factors associated with HBV and HCV infection among subjects with type 2 diabetes from South India	133
<i>I. Dicembrini, B. Tomberli, B. Nreu, G.I. Baldereschi, F. Fanelli, E. Mannucci and M. Monami</i> Peripheral artery disease and amputations with Sodium-Glucose co-Transporter-2 (SGLT-2) inhibitors: A meta-analysis of randomized controlled trials	138
<i>R. Haider, K. Hyun, N.W. Cheung, J. Redfern, A. Thiagalingam and C.K. Chow</i> Effect of lifestyle focused text messaging on risk factor modification in patients with diabetes and coronary heart disease: A sub-analysis of the TEXT ME study	184
International Diabetes Federation	
How can advocates, healthcare professionals and policy makers join forces to fight CVD in people with diabetes?	196
Commentaries	
<i>G. Schernthaner, C. Wanner, D. Jurišić-Eržen, C. Guja, J. Gumprecht, I.R. Jarek-Martynowa, A. Karasik, N. Lalić, B.N. Mankovsky, M. Prázný, T. Tankova, A. Tsur, T.C. Wascher and I. Wittmann</i> CARMELINA: An important piece of the DPP-4 inhibitor CVOT puzzle	30
<i>D. Giugliano, M.I. Maiorino, M. Longo and K. Esposito</i> Are gliflozins the new statins for diabetes?	191
<i>K.W. Hansen</i> Converting haemoglobin A1c and average glucose. Time to change?	194
Brief report	
<i>J.B. Farinha, W. Boff, G.C. dos Santos, F.P. Boeno, T.R. Ramis, A.F. Vieira, R.C.O. Macedo, J. Rodrigues-Krause and A. Reischak-Oliveira</i> Acute glycemic responses along 10-week high-intensity training protocols in type 1 diabetes patients	111