



Durability of antidepressant response to repetitive transcranial magnetic stimulation: Systematic review and meta-analysis

Suhan Senova^{a, b, c, 1}, Gonçalo Cotovio^{a, d, e, h, 1}, Alvaro Pascual-Leone^{f, g}, Albino J. Oliveira-Maia^{a, d, e, h, *}

^a Champalimaud Clinical Centre, Champalimaud Centre for the Unknown, Lisboa, Portugal

^b Neurosurgery and PePsy Departments, Assistance Publique-Hôpitaux de Paris (APHP), Groupe Henri-Mondor Albert-Chenevier, Créteil, France

^c Equipe 14, U955 INSERM, Institut Mondor de Recherche Biomedicale and Faculté de Médecine, Université Paris Est, Créteil, France

^d Department of Psychiatry and Mental Health, Centro Hospitalar de Lisboa Ocidental, Lisboa, Portugal

^e NOVA Medical School | Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisboa, Portugal

^f Berenson-Allen Center for Noninvasive Brain Stimulation, Division of Cognitive Neurology, Department of Neurology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

^g Institut Guttmann de Neurorehabilitación, Universitat Autònoma de Barcelona, Badalona, Barcelona, Spain

^h Champalimaud Research, Champalimaud Centre for the Unknown, Lisboa, Portugal

ARTICLE INFO

Article history:

Received 5 April 2018

Received in revised form

21 September 2018

Accepted 1 October 2018

Available online 2 October 2018

Keywords:

Depression

Repetitive transcranial magnetic

stimulation

Durability

Relapse

Meta-analysis

ABSTRACT

Background: The therapeutic options for treatment-resistant depression (TRD) encompass a range of neuromodulatory techniques, including repetitive transcranial magnetic stimulation (rTMS). While rTMS is safe and has documented short-term efficacy, durability of antidepressant effects is poorly established.

Objective: Assess existing evidence regarding durability of rTMS-induced antidepressant response.

Methods: We performed a systematic review of studies reporting antidepressant outcome measures collected three or more months after the end of an induction course of rTMS for depression. Among responders to the induction course, we used a meta-analytic approach to assess response rates at 3 (m3), 6 (m6) or 12 (m12) months after induction, and studied predictors of responder rates using meta-regression.

Results: Nineteen studies published between 2002 and 2018 were included. Eighteen were eligible for analysis at m3 (732 patients) and m6 (695 patients) and 9 at m12 (247 patients). Among initial responders, 66.5% sustained response at m3 (95% CI = 57.1–74.8%, $I^2 = 27.6\%$), 52.9% at m6 (95% CI = 40.3–65%, $I^2 = 0\%$), and 46.3% at m12 (95% CI = 32.6–60.7%, $I^2 = 0\%$), in the absence of any major bias. Random-effects meta-regressions further demonstrated that a higher proportion of women, as well as receipt of maintenance treatment, predicted higher responder rates at specific time-points.

Conclusions: rTMS is a durable treatment for depression, with sustained responder rates of 50% up to 1 year after a successful induction course of treatment. Maintenance treatment may enhance the durability of the antidepressant effects of rTMS, and should be considered in clinical practice, as well as systematically explored in future clinical trials.

© 2018 Elsevier Inc. All rights reserved.

Introduction

Antidepressant medication and psychotherapy are first line treatments for patients suffering from major depression [1]. In case of insufficient benefit from, or intolerance to, medication, patients may be offered a number of neuromodulatory options, namely electroconvulsive therapy (ECT), repetitive transcranial magnetic

stimulation (rTMS) and vagal nerve stimulation (VNS). Transcranial direct current stimulation (tDCS) and deep brain stimulation (DBS) are still investigational [2–6]. Choosing among these options requires careful weighing of their characteristics, namely durability of antidepressant effects, of particular importance since these patients typically suffer from multiple depressive episodes across their lifetime [7,8]. However, durability of antidepressant effects of rTMS has been insufficiently examined to date.

Efficacy of dorsolateral prefrontal cortex (DLPFC) rTMS for major depression has been documented by a large number of studies, including multicenter trials, and analyzed in several meta-analyses (e.g. Refs. [4,9–12]). Efficacy has been demonstrated for both

* Corresponding author. Champalimaud Research and Clinical Centre, Champalimaud Centre for the Unknown, Avenida Brasília, 1400-038, Lisboa, Portugal.

E-mail address: albino.maia@neuro.fchampalimaud.org (A.J. Oliveira-Maia).

¹ equally contributing authors.

medication-free and medicated patients [13–17], and the United States Food and Drug Administration (FDA) has formally cleared four different devices for rTMS treatment of patients with prior treatment-failure for depression. The protocol leading to initial FDA clearance was based on the application of daily treatment sessions, 5 days a week, for up to six weeks, with stimulation delivered over the left hemisphere at high frequency (typically 10 Hz or higher) and high intensity (120% of the resting motor threshold) [18,19]. There is less evidence to support rTMS protocols using lower stimulation frequency or stimulation intensities [20].

Durability of antidepressant effects after rTMS induction has been assessed on short timescales, typically no longer than 3 months after treatment [14,21–23]. A previous meta-analysis including randomized studies with follow-up phases ranging from 2 to 16 weeks, none of which used rTMS maintenance treatment, concluded that rTMS has a small antidepressant effect during follow-up [22]. Meta-analyses taking into consideration longer follow-up phases, and considering options for maintenance treatment, have not been reported, probably because there are very few randomized controlled trials (RCT) with long-term follow-up. Indeed, in RCT's for major depressive disorder (MDD), sustained blinding for long periods after the end of treatment raises difficult logistic and ethical questions. However, considering the high relapse rate for major depressive disorder [7], understanding treatment effects for only three months is not sufficiently informative. This is particularly important when considering the emergence of chronic neurostimulation treatment options for treatment resistant depression (TRD), such as VNS, DBS [5,24,25] and epidural prefrontal cortical stimulation [26], that offer the potential for prolonged efficacy.

Nevertheless, several open-label rTMS studies have retained patients for very long follow-up phases, of up to 6 years [27–29]. A meta-analysis including such studies, similar to what has been performed for electroconvulsive therapy [30], could thus begin to provide quantitative responses for the question of long-term durability of the antidepressant effects of rTMS [31], as well as explore the best alternatives for relapse prevention after successful rTMS [8,32]. Thus, the primary objective of this meta-analysis was to provide a systematic overview of the existing evidence regarding post-rTMS durability of response for patients followed for at least 3 months after acute treatment. The secondary goal was to evaluate the impact of maintenance rTMS treatment on durability of antidepressant effects, since this has been proposed as a potential alternative for relapse prevention, but its efficacy is not clearly established [33]. Considering both the previously mentioned meta-analysis for short-term follow-up, and trials including a longer follow-up period, we hypothesized that the efficacy of acute rTMS decreases over time, but is prolonged by rTMS maintenance treatment.

Material & methods

This systematic review and meta-analysis was performed according to PRISMA Guidelines [34].

Search strategy

An electronic literature search in PubMed and Web of Science was performed, including papers published up to June 2018. Mesh terms used are described in [supplementary table 1](#). Two authors (SS and GC) independently eliminated ineligible reports in sequential steps, judging first by title and abstract review, followed by full-text screening. Hand-searches of the reference sections of prior reviews, as well as of previously identified eligible studies, were carried out to identify additional eligible studies. Inconsistencies in eligibility assessments were resolved by consensus, when necessary with

participation of a third author. Quality assessment for each study was performed with the Newcastle-Ottawa Quality Assessment Scale for Cohort Studies [35].

Study eligibility criteria

Inclusion criteria

- (1) prospective or retrospective study reported in a peer-reviewed publication;
- (2) use of rTMS induction for treatment of a depressive episode (unipolar or bipolar) diagnosed by clinical judgment or formal diagnostic criteria (e.g., DSM-IV);
- (3) treatment response defined consistently in each study through clinical judgment, use of formal diagnostic criteria and/or clinician-rated depression severity rating scales (Hamilton Depression Rating Scale – HAM-D; Cognitive Global Impression Scale – CGI);
- (4) patients considered to be rTMS responders or remitters after rTMS induction were monitored at one or more time points, at least 3 months after the end of rTMS induction;
- (5) responder rates at each time-point were defined by the investigators of the original study and reported as the percentage of the initial responder or remitter sample, or could be directly extracted from the data in the manuscript.

Exclusion criteria

- (1) case studies or case series with less than 4 patients;
- (2) inclusion of non-affective psychosis, dementia, neurological disease, alcoholism or unstable medical conditions among the patient sample;
- (3) inclusion of patients younger than 18 years;
- (4) review or meta-analysis not reporting original data.

Data extraction

Data were extracted from each selected manuscript, using the text, tables, and/or figures, independently by three authors of this meta-analysis. Discrepancies were resolved through consensus, prior to data analysis. The primary outcome was the responder rate and the endpoints were three moments after rTMS induction treatment: m3 (three to four months), m6 (five to six months) and m12 (ten to twelve months). Response was defined in each study according to specific criterion chosen by the authors, and expressed at each endpoint relatively to the number of patients who responded to induction rTMS and entered the follow-up phase. An intention-to-treat analysis approach was chosen, such that patients lost to follow-up were considered as non-responders.

Additional data were extracted to assess potential predictors of responder rates. Age and gender distributions were extracted as potential demographic predictors of response. Clinical variables of interest were depression severity prior to rTMS induction, duration of the affective disorder, number of previous episodes, duration of the current episode and percentage of patients with bipolar depression. Inclusion of patients with TRD was also extracted as a clinical variable. Since TRD is defined variably according to different criteria [36], for the purposes of this meta-analysis we specified TRD according to stage II from the Thase & Rush criteria [37]. Parameters of the rTMS induction treatment were also extracted: intensity, rTMS frequency, number of pulses per treatment and total number of rTMS treatments. Other treatment-related variables were the study-defined criterion of response to rTMS induction, percentage of patients receiving left DLPFC rTMS, and delivery of maintenance

rTMS treatment. For this study, maintenance rTMS was defined as any rTMS session delivered after the induction cycle.

Meta-analyses

Statistical analyses were performed using Comprehensive Meta Analysis Version 3. Due to differences in study designs and patient populations, mean responder rates with 95% confidence intervals (CIs) were calculated with a random-effects model. I^2 statistic was performed to assess heterogeneity. Univariate meta-regression analyses were performed to identify predictors of responder rates. Subgroup analyses were performed at m3, m6 and m12, to compute mean responder rates with 95% CIs according to maintenance treatment, namely among studies exclusively with, versus exclusively without, maintenance treatment.

Publication bias analysis

Publication bias was assessed by visual inspection of funnel plots of standard error vs. logit of responder rates, whenever more than 10 studies were available. Duval and Tweedie's Trim-and-Fill analysis was used to test if the funnel plot was symmetrical around the overall mean weighted responder rates of all studies. To test if studies with lower effect sizes differed systematically from studies with higher effect sizes we used the Begg and Mazumdar Rank Order Correlation (Kendall's tau b) between the standardized effect sizes and the standard error of the mean (SEM) in each study, as well as the Egger's regression of $1/SEM$ (predictor) on the standardized effect sizes.

Results

Search results

In initial literature searches, 207 articles were found and screened for eligibility, 30 of which were chosen for full-text inspection after title and abstract review. Among the references quoted in these articles, 32 more were selected for full-text inspection. Finally, 23 articles published between 2002 and 2018 were included in the systematic review (Fig. 1): 3 RCT [33,38,39], 14 open-label prospective studies [28,40–50,52,53] and 2 retrospective studies [54,55]. For one RCT [39] only the open label extension could be included.

Four studies [51,56–58] were not considered for quantitative review since response and relapse rates for responders to initial course of rTMS could not be extracted. Among the remaining 19 studies included in quantitative review, quality was fair to good (Supplementary Table 2). Four studies included only MDD patients with TRD, defined as non-responders to two or more courses of adequate pharmacotherapy [39,40,49,53], while 9 studies included MDD patients both with and without TRD [28,33,41,43,44,46,47,50,54]. The 6 remaining studies included patients with bipolar depression [38,42,45,48,52,55], two of which including only patients with TRD [38,52]. The criterion to define responders was also variable, as is further detailed in Table 1 and Supplementary Table 3.

One of the central objectives of the quantitative synthesis was regarding the effects of maintenance treatment. Thus, one of the open-label prospective studies [52] and one RCT [38], reporting data separately for maintenance and non-maintenance groups, were split accordingly, and considered as two distinct studies for meta-analyses. Among the studies delivering rTMS maintenance treatments, the design of the maintenance protocols was highly variable. In 7 studies, rTMS maintenance was administered according to planned treatment schemes, ranging from weekly sessions [48] to clusters of 5 sessions administered in two and one-half

days every month [45], as well as extended taper [38,40,46,52,55] (Supplementary Table 4). In three studies, rTMS maintenance was administered as 'rescue' treatment, after evidence of symptom worsening [28,43,47]. Finally, in one RCT [33], while all patients were offered symptomatic re-introduction of rTMS, a subgroup of patients was randomized to receive additional planned maintenance, with a once-monthly session. Since this frequency of rTMS maintenance is well below the standard in other studies (Supplementary Table 4), both groups were considered together and as receiving symptomatic re-introduction of rTMS.

Responder rates at m3, m6 and m12

After a successful index rTMS induction course, among 732 patients from 18 studies, 66.5% (95% CI = 57.1–74.8%, $I^2 = 27.6\%$) were still responders at m3 (Fig. 2a). At m6, among 695 patients from 18 studies, 52.9% (95% CI = 40.3–65%, $I^2 = 0\%$) were still responders (Fig. 2b). At m12, among 247 patients from 9 studies, 46.3% (95% CI = 32.6–60.7%, $I^2 = 0\%$) were still responders (Fig. 2c). While, at m3 and m6, cumulative analyses suggested a trend for responder rates to be higher for older studies (Supplementary Figures 1a and 2a), one-study removed analyses showed the final results were not dependent on any particular study at any time-point (Supplementary Figure 1b, 2b and 3b). Heterogeneity of the studies was low to moderate for all time-points.

Several sensitivity analyses were conducted to verify the nature of these findings when excluding studies according to several different criteria. One study [33] was identified as an outlier with a higher risk of bias due to presenting the highest loss to follow-up at all time points (>33.3% at m3, >50% at m6 and >66.6% at m12; Supplementary Table 5), as well as the lowest response criterion to induction rTMS (25%; Table 1). Another study was considered as potentially problematic since responders were defined according to clinical judgment only, without referring to any clinical score of depression [49]. We thus repeated the meta-analysis after excluding these trials (Supplementary Figure 6b). Response criteria used by the authors to include responders in the follow-up phase could also influence the results. In order to address this potential bias, we performed an additional sensitivity analysis using only studies defining response as at least 50% reduction in the HAM-D score (Supplementary Figure 6c). Furthermore, and despite the fact that the percentage of patients with bipolar disorder in each study was not found to be a predictor of response rate at any time point (see section on predictors of responder rates), we conducted sensitivity analyses considering studies that included only patients with major depressive episode (Supplementary Figure 6d) or studies that included only patients with bipolar disorder (Supplementary Figure 6e). Finally, sensitivity analyses were conducted considering only patients with (Supplementary Figure 6f) or without (Supplementary Figure 6g) chronic depression, defined according to the duration of the current depressive episode. When comparing with the full dataset (Supplementary Figure 6a) these several sensitivity analyses rendered similar results.

Estimation of bias

While at m3 and m6 we gathered information from more than 10 studies, allowing for assessments of bias, at m12 only 9 studies were found and assessment of bias was not possible. For both m3 and m6 the funnel plot was symmetrical at visual inspection (Fig. 3a and b), the Begg and Mazumdar's tests and the Egger's tests were not significant ($p > 0.4$ for both tests), and there was no imputed study after the Duval and Tweedie's Trim and Fill analysis. Thus, overall, our meta-analysis did not suffer from obvious publication bias.




Fig. 1. Article selection flow chart according to PRISMA Statement [34]. *records extracted from the references of full-text articles screening.

Predictors of responder rates at m3 and m6

Univariate meta-regressions with random-effects analyses were performed to assess potential predictors of responder rates at m3 and m6 (Table 2, Supplementary Figures 4 and 5). Such predictors were not assessed at m12, due to the low number of studies at this time-point. The percentage of women included in each study was a significant positive predictor of being a responder at m3 ($p = 0.009$, effect size: 0.04, 95% CI = 0.01–0.07, Z-value: 2.6) and m6 ($p = 0.02$, effect size: 0.04, 95% CI = 0.007–0.07, Z-value: 2.37). There was a trend for lower response criterion to induction rTMS to be a predictor of worse response rate both at m3 ($p = 0.07$) and m6 (0.05), and for higher quality of studies to be a predictor of worse response rate at m3 ($p = 0.05$). Finally, maintenance treatment was a significant positive predictor of response rates at m3 ($p = 0.03$, effect size: 0.91, 95% CI = 0.088–1.74, Z-value: 2.17) and m6 ($p = 0.02$, effect size: 0.95, 95% CI = 0.12–1.79, Z-value: 2.24). The remaining variables, including rTMS protocol parameters in the acute treatment such as stimulation location (Left DL-PFC), % of motor threshold (MT), pulses/day, frequency and induction days, were either not significant predictors of responder rates at m3 and m6, or there was not enough data to assess their effects using meta-regression (Table 2).

Subgroup analyses

Since the question of maintenance treatment is of crucial clinical relevance, we performed subgroup analyses to compute mean responder rates separately for patients receiving or not receiving maintenance rTMS at m3 and m6. At m3, responder rate was 35.8% higher for the 383 patients from 8 studies receiving maintenance (76.2%, 95% CI = 63–85.8%, $I^2 = 0\%$) than for the 349 patients from 8 studies that did not receive maintenance (56.1%, 95% CI = 43.6–67.8%, $I^2 = 38.7\%$), while at m6, responder rate was 58.7% higher for the 417 patients receiving maintenance in 10 studies (61.1%, 95% CI = 49.8–71.3%, $I^2 = 0\%$) than for the 278 patients from 8 studies that did not receive maintenance (38.5%, 95% CI = 21.9–58.3%, $I^2 = 0\%$). Analyses comparing planned and symptomatic re-introduction of rTMS schemes did not demonstrate any significant differences between the two approaches (Supplementary Table 4).

Discussion

In this systematic review and meta-analysis, we describe relevant findings with regards to durability of the antidepressant effects of rTMS treatment, among patients who respond to an acute course of rTMS. Indeed, 66.5% of these patients sustained response

Table 1

Demographic, clinical and methodological data extracted from each study included in the quantitative review.

Study				Demographic		Diagnosis		Disease Characteristics (Baseline)				rTMS Parameters					Response Criteria		Maintenance	
First Author	Year	Type	n	Age (y)	Fem.(%)	BD (%)	TRD	HAMD	Dis.Dur. (y)	Epis. Dur. (m)	Prior Epis. (n)	Left DL-PFC (%)	MT (%)	Pulses/day (n; L/R)	Freq. (Hz; L/R)	Ind. Days (n)	Scale	Red.(%)	TMS	Type
Dannon [41]	2002	Pros.	23	56.8	66.7	0	No			10	3.2	100	90	1200	10	20	HAMD-17	≥60	No	
Boutros [39]	2002	RCT ^b	5	48.8	20	0	Yes	36.6				100	80	800	20	10	HAMD-25	>30	No	
Li [48]	2004	Pros.	7	44.9	71.4	100	No	33.7		20.4		100	110	1600	5	10	HAMD	>50 ^a	Yes	Pl.
Benadhira [40]	2005	Pros.	8			0	Yes					100	80	1600	10	16	HAMD	>50	Yes	Pl.
O'Reardon [28]	2005	Pros.	10	49.6	60	0	No					100	100				HAMD-17	>50	Yes	Sym.
Eranti [44]	2007	Pros.	4			0	No					100	110	1000	10	15	HAMD-17	>50	No	
Cohen [54]	2009	Retr.	204	43.3	51	0	No	22		14.8	2.2	12					HAMD	n.d.	No	
López-Ibor [49]	2010	Pros.	26			0	Yes					100	70		20	10	Clinical Judgement		No	
Janicak [47]	2010	Pros.	99	49.1	53.5	0	No	29.0		12.7		100	120	3000	10		HAMD-17	≥25	Yes	Sym.
Rosenberg [53]	2011	Pros.	8	46.6	50	0	Yes	28.8	19.9	37.6	1.2						HAMD-24	≥50	No	
Dell'osso [42]	2011	Pros.	6			100	No	21.2				0	110	300	1	15	HAMD-21	>50	No	
Mantovani [50]	2012	Pros.	55			0	No										HAMD-24	≥50	No	
Richieri (M) [52]	2013	Pros.	37	52.8	64.9	19	Yes		15.1			37.8	120	2000/720	10/1	20	Sy.	>50	Yes	Pl.
Richieri (NM) [52]	2013	Pros.	22	54.5	36.4	36	Yes		16.7			72.7	120	2000/720	10/1	20	Sy.	>50	No	
Connolly [55]	2013	Retr.	42			>0 ^c	No										CGI	≥50	Yes	Pl.
Fitzgerald [45]	2013	Pros.	35	44.8	77.1	26	No	22.9									HAMD-17	>50	Yes	Pl.
Dunner [43]	2014	Pros.	120			0	No										CGI-S	≤3	Yes	Sym.
Harel [46]	2014	Pros.	12			0	No						120	1680	20	20	HAMD-21	>50	Yes	Pl.
Philip [33]	2016	RCT ^b	49	48.6	61.2	0	No	23.8				100	120	3000	10	30	HAMD-21	>25	Yes	Sym.
Benadhira (M) [38]	2017	RCT	10	50.8	50	n.c.	Yes	20.3				100	110	2000	10	20	HD-17	>50	Yes	Pl.
Benadhira (NM) [38]	2017	RCT	7	56	83.3	n.c.	Yes	20.7				100	110	2000	10	20	HD-17	>50	No	

BD: bipolar disorder; CGI: Clinician-Reported Clinical Global Impressions-; CGI-S: Clinician-Reported Clinical Global Impressions-Severity of Illness scale; Dis. Dur.: disease duration; Epis. Dur.: episode duration; Fem. (%): % of women; HAMD: Hamilton depression score; M: maintenance; n: number of included patients; n.c.: not computable; n; d.: not determined; NM: non maintenance; Pl.: planned maintenance; Prior Epis.: prior episodes; Pros.: prospective studies; RCT: randomized controlled studies; Red.: reduction; Retr.: retrospective studies; Sy: symptoms; Sym.: symptomatic re-introduction of rTMS; TRD: treatment resistant depression; y: year. Study type classifications were performed according to previous definitions [59,60].

^a Despite the absence of predefined response criterion for the inclusion in the follow-up phase, all the patients had a >50% reduction from the baseline HAMD score when entering the maintenance phase according to the table of results displayed in the article.

^b One of the randomized studies comparing two groups of patients was [33]. However, in this study, patients of both groups received rTMS maintenance as needed (based on clinical observation), while scheduled rTMS maintenance was the randomized intervention. For this reason, both groups were recipients of rTMS maintenance, as defined for this manuscript, and we thus considered the data from the two groups jointly, and classified [33] as a prospective study. While [39] also describes results from a RCT of acute rTMS, the data included in this meta-analysis are the results from an open-label extension of the active arm of the RCT.

^c Bipolar patients were included but their proportion could not be determined.




Fig. 2. Meta-analyses. Meta-analyses of Responder rates at 3 months follow-up (m3, panel a), 6 months follow-up (m6, panel b) and 12 months follow-up (m12, panel c), among patients benefiting from an acute course of rTMS. The analyses show that there are sustained responder rates to rTMS, of up to 67%, from 3 months to 1 year after an initial treatment course. CI: confidence interval; M: study arm including only patients with maintenance rTMS; NM: study arm including only patients without maintenance rTMS.

after 3 months, with responder rates decreasing progressively across time, and 46.3% maintaining response 1 year after induction treatment. These results expand the conclusions drawn from a previous meta-analysis [22] with shorter follow-up periods, of only up to 16 weeks. Here we show that the benefit of rTMS might last up to 1 year in close to half of the patients who respond to acute treatment. Furthermore, we found that studies including more women had higher responder rates at 3 and 6 months, in confirmation of the findings described by Kedzior and colleagues [10].

Nevertheless, since results were not reported separately according to gender, we cannot conclude that women have a better outcome than men, and suggest that future studies report outcomes by gender so this finding could be better explored. Importantly, rTMS maintenance treatment, when applied to those responding to acute rTMS treatment, was associated with higher sustained responder rates after 3 months, and especially after 6 months, which were respectively 35.8% and 58.7% greater than those reported in the absence of rTMS maintenance. This could reflect a more important




Fig. 3. Assessment of bias. Assessment of potential bias of meta-analyses of Responder rates at 3 months follow-up (m3, panel a) and 6 months (m6, panel b) among patients benefitting from an acute course of rTMS. In each panel, a funnel plot of the standard error of responder rate over logit of the responder rate, as well as the results for the Egger's test and Begg & Mazumdar's test, are shown.

decline of responder rates in patients who do not receive rTMS maintenance treatment, for whom response rate was 31.4% lower at m6 relative to m3, than in those who did receive rTMS maintenance, where response rate was only 19.8% lower at m6 relative to m3. These findings, which have not been explored in previous meta-analyses, suggest that rTMS maintenance should be systematically considered for patients who respond to an initial treatment course of rTMS for depression, to enhance the chances to maintain the benefits of rTMS for a longer period.

While this meta-analysis suggests that there is value in maintenance rTMS, it provides limited insight regarding the protocol

that should be used, specifically the duration and frequency of treatments, as well as optimal stimulation parameters. Among the 11 studies in which rTMS maintenance was offered, 4 described protocols including symptomatic re-introduction of rTMS schedules [28,33,43,47], whereas the remaining 7 were performed with planned maintenance schedules [38,40,45,46,48,52,55]. The available data did not suggest superiority for any of the two types of approach. Furthermore, the design of planned strategies varied significantly across studies, regarding both frequency, ranging from twice-weekly [57] to monthly [55], as well as duration, ranging from 3 months [40] to up to 2 years [45]. The methods for

Table 2

Effect sizes with 95% CI and p-values for moderators of response rates at 3 months (m3) and 6 months (m6) follow-up, calculated using univariate meta-regression.

	All Studies					
	M3			M6		
	Effect Size	95% CI	p-value	Effect Size	95% CI	p-value
Maintenance rTMS	0.91	0.088–1.74	0.03	0.96	0.12–1.79	0.03
% of Patients with Bipolar Depression	0.75	–1.75–3.24	0.56	0.60	–1.28–2.49	0.53
Response Criterion	0.04	–0.004–0.09	0.07	0.04	0.0006–0.07	0.05
Age	–0.03	–0.15–0.09	0.63	0.03	–0.10–0.15	0.7
% of Women	0.04	0.01–0.07	0.009	0.04	0.007–0.08	0.02
Baseline Clinical Score	–0.05	–0.18	0.46	–0.004	–0.14–0.13	0.95
Episode Duration	n.e.d.	n.e.d.	n.e.d.	n.e.d.	n.e.d.	n.e.d.
% of Patients with Left DLPFC Stimulation	0.0009	–0.01–0.02	0.9	0.004	–0.01–0.01	0.59
Motor Threshold	–0.009	–0.04–0.02	0.61	–0.007	–0.05–0.04	0.76
Number of Pulses/day	–0.0001	–0.001–0.0009	0.85	–0.0001	–0.0007–0.0005	0.77
Frequency of Stimulation	–0.05	–0.17–0.08	0.44	–0.60	–0.17–0.04	0.25
Number of Induction Days	–0.04	–0.15–0.07	0.46	–0.05	–0.15–0.05	0.34
Quality of the Study	–0.27	–0.54–0.002	0.05	–0.27	–0.59–0.05	0.1

DL-PFC: dorsolateral prefrontal cortex; rTMS: repeated transcranial magnetic stimulation; y: year.

n.e.d.: not enough data were gathered to run metaregressions on episode duration, disease duration and on number of previous episodes of depression.

n.a.: not applicable.

symptomatic re-introduction of rTMS were also not fully described, namely with regards to the exact criteria for re-introduction of rTMS. Ultimately, there was too much variability, and insufficient replication, to allow for clear meta-analytic comparisons. Thus, a definitive maintenance strategy or treatment schedule cannot be fully defined at this moment. Nevertheless, we believe these results highlight the need for controlled studies of symptomatic re-introduction and/or planned maintenance strategies for maintenance rTMS, that could provide further support not only for rTMS as a viable long-term treatment regimen for depressive disorders, but also help define which strategies should be used.

Our results are in line with what has been reported for durability of other neurostimulation approaches to treat MDD or TRD [61]. A meta-analysis assessing the relapse rate after a successful course of ECT in MDD patients showed that close to 40% of patients, both with and without maintenance ECT, relapsed after 6 months [30]. For tDCS 2 RCTs have assessed the long-term effectiveness of tDCS and both reported a relapse rate of approximately 50% at m6 follow-ups [2]. Regarding invasive neuromodulation, where stimulation is typically delivered chronically, VNS is associated with a 52% response rate after 6 months [62], while DBS of the ventral capsule/ventral striatum is associated with response rates of 43.7% at m3 and 40% at m6 follow-up [63]. Consistently, in a 2014 meta-analysis, Berlim and colleagues found that the 12-month response rate following DBS treatment was 39.9%. While these numbers are similar to those reported here, it is important to underline that patient populations in the several studies are not necessarily equivalent. Importantly, the patients included in our meta-analysis were not exclusively TRD patients, while all patients offered VNS or DBS suffer from TRD. Moreover for VNS or DBS studies response rates were computed for all patients included at the beginning of the studies, whereas in our and other meta-analyses [30] response rates during follow-up were computed for responders to an acute course of treatment.

In the assessment of the included publications, we did not find evidence of relevant publication bias, supporting the validity of our results. However, the data available for this study suffers from other potential limitations. First and foremost, while our inclusion and exclusion criteria were not overly restrictive, the number of studies with long enough follow-up phases was low. Furthermore, some did not report responder or relapse rates, and thus could not be considered for meta-analyses. Importantly, most studies reported data for only one or two time-points, such that the data reported for each time-point of the meta-analysis does not represent the same group of studies. Heterogeneity of study methodology or study population is also of concern. However, sensitivity analyses removing outlier studies or studies with atypical antidepressant response criteria (i.e., anything other than HAM-D $\geq 50\%$), as well as comparing studies including specific patient subgroups, essentially confirmed the findings in the full sample (supplementary fig 6). Importantly, we could not draw any strong conclusion regarding the durability of response to rTMS for MDD versus bipolar patients (supplementary Fig. 6d and 6e) nor for patients with versus without chronic depression (supplementary Fig. 6f and 6g). Future studies systematically and directly comparing these subgroups of patients are required to address these questions. Nevertheless, and while this study does not provide level 1 evidence given limitations in the available literature, it provides clear directions for additional well designed rTMS studies, for well-defined groups of patients, with at least 3 months' post-treatment follow-up, and sound reporting of outcome measures accordingly to common reported criteria, at standard time-points.

Another important limitation results from the fact that our analyses included almost exclusively open label and naturalistic studies. We believe this reflects important ethical and logistic

challenges in the conduction of long-term randomized and sham-controlled studies among depressed patients. In fact, one of the studies used in the meta-analyses described a randomized protocol, testing the efficacy of a planned rTMS maintenance protocol to prevent relapse [33], offered in addition to symptomatic re-introduction of rTMS that was available for all patients, and in the absence of a sham stimulation intervention for the control arm of the study. This limited the use of this study design to address the relevance of maintenance treatment, given that both study arms effectively received maintenance. Thus, we remained limited in the ability of extracting causal relationships from the data described here, since there is insufficient control for potential confounders. For example, it is possible that, during follow-up/maintenance phases, new antidepressants could be initiated, which could contribute towards the overall duration of the antidepressant response to induction rTMS and, albeit to a lesser degree, towards the advantages associated with maintenance rTMS. For these reasons, we strongly believe that studies with a RCT design are necessary to confirm the findings reported here, most importantly, to confirm the utility and define the optimal strategy for rTMS maintenance treatment. We expect that the findings reported here could help refine the design of future clinical trials in this field.

In conclusion, a systematic review and meta-analysis of the available literature suggests that rTMS is a durable treatment for depression, with sustained responder rates of 46–67%, from 3 months to 1 year after a successful initial treatment course. Of critical importance for current rTMS practice in depression treatment, maintenance rTMS delivered to those that respond to induction rTMS was found to enhance the rates of sustained responders across time-points, mainly 6 months after rTMS induction, when responder rates were almost 60% higher than among patients that did not receive maintenance rTMS.

Funding

SS was funded by *Assistance Publique des Hôpitaux de Paris*. APL was supported in part by the Berenson-Allen Foundation, the Sidney R. Baer, Jr. Foundation, the National Institutes of Health (R21 MH099196), and Harvard Catalyst | The Harvard Clinical and Translational Science Center (NCRR and the NCATS NIH, UL1 RR025758). Fundação para a Ciência e a Tecnologia funded GC through a Doctoral Scholarship (SFRH/BD/130210/2017), and AJO-M through a Junior Research and Career Development Award from the Harvard Medical School - Portugal Program (HMSP-ICJ/0020/2011). The content of this study is solely the responsibility of the authors and does not necessarily represent the official views of the Fundação para a Ciência e Tecnologia, Harvard Catalyst, Harvard University and its affiliated academic health care centers, the National Institutes of Health, or the Sidney R. Baer Jr. Foundation.

Declarations of interest

APL serves on the scientific advisory boards for Nexstim, NeuroRx, Starlab Neuroscience, Neuroelectronics, Axilum Robotics, and Neosync; and is listed as an inventor on several issued and pending patents on the real-time integration of transcranial magnetic stimulation with electroencephalography and magnetic resonance imaging. SS, GC and AJO-M report no competing interests.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brs.2018.10.001>.

References

- [1] Bauer M, Pfennig A, Severus E, Whybrow P, Angst J, Möller H. Task Force on Unipolar Depressive Disorders. World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for biological treatment of unipolar depressive disorders, part 1: update 2013 on the acute and continuation treatment of unipolar depressive disorders. *World J Biol Psychiatr* 2013;14(5):334–85.
- [2] Brunoni AR, Moffa AH, Fregni F, Palm U, Padberg F, Blumberger DM, et al. Transcranial direct current stimulation for acute major depressive episodes: meta-analysis of individual patient data. *Br J Psychiatr* 2016;208(6):522–31.
- [3] Cristancho P, Cristancho MA, Baltuch GH, Thase ME, O'Reardon JP. Effectiveness and safety of vagus nerve stimulation for severe treatment-resistant major depression in clinical practice after FDA approval: outcomes at 1 year. *J Clin Psychiatr* 2011;72(10):1376–82.
- [4] Kedzior KK, Reitz SK. Short-term efficacy of repetitive transcranial magnetic stimulation (rTMS) in depression—reanalysis of data from meta-analyses up to 2010. *BMC psychiatry* 2014b;2(1):39.
- [5] Mayberg HS, Lozano AM, Voon V, McNeely HE, Seminowicz D, Hamani C, et al. Deep brain stimulation for treatment-resistant depression. *Neuron* 2005;45(5):651–60.
- [6] Song G-M, Tian X, Shuai T, Yi L-J, Zeng Z, Liu S, et al. Treatment of adults with treatment-resistant depression: electroconvulsive therapy plus antidepressant or electroconvulsive therapy alone? Evidence from an indirect comparison meta-analysis. *Medicine* 2015;94(26): e1052.
- [7] Keller MB, Klerman GL, Lavori PW, Coryell W, Endicott J, Taylor J. Long-term outcome of episodes of major depression: clinical and public health significance. *J Am Med Assoc* 1984;252(6):788–92.
- [8] Kelly MS, Oliveira-Maia AJ, Bernstein M, Stern AP, Press DZ, Pascual-Leone A, et al. Initial response to transcranial magnetic stimulation treatment for depression predicts subsequent response. *J Neuropsychiatr Clin Neurosci* 2016;29(2):179–82.
- [9] Brunoni AR, Chaimani A, Moffa AH, Razza LB, Gattaz WF, Daskalakis ZJ, et al. Repetitive transcranial magnetic stimulation for the acute treatment of major depressive episodes: a systematic review with network meta-analysis. *JAMA psychiatry* 2017;74(2):143–52.
- [10] Kedzior KK, Azorina V, Reitz SK. More female patients and fewer stimuli per session are associated with the short-term antidepressant properties of repetitive transcranial magnetic stimulation (rTMS): a meta-analysis of 54 sham-controlled studies published between 1997–2013. *Neuropsychiatric Dis Treat* 2014a;10:727–56.
- [11] Lepping P, Schönfeldt-Lecuona C, Sambhi R, Lanka S, Lane S, Whittington R, et al. A systematic review of the clinical relevance of repetitive transcranial magnetic stimulation. *Acta Psychiatr Scand* 2014;130(5):326–41.
- [12] Liu B, Zhang Y, Zhang L, Li L. Repetitive transcranial magnetic stimulation as an augmentative strategy for treatment-resistant depression, a meta-analysis of randomized, double-blind and sham-controlled study. *BMC Psychiatr* 2014;14(1):342.
- [13] Berlim MT, Van den Eynde F, Daskalakis ZJ. Clinically meaningful efficacy and acceptability of low-frequency repetitive transcranial magnetic stimulation (rTMS) for treating primary major depression: a meta-analysis of randomized, double-blind and sham-controlled trials. *Neuropsychopharmacology* 2013a;38(4):543–51.
- [14] Berlim MT, Van den Eynde F, Daskalakis ZJ. High-frequency repetitive transcranial magnetic stimulation accelerates and enhances the clinical response to antidepressants in major depression: a meta-analysis of randomized, double-blind, and sham-controlled trials. *J Clin Psychiatr* 2013b;74(2):e122–9.
- [15] Burt T, Lisanby SH, Sackeim HA. Neuropsychiatric applications of transcranial magnetic stimulation: a meta analysis. *Int J Neuropsychopharmacol* 2002;5(1):73–103.
- [16] Herrmann LL, Ebmeier KP. Factors modifying the efficacy of transcranial magnetic stimulation in the treatment of depression: a review. *J Clin Psychiatr* 2006;67(12):1870–6.
- [17] Slotema CW, Dirk Blom J, Hoek HW, Sommer IE. Should we expand the toolbox of psychiatric treatment methods to include Repetitive Transcranial Magnetic Stimulation (rTMS)? A meta-analysis of the efficacy of rTMS in psychiatric disorders. *J Clin Psychiatr* 2010;71(7):873.
- [18] Avery DH, Holtzheimer 3rd PE, Fawaz W, Russo J, Neumaier J, Dunner DL, et al. A controlled study of repetitive transcranial magnetic stimulation in medication-resistant major depression. *Biol Psychiatry* 2006;59(2):187–94.
- [19] Stern WM, Tormos JM, Press DZ, Pearlman C, Pascual-Leone A. Antidepressant effects of high and low frequency repetitive transcranial magnetic stimulation to the dorsolateral prefrontal cortex: a double-blind, randomized, placebo-controlled trial. *J Neuropsychiatry* 2007;19(2):179–86.
- [20] Xie J, Chen J, Wei Q. Repetitive transcranial magnetic stimulation versus electroconvulsive therapy for major depression: a meta-analysis of stimulus parameter effects. *Neurol Res* 2013;35(10):1084–91.
- [21] Allan CL, Herrmann LL, Ebmeier KP. Transcranial magnetic stimulation in the management of mood disorders. *Neuropsychobiology* 2011;64(3):163–9.
- [22] Kedzior KK, Reitz SK, Azorina V, Loo C. Durability of the antidepressant effect of the high-frequency repetitive transcranial magnetic stimulation (rTMS) in the absence of maintenance treatment in major depression: a systematic review and meta-analysis of 16 double-blind, randomized, sham-controlled trials. *Depress Anxiety* 2015;32(3):193–203.
- [23] Martin JLR, Barbanjo MJ, Schlaepfer TE, Thompson E, Pérez V, Kulisevsky J. Repetitive transcranial magnetic stimulation for the treatment of depression. *Br J Psychiatr* 2003;182(6):480–91.
- [24] Hamani C, Mayberg H, Snyder B, Giacobbe P, Kennedy S, Lozano AM. Deep brain stimulation of the subcallosal cingulate gyrus for depression: anatomical location of active contacts in clinical responders and a suggested guideline for targeting: clinical article. *J Neurosurg* 2009;111(6):1209–15.
- [25] Lozano AM, Mayberg HS, Giacobbe P, Hamani C, Craddock RC, Kennedy SH. Subcallosal cingulate gyrus deep brain stimulation for treatment-resistant depression. *Biol Psychiatry* 2008;64(6):461–7.
- [26] Williams NR, Short EB, Hopkins T, Bentzley BS, Sahlem GL, Pannu J, et al. Five-year follow-up of bilateral epidural prefrontal cortical stimulation for treatment-resistant depression. *Brain stimulation* 2016;9(6):897–904.
- [27] Dunner DL, Aaronson ST, Sackeim HA, Janicak PG, Carpenter LL, Boyadjis T, et al. A multisite, naturalistic, observational study of transcranial magnetic stimulation for patients with pharmacoresistant major depressive disorder: durability of benefit over a 1-year follow-up period. *J Clin Psychiatr* 2014b;75(12):1394–401.
- [28] O'Reardon JP, Blumner KH, Peshek AD, Pradilla RR, Pimiento PC. Long-term maintenance therapy for major depressive disorder with rTMS. *J Clin Psychiatr* 2005;66(12):1524–8.
- [29] Rapinesi C, Bersani FS, Kotzalidis GD, Imperatori C, Del Casale A, Di Pietro S, et al. Maintenance deep transcranial magnetic stimulation sessions are associated with reduced depressive relapses in patients with unipolar or bipolar depression. *Front Neurol* 2015a;6:16.
- [30] Jelovac A, Kolshus E, McLoughlin DM. Relapse following successful electroconvulsive therapy for major depression: a meta-analysis. *Neuropsychopharmacology* 2013;38(12):2467–74.
- [31] Bermudes RA, Lanocha KI, Janicak PG. Transcranial magnetic stimulation: clinical applications for psychiatric practice. *American Psychiatric Pub*; 2017.
- [32] Loo CK, McFarquhar TF, Mitchell PB. A review of the safety of repetitive transcranial magnetic stimulation as a clinical treatment for depression. *Int J Neuropsychopharmacol* 2008;11(1):131–47.
- [33] Philip NS, Dunner DL, Dowd SM, Aaronson ST, Brock DG, Carpenter LL, et al. Can medication free, treatment-resistant, depressed patients who initially respond to TMS be maintained off medications? A prospective, 12-month multisite randomized pilot study. *Brain stimulation* 2016;9(2):251–7.
- [34] Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7): e1000097.
- [35] Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. Newcastle-Ottawa quality assessment scale cohort studies. 2014.
- [36] Berlim MT, Turecki G. Definition, assessment, and staging of treatment-resistant refractory major depression: a review of current concepts and methods. *Can J Psychiatr* 2007;52(1):46–54.
- [37] Thase ME, Rush AJ. When at first you don't succeed: sequential strategies for antidepressant nonresponders. *J Clin Psychiatr* 1997;58(Suppl 13):23–9.
- [38] Benadhira R, Thomas F, Bouaziz N, Braha S, Andrianisaina PS, Isaac C, et al. A randomized, sham-controlled study of maintenance rTMS for treatment-resistant depression (TRD). *Psychiatr Res* 2017;258:226–33.
- [39] Boutros NN, Gueorguieva R, Hoffman RE, Oren DA, Feingold A, Berman RM. Lack of a therapeutic effect of a 2-week sub-threshold transcranial magnetic stimulation course for treatment-resistant depression. *Psychiatr Res* 2002;113(3):245–54.
- [40] Benadhira R, Saba G, Samaan A, Dumortier G, Lipski H, Gastal D, et al. Transcranial magnetic stimulation for refractory depression. *Am J Psychiatry* 2005;162(1): 193–193.
- [41] Dannon PN, Dolberg OT, Schreiber S, Grunhaus L. Three and six-month outcome following courses of either ECT or rTMS in a population of severely depressed individuals—preliminary report. *Biol Psychiatry* 2002;51(8):687–90.
- [42] Dell'Osso B, D'urso N, Castellano F, Ciabatti M, Altamura AC. Long-term efficacy after acute augmentative repetitive transcranial magnetic stimulation in bipolar depression: a 1-year follow-up study. *J ECT* 2011;27(2):141–4.
- [43] Dunner DL, Aaronson ST, Sackeim HA, Janicak PG, Carpenter LL, Boyadjis T, et al. A multisite, naturalistic, observational study of transcranial magnetic stimulation for patients with pharmacoresistant major depressive disorder: durability of benefit over a 1-year follow-up period. *J Clin Psychiatr* 2014a;75(12):1394–401.
- [44] Eranti S, Mogg A, Pluck G, Landau S, Purvis R, Brown RG, et al. A randomized, controlled trial with 6-month follow-up of repetitive transcranial magnetic stimulation and electroconvulsive therapy for severe depression. *Am J Psychiatry* 2007;164(1):73–81.
- [45] Fitzgerald PB, Grace N, Hoy KE, Bailey M, Daskalakis ZJ. An open label trial of clustered maintenance rTMS for patients with refractory depression. *Brain stimulation* 2013;6(3):292–7.
- [46] Harel EV, Rabany L, Deutsch L, Bloch Y, Zangen A, Levkovitz Y. H-coil repetitive transcranial magnetic stimulation for treatment resistant major depressive disorder: an 18-week continuation safety and feasibility study. *World J Biol Psychiatr* 2014;15(4):298–306.
- [47] Janicak PG, Nahas Z, Lisanby SH, Solvason HB, Sampson SM, McDonald WM, et al. Durability of clinical benefit with transcranial magnetic stimulation (TMS) in the treatment of pharmacoresistant major depression: assessment of relapse during a 6-month, multisite, open-label study. *Brain stimulation* 2010;3(4):187–99.

- [48] Li X, Nahas Z, Anderson B, Kozel FA, George MS. Can left prefrontal rTMS be used as a maintenance treatment for bipolar depression? *Depress Anxiety* 2004;20(2):98–100.
- [49] López-Ibor Aliño J, Pastrana Jiménez J, Cisneros Flores S, López-Ibor Alcocer M. Eficacia de la estimulación magnética transcraneal en depresión. Estudio naturalístico. *Actas Esp Psiquiatr* 2010;38(2):87–93.
- [50] Mantovani A, Pavlicova M, Avery D, Nahas Z, McDonald WM, Wajdik CD, et al. Long-term efficacy of repeated daily prefrontal transcranial magnetic stimulation (TMS) in treatment-resistant depression. *Depress Anxiety* 2012;29(10):883–90.
- [51] Rapinesi C, Curto M, Kotzalidis GD, Del Casale A, Serata D, Ferri VR, et al. Antidepressant effectiveness of deep transcranial magnetic stimulation (dTMS) in patients with major depressive disorder (MDD) with or without alcohol use disorders (AUDs): a 6-month, open label, follow-up study. *J Affect Disord* 2015b;174:57–63.
- [52] Richieri R, Guedj E, Michel P, Loundou A, Auquier P, Lançon C, et al. Maintenance transcranial magnetic stimulation reduces depression relapse: a propensity-adjusted analysis. *J Affect Disord* 2013;151(1):129–35.
- [53] Rosenberg O, Isserles M, Levkovitz Y, Kotler M, Zangen A, Dannon P. Effectiveness of a second deep TMS in depression: a brief report. *Prog Neuro Psychopharmacol Biol Psychiatr* 2011;35(4):1041–4.
- [54] Cohen RB, Boggio PS, Fregni F. Risk factors for relapse after remission with repetitive transcranial magnetic stimulation for the treatment of depression. *Depress Anxiety* 2009;26(7):682–8.
- [55] Connolly KR, Helmer A, Cristancho MA, Cristancho P, O'Reardon JP. Effectiveness of transcranial magnetic stimulation in clinical practice post-FDA approval in the United States: results observed with the first 100 consecutive cases of depression at an academic medical center. *J Clin Psychiatr* 2012;73(4):e567–73.
- [56] Koerselman F, Laman DM, van Duijn H, Van Duijn M, Willems M. A 3-month, follow-up, randomized, placebo-controlled study of repetitive transcranial magnetic stimulation in depression. *J Clin Psychiatr* 2004;65(10):1323–8.
- [57] Levkovitz Y, Isserles M, Padberg F, Lisanby SH, Bystritsky A, Xia G, et al. Efficacy and safety of deep transcranial magnetic stimulation for major depression: a prospective multicenter randomized controlled trial. *World Psychiatr* 2015;14(1):64–73.
- [58] Mogg A, Pluck G, Eranti S, Landau S, Purvis R, Brown R, et al. A randomized controlled trial with 4-month follow-up of adjunctive repetitive transcranial magnetic stimulation of the left prefrontal cortex for depression. *Psychol Med* 2008;38(03):323–33.
- [59] Mann C. Observational research methods. Research design II: cohort, cross sectional, and case-control studies. *Emerg Med J* 2003;20(1):54–60.
- [60] Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMC Med* 2010;8(1):18.
- [61] Akhtar H, Bukhari F, Nazir M, Anwar MN, Shahzad A. Therapeutic efficacy of neurostimulation for depression: techniques, current modalities, and future challenges. *Neuroscience bulletin* 2016;32(1):115–26.
- [62] Marangell LB, Rush AJ, George MS, Sackeim HA, Johnson CR, Husain MM, et al. Vagus nerve stimulation (VNS) for major depressive episodes: one year outcomes. *Biol Psychiatry* 2002;51(4):280–7.
- [63] Malone DA, Dougherty DD, Rezai AR, Carpenter LL, Friehs GM, Eskandar EN, et al. Deep brain stimulation of the ventral capsule/ventral striatum for treatment-resistant depression. *Biol Psychiatry* 2009;65(4):267–75.