

Dysfunctional Neural Processes Underlying Context Processing Deficits in Schizophrenia

Seung Suk Kang, Angus W. MacDonald, and Scott R. Sponheim

ABSTRACT

BACKGROUND: People with schizophrenia (PSZ) have profound deficits in context processing, an executive process that guides adaptive behaviors according to goals and stored contextual information. Although various neural processes are involved in context processing and are affected in PSZ, the core underlying neural dysfunction is unclear.

METHODS: To determine the relative importance of neural dysfunctions within prefrontal cognitive control, sensory activity, and motor activity to context processing deficits in PSZ, we examined event-related potentials (ERPs) in 60 PSZ and 51 healthy control subjects during an optimal context processing task. We also analyzed the Ex-Gaussian reaction time distribution to examine abnormalities in motor control variability in PSZ.

RESULTS: Compared with healthy control subjects, PSZ had lower response accuracy and greater variability in their normal reaction times during high context processing demands. Latencies of normal and slow responses were generally increased in PSZ. High context processing-related reductions in frontal ERPs were indicative of specific deficits in proactive and reactive cognitive controls in PSZ, while ERPs associated with visual and motor processes were reduced regardless of context processing demands, indicating generalized visuomotor deficits. In contrast to previous studies, we found that diminished frontal responses reflective of proactive control of the contextual cue, rather than visual responses of cue encoding, predicted response accuracy deficits in PSZ. In addition, probe-related ERP components of motor preparation, prefrontal reactive control, and frontomotor interaction predicted Ex-Gaussian indices of reaction time instability in PSZ.

CONCLUSIONS: Prefrontal proactive and reactive control deficits associated with failures in using mental representation likely underlie context processing deficits in PSZ.

Keywords: Context processing, ERPs, Prefrontal dysfunction, Reaction time variability, Schizophrenia, Visuomotor deficits

<https://doi.org/10.1016/j.bpsc.2019.03.012>

A deficit in context processing and the associated prefrontal network dysfunction has been identified as central to schizophrenia (1). Context processing involves the control of behaviors according to goals or other stored abstract representations that reflect contextual information (e.g., rules) (2) and is considered an executive function related to goal representation. Experimental studies using tasks that require updating of response rules based on previous cues (e.g., AX version of the continuous performance task [AX-CPT]) have demonstrated that people with schizophrenia (PSZ) show deficits in using contextual cue information to guide responses (3,4). Context processing deficits were observed from the onset of the disease (5) and were predictive of the long-term outcome (6). In addition, the deficits have been identified as a specific deficit (i.e., not attributable to generalized cognitive impairments) that is associated with genetic liability for schizophrenia (3,4) and therefore may reflect pathology central to the disorder (1,5,7). Thus, it is imperative to determine the underlying abnormal neural mechanisms in PSZ. Context processing requires coordination among various neural

processes, including sensory encoding of context information, cognitive controls to guide sensory-motor processes, and preparation and execution of goal-relevant motor responses. Abnormalities in all these processes have been recognized in PSZ (8–10), so the process most responsible for context processing deficits in PSZ remains controversial (11).

The control elements of context processing have been conceptualized according to the dual mechanisms of control theory, which specifies two distinct prefrontal cognitive control modes with different temporal dynamics (12,13). The proactive control mode refers to the early selection of goal-relevant information to update contextual rules or active maintenance of such information in working memory (WM) prior to a behavioral response (e.g., contextual cue processing in the AX-CPT), which in turn biases attentional and perceptual neural processes toward goal-directed behaviors. In contrast, the reactive control mode encompasses functions that are recruited for the detection and resolution of behavioral conflicts at the time a response is required (e.g., response control in Stroop task). Functional magnetic resonance imaging (fMRI) studies

Context Processing Deficits in Schizophrenia

comparing neural responses between the AX-CPT and Stroop task (14) and those between cue and probe periods of the AX-CPT (15) revealed that PSZ had prefrontal cortical responses consistent with deficient proactive control but relatively intact reactive control. However, the AX-CPT also requires reactive control, and the slow time course of hemodynamic responses in fMRI provides incomplete differentiation of cue-related versus probe-related neural responses. A recent electroencephalogram (EEG) study using the AX-CPT identified deficient midline theta oscillations associated with specific deficits in proactive control and generalized deficits in reactive control in PSZ (16). The disproportionate frequencies of AX/AY/BX/BY conditions in the task, however, might lead to confounding effects of neural responses associated with probability in addition to those of context processing. Therefore, the use of electrophysiological measurements and an optimal task without biases toward a specific condition will improve the functional neural measurements of proactive and reactive control processes to better examine the nature of the context processing deficit in schizophrenia.

In contrast to a prefrontal cognitive control impairment in schizophrenia, recent event-related potential (ERP) studies implied that failures in visual systems accounted for context processing (11) and WM (17) impairments in schizophrenia. In particular, Dias *et al.* (11) used the AX-CPT to show that deficient occipital responses for encoding contextual information better explained context processing impairments in PSZ than prefrontal responses. They also found that PSZ had intact top-down modulation of visual processing despite overall reductions in the visual responses, suggesting that early visual stream deficits rather than prefrontal top-down control are the primary cause of deficits in context processing and WM in schizophrenia. However, the task used in this work had much greater visual processing demands than commonly used; therefore, it is necessary to test context processing in PSZ without biases toward a specific neural process.

Context processing deficits in PSZ might also be affected by psychomotor deficits that have been clinically documented (18,19) and demonstrated with reaction time (RT) and ERP motor components such as lateralized readiness potential (LRP) (20–22). In this regard, analysis of intraindividual variability (i.e., trial-by-trial fluctuation) in RTs may provide rich information on instability in cognitive processes probing pathophysiology in PSZ (23) unavailable to conventional mean RT analyses (24,25). As illustrated in Figure 1C, an Ex-Gaussian distribution analysis of RT decomposes a typical positively skewed RT distribution into the parameters that estimate one's normal motor speed (μ : mean of Gaussian component), intraindividual RT variability of normal responses (σ : SD of Gaussian component) that reflect trial-by-trial alterations in response processes, and finally an attentional lapse component that contributes to slow responses (τ : mean of exponential component) (24). Investigation of motor cortical activities along with the RT variability indices might further capture a problem with efficiently processing information during context processing in PSZ.

The goal of the current study was to identify behavioral features of context processing deficits and determine relative impairments in neural processes related to prefrontal cognitive controls, sensory and motor activities, and their contributions

to context processing deficits in PSZ. To identify the specific deficits in cognitive and neural processes in PSZ, we used the Stimulus Response Reversal Task (SRRT), which measures context processing without biases toward specific condition or neural processes (i.e., equivalent probabilities of conditions and sufficient stimulus duration for visual encoding). The SRRT requires both proactive controls to update response rules and reactive controls to initiate a goal-relevant response, overriding an interference of the automatic response triggered by the probe stimulus. We also used ERP components to isolate the timing of neural responses associated with visual, cognitive control, and motor functions in context processing (see Figure 1B): 1) visual ERP components, including cue and probe period visual N1 (cvN1 and pvN1, respectively) that reflect ventral occipital-temporal cortical functions of visual encoding (11,26,27); 2) frontal cognitive components of negative potentials, including cue-locked frontal N2 (cfN2) and pre-response frontal negativity (prfN) that are known to reflect prefrontal proactive control updating contextual information (11,28) and prefrontal reactive control activity for conflict resolution (29,30), respectively; and 3) LRP reflecting preparatory motor processes (31). Along with a response accuracy (i.e., %Correct), we analyzed the three Ex-Gaussian RT parameters to examine various aspects of disturbances in motor control during context processing. We tested whether PSZ had differential deficits for context processing in ERP components reflecting proactive/reactive controls and visual and motor processes. We hypothesized that ERP indices of prefrontal proactive and reactive controls rather than visual and motor processing would primarily explain performance deficits (i.e., low accuracy and high latency variability in behavioral responses) in high context processing (HCP) load in PSZ.

METHODS AND MATERIALS


Participants

In total, 60 PSZ and 51 healthy control subjects (HCs) participated. The demographic and clinical characteristics of the participants are summarized in Supplemental Table S1. Patients were recruited from hospitals, clinics, residential facilities, and community support programs for individuals with severe mental disorders in the Minneapolis–St. Paul metropolitan area of Minnesota. Diagnostic information was obtained by a trained interviewer completing the Structured Clinical Interview for DSM-IV-TR Axis I Disorders (32) and the Psychosis Module of the Diagnostic Interview for Genetic Studies (33) with each participant. Only patients who met DSM-IV TR criteria for schizophrenia based on consensus diagnosis of two raters were included in this report. HCs were recruited through advertising broadly in the metropolitan community. Inclusion/exclusion criteria are described extensively elsewhere (34). The Minneapolis VA Medical Center and University of Minnesota Institutional Review Boards approved the protocol.


SRRT and Evaluation of Task Performances

In the SRRT, subjects were asked to select responses to a directional probe word (“Left” or “Right”) according to a contextual cue (a red or green rectangle) presented prior to the

A Procedures of SRRT



B Cognitive Processes and ERP Components in SRRT ERP Paradigm



C Ex-Gaussian RT Distribution Analysis




Figure 1. (A) Schematic of Stimulus Response Reversal Task (SRRT) with low context processing (LCP) and high context processing (HCP) trial types. In this task, subjects were asked to select responses to a directional probe word (“Left” or “Right”) according to a contextual cue (a red or green rectangle) presented prior to the probe. Each trial began with a central cross fixation, which was followed by a green or red square cue that was presented for 650 ms. After a 3000- to 4000-ms delay, a probe word (“Right” or “Left”) was presented for 650 ms. The next trial began 2000 ms after probe stimuli offset. (B) Diagram depicting the sequence of cognitive processes and event-related potential (ERP) components in the SRRT ERP paradigm that consisted of context processing during the task. (C) Example of Ex-Gaussian reaction time (RT) distribution from a subject’s correct response latency of SRRT HCP trials. Ex-Gaussian RT distribution is a convolution of the two additive processes (i.e., Gaussian and exponential processes). The μ and σ represent the mean and SD, respectively, of the normal (Gaussian) component of RT distribution, reflecting the mean speed and variability of normal responses. Thus, a large σ indicates increased trial-to-trial jittering of usual responses that might be caused by instability of motor processes. On the other hand, τ represents the mean of the exponential component of RT distribution, reflecting the mean of extremely slow responses (i.e., outlier responses). Thus, a larger τ indicates a longer and thicker tail of exponential distributions of slow responses that often occur due to attentional lapse. The mean and SD of the Ex-Gaussian distribution are equal to $\mu + \tau$ and $\sqrt{\sigma^2 + \tau^2}$, respectively. ITI, intertrial interval.

probe. In the low context processing (LCP; green cue) condition, subjects needed to respond with the thumb of the hand indicated by a probe. In the HCP (red cue) condition, they

needed to respond with the thumb of the opposite hand. The HCP and LCP conditions were intermixed with the equivalent frequency in a block, which consisted of 71 test trials. Subjects

Context Processing Deficits in Schizophrenia

completed three blocks of SRRT trials (see Figure 1A for details of a trial sequence and timings of the SRRT). We computed the %Correct index of response accuracy and estimated μ , σ , and τ parameters of an Ex-Gaussian distribution for central tendency and variability of RT during context processing using only correct trials (see Figure 1C for details). To this end, we used the *retimes* R package (<https://cran.r-project.org/web/packages/retimes/>) that uses methods described by Heathcote (35).

EEG Recordings and Processes

EEG data were collected at a sampling rate of 1024 Hz using a 64- or 128-channel BioSemi ActiveTwo EEG system (<http://www.biosemi.com>) while participants performed the SRRT. All EEG processing was conducted using MATLAB (The Mathworks Inc., Natick, MA), including low-pass filtering (256 Hz cutoff frequency), high-pass filtering (0.5 Hz cutoff frequency), downsampling to 256 Hz using the *antialiasing* resample function, and referencing to linked earlobe signal. Noisy electrode signals were identified via visual inspection and excluded from processing. EEG was segmented to three types of epochs: time-locked to cue onsets (−500 to 2000 ms), probe onsets (−500 to 1500 ms), and response onsets (−1100 to 100 ms). Atypical noisy time segments and physiological and electrode-specific artifacts were removed using a custom MATLAB-based pipeline using independent component analysis (see Supplement for details). After independent component analysis denoising, excluded bad electrode signals were interpolated using the spherical spline method (36) with optimal interpolation parameters (37). Finally, all scalp EEG signals were referenced to the average signal to produce an unbiased topographical distribution of EEG signals, and then 64- and 128-channel EEG data were converted to an 81-channel 10-10 EEG standard montage using BESA (<https://www.besa.de>) to produce topographical distributions of ERP components with the same montage.

ERP Analyses

Cue-, probe-, and response-locked ERPs were computed by averaging corresponding EEG epochs filtered at 30-Hz low pass for trials with correct responses. Baseline correction was conducted by subtracting mean amplitudes of 100 ms prestimulus time window (i.e., precue 100 ms for cue-locked ERPs and preprobe 100 ms for probe- and response-locked ERPs). Spatiotemporal windows for ERP quantification and analyses were determined based on previous studies with minor modification to account for ERP differences due to differences in the stimuli and temporal structure of the tasks. Visual ERP components (cvN1 and pvN1) were quantified at bilateral occipitotemporal sites (10,26). Frontal cognitive components of cfN2 (11,28) and prfN (29) were quantified at Fz and FCz, respectively, along with bilateral frontal sites (F3 and F4) where their HCP–LCP amplitude differences were maximal. We included the bilateral occipital and frontal electrodes to test whether PSZ had lateralized deficits in the neural processes as reported in neuroimaging studies (5,38). Finally, the LRP being typically most obvious around 100 ms prior to a response in bilateral central electrodes (C3 and C4) was quantified using the formula isolating interhemispheric differences in motor

potentials (31). The onset latency of the LRP was determined as the time point at which the amplitude reached 50% of its peak amplitude (21). To minimize high-frequency signals causing measurement error in the LRP onset latency, we applied an 8-Hz low-pass filter to the LRP. See details of ERP analyses in the Supplement.

Statistical Analyses

For behavioral performance and ERP components, repeated-measures analyses of variance with a between-group factor of diagnostic group (PSZ or HCs) and a within-group factor of cue condition (HCP or LCP) were conducted for dependent variables of behavioral indices and each ERP component. Any significant interaction effect was further analyzed with analysis of variance and *t* tests. Effect sizes were estimated using η^2 and Cohen's *d* formulas. To identify neural processes predictive of context processing abilities, we conducted multiple regression analyses with predictors being amplitudes of the ERP component amplitudes and the dependent variable being one of the four task performance indices (i.e., %Correct, μ , σ , or τ). To control problems due to multicollinearity among the predictors (39,40), we conducted ridge regression analyses using the *ridge* R package (<https://cran.r-project.org/web/packages/ridge/index.html>) that automatically determines an optimal ridge regularization parameter and provides significance tests of predictors (41,42) (see Supplement for details). To test whether the cognitive and neural processes during the SRRT predict schizophrenic symptom severity, we also conducted ridge regression analyses with the predictors and dependent variables of Brief Psychiatric Rating Scale symptom scores.

RESULTS

Behavioral Performance

Repeated-measures analysis of variance results of performance indices are depicted in Figure 2 and summarized in Supplemental Table S3. There was a group-by-condition interaction effect on response accuracy (%Correct), indicating differential performance deficits of PSZ (i.e., lower accuracy in the HCP condition [$t_{109} = -2.33, p = .022, d = -0.44$] but, importantly, not in the LCP condition [$t_{109} = -1.53, p = .129, d = -0.29$]) (see Figure 2). Similarly, σ had a group-by-condition interaction characterized by a significant group difference only in the HCP condition (HCP: $t_{109} = 4.45, p < 10^{-4}, d = 0.85$; LCP: $t_{109} = 1.53, p = .130, d = 0.29$). Unlike HCs, PSZ had larger σ for HCP trials than for LCP trials ($t_{58} = 2.12, p = .039, d = 0.28$), indicating instability of controlled responses in PSZ for HCP. In contrast, μ and τ had group and condition effects but no group-by-condition interaction, indicating longer latency of PSZ than of HCs in both normal and slow responses regardless of the context processing load. Although the group-by-condition interaction effect was not significant in τ , only PSZ had significantly larger τ in the LCP condition than in the HCP condition ($t_{58} = 2.42, p = .019, d = 0.31$).

Visual ERP Components

The cvN1 amplitude had no group difference in any condition and electrode but had a condition difference (LCP > HCP)




Figure 2. Response accuracy (%Correct) and the three parameters of exponentially modified Gaussian reaction time (RT) distribution of Stimulus Response Reversal Task. People with schizophrenia (PSZ) had response accuracy deficits only in the high context processing (HCP) condition (A), suggesting a deficit specific to trials requiring controlled motor responses. Regardless of context-processing load, PSZ had longer mean latency than healthy control subjects (HCs) in normal (μ) (B) and slow (τ) responses (D) that were differentiated by deconvolution of an RT distribution into a Gaussian distribution and an exponential RT distribution, indicating generalized problems in motor control in PSZ. In contrast, they had larger variability in normal responses (σ) (C) than HCs only for high context processing trials, suggesting that PSZ had specific deficits of unstable motor controls especially on trials where they should override a pre-potent response and initiate a context-relevant motor response. Note that one patient with schizophrenia who had difficulty in understanding how to perform the task and extremely low response accuracy was excluded from performance analyses. * $p < .05$, *** $p < .0005$. LCP, low context processing.

(Figure 3), perhaps because the green cue had higher luminance than the red cue (43). It also had significant group-by-electrode and trend-level group-by-condition-by-electrode interaction effects, which were explained by differential condition-by-electrode interaction effects in the two groups; HCs had the condition effect only in P7 electrode ($t_{51} = 2.50$, $p = .016$), while PSZ had the condition effect only in P8 electrode ($t_{56} = 2.24$, $p = .029$). The pvN1 peak amplitude had significant group (PSZ < HCs) and condition (HCP > LCP) effects but had no other effect. The pvN1 peak latency had a group effect ($F_{1,109} = 7.76$, $p = .006$, $\eta^2 = .07$; PSZ > HCs) and an electrode effect ($F_{1,109} = 5.60$, $p = .020$, $\eta^2 = .05$; PO7 > PO8) while cvN1 peak latency had no such effects. These findings suggest that impairments in early visual processes in PSZ did not interact with the context processing demands of the task.

Proactive and Reactive Control ERP Components

The cfN2 amplitudes had group (PSZ < HCs), electrode (Fz > F3 > F4), group-by-condition interaction, and group-by-condition-by-electrode interaction effects (Figure 4). Follow-up tests found a significant group difference in the HCP condition ($F_{1,107} = 8.02$, $p = .006$, $\eta^2 = .07$) but not in the LCP condition ($F_{1,107} = 2.47$, $p = .119$, $\eta^2 = .02$). Such a differential group difference (i.e., group-by-condition interaction) was observed in Fz electrode ($F_{1,107} = 4.87$, $p = .030$, $\eta^2 = .04$) and F3 electrode ($F_{1,107} = 9.49$, $p = .003$, $\eta^2 = .08$) but not in F4 electrode ($F_{1,107} = 0.01$, $p = .937$, $\eta^2 < .01$). The prfN amplitude had group (PSZ < HCs), condition (LCP > HCP), and trend-level electrode (FCz > F3 > F4) effects. Group-by-condition and group-by-electrode interaction effects were also significant for prfN. Follow-up analyses found prfN group differences

in the HCP condition ($F_{1,108} = 12.84$, $p = .001$, $\eta^2 = .11$) but not in the LCP condition ($F_{1,108} = 2.80$, $p = .093$, $\eta^2 = .03$). In addition, prfN group difference were significant at FCz electrode ($F_{1,108} = 16.96$, $p < 10^{-4}$, $\eta^2 = .14$) and F3 electrode ($F_{1,108} = 4.74$, $p = .032$, $\eta^2 = .04$) but not at F4 electrode ($F_{1,108} = 0.92$, $p = .340$, $\eta^2 = .01$).

Motor ERP Component

LRP amplitude was not different between conditions but was largely reduced in PSZ compared with HCs regardless of context processing loads (Figure 5). LRP onset latency had no group effect ($F_{1,101} = 1.11$, $p = .294$, $\eta^2 = .01$) but showed a trend-level condition effect ($F_{1,101} = 3.59$, $p = .061$, $\eta^2 = .03$; HCP > LCP) and a significant group-by-condition interaction effect ($F_{1,101} = 5.32$, $p = .023$, $\eta^2 = .05$). Follow-up analysis found that PSZ had longer LRP onset latency than HCs in the HCP condition ($t_{101} = 2.14$, $p = .035$, $d = 0.42$) but not in the LCP condition ($t_{101} = 0.42$, $p = .677$, $d = 0.08$). To further examine whether this HCP-specific LRP problem relates to dysfunctional prefrontal-motor coordination for context-dependent motor control, we examined correlations between motor LRP and cognitive prfN amplitudes. We found that HCs had significant correlation between LRP and prfN amplitudes in the HCP condition ($r_{51} = .35$, $p = .011$) but not in the LCP condition ($r_{51} = .16$, $p = .261$). PSZ showed no correlation in either condition (Figure 5D). Given the group-by-condition interaction pattern of the correlation (i.e., significant group difference in the correlation for the HCP condition [$z = 2.19$, $p = .029$] but not for the LCP condition [$z = 1.49$, $p = .136$]), the interaction between LRP and prfN might be indicative of specific deficits in prefrontal-motor coordination for reactive control in PSZ.




Figure 3. Visual event-related potential components for cue (left) and probe (right) stimuli. **(A, B)** Topographic distributions of the visual N1 components in response to cue and probe stimuli (cvN1 [left] and pvN1 [right]) in healthy control subjects (HCs) and people with schizophrenia (PSZ), which were quantified by mean amplitudes within the time windows (130–230 ms after stimulus onsets) for the low context processing (LCP) and high context processing (HCP) conditions as well as the HCP–LCP difference. The cvN1 was maximal at P7 and P8, while the pvN1 was maximal at PO7 and PO8. **(C, D)** Event-related potential waveforms time-locked to the onsets of visual stimuli, where the time windows used for the peak amplitude and latency quantification are marked by shaded areas (130–230 ms after onsets of the visual stimuli). **(E, F)** Peak amplitudes of the cue and probe N1 components measured in P7/P8 and PO7/PO8 electrodes. No group difference was observed in the cvN1, which was larger for LCP cues (green) than for HCP cues (red); HCs had the significant difference between cue types at P7, while PSZ showed the cue type effect at P8. The pvN1 was diminished in PSZ compared with HCs, especially in PO8, but both groups had larger pvN1 amplitudes in the HCP condition. * $p < 1.0$, * $p < .05$, ** $p < .005$.

Relationships Among ERP Components, Task Performance Indices, and Schizophrenic Symptoms

In the ridge regression analyses with dependent variables of performance indices and predictors of ERP components, we included the interaction term between the response period prefrontal and motor ERPs (i.e., the product of prfN and LRP amplitudes [$\text{prfN} \times \text{LRP}$]) as a predictor of the frontomotor coordination process of reactive control. Only the ERP component of prefrontal proactive control (cfN2) predicted performance accuracy. The mean and variability of normal responses (μ and σ) were primarily predicted by LRP amplitude and the frontomotor interaction component ($\text{prfN} \times \text{LRP}$). Lastly, the mean of slow response latencies (τ) was primarily predicted by the prefrontal reactive control component (prfN). In the regressions predicting Brief Psychiatric Rating Scale

scores, we also found that $\text{prfN} \times \text{LRP}$ predicted severity of overall and disorganization symptoms of schizophrenia (see Supplemental Tables S4 and S5 and Supplemental Figure S2 for details).

DISCUSSION

This study revealed that PSZ showed poor sensitivity of frontal brain responses to contextual information, which predicted poor use of context to guide behavioral responses. Particularly in situations that required overriding prepotent responses based on contextual information (HCP condition), PSZ had lower accuracy and larger RT variability (σ) than HCs, but not when such control demands were low (LCP condition). In contrast, PSZ also had increased latency in normal and slow responses (μ and τ) regardless of context processing load,




Figure 4. Cue period frontal N2 event-related potential component (cfN2) representing prefrontal proactive control (left) and prereshponse frontal negativity (prfN) component representing prefrontal reactive control (right). **(A, B)** Topographic distributions of cfN2 and prfN components in healthy control subjects (HCs) and people with schizophrenia (PSZ), which were quantified by mean amplitudes within the time windows (cfN2: 220–350 ms after cue onsets; prfN: 500–200 ms before response onsets) for the low context processing (LCP) and high context processing (HCP) conditions as well as the HCP–LCP difference. The cfN2 was quantified at F3, Fz, and F4 sites, while the prfN was quantified at F3, FCz, and F4 sites. **(C, D)** Event-related potential waveforms time-locked to the cue and response onsets, where the time windows used for the mean amplitudes are marked by shaded areas. **(E, F)** Mean amplitudes of the cfN2 and prfN components were larger for the HCP condition than for the LCP condition. The components were diminished in PSZ compared with HCs, but only in the HCP condition. These differences were observed at F3 and Fz sites (cfN2) or at FCz site (prfN). The condition effect was absent (cfN2) or diminished (prfN) in PSZ compared with HCs. These results suggested that PSZ had deficits in prefrontal proactive and reactive controls that were specific to HCP demands. * $p < .05$, ** $p < .005$, *** $p < .0005$.

suggesting additional generalized deficits in motor preparation and attentional vigilance. During encoding of contextual cues, PSZ had intact visual responses (cvN1) but reduced frontal cognitive responses (cfN2), pointing to deficient prefrontal proactive control processes but normal perception of simple and clearly discernible visual cue stimuli. In response to probe letters that followed cues, PSZ had diminished and delayed visual responses (pvN1) and reduced frontal responses prior to button presses (prfN), indicating deficits in visual processing of letter stimuli and prefrontal reactive control. PSZ also exhibited abnormal motor preparation processes that were associated with the speed of button presses to probes. In both the LCP and HCP conditions, PSZ had diminished motor potentials

(LRPs) leading up to a behavioral response, but they showed a delayed LRP onset latency only for HCP trials, suggesting problems in controlled motor processes. None of the task performance and ERP indices was explained by antipsychotic medication dosage (see Supplement).

Unlike previous reports (11) emphasizing sensory processing deficits (i.e., the bottom-up hypothesis of schizophrenia) (44), the current study revealed that response accuracy during a context processing task was predicted by prefrontal proactive control (cfN2) rather than neural responses for visual encoding of contextual cues (cvN1). Consistent with our recent findings that encoding period theta frequency neural responses in prefrontal cortices rather than visual cortices was




Figure 5. Lateralized readiness potentials (LRPs) time-locked to button response onsets and measured as the difference between electrodes over the motor cortex of left and right hemispheres (C3 and C4). **(A)** LRPs were maximal around 100 ms prior to response onsets. Two time windows marked by the two shaded areas were used for LRP quantification for the mean amplitudes (50–200 ms before responses) and the onset latency (50–350 ms before responses) because the onset latency was often substantially earlier than the ranges of the peak LRP amplitudes, as observed in previous studies (19). The dashed lines represent LRP onset latency indicating the time point when the LRP reached 50% of its maximum amplitude. **(B)** People with schizophrenia (PSZ) had largely diminished LRP amplitudes regardless of the task condition. **(C)** PSZ had longer LRP onset latency than healthy control subjects (HCs) only in the high context processing (HCP) condition. **(D)** The relationships between pre-response frontal negativity (prfN) event-related potential component representing prefrontal reactive control and LRP component representing preparatory motor cortical activity during the low context processing (LCP) and HCP conditions in two groups. HCs had significant positive correlation between prfN and LRP amplitudes in the HCP condition ($r_{51} = .35$, $p = .01$) but not in the LCP condition ($r_{51} = .16$, $p = .261$), while PSZ failed to show an association in either context processing condition (HCP: $r_{52} = -.08$, $p = .582$; LCP: $r_{52} = -.14$, $p = .340$). These results suggested that HCs might have increased prefrontal-motor coordination underlying reactive control for context processing in HCP trials, while PSZ failed to have such coordination.

predictive of WM deficits in PSZ (45), these results suggest that schizophrenia is associated with failures in context cue encoding owing to prefrontal proactive control deficits rather than simple visual deficits. Given the differential deficits of PSZ in cfN2, prfN, and LRP onset latency for HCP, proactive control for rule updating and reactive control of motor processes are the primary specific deficits underlying context processing deficits in schizophrenia.

Proactive and Reactive Control Deficits in PSZ

A previous ERP study using AX-CPT identified coactivations in the bilateral dorsolateral prefrontal and parietal cortices as the neural sources of frontal N2 component during a cue period (46). Therefore, the cfN2 during the SRRT might reflect dorsolateral prefrontal cortex–parietal activations for proactive control and the updating of WM for contextual rules. fMRI studies using the SRRT have shown activations in middle frontal, anterior cingulate, inferior frontal, and inferior parietal cortices for HCP trials (47,48). Given the bilateral-frontal and frontocentral topography of prfN, it might reflect coactivations in the dorsolateral prefrontal-inferior frontal-anterior cingulate

cortices network underlying reactive control for detection and resolution of response conflict by inhibiting a prepotent motor response (49,50) and initiating a correct response according to contextual rules stored in WM (51). The context-dependent deficits in the cfN2 and prfN components in PSZ suggest that they reflect prefrontal-specific deficits involving both proactive and reactive controls. These findings partly replicate the previous AX-CPT fMRI findings that PSZ have deficient activations for proactive control (cue period) but hyper-activations for reactive control (probe period) in the prefrontal-parietal network (15). The hyper- and hypofrontality findings for reactive control in the AX-CPT and SRRT might be due to differences in task demands (52); the SRRT requires more reactive controls than the AX-CPT, not only inhibition of a prepotent automatic response but also initiation of a goal-relevant response.

In contrast to consistent findings of proactive control deficits, findings of reactive controls in PSZ are somewhat inconsistent. Unlike the EEG findings of reactive control deficits in the current SRRT and recent AX-CPT (16) studies, recent behavioral findings of experimental studies using

computerized conflict resolution tasks (i.e., the Stroop and Flanker tasks) (53–55) suggested intact reactive control abilities in PSZ. However, such tasks have the same rules requiring selective attention to task-relevant features of stimuli, while context processing tasks require using current mental representation of contextual rules varying across trials. Therefore, reactive control deficits in PSZ might arise from their difficulty in using mental representations rather than selective attention.

Visual Processing Deficits and Top-Down Visual Modulation in PSZ

We only partly replicated previous findings of visual encoding deficits in PSZ (11,17): intact visual responses to contextual cues but deficient visual responses to probes. This might be due to differences in the visual stimuli used for the cue. We used a simple visual cue (a rectangle in red/green color), while the previous studies used a letter (11) or an abstract shape (17). It appears that PSZ had visual encoding deficits for complex shapes but not for simple and easily discernible objects. Furthermore, PSZ might have had more difficulty in visual encoding of a letter that was presented for only 100 ms in the previous study (11), which challenged subsequent attentional processes. This might explain previous findings of associations of performance deficits with visual encoding deficits rather than prefrontal deficits. Our results show that both groups had larger pvN1 amplitudes in HCP trials than in LCP trials, suggesting that the prefrontal proactive control enhanced visual neuronal activity for controlled responses to probes. As indicated by the results from the supplementary analyses of cfN2-pvN1 ERP pairs (see Supplemental Figure S3) that support the previous finding (11), PSZ might have relatively intact top-down modulation of visual processing.

Contribution of Deficits in Controlled Motor Processes to Context Processing Deficits in PSZ

Consistent with LRP findings during Flanker tasks (56), PSZ had diminished LRP amplitudes regardless of response conflicts but increased LRP onset latency only for HCP trials, suggesting generalized and specific deficits of motor processes in PSZ. The onset latency of response-locked LRP represents amount of time between the onset of preparatory motor activities and the execution of the response. Therefore, the increased LRP onset latency might reflect deficient prefrontal reactive control of motor cortical activity to initiate a goal-relevant motor response in PSZ.

In line with the LRP findings, we also found general and specific abnormalities in Ex-Gaussian RT indices in PSZ, replicating recent findings of increased σ and τ in first-episode PSZ (57) that indicated unstable and dysregulated motor processes in schizophrenia. PSZ had HCP-specific abnormality in the RT variability index σ , which was primarily predicted by the interaction between prfN and LRP (prfN \times LRP). Consistent with a recent fMRI finding that smaller σ was predicted by larger activities of dorsal anterior cingulate cortex (58) that might be the primary source of prfN, these findings suggest that deficient prefrontal-motor coordination for reactive control may cause unstable motor processes

characterized by large trial-by-trial fluctuations in the latency of controlled motor responses. PSZ also had large increases in slow response latency (τ) which was predicted by smaller prfN and prfN \times LRP. Large τ has often been associated with attentional lapse (57,59), and recent fMRI studies found that attentional lapse is associated with diminished suppression of default mode brain network activities (60,61). Therefore, our finding of larger τ for LCP trials than for HCP trials in PSZ might indicate their increased attentional lapse associated with not only deficient prefrontal reactive control but also inability to suppress spontaneous activities of the default mode brain network that tends to be larger in the easier trials, leading to more mind wandering (62). In addition, prfN \times LRP predicted severity of overall and disorganization symptoms of schizophrenia. Given the recent fMRI finding that abnormal prefrontal-motor functional connectivity predicted overall schizophrenia symptom severity (63), the results suggested that failures in prefrontal-motor coordination underlying context processing deficits in PSZ relates to various schizophrenic symptoms as well as their cognitive deficits.

In summary, using an optimal task, RT variability analyses, and high temporal resolution ERP indices, the current study identified specific and generalized deficits in neural processes underlying context processing deficits in PSZ. Inaccuracy and instability of motor control predicted by deficient proactive prefrontal and reactive prefrontal-motor control activities in the HCP condition indicated their specific context processing deficits associated with failures in using mental representations of varying contextual rules. In contrast, delayed response latencies and reduced visual and motor cortical activities observed regardless of task demands suggested generalized deficits in visuomotor processes in schizophrenia. The identification of the specific and generalized cognitive/neural deficits may provide guidelines for future mechanistic studies and alternative interventions (e.g., cognitive remediation) of schizophrenia. Future studies analyzing interregional neural coordination may further clarify cortical dysintegration mechanisms underlying context processing deficits in schizophrenia.

ACKNOWLEDGMENTS AND DISCLOSURES

This work was supported by Grant No. CSMRF I01CX000227 from the Veterans Health Administration (to SRS) and Grants Nos. R24MH069675 and R03MH106831 from the National Institute of Mental Health (to SRS) as well as by the Mental Health Patient Service Line at the Minneapolis VA Medical Center.

We are grateful to the participants and also for the efforts of Jessica Johns.

The authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the Department of Biomedical Sciences (SSK), University of Missouri–Kansas City, Kansas City, Missouri; Department of Psychiatry (AWM, SRS) and Department of Psychology (AWM, SRS), University of Minnesota, and Veterans Affairs Health Care System (SRS), Minneapolis, Minnesota.

Address correspondence to Seung Suk Kang, Ph.D., 2411 Holmes Street, Kansas City, MO 64108; E-mail: kangseung@umkc.edu.

Received Mar 1, 2019; accepted Mar 20, 2019.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.bpsc.2019.03.012>.

REFERENCES

- MacDonald A, Carter CS, Kerns JG, Ursu S, Barch D, Holmes A, *et al.* (2005): Specificity of prefrontal dysfunction and context processing deficits to schizophrenia in a never medicated first-episode psychotic sample. *Am J Psychiatry* 162:475–484.
- Cohen JD, Servanschauber D (1992): Context, cortex, and dopamine: A connectionist approach to behavior and biology in schizophrenia. *Psychol Rev* 99:45–77.
- Snitz BE, MacDonald AW, Carter CS (2005): Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: A meta-analytic review of putative endophenotypes. *Schizophr Bull* 32: 179–194.
- MacDonald AW, Pogue-Geile MF, Johnson MK, Carter CS (2003): A specific deficit in context processing in the unaffected siblings of patients with schizophrenia. *Arch Gen Psychiatry* 60:57–65.
- Barch DM, Carter CS, Braver TS, Sabb FW, MacDonald A, Noll DC, Cohen JD (2001): Selective deficits in prefrontal cortex function in medication-naïve patients with schizophrenia. *Arch Gen Psychiatry* 58:280–288.
- Green MF (2006): Cognitive impairment and functional outcome in schizophrenia and bipolar disorder. *J Clin Psychiatry* 67(Suppl 9):3–8; discussion 36–42.
- Barber AD, Carter CS (2005): Cognitive control involved in overcoming prepotent response tendencies and switching between tasks. *Cereb Cortex* 15:899–912.
- Butler PD, Silverstein SM, Dakin SC (2008): Visual perception and its impairment in schizophrenia. *Biol Psychiatry* 64:40–47.
- Luck SJ, Gold JM (2008): The construct of attention in schizophrenia. *Biol Psychiatry* 64:34–39.
- Walker EF, Savoie T, Davis D (1994): Neuromotor precursors of schizophrenia. *Schizophr Bull* 20:441–451.
- Dias EC, Butler PD, Hoptman MJ, Javitt DC (2011): Early sensory contributions to contextual encoding deficits in schizophrenia. *Arch Gen Psychiatry* 68:654–664.
- Braver TS (2012): The variable nature of cognitive control: A dual mechanisms framework. *Trends Cogn Sci* 16:106–113.
- Braver TS, Paxton JL, Locke HS, Barch DM (2009): Flexible neural mechanisms of cognitive control within human prefrontal cortex. *Proc Natl Acad Sci U S A* 106:7351–7356.
- Lesh TA, Westphal AJ, Niendam TA, Yoon JH, Minzenberg MJ, Ragland JD, *et al.* (2013): Proactive and reactive cognitive control and dorsolateral prefrontal cortex dysfunction in first episode schizophrenia. *NeuroImage Clin* 2:590–599.
- Edwards BG, Barch DM, Braver TS (2010): Improving prefrontal cortex function in schizophrenia through focused training of cognitive control. *Front Hum Neurosci* 4:32.
- Ryman SG, Cavanagh JF, Wertz CJ, Shaff NA, Dodd AB, Stevens B, *et al.* (2018): Impaired midline theta power and connectivity during proactive cognitive control in schizophrenia. *Biol Psychiatry* 84:675–683.
- Haenschel C, Bittner RA, Haertling F, Rotarska-Jagiela A, Maurer K, Singer W, Linden DEJ (2007): Contribution of impaired early-stage visual processing to working memory dysfunction in adolescents with schizophrenia. *Arch Gen Psychiatry* 64:1229–1240.
- Wolff AL, O'Driscoll GA (1999): Motor deficits and schizophrenia: The evidence from neuroleptic-naïve patients and populations at risk. *J Psychiatry Neurosci* 24:304–314.
- Pappa S, Dazzan P (2009): Spontaneous movement disorders in antipsychotic-naïve patients with first-episode psychoses: A systematic review. *Psychol Med* 39:1065–1076.
- Luck SJ, Kappenman ES, Fuller RL, Robinson B, Summerfelt A, Gold JM (2009): Impaired response selection in schizophrenia: Evidence from the P3 wave and the lateralized readiness potential. *Psychophysiology* 46:776–786.
- Kappenman ES, Luck SJ, Kring AM, Lesh TA, Mangun GR, Niendam T, *et al.* (2016): Electrophysiological evidence for impaired control of motor output in schizophrenia. *Cereb Cortex* 26:1891–1899.
- Mathalon DH, Fedor M, Faustman WO, Gray M, Askari N, Ford JM (2002): Response-monitoring dysfunction in schizophrenia: An event-related brain potential study. *J Abnorm Psychol* 111:22–41.
- MacDonald SWS, Nyberg L, Bäckman L (2006): Intra-individual variability in behavior: Links to brain structure, neurotransmission and neuronal activity. *Trends Neurosci* 29:474–480.
- Palmer EM, Horowitz TS, Torralba A, Wolfe JM (2011): What are the shapes of response time distributions in visual search? *J Exp Psychol Hum Percept Perform* 37:58–71.
- Van Zandt T (2000): How to fit a response time distribution. *Psychon Bull Rev* 7:424–465.
- Nobre AC, Allison T, McCarthy G (1998): Modulation of human extrastriate visual processing by selective attention to colours and words. *Brain* 121:1357–1368.
- Di Russo F, Martínez A, Sereno MI, Pitzalis S, Hillyard SA (2002): Cortical sources of the early components of the visual evoked potential. *Hum Brain Mapp* 15:95–111.
- Patel SH, Azzam PN (2005): Characterization of N200 and P300: Selected studies of the event-related potential. *Int J Med Sci* 2:147–154.
- Czernochowski D (2015): ERPs dissociate proactive and reactive control: Evidence from a task-switching paradigm with informative and uninformative cues. *Cogn Affect Behav Neurosci* 15:117–131.
- Czernochowski D, Nessler D, Friedman D (2010): On why not to rush older adults: Relying on reactive cognitive control can effectively reduce errors at the expense of slowed responses. *Psychophysiology* 47:637–646.
- Eimer M (1998): The lateralized readiness potential as an on-line measure of central response activation processes. *Behav Res Methods Instrum Comput* 30:146–156.
- First M, Spitzer R, Gibbon M, Williams J (2001): Structured Clinical Interview for DSM-IV-TR Axis I Disorders. New York: Biometrics Research, New York State Psychiatric Institute.
- Nurnberger JI, Blehar MC, Kaufmann CA, York-cooler C, Simpson SG, Harkavy-friedman J, *et al.* (1994): Diagnostic Interview for Genetic Studies. *Arch Gen Psychiatry* 51:849–859.
- Goghari VM, MacDonald AW, Sponheim SR (2011): Temporal lobe structures and facial emotion recognition in schizophrenia patients and nonpsychotic relatives. *Schizophr Bull* 37:1281–1294.
- Heathcote A (1996): RTSYS: A DOS application for the analysis of reaction time data. *Behav Res Methods Instrum Comput* 28:427–445.
- Perrin F, Pernier J, Bertrand O, Echallier JF (1989): Spherical splines for scalp potential and current density mapping. *Electroencephalogr Clin Neurophysiol* 72:184–187.
- Kang SS, Lano TJ, Sponheim SR (2015): Distortions in EEG interregional phase synchrony by spherical spline interpolation: Causes and remedies. *Neuropsychiatr Electrophysiol* 1:9.
- MacDonald AW, Carter CS (2003): Event-related fMRI study of context processing in dorsolateral prefrontal cortex of patients with schizophrenia. *J Abnorm Psychol* 112:689–697.
- Hoerl AE, Kennard RW (1970): Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics* 12:55–67.
- García CB, García J, López Martín MM, Salmerón R (2015): Collinearity: Revisiting the variance inflation factor in ridge regression. *J Appl Stat* 42:648–661.
- Cule E, Vineis P, De Iorio M (2011): Significance testing in ridge regression for genetic data. *BMC Bioinformatics* 12:372.
- Cule E, De Iorio M (2012): A semi-automatic method to guide the choice of ridge parameter in ridge regression. arXiv:1205.0686. Available at: <https://arxiv.org/pdf/1205.0686>.
- Johannes S, Münte TF, Heinze HJ, Mangun GR (1995): Luminance and spatial attention effects on early visual processing. *Brain Res Cogn Brain Res* 2:189–205.
- Javitt DC (2009): When doors of perception close: Bottom-up models of disrupted cognition in schizophrenia. *Annu Rev Clin Psychol* 5:249–275.
- Kang SS, MacDonald AW, Chafee MV, Im C-H, Bernat EM, Davenport ND, Sponheim SR (2018): Abnormal cortical neural synchrony during working memory in schizophrenia. *Clin Neurophysiol* 129:210–221.
- Dias EC, Foxe JJ, Javitt DC (2003): Changing plans: A high density electrical mapping study of cortical control. *Cereb Cortex* 13:701–715.

47. Goghari VM, MacDonald AW (2008): Effects of varying the experimental design of a cognitive control paradigm on behavioral and functional imaging outcome measures. *J Cogn Neurosci* 20:20–35.
48. Goghari VM, MacDonald AW (2009): The neural basis of cognitive control: Response selection and inhibition. *Brain Cogn* 71:72–83.
49. Botvinick MM, Braver TS, Barch DM, Carter CS, Cohen JD (2001): Conflict monitoring and cognitive control. *Psychol Rev* 108:624–652.
50. Levy BJ, Wagner AD (2011): Cognitive control and right ventrolateral prefrontal cortex: Reflexive reorienting, motor inhibition, and action updating. *Ann N Y Acad Sci* 1224:40–62.
51. Haddon JE, Killcross S (2006): Prefrontal cortex lesions disrupt the contextual control of response conflict. *J Neurosci* 26:2933–2940.
52. Callicott JH, Mattay VS, Verchinski BA, Marenco S, Egan MF, Weinberger DR (2003): Complexity of prefrontal cortical dysfunction in schizophrenia: More than up or down. *Am J Psychiatry* 160:2209–2215.
53. Smid H, Bruggeman R, Martens S (2016): Normal cognitive conflict resolution in psychosis patients with and without schizophrenia. *J Abnorm Psychol* 125:88–103.
54. Westerhausen R, Kompus K, Hugdahl K (2013): Unaffected control of distractor interference in schizophrenia: A meta-analysis of incompatibility slowing in flanker tasks. *J Psychiatr Res* 47:246–251.
55. Westerhausen R, Kompus K, Hugdahl K (2011): Impaired cognitive inhibition in schizophrenia: A meta-analysis of the Stroop interference effect. *Schizophr Res* 133:172–181.
56. Kappenman ES, Kaiser ST, Robinson BM, Morris SE, Hahn B, Beck VM, *et al.* (2012): Response activation impairments in schizophrenia: Evidence from the lateralized readiness potential. *Psychophysiology* 49:73–84.
57. Fassbender C, Scangos K, Lesh TA, Carter CS (2014): RT distributional analysis of cognitive-control-related brain activity in first-episode schizophrenia. *Cogn Affect Behav Neurosci* 14:175–188.
58. van Belle J, van Raalten T, Bos DJ, Zandbelt BB, Oranje B, Durston S (2015): Capturing the dynamics of response variability in the brain in ADHD. *NeuroImage Clin* 7:132–141.
59. Epstein JN, Conners CK, Hervey AS, Tonev ST, Arnold LE, Abikoff HB, *et al.* (2006): Assessing medication effects in the MTA study using neuropsychological outcomes. *J Child Psychol Psychiatry* 47:446–456.
60. Fassbender C, Zhang H, Buzy WM, Cortes CR, Mizuiri D, Beckett L, Schweitzer JB (2009): A lack of default network suppression is linked to increased distractibility in ADHD. *Brain Res* 1273:114–128.
61. Weissman DH, Roberts KC, Visscher KM, Woldorff MG (2006): The neural bases of momentary lapses in attention. *Nat Neurosci* 9:971–978.
62. Mason MF, Norton MI, Van Horn JD, Wegner DM, Grafton ST, Macrae CN (2007): Wandering minds: The default network and stimulus independent thought. *Science* 315:393–395.
63. Du X, Choa F-S, Chiappelli J, Wisner KM, Wittenberg G, Adhikari B, *et al.* (2019): Aberrant middle prefrontal-motor cortex connectivity mediates motor inhibitory biomarker in schizophrenia. *Biol Psychiatry* 85:49–59.