




Contents lists available at ScienceDirect

Sleep Health

Journal of the National Sleep Foundation

journal homepage: sleephealthjournal.org

Effect of positive airway pressure therapy in children with obstructive sleep apnea syndrome: does positive airway pressure use reduce pedestrian injury risk? ☆☆☆

Kristin T. Avis, PhD, CBSM^a, Karen L. Gamble, PhD^b, David C Schwebel, PhD^{c,*}

^a University of Alabama at Birmingham, Department of Pediatrics, Birmingham, AL

^b University of Alabama at Birmingham, Department of Psychiatry, Birmingham, AL

^c University of Alabama at Birmingham, Department of Psychology, Birmingham, AL

ARTICLE INFO

Article history:

Received 31 July 2018

Received in revised form 16 November 2018

Accepted 13 December 2018

Keywords:

Pedestrian safety

Children

Injury risk

Obstructive sleep apnea syndrome

PAP

ABSTRACT

Introduction: Treatment with positive airway pressure (PAP) therapy reduces injury risk among adults with obstructive sleep apnea syndrome (OSAS), but the effect of PAP therapy on children's injury risk is unknown. This study investigated whether treatment of OSAS with PAP reduces children's pedestrian injury risk in a virtual reality pedestrian environment.

Methods: Forty-two children ages 8–16 years with OSAS were enrolled upon diagnosis by polysomnography. Children crossed a simulated street several times upon enrollment, prior to PAP treatment, and again after 3 months of PAP therapy. Children underwent sleep studies at all time points.

Results: Children adherent with PAP had a significant reduction in hits by a virtual vehicle ($P < .01$) and less time to contact with oncoming vehicles ($P < .01$) following treatment. Those who were nonadherent did not show improved safety. There was no change in attention to oncoming traffic.

Conclusions: OSAS may have significant consequences on children's daytime functioning in a critical domain of personal safety: pedestrian skills. In pedestrian simulation, children with OSAS adherent to PAP therapy showed improvement in pedestrian safety and had fewer collisions with a virtual vehicle following treatment. Results highlight need for heightened awareness of the real-world benefits of treatment for pediatric sleep disorders.

Level of evidence: Level II Therapeutic Study.

© 2019 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.

Annually, 6300 American pedestrians are killed and 190,000 others are injured, and a disproportionate number of injured and killed pedestrians are children.¹ Not surprisingly, prevention of pediatric pedestrian injury is targeted as a national public health priority.² Many factors contribute to unintentional pedestrian injury. Among them are cognitive skills of the pedestrian, including reaction time,

impulsivity, risk taking, attention, and decision making.^{3,5,6} These same characteristics that influence pedestrian safety are negatively influenced by sleepiness, both from sleep deprivation/insufficient sleep and from sleep disorders such as obstructive sleep apnea syndrome (OSAS).^{7–13}

OSAS is a common pediatric sleep-related breathing disorder, with estimated prevalence of 1%–5% among nonobese children and 25%–40% among obese children.^{14,15} In adults, OSAS has negative consequences on mood and behavioral regulation and also on neurocognitive functioning, including vigilance, attention, problem solving, and visual and motor functioning.^{16,17} Adults with OSAS are at high risk for human error and mental inefficiency, and are more likely to have inattention, risk taking, and daytime sleepiness resulting in motor vehicle crashes and occupational injury.^{6–8,16–18}

The standard treatment for adult OSAS is positive airway pressure therapy (PAP).^{19–21} PAP consists of patients wearing a mask interface covering the nose or nose and mouth connected to a machine by a

☆ Acknowledgement of funding source: The project was supported primarily by the RESMED Foundation and also by Award Number R01HD058573 from the Eunice Kennedy Shriver National Institute of Child Health & Human Development. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Eunice Kennedy Shriver National Institute of Child Health & Human Development, the National Institutes of Health, or RESMED foundation.

☆☆ Conflict of interest: The authors have no conflicts of interest to disclose.

* Corresponding author at: Department of Psychology, University of Alabama at Birmingham, 1300 University Blvd, CH 415, Birmingham, AL 35294. Tel.: +1 205 934 8745.

E-mail address: schwebel@uab.edu (D.C. Schwebel).

tube. The machine blows a determined amount of pressure into the airway functioning as a “stent” to keep the airway patent. There is evidence that treatment of adult OSAS with PAP improves attention, cognitive functioning, and sleepiness and reduces the risk of motor vehicle crashes.^{22–25} However, the effect of PAP in children with OSAS who are not yet old enough to drive, and thus engage in traffic primarily as pedestrians, is unknown.

Pedestrian behavior, although enacted subconsciously by most adults, is a highly complex cognitive and perceptual task. A safe pedestrian must simultaneously process several pieces of information, interpret their meaning, and make a decision to cross the street when a safe opportunity arises. These tasks must occur very quickly. Impulse control, attention, and reaction time are cognitive skills still developing among children that may impact pedestrian safety,^{3–5,26–27} and these same cognitive impairments are reported among children with untreated OSAS who engage in laboratory tasks and questionnaires.^{9–12,28} Thus, children with OSAS may have deficits in pedestrian safety. In fact, previous research suggests that children with OSAS were at higher risk for pedestrian injury than healthy peers of the same age, race, sex, and estimated household income.²⁹ The present study was designed to extend that result by testing whether treating OSAS may improve pedestrian safety among children with OSAS.

Specifically, we recruited children with OSAS to engage in a virtual reality pedestrian environment in 2 conditions: when OSAS was initially diagnosed but untreated and after OSAS was controlled for 3 months using treatment with PAP. We hypothesized that children with OSAS would be safer pedestrians after adhering to treatment with PAP vs when untreated or after nonadherent treatment.

Methods

Participants and recruitment

Using a within-subjects design, forty-two 8- to 16-year-olds with OSAS were enrolled. Diagnosis and recruitment occurred in the Pediatric Sleep Disorders Center at Children's of Alabama the morning following a diagnostic sleep study. All participating children met International Classification of Sleep Disorders-2 diagnostic criteria for OSAS based on diagnostic assessments that included nocturnal polysomnography using Sandman 9.2 technology software (Embla, Broomfield, CO) and thorough clinical evaluation from 1 of 2 attending board-certified sleep specialists early in the morning after the overnight nocturnal polysomnography. Standard polysomnography consisted of electroencephalogram, electromyogram, electrooculogram (right, left), arterial oxygen saturation (SaO₂) oximeter pulse wave form, end-tidal carbon dioxide tension, nasal pressure monitoring, oronasal flow using thermistor, and thoracic and abdominal wall motion. Standard pediatric scoring was used for respiratory events.³⁰ Diagnosis of OSAS was defined as having an apnea/hypopnea index (AHI) ≥ 1.5 per hour.^{31–34} All children were treated with continuous PAP rather than bilevel PAP. Exclusion criteria for the clinical sample included cognitive or physical disabilities that prevented participation in the experimental protocol (eg, mental retardation, blindness, use of a wheelchair), comorbid medical or neurologic conditions, or antipsychotic medication use. No children were excluded.

General procedure

We recruited to the study children diagnosed with OSAS requiring and naive to PAP therapy. As part of routine care, PAP was prescribed as follows: (1) after surgical intervention (tonsillectomy/adenoidectomy), follow-up sleep study determined OSAS was still present, or (2) at the diagnostic study when surgical intervention was not an option (ie, surgery was done at any point prior to initial sleep study and no regrowth of tonsillar/adenoidal tissue has

occurred, persistent OSAS likely due to obesity). Following PAP prescription, an initial research visit was scheduled that included informed consent procedures and evaluation in the pedestrian virtual reality environment.

Following the initial research appointment, the family returned to the Pediatric Sleep Disorders Centers for routine care, which consisted of (1) PAP education, (2) behavioral techniques to aid in adjusting to PAP regimen, and (3) meeting with a home durable equipment representative who fit the child for an appropriate face mask and provided the PAP machine to the family. The child practiced for 1 week to adjust to the new regimen. One week after the baseline visit, children returned to the sleep laboratory for an overnight sleep study to titrate their PAP to the optimal pressure according to titration manual guidelines. After 3 months of use of PAP at home, the family returned for a posttreatment overnight sleep study (conducted while the child was wearing the PAP) to document use of and adequate control of OSAS with PAP and to obtain an objective check of adherence from a downloadable card housed in the machine. The morning following the final overnight visit, the family completed a second research appointment evaluating pedestrian safety in the virtual reality environment; this occurred after 3 months treatment of PAP.

Measures

Adherence

Our objective was to evaluate injury outcomes in participants who were adherent. Adherence was monitored with downloadable computerized use meters in the PAP machine that document hours of use each night. Following standard practice in the literature, the minimal criterion for category of “adherent” was greater than or equal to average of 4 hours use per night.^{35–36}

Sleepiness

To verify sleepiness, we sampled level of sleepiness using the Epworth Sleepiness Scale modified for children.³⁷ The Epworth Sleepiness Scale assesses daytime sleepiness and is modified for children to rate the propensity to fall asleep in 8 situations in which a child may be likely to fall asleep.⁹ Response options are “would never doze,” slight chance of dozing,” “moderate chance of dozing,” and “high chance of dozing.” The questionnaire, originally designed for adults, is used in studies with children with OSAS with slight variations in terminology on 2 questions: (a) the mention of alcohol is removed in question 7, and (b) question 8 was modified to indicate the child is a passenger in a car.⁹

Virtual reality pedestrian environment

Details of the virtual reality pedestrian environment (VRPE), including validation data demonstrating behavior in the virtual world corresponds with behavior in real pedestrian environments among both children and adults, are available elsewhere.²⁶ Briefly, the environment has demonstrated convergent, internal, and face validity and has been used in several studies showing impaired pedestrian behavior under a variety of conditions and situations among both children and adults.^{3,26,38}

While participating in the virtual environment task, children stood on a wooden simulated curb and viewed the virtual pedestrian environment on 3 consecutive monitors arranged in a semicircle in front of them. Children were immersed in the virtual environment as they watched vehicles pass bidirectionally on the screens and heard environmental and traffic noise through speakers in the room. After deciding it was safe to cross, they stepped off the curb onto a pressure plate connected to the computer, and a sex-matched avatar was then activated to cross the street. The avatar's walking speed in the VRPE matched children's walking speed, which was evaluated prior to the VRPE task in a separate location. If the avatar safely reached the other

side, children heard 1 of 2 positive messages such as “Yes! Great job!” If the child made it across safely but was close to being hit by a car, the child heard, “Whoa! That was close!” If the child was struck by one of the cars, they heard, “Uh oh, you should try that again.” Thus, the child was immersed into a virtual world while deciding when it was safe to cross. After choosing to cross, the world became third person, and the child witnessed the safety (or danger) of the crossing. During the experimental visits, children performed 10 practice trials to reduce learning effects and then engaged in 12 virtual street crossings. Behavior in the 12 crossings was used for analysis.

Measures of crossing behavior

We considered 3 pedestrian outcome measures. First, we looked at overall risk of simulated injury. We also considered 2 aspects of crossing behavior: looking at traffic and time to contact by an oncoming vehicle after entering a traffic gap.

- 1 Simulated injury (collisions with virtual vehicles): “Hits” were any direct collisions between the virtual pedestrian and a vehicle.
- 2 Looking at traffic: To evaluate attention to traffic, looks toward traffic were tallied by head-tracking equipment (Trackir4:Pro, NaturalPoint Inc, Corvallis, OR) that monitored participants’ visual attention to traffic from the left and right. We summed the number of times participants looked left plus the number of times they looked right while waiting to cross, divided by the average wait time in seconds.
- 3 Time to contact by an oncoming vehicle: The time to contact by an oncoming vehicle was the shortest latency (in seconds) between the child and an oncoming vehicle during the time the child was in the crosswalk. Shorter time to contact by an oncoming vehicle indicates selection of a riskier traffic gap to cross within.

Data analysis plan

Descriptive statistics were considered first to examine the distribution of scores and ensure that assumptions of inferential analyses were met. We also examined descriptively the number of children who were adherent and nonadherent to the PAP treatment. To address the primary hypothesis that children with OSAS would engage more safely as pedestrians after adhering to treatment with PAP vs when untreated or after nonadherent treatment, we constructed mixed-model, ordinal logistic regressions run using the generalized estimating equations procedure to compare pre- and posttreatment hits by simulated traffic in a stratified manner, with one model constructed among children who were adherent to PAP and a second among children who were nonadherent. Given our results that the regression was statistically significant among adherent children but not among nonadherent children, we proceeded to evaluate secondary pedestrian behavior outcomes, time to contact, and attention to traffic among only the adherent group. Paired-sample *t* tests compared behavior pre- and posttreatment among those children. All analyses were conducted using SPSS, version 22 (IBM Corp, Armonk, NY). Significance was ascribed at $P < .05$.

Results

Forty-two children with OSAS were enrolled. One child was excluded who could not be provided with a machine due to inadequate insurance coverage, and 12 children were lost to follow-up over the 3-month time period. Analyses comparing those lost to follow-up to those who completed the study found that children lost to follow-up were more likely to be male (92% vs 59%) but otherwise were similar on demographic, body mass index (BMI), and baseline sleep measures.

Table 1 presents data for the sample of 29 children who completed the study, both overall and divided into adherent and

nonadherent subgroups. Of the 29 participants, 20 children were evaluated at both visits, had usable data, and did not have persisting OSAS. One child was excluded because the downloadable card malfunctioned, and 8 children had persisting OSAS greater than 1.5 AHI and so were categorized as inadequately treated due to poor titration. Based on the criteria of PAP use for an average of greater than 4 hours a night, 11 of the 20 included children were adherent to the PAP therapy and 9 were nonadherent. The nonadherent group was somewhat older than the adherent group, but otherwise, the 2 groups were similar on all variables tested.

To test the primary hypothesis, stratified mixed-model, ordinal logistic regression analyses indicated that, following treatment, children adherent and adequately treated with PAP experienced a significant reduction in hits by simulated vehicles in the VRPE (Wald $\chi^2[1] = 6.86, P = .009$, Fig. 1) but children not adherent to treatment with PAP did not experience a significant reduction in hits by simulated vehicles (Wald $\chi^2[1] = 1.47, P = .23$). A follow-up mixed binomial regression indicated that, when untreated, children were 12 times more likely to have a simulated hit by a virtual vehicle than when adherent to PAP (Wald $\chi^2[1] = 5.61$, odds ratio 12.00, 95% confidence interval 1.12–128.84, $P = .018$).

Given these results, we next examined secondary pedestrian safety outcomes at baseline and posttreatment among children adherent to treatment. As shown in Table 2, children adherent to treatment showed a significant improvement in time to contact with oncoming vehicles posttreatment ($t[10] = 4.05, P < .01$), indicating they had left more time between themselves and oncoming traffic following treatment. There was not a significant change posttreatment in looking at oncoming traffic ($t[10] = -.63, P = .55$).

Discussion

Our results suggest that children with OSAS who were treated with PAP therapy experienced significant improvement in their pedestrian safety, as measured in a virtual pedestrian environment, following adherence to PAP therapy. In contrast to children who were not adherent, we found that adherent children experienced a significant reduction in collisions with virtual vehicles. We also saw an improvement in leaving more time before contact with oncoming traffic among adherent children.

Results reinforce the fact that pediatric OSAS causes significant cognitive impairment that manifests itself in multiple applied domains, including pedestrian safety, and that adherent treatment with PAP therapy can produce positive change in functioning. Crossing streets safely requires simultaneous processing of several pieces of information and therefore is a highly complex cognitive and perceptual task. OSAS may impair relevant cognitive components.²⁸ When children with OSAS were adherent to PAP, they seemed to regain cognitive capacity and engaged more safely in the virtual street environment, functioning closer to what their age and development status would predict. When they were untreated at baseline, their pedestrian safety was compromised.

Consistent with other research, we did not find that looking at traffic was impaired or improved with PAP therapy.^{29,39} A possible explanation for this finding is that when they are sleepy, children are still able to follow simple and rote rules learned when they were young, such as looking both ways before crossing, but they may not fully process cognitively the complex environment they perceive or select safe traffic gaps to cross within. This hypothesis corresponds with results from research investigating pedestrian safety in untreated pediatric sleep disorders of central hypersomnia and untreated OSAS, in which children looked both ways at traffic but did not process the crossing properly and were struck by virtual vehicles.^{29,39}

Table 1
Demographic, BMI, and sleep characteristics of sample at baseline

Characteristic	Completers (n = 29)	Adherent (n = 11)	Nonadherent (n = 9)	Test statistic ^a
Sex, % male	58.6	54.5	55.6	<i>P</i> = .658
Age (y), mean (SD)	11.9 (2.7)	10.7 (2.2)	14.7 (1.1)	<i>t</i> = 5.1**
Ethnicity, % white	42.9	54.5	50.0	<i>P</i> = .605
Household income, % below \$20,000	30.8	45.5	42.9	<i>P</i> = .648
ESS total, mean (SD) ^b	12.7 (5.4)	13.2 (5.1)	16.4 (4.6)	<i>t</i> = 1.4
BMI, mean (SD) (age and sex adjusted)	36.1 (11.9)	37.7 (11.1)	38.4 (14.3)	<i>F</i> _{1,16} = 0.01
BMI z score, mean (SD)	2.29 (0.16)	MED = 2.5	MED = 2.9	<i>U</i> = 26.0
Polysomnography data				
Total sleep time (min), mean (SD)	445.3 (54.2)	455.0 (51.1)	445.2 (27.9)	<i>t</i> = 0.544
Sleep efficiency %, mean (SD)	85.4 (10.6)	MED = 93.6	MED = 89.9	<i>U</i> = 53.5
Sleep latency (min), mean (SD)	16.4 (15.2)	MED = 12.5	MED = 7.0	<i>U</i> = 67.0
Awakenings (count), mean (SD)	24.0 (11.4)	21.0 (11.5)	28.2 (11.7)	<i>t</i> = 1.4
Stage 1 sleep (%), mean (SD)	8.0 (5.0)	6.5 (3.9)	10.9 (5.5)	<i>t</i> = 2.1
Stage 2 sleep (%), mean (SD)	49.4 (10.0)	48.0 (11.1)	50.4 (10.1)	<i>t</i> = 0.5
Stage 3 sleep (%), mean (SD)	23.6 (9.3)	25.2 (10.3)	19.7 (8.2)	<i>t</i> = 1.3
REM sleep (%), mean (SD)	18.9 (6.4)	20.3 (6.2)	19.0 (6.8)	<i>t</i> = 0.4
AHI, mean (SD)	10.7 (15.0)	MED = 3.0	MED = 5.7	<i>U</i> = 35.5

ESS, Epworth Sleepiness Scale; MED, median.

^a Test statistic compares adherent to nonadherent groups. Fisher exact test *P* values presented for categorical variables; Independent samples *t* or Mann-Whitney *U* statistics presented for continuous variables, depending on distribution of outcome variable; and analysis of covariance used for BMI to covary age and sex.

^b n = 10 for adherent group and n = 8 for nonadherent group on this variable due to missing data.

** *P* < .01.

Our findings have implications for clinical practice. Although a few pediatric studies have documented cognitive improvement on laboratory tasks following PAP therapy,^{40–42} this study extends findings to the applied outcome of unintentional pedestrian injury risk. In adults with OSAS, PAP therapy improves occupational safety, driving performance, and injury risk following PAP therapy.^{22–24} This study offers initial evidence that pediatric functioning concords with that of adults and PAP therapy has the potential to reduce injury risk among children engaging in what is perhaps the most common mode of transportation where children influence their own safety: pedestrian injury.

We note several study limitations. First, we recruited children with OSAS from a clinically referred group of children that may represent a group with more severe symptoms that prompt a referral than the larger community sample of children with OSAS. Second, although we chose 3 months as an outcome point based on prior research,^{41–42} the minimal amount of daily PAP use required for improvement in safety and the length of time needed to wear PAP before safety improvement is observed remain unknown. Third, our study suffered from attrition over the 3-month time period. Fourth, we used a noncounterbalanced within-subjects research design. Ethical considerations prohibited counterbalancing or inclusion of a

clinical control group, but future research might consider alternative research designs to control for the possibility of practice effects and the simple cognitive-perceptual development that occurs in children over a 3-month time period and might influence pedestrian behavior and skill. Finally, the findings are limited to pedestrian injury. Future research should consider other aspects of health and safety among children with OSAS, other treatments that might reduce injury risk in children with OSAS, and possible countermeasures that could be implemented to promote increased safety.

Disclosure

KTA received support from RESMED foundation. All data collection, statistical analysis, and manuscript writing were performed by the investigators independent of RESMED foundation. The authors have no financial relationships relevant to this article to disclose.

Acknowledgments

We thank Carole L Marcus for her guidance as a consultant on the initial study design and Anna Johnston, research coordinator for the




Fig. 1. Reduction in hits by a virtual vehicle in children with OSAS adherent to PAP.

Table 2
Pedestrian outcome measures in children with OSAS adherent to PAP

Pedestrian risk outcome measure	Pre M (SD)	Post M (SD)	t	df	P
Time to contact	3.45 (0.61)	4.43 (0.68)	4.05	10	.002
Looking at traffic	27.60 (9.97)	30.27 (8.54)	0.63	10	.546

UAB Youth Safety Lab. We thank Digital Artefacts for their development and support of the virtual reality system.

References

- Centers for Disease Control and Prevention. WISQARS™ (Web-based Injury Statistics Query and Reporting System). Retrieved 7/18/18 from <https://www.cdc.gov/injury/wisqars/index.html>; 2018.
- National Highway Traffic Safety Administration. Pedestrian safety. Retrieved 7/18/18 from <https://www.nhtsa.gov/road-safety/pedestrian-safety>; 2018.
- Schwebel DC, Gaines J, Severson J. Validation of virtual reality as a tool to understand and prevent child pedestrian injury. *Accid Anal Prev*. 2008;40:1394–1400.
- Schwebel DC, Barton BK. Contributions of multiple risk factors to child injury. *J Pediatr Psychol*. 2005;30(7):553–561.
- Thompson JA, Tolmie AK, Foot HC, Sarvary PH, Whelan KM. Influence of virtual reality on the roadside crossing judgments of child pedestrians. *J Exp Psychol Appl*. 2005;11(3):175–186.
- Dinges DF. An overview of sleepiness and accidents. *J Sleep Res*. 1995;4:4–14.
- Findley L. Vigilance and automobile accidents in patients with sleep apnea or narcolepsy. *Chest*. 1995;108(3):619–624.
- American Thoracic Society. sleep apnea, sleepiness, and driving risk. *Am J Respir Crit Care Med*. 1994;50:1463–1473.
- Melendres MC, Lutz JM, Rubin ED, Marcus CL. Daytime sleepiness and hyperactivity in children suspected with sleep-disordered breathing. *Pediatrics*. 2004;114:768–775.
- Beebe DW, Groesz L, Wells C, et al. The neuropsychological effects of obstructive sleep apnea: a meta-analysis of norm referenced and case controlled data. *Sleep*. 2003;26(3):298–307.
- Beebe DW, Wells CT, Jeffries J, Chini B, Kalra M, Amin R. Neuropsychological effects of pediatric obstructive sleep apnea. *J Int Neuropsychol Soc*. 2004;10(7):962–975.
- Beebe DW. Neurobehavioral morbidity associated with disordered breathing during sleep in children: a comprehensive review. *Sleep*. 2006;29(9):1115–1134.
- Redline S, Tishler PV, Schluchter M, Aylor J, Clark K, Graham G. Risk factors for sleep-disordered breathing in children. Associations with obesity, race, and respiratory problems. *Am J Respir Crit Care Med*. 1999;159:1527–1532.
- Marcus CL, Brooks LJ, Ward SD, et al. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*. 2012;130:1–42.
- Alonso-Alvarez ML, Cordero-Guevara JA, Teran-Santos J, et al. Obstructive sleep apnea in obese community-dwelling children: the NANOS study. *Sleep*. 2014;37:943–949.
- Lal C, Strange C, Bachman D. Neurocognitive impairment in obstructive sleep apnea. *Chest*. 2012;14:1601–1610.
- Treager S, Reston J, Schoelles K, Phillips B. Obstructive sleep apnea and risk of motor vehicle crash. *J Clin Sleep Med*. 2009;5:573–581.
- Aksan N, Marini R, Tippin J, et al. Driving performance and driver state in obstructive sleep apnea: what changes with positive airway pressure? *Proc Int Driv Symp Hum Factors Driv Assess Train Veh Des*. 2017:9–15.
- Gay P, Weaver T, Loube D, Iber C. Evaluation of positive airway pressure treatment for sleep related breathing disorders in adults. *Sleep*. 2006;29:381–401.
- Morgenthaler TI, Kapen S, Lee-Chiong T, et al. Practice parameters for the medical therapy of obstructive sleep apnea. *Sleep*. 2006;29(8):1031–1035.
- Kushida CA, Littner MR, Hirshkowitz M, et al. Practice parameters for the use of continuous and bilevel positive airway pressure devices to treat adult patients with sleep-related breathing disorders. *Sleep*. 2006;29:375–380.
- Tregear TS, Reston J, Schoelles K, Phillips B. Continuous positive airway pressure reduces risk of motor vehicle crash among drivers with obstructive sleep apnea. *Sleep*. 2010;33(10):1373–1380.
- George CF. Reduction in motor vehicle collisions following treatment of sleep apnea with nasal CPAP. *Thorax*. 2001;56:508–512.
- Findley L, Smith C, Hooper J, Dineen M, Surratt PM. Treatment with nasal CPAP decreases automobile accidents in patients with sleep apnea. *Am J Respir Crit Care Med*. 2000;161:857–859.
- Sanchez AI, Martinez P, Miro E, Bardwell WA, Buela-Casal GCPAP. Behavioral therapies in patients with obstructive sleep apnea: effects on daytime sleepiness, mood, and cognitive function. *Sleep Med Rev*. 2009;13:223–233.
- Schwebel DC, Davis AL, O'Neal EE. Child pedestrian injury: a review of behavioral risks and preventive strategies. *Am J Lifestyle Med*. 2012;6(4):292–302.
- Barton BK. Integrating selective attention into developmental pedestrian safety research. *Can Psychol*. 2006;47:203–210.
- Cardoso TSG, Pompeia S, Miranda MC. Cognitive and behavioral effects of obstructive sleep apnea syndrome in children: a systematic literature review. *Sleep Med*. 2018;46:46–55.
- Avis K, Gamble KL, Schwebel DC. Obstructive sleep apnea syndrome increases pedestrian injury risk in children. *J Pediatr*. 2015;16:109–114.
- American Academy of Sleep Medicine. International Classification of Sleep Disorders. Diagnostic and Coding Manual. 2nd ed. Westchester, IL: American Academy of Sleep Medicine; 2005.
- Berry RB, Budhiraja R, Gottlieb DJ, et al. Rules for scoring respiratory events in sleep: update of the 2007 AASM Manual for the Scoring of Sleep and Associated Events. *J Clin Sleep Med*. 2012;8(5):597–619.
- Witmans MB, Keens TG, Davidson Ward SL, Marcus CL. Obstructive hypopneas in children and adolescents: normal values. *Am J Respir Crit Care Med*. 2003;168:1540.
- Treager N, Schultz B, Pollock AN, Mason T, Marcus CL, Arens R. Polysomnographic values in children 2–9 years old: additional data and review of the literature. *Pediatr Pulmonol*. 2005;40:22–30.
- Uliel S, Tauman R, Greenfeld M, Sivan Y. Normal polysomnographic respiratory values in children and adolescents. *Chest*. 2004;125:872–878.
- King MS, Xanthopoulos MS, Marcus CL. Improving positive airway pressure in children. *Sleep Med Clin*. 2014;9:219–234.
- Pepin JL, Krieger J, Rodenstein D, et al. Effective compliance during the first 3 months of continuous positive airway pressure. A European prospective study of 121 patients. *Am J Respir Crit Care Med*. 1999;160:1124–1129.
- Johns MW. A New method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep*. 1991;14:540–545.
- Stavrinou D, Biasini FJ, Fine PR, Hodgins JB, Khatri S, Mrug S, Schwebel DC. Mediating factors associated with pedestrian injury in children with attention-deficit disorder. *Pediatrics*. 2011;128(2):296–302.
- Avis KT, Gamble KL, Schwebel DC. Does excessive daytime sleepiness affect children's pedestrian safety? *Sleep*. 2014;37(2):283–287.
- Marcus CL, Rosen G, Ward SL, Halbower AV, Sterni L, et al. Adherence to and effectiveness of positive airway pressure therapy in children with obstructive sleep apnea. *Pediatrics*. 2006;117:e442–e451.
- Marcus CL, Radcliffe J, Konstantinopoulou S, Beck S, Cornaglia M, et al. Effects of positive airway pressure therapy on neurobehavioral outcomes in children with obstructive sleep apnea. *Am J Respir Crit Care Med*. 2012;185:998–1003.
- Beebe DW, Byars KC. Adolescents with obstructive sleep apnea adhere poorly to positive airway pressure (PAP) but users show improved attention and school performance. *PLoS One*. 2011;6(3):e16924. <https://doi.org/10.1371/journal.pone.0016924>.