



## Original article

## Effects of long-term consumption of broccoli sprouts on inflammatory markers in overweight subjects



María Teresa López-Chillón<sup>a</sup>, Carmen Carazo-Díaz<sup>b</sup>, David Prieto-Merino<sup>b, c</sup>,  
Pilar Zafrilla<sup>a</sup>, Diego A. Moreno<sup>d, \*</sup>, Débora Villaño<sup>a</sup>

<sup>a</sup> Universidad Católica San Antonio de Murcia (UCAM), Department of Pharmacy, Faculty of Health Sciences, Campus de los Jerónimos 30107 Guadalupe, Murcia, Spain

<sup>b</sup> Applied Statistical Methods in Medical Research Group, Catholic University of Murcia (UCAM), Murcia, Spain

<sup>c</sup> Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London, United Kingdom

<sup>d</sup> CEBAS-CSIC, Department of Food Science and Technology, Phytochemistry and Healthy Foods Lab, Research Group on Quality, Safety and Bioactivity of Plant Foods. Campus de Espinardo - 25, E-30100 Espinardo, Murcia, Spain

## ARTICLE INFO

## Article history:

Received 22 February 2018

Accepted 6 March 2018

## Keywords:

Glucosinolates  
Brassica oleracea  
Sulforaphane  
IL-6  
Bioavailability  
Inflammation

## SUMMARY

**Background & aims:** Broccoli sprouts represent an interesting choice of healthy food product as they are rich in glucosinolates and their cognate bioactive metabolites, isothiocyanates able to counteract the negative effects of diverse pathologies. As obesity is linked to an inflammatory component, the aim of the study was to evaluate the anti-inflammatory action of broccoli sprouts in overweight adult subjects.

**Methods:** An *in vivo* controlled study was performed in 40 healthy overweight subjects (ClinicalTrials.gov ID NCT 03390855). Treatment phase consisted on the consumption of broccoli sprouts (30 g/day) during 10 weeks and the follow-up phase of 10 weeks of normal diet without consumption of these broccoli sprouts. Anthropometric parameters as body fat mass, body weight, and BMI were determined. Inflammation status was assessed by measuring levels of TNF- $\alpha$ , IL-6, IL-1 $\beta$  and C-reactive protein.

**Results:** IL-6 levels significantly decreased (mean values from 4.76 pg/mL to 2.11 pg/mL with 70 days of broccoli consumption,  $p < 0.001$ ) and during control phase the inflammatory levels were maintained at low grade (mean values from 1.20 pg/mL to 2.66 pg/mL,  $p < 0.001$ ). C-reactive protein significantly decreased as well.

**Conclusions:** This study represents an advance in intervention studies as the broccoli sprouts were included in a daily dietary pattern in quantities that reflect a real consumption. Further studies are necessary to elucidate the role of this healthy rich and nutritious food product, but these promising results support the current evidence on the healthy properties of *Brassica* varieties.

© 2018 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

## 1. Introduction

Nowadays there is an increasing demand by consumers on healthy food products prepared in convenient forms, simple to use and not containing additives. In this sense, broccoli sprouts (*Brassica oleracea* var. *italica*) are a good option, as they are the main dietary source of glucosinolates, nitrogen-sulphur compounds, and phenolic derivatives (flavonoid glycosides, and hydroxycinnamic acids), vitamins A, C, E, K and minerals [1]. Glucosinolates are derivatives from amino acids and particularly abundant in *Brassica*

species and are believed to be responsible of the biological effects attributed to these vegetables. Cultivar conditions can affect and improve the biosynthesis of these secondary metabolites in the plant, favouring their storage in the sprouts. Broccoli sprouts contain 10- to 50-fold higher concentrations of glucosinolates than the mature broccoli [2].

Isothiocyanates are degradation products from glucosinolates formed by the hydrolytic action of myrosinase vegetable enzyme when the vegetable structure is crushed, e.g. during mastication process. Hydrolysis can also be performed by gastrointestinal enzymes during digestion; the metabolic conversion of glucosinolates to isothiocyanates is a crucial step for the benefits observed with the consumption of *Brassica* vegetables [3]. Most of these reported effects are linked to anti-cancer properties and epidemiological

\* Corresponding author.

E-mail address: [dmoreno@cebas.csic.es](mailto:dmoreno@cebas.csic.es) (D.A. Moreno).

studies have evidenced a decrease in the risk of cancer with the intake of cruciferous foods [4,5]. Human clinical studies have mainly focused on these antitumoral activities, with mechanisms including the upregulation of phase II detoxification enzymes, as well as the direct action on cell cycle, causing apoptosis of cancer cells [6] [7]. There is less evidence on the anti-inflammatory properties of cruciferous vegetables in humans.

A non-communicable disease of high prevalence in Western societies is obesity. Nowadays it is assumed that obesity is a condition characterized by a chronic low-grade inflammation status. Levels of pro-inflammatory mediators as IL-6, TNF- $\alpha$  and C-reactive protein are increased with the higher visceral adiposity and all of them are risk factors for cardiovascular diseases, metabolic syndrome and diabetes [8]. The permanent low-grade of inflammation is a response of the organism to the imbalance in the nutrient due to energy excess and it is manifested in the development of insulin resistance and type 2 diabetes [9]. Dietary interventions able to reduce the inflammatory conditions linked to obesity may improve health conditions of overweight population.

The hypothesis of our research is that broccoli sprouts are able to reduce the inflammatory status in overweight subjects due to their content in phytochemicals, mainly glucosinolates. Hence, the aim of the study is to evaluate the changes in markers of inflammation in overweight subjects after 10-week ingestion of broccoli sprouts as well as to assess the bioavailability of glucosinolates to correlate the changes observed.

## 2. Material and methods

### 2.1. Study design

We performed an interventional follow-up study to evaluate the effect of the daily consumption of broccoli sprouts during 10 weeks (70 days).

The study was performed according with the Helsinki Declaration of Human Studies and approved by the Ethical Committee of the Catholic University of Murcia as well as the Bioethics Sub-Committee of the CSIC' Department of Ethics for the AGL-2013-46247-P project registered also at ClinicalTrials.gov ID NCT 03390855. Volunteers (n = 40; 21 men, 19 women) were recruited in the Catholic University of Murcia (UCAM) and all of them were informed on the characteristics of the study and they signed the written-informed consent. The inclusion criteria were: BMI within the overweight range according to the World Health Organization criteria (24.9–29.9 kg/m<sup>2</sup>), aged 35–55 years, taking no vitamins, supplements or medication during the previous two months; no-smoking. The exclusion criteria were: diagnosed diseases as hypertension and cardiovascular pathologies, diabetes, hepatic, gastrointestinal and renal diseases, as well as the intake of drugs related to these pathologies, vegetarian diet, pregnancy or breast-feeding. Dietetic and life style habits were recorded from all participants.

There were no drop-outs during the whole period of study and no adverse effects were reported due to the broccoli sprout ingestion. Physical parameters of the volunteer at the beginning of the study are listed in Table 1.

One week before the beginning of the intervention period, subjects were asked to avoid the consumption of *Brassica* vegetables (broccoli, radish, cauliflower, Brussel sprouts, mustards, among others) and their derived products, and to follow a well-balanced diet (based on Mediterranean diet), with no other food restriction criteria. These dietary instructions were maintained during the entire period of study. Besides, they were requested to record any sign of adverse effect, illness or deviation of the experimental diet. The subjects maintained their usual lifestyles during the study.

On the first day, participants were given the portions of fresh broccoli sprouts to be taken for the whole week (7 trays of broccoli sprouts of 30 g each) and each week they had an appointment to provide them the fresh products. The intervention consisted on a 10-week period which included daily consumption of a portion (30 g) of raw, fresh broccoli sprouts. This amount is consistent with a half-serving [11]. Subjects were instructed to ingest 1 tray per day and to keep the trays refrigerated (4 °C) at home. The intake of the broccoli sprouts was included in their normal daily diet and no specific time of consumption was established, with the only limitation of avoid cooking of the sprouts and to consume them fresh. Cooking procedures can affect the content of glucosinolates as well as their bioavailability [12,13] and therefore some recipes were provided to the participants to facilitate the intake of the sprouts without affecting the phytochemical composition and absorption. We gave instructions to the volunteer of not cooking broccoli sprouts but of consuming them in raw manner. They included the sprouts in vegetable salads, cold pasta salads or in cold sandwiches with different combinations of the following ingredients: cheese, ham, tomato, lettuce, grilled pork or grilled vegetables (in both cases the sprouts should be added after the grill and once the ingredient was lukewarm). It could also be included in burgers, as the “Californian style burger”, which ingredients are broccoli sprouts, bacon, avocado, tomato, burger and bread. Other recipes include elaboration of “Gazpacho” (Spanish cold soup made with vegetables as tomato, cucumber, green pepper, onion, oil, vinegar and salt), or mixed with smashed potatoes with melted cheese, or mixed with bread and spreadable cheese. In all cases it was important to use in cold or warm temperatures so that the glucosinolates were preserved.

After the intervention period, a follow-up recovery period for all subjects continued for other 90 days with no ingestion of broccoli sprouts.

Fasting blood samples and 24-h urine samples were taken on day 0 (D0: just before starting the intervention), day 70 (D70: end of intervention period), day 90 (D90: 20 days after end of intervention) and 160 (D160: 90 days after end of intervention). Blood samples were collected from each subject by venipuncture from the antecubital vein; 3 mL were placed in heparin tubes and centrifuged at 10,000 rpm for 10 min at 4 °C. Plasma was aliquoted and stored at –80 °C until analysis. Analysis were performed once each period was finished and in the same batch to minimize analytical variations. The total volume of the 24 h-urine was recorded to calculate the absolute amounts of the compounds and metabolites excreted in the study period and aliquots were frozen at –80 °C for further analysis. Body weight and percentage of fat mass were measured as well and BMI calculated in each sampling time point.

### 2.2. Broccoli sprouts

Raw, fresh broccoli sprouts (*B. oleracea* var. *italica*) were supplied by Aquaporins&Ingredients, S.L (Alcantarilla, Murcia, Spain). The sprouts were biostimulated with methyl jasmonate 250  $\mu$ M, for 4 days previous to delivery, in order to increase up to 2-fold levels the production of bioactive compounds, according to a protocol

**Table 1**  
Baseline characteristics of volunteer (n = 40; 21 men, 19 women).

Variable	Mean $\pm$ standard deviation
Age (years)	46 $\pm$ 6
Height (m)	1.72 $\pm$ 0.08
Weight (kg)	85.8 $\pm$ 16.7
BMI (kg/m <sup>2</sup> )	28.9 $\pm$ 4.0
Body fat mass (%)	30.34 $\pm$ 7.54

previously validated [10]. In that study we performed some tests on elicitation and seed priming to enrich the broccoli sprouts in glucosinolates. We used the elicitor methyl jasmonate (MeJA) by priming the seeds as well as by spraying daily over the cotyledons from day 4–7 of germination. We observed that MeJA at concentrations of 250  $\mu\text{M}$  act as stressor in the plant and enhances the biosynthesis of the phytochemicals glucosinolates. Compared to control plants without MeJA treatment, the content of compounds as the aliphatic glucosinolate glucoraphanin was enhanced up to a 70% and similar increases were observed with glucoiberin or glucobrassicin. In this way, we improved the content of these health-promoting compounds. Other nutritional facts did not change with these treatments.

Three trays of sprouts were collected once a week during the study, frozen and lyophilized prior to analysis on glucosinolates and isothiocyanates, as previously described [2]. The phytochemicals (glucosinolates) provided by the broccoli sprouts are summarized in Table 2.

### 2.3. Biochemical analysis

Markers of inflammation as IL-6, C-reactive protein, IL-1 $\beta$  and TNF- $\alpha$  in plasma were determined in our laboratory using high-sensitivity ELISA kits. ichroma™ hsCRP kits were purchased from Bioditech Meed Inc.'s. Human IL-6 ELISA high sensitivity kits were from BioVendor. Human IL-1 $\beta$  high sensitivity ELISA kits and human TNF- $\alpha$  high sensitivity ELISA kits were acquired from IBL International GmbH Instrumental.

### 2.4. Analysis of glucosinolates and isothiocyanates in urine and plasma

Levels of glucosinolates, isothiocyanates and their metabolites (GRA, IB, SFN, SFN-GSH, SFN-NAC, SFN-CYS, I3C, 3,3-DIM) were measured in urine by a rapid, sensitive and high throughput UHPLC-QqQ-MS/MS [14]. All LC-MS grade solvents were obtained from J.T. Baker (Phillipsburg, New Jersey, USA). The standards of Sulforaphane, SFN-glutathione, SFN-cysteine and SFN-N-acetylcysteine (SFN, SFN-GSH, SFN-CYS, SFN-NAC, respectively) and Iberin (IB), Indole-3-carbinol (I3C) and 3,3-Diindolyl-methane (DIM\_3\_3) were from SantaCruz Biotech (CA, USA). Glucoraphanin (GRA) was obtained from Phytoplan (Diehm & Neuberger GmbH, Heidelberg, Germany).

Urine and plasma samples were extracted using SPE Strata-X cartridges (33  $\mu\text{m}$  Polymeric Strong Cation) following manufacturer's instructions (Phenomenex, Torrance, CA, USA). The cartridges were preconditioned with 2 mL of methanol and equilibrated with 2 mL of water:formic acid (98:2, v/v). Samples (400  $\mu\text{L}$ ) were diluted with 2 mL water:formic acid (98:2, v/v) and

loaded into the column. SPE cartridges were washed with 2 mL water:formic acid (98:2, v/v). Elution of target metabolites was performed with 1 mL of methanol/formic acid (98:2, v/v). Samples eluted were dried using a SpeedVac concentrator (Savant SPD121P, Thermo Scientific, Massachusetts, USA). The extracts were reconstituted in 200  $\mu\text{L}$  of mobile phases A/B (90:10, v/v) and filtered with PTFE 0.22  $\mu\text{m}$  filters. Chromatographic separation was carried out using a ZORBAX Eclipse Plus C-18 (2.1  $\times$  50 mm, 1.8  $\mu\text{m}$ ) (Agilent Technologies) and the mobile phases employed were: solvent A ammonium acetate, 13 mM (pH 4 with acetic acid) and solvent B acetonitrile/acetic acid (99.9:0.1 v/v) as previously described [15]. Twenty microlitres of each sample were acquired in a Agilent Technologies UHPLC-1290 Series coupled to a 6460 QqQ-MS/MS (Agilent Technologies, Waldbronn, Germany). Compounds were identified and quantified using MRM transitions and positive or negative ESI mode for confirmation of the target analytes, compared to available external standards [14]. Standard curves were prepared freshly every day of analysis.

### 2.5. Statistical analysis

Continuous variables were summarised with means and standard deviations while qualitative variables were summarised with proportions. We estimated the relative change on the continuous biomarkers (weight, BMI, body fat, IL\_6, C-reactive protein and 3,3-DIM), between the different periods (before and after Broccoli ingestion) using linear regression models. Additional models were run controlling for age and sex in case these variables were confounding the effect of the broccoli. We then compared the proportion of samples with detectable metabolites in sulforaphane pathway (SFN, SFN-CYS and SFN-NAC) between visits. We studied the association between changes observed in body fat mass, IL6 and C-reactive protein (outcome variables) and changes in metabolites 3,3-DIM, SFN-NAC, SFN-CYS and SFN (explanatory variables) using linear regression models. Data were analysed using R (3.4.1 version) software package.

## 3. Results

### 3.1. Bioactive compounds in broccoli sprouts

Broccoli sprouts from each of the 10 weeks of the study were characterized for their glucosinolate (GLS) contents (Table 2). Results are presented as the serving portion (30 g) consumed daily by the volunteers. The major aliphatic glucosinolate in broccoli sprout detected was glucoraphanin (GRA, 4-methyl-sulphonylbutyl glucosinolate) and the major indolic glucosinolate detected was neoglucobrassicin (NGB, 1-methoxy-3-indolylmethyl glucosinolate). Total concentration of aliphatic glucosinolates was 80.50 mg/30 g f.w., equivalent to 6.22  $\mu\text{mol/g}$  fresh weight or 65.47  $\mu\text{mol/g}$  dry weight. This concentration was two-fold higher than indolic glucosinolates (40.62 mg/30 g f.w., equivalent to 2.88  $\mu\text{mol/g}$  fresh weight or 30.32  $\mu\text{mol/g}$  dry weight). Volunteers consumed an average of 51 mg (117  $\mu\text{mol}$ ) and 20 mg (42  $\mu\text{mol}$ ) of glucoraphanin and neoglucobrassicin, respectively, on a daily basis, during the 70 days of the dietary intervention.

### 3.2. Biological effects of broccoli sprout on markers of inflammation and body composition

Baseline characteristics of volunteer are described in Table 1. Changes on plasma concentration of biomarkers at different time points of the intervention are shown in Table 3. Day 0 and day 70 refers to the first and last days of the broccoli ingestion,

**Table 2**  
Glucosinolate contents in broccoli sprouts daily portions (mg/30 g F.W.).

	Mean $\pm$ Standard deviation (n = 3)
Glucoiberin (GIB)	19.28 $\pm$ 0.98
Glucoraphanin (GRA)	51.08 $\pm$ 1.06
4-Hydroxyglucobrassicin (HGB)	3.67 $\pm$ 0.41
Glucoerucin (GER)	10.14 $\pm$ 1.20
Glucobrassicin (GBS)	9.69 $\pm$ 0.95
4-Methoxyglucobrassicin (MBG)	7.14 $\pm$ 0.61
Neoglucobrassicin (NGB)	20.11 $\pm$ 1.66
Aliphatic Glucosinolates ( $\Sigma$ )	80.50 $\pm$ 2.18
Indolic Glucosinolates ( $\Sigma$ )	40.62 $\pm$ 2.07
Total ( $\Sigma$ )	121.11 $\pm$ 4.00

respectively. The days 90 and 160 refers to 20 days and 90 days of follow-up upon the broccoli dosage period, respectively.

The evolution of mean values of continuous variables are shown in Table 4. The metabolite 3,3'-diindolylmethane (3,3-DIM) was included in the statistical analysis as it was detected in all volunteers, at concentrations higher than limit of quantification, hence, for statistical purposes, it was treated as a continuous variable. Evolution of ratios of continuous variables in each visit are shown in Fig. 1.

No significant changes were observed in weight and BMI. By contrast, body fat mass slightly decreased significantly after 70 days of broccoli consumption (ratio = 0.947, P-value = 0.02586) and returned to basal levels at day 90, a state that was maintained until day 160 (P-value = 0.94899 y P-value = 0.07644).

Plasma interleukine-6 (IL-6) concentrations decreased significantly (by 38%) after 70 days of broccoli ingestion as well, respect to basal value (ratio = 0.381, P-value < 0.00001). Moreover, these lower levels continue to significantly decrease after 20 days of ceasing broccoli ingestion (ratio = 0.195, P-value < 0.00001). At 90 days of the follow up period (day 160), levels returned somewhat but without returning to baseline values (ratio = 0.472, P-value = 0.00000). Figure 2 illustrates how the changes in IL-6, during broccoli intake, depends on baseline values. The negative slope of the regression line indicates that volunteer with higher concentrations at baseline tend to lose more concentration of this biomarker. Decreases in C-reactive protein were also observed, during broccoli ingestion period (ratio = 0.592, P-value = 0.00915). Shortly in the follow-up period, levels returned to baseline conditions (P-value = 0.92162 and P-value = 0.72756, at 90 and 160 days, respectively).

These results did not substantially change when we repeated the regression models adjusting for age and sex. TNF $\alpha$  and IL-1 $\beta$  were detected in a small number of samples and most of them below the limit of quantification, hence no valid conclusions can be inferred and data have not been considered for statistical purposes.

### 3.3. Bioavailability and metabolism of glucosinolates and isothiocyanates

Some glucosinolates and isothiocyanates, as glucoraphanin, glucoiberin, iberin, glucoerucin, erucin and glucobrassicin were absent in the urine samples. Indole-3-carbinol was detected only after broccoli ingestion in low quantities and in 50% of samples. In contrast, the metabolite 3,3'-diindolylmethane (3,3-DIM) was detected and quantified in all volunteers and for statistical purposes it was treated as a continuous variable. It increased significantly during broccoli ingestion (ratio = 1.947, P-value < 0.00001). Shortly in the follow-up period, levels returned to baseline conditions (P-value = 0.10484 and P-value = 0.12312, at 90 and 160 days, respectively).

Metabolites from sulfuraphane pathway are present in 24 h-urine samples (Table 3); the metabolite at higher amount was SFN-

**Table 4**  
Evolution of mean values (ratios) on the time points.<sup>a</sup>

Variable	Mean	Ratio	95% CI	P-value
Weight (kg)	86.141			
From D0 to D70	86.346	1.002	(0.997–1.008)	0.38388
From D0 to D90	85.983	0.998	(0.987–1.010)	0.74963
From D0 to D160	86.424	1.003	(0.994–1.012)	0.45589
Body mass index (kg/m <sup>2</sup> )	28.877			
From D0 to D70	28.945	1.002	(0.997–1.008)	0.39007
From D0 to D90	28.797	0.997	(0.986–1.009)	0.62441
From D0 to D160	28.971	1.003	(0.994–1.012)	0.46125
Body fat mass (%)	28.834			
From D0 to D70	27.298	0.947	(0.903–0.993)	<b>0.02586</b>
From D0 to D90	28.795	0.999	(0.956–1.043)	0.94899
From D0 to D160	29.955	1.039	(0.996–1.084)	0.07644
IL-6 (pg/mL)	4.594			
From D0 to D70	1.748	0.381	(0.298–0.486)	<b>&lt;0.00001</b>
From D0 to D90	0.896	0.195	(0.149–0.255)	<b>&lt;0.00001</b>
From D0 to D160	2.170	0.472	(0.366–0.609)	<b>&lt;0.00001</b>
C-reactive protein ( $\mu$ g/mL)	1.431			
From D0 to D70	0.847	0.592	(0.405–0.865)	<b>0.00915</b>
From D0 to D90	1.459	1.020	(0.677–1.536)	0.92162
From D0 to D160	1.553	1.085	(0.665–1.771)	0.72756
DIM_3_3 ( $\mu$ M)	0.334			
From D0 to D70	0.650	1.947	(1.705–2.223)	<b>&lt;0.00001</b>
From D0 to D90	0.335	0.757	(0.539–1.063)	0.10484
From D0 to D160	0.376	0.850	(0.689–1.048)	0.12312

Bold are significant at P < 0.05 and highly significant at P < 0.01.

<sup>a</sup> Data are adjusted by baseline levels.



NAC (mean concentration 2.0301  $\mu$ M, corresponding to 3.21  $\mu$ mol/24 h), whereas SFN was the compound with the lowest excretion (0.543  $\mu$ M, corresponding to 0.77  $\mu$ mol/24 h). The sum of SFN, SFN-NAC and SFN-CYS was ~5.11  $\mu$ mol/24 h. Considering an amount of GRA of 117  $\mu$ mol by serving, a 4% on average was metabolized through mercapturic acid pathway.

Figure 3 shows the proportion of individuals in which the metabolites have been detected and quantified at each visit and Fig. 4 shows the differences of these proportions from baseline with their confidence intervals. The percentage of individuals where SFN-NAC is detected increases significantly during broccoli intervention (45% increase; P-value = 0.00001). Afterwards, the percentage diminishes although it is statistically different from baseline (32.5% difference; P-value = 0.00303). At 160 days, no significant differences from baseline are observed (P-value = 0.07139). Similar behaviour is detected with SFN-CYS and SFN. Percentages increased during broccoli ingestion (67.5% increases in SFN-CYS P-value = 0.0000; 82.5% increases in SFN; P-value < 0.0001). Afterwards, the percentages of individuals detected decreased to baseline conditions (P-value = 0.43858 and P = value 0.26355 for SFN-CYS and SFN, respectively). This behaviour is maintained for the longer period at 160 days (P-value = 0.29330 and P = value 0.73532, respectively). SFN-GSH was detected in very few samples of volunteer during broccoli ingestion (data not included), hence, it has not been considered for statistical purposes.

**Table 3**

Changes observed with broccoli treatment as well as during follow-up period. Values are expressed as mean (confidence interval 95%).

Variable	Day 0	Day 70	Day 90	Day 160
Weight (kg)	85.79 (80.38–91.20)	85.69 (80.42–90.95)	83.83 (79.47–88.18)	84.04 (79.66–88.41)
BMI (kg/m <sup>2</sup> )	28.88 (27.56–30.20)	28.93 (27.63–30.23)	28.49 (27.40–29.57)	28.60 (27.52–29.68)
Body fat mass (%)	30.34 (27.29–33.39)	29.32 (26.93–31.71)	30.29 (27.87–32.72)	32.09 (29.69–34.49)
IL-6 (pg/mL)	4.76 (4.21–5.31)	2.11 (1.61–2.61)	1.20 (0.88–1.52)	2.66 (1.89–3.44)
C-reactive protein ( $\mu$ g/mL)	2.42 (1.45–3.40)	1.52 (0.70–2.34)	1.92 (1.02–2.82)	2.32 (1.07–3.56)
SFN-NAC ( $\mu$ M)	0.193 (0.00–0.41)	2.301 (1.85–2.75)	0.023 (0.01–0.04)	0.094 (0.00–0.19)
SFN-CYS ( $\mu$ M)	0.116 (0.00–0.26)	0.800 (0.57–1.03)	0.078 (0.00–0.22)	0.081 (0.00–0.19)
3,3-DIM ( $\mu$ M)	0.484 (0.38–0.59)	0.707 (0.61–0.80)	0.449 (0.33–0.57)	0.461 (0.36–0.56)
SFN ( $\mu$ M)	0.098 (0.00–0.23)	0.543 (0.40–0.69)	0.038 (0.00–0.13)	0.022 (0.01–0.03)




**Fig. 1.** Evolution of ratios of continuous variables in each visit according to model (1): a) changes in weight, BMI and body fat mass; b) changes in IL-6 and C-reactive protein levels.


The decrease in IL-6 levels was significantly related to the increase in 24 h-urine SFN levels ( $p = 0.03319$ ). In case of C-reactive protein, the decrease was significantly related to the increases in 24 h-urine SFN-NAC ( $p = 0.04783$ ) and SFN-CYS ( $p = 0.04116$ ). (Supp. Table-6).

**4. Discussion**

We conducted a human intervention study to test whether regular consumption of broccoli sprouts improves inflammatory biomarkers in overweight subjects. Adipose tissue is related to higher secretion of pro-inflammatory cytokines as  $TNF-\alpha$  and IL-6 and elevated levels of these proteins have been described in overweight individuals [16,17]. These proteins are linked to several disease states [18] and C-reactive is an important predictive marker



**Fig. 2.** Example of how changes in IL-6 during broccoli intake depends on baseline values of IL-6 (variables log-transformed).



**Fig. 3.** Proportion of individuals and 95% IC of metabolites detected at each visit.

of cardiovascular events [19]; hence the reduction of their levels with dietary intervention could contribute to a better prognosis on obesity-associated disorders. In our study we observed a noticeable anti-inflammatory effect with the ingestion of broccoli sprouts, with a significant reduction by 38% and 59% in IL-6 and C-reactive protein concentrations, respectively.

Clinical studies on human participants on the anti-inflammatory properties of *Brassica* products are scarce. Our research group has previously described a significant decrease on markers of inflammatory processes, as the metabolites tetranor-PGEM (from prostaglandins  $E_1$  and  $E_2$ ) and  $11\beta$ -PGF $2\alpha$  (from prostaglandin D2) after consumption of a single portion of broccoli sprouts [20]. Other authors have reported decreases on C-reactive protein levels by 48% after 10-day broccoli intake (250 g/day) in smokers, confirming our results; however, no changes on IL-6 levels were detected [21]. Decreases in IL-6 and C-reactive protein were also observed after 14




Fig. 4. Changes in binary variables over periods.

days of cruciferous consumption [22], but the amounts used (7 g/kg body weight, 14 g/kg body weight) far exceeded those of our experiment.

Differences in population studied, study design, type of *Brassica* or amount of product consumed, could explain the different results observed. Our broccoli sprouts contained significant quantities of aliphatic glucosinolates as glucoraphanin, glucoiberin and glucoerucin, which derive from the aminoacid methionine, as well as indolic glucosinolates as methoxy and hydroxy derivatives of glucobrassicin, that derive from the aminoacid tryptophan. Broccoli sprouts are especially rich in glucoraphanin (up to 10-fold above adult organ (inflorescence) levels) that drop with the plant growing, as the plant material increases without concomitant synthesis of glucoraphanin [23]. Hence, potential beneficial concentrations are easier to achieve with dietary quantities of sprouts vs broccoli heads (inflorescences).

Levels of glucosinolates and their metabolites isothiocyanates were measured in 24 h-urine by UHPLC-MS/MS in order to ascertain the consumption of broccoli and with the aim to find out if any metabolite is related to the changes in the biochemical parameters observed. We did not observe significant levels of intact glucosinolates in 24 h-urine samples, being explainable as these compounds suffer extensive modifications prior to absorption in the gut. They are present in the intact plant as glucosides and, upon tissue damage, the enzyme myrosinase catalyses their rapid hydrolysis of the glucose moiety [24]. The aglycone of each glucosinolate suffers further hydrolytic metabolism to isothiocyanate in the gastrointestinal tract by gut microbiota; these compounds are then absorbed by enterocytes and distributed systemically [25]. In particular, the isothiocyanate sulforaphane (1-isothiocyanate-4-methyl-sulfinylbutane) is formed from the glucosinolate glucoraphanin (4-methyl-sulphanylbutyl glucosinolate). Therefore, isothiocyanates are the compounds mainly present in human tissues to which can be attributed the biological activities.

In humans, the isothiocyanates are metabolized via the mercapturic acid pathway. Conjugation with glutathione is catalysed by glutathione transferase and GSH-conjugates are metabolized rendering SFN-CYS and SFN-NAC. It has been proven that polymorphisms of these enzymes have a significant impact on ITC

metabolism [26]. From the results described, the metabolites SFN-NAC, SFN-CYS and SFN can be considered as good markers of ingestion, as their presence is related only with the broccoli period, indicating the compliance of the experimental diet.

Concerning indole glucosinolates, indole-3-carbinol (I3C) is released by hydrolysis of glucobrassicin (3-indolyl-methylglucosinolate) by myrosinase action. This type of indole glucosinolates are present in seeds, mature plant and some sprouts cultivars, but are not commonly present in all *Brassica* varieties [27,28]. After ingestion, I3C is modified by the acidic pH in the stomach and dimerizes to 3,3'-diindolylmethane (3,3-DIM) [29]. Hence, the presence of 3,3-DIM in the 24 h-urine samples is related to the metabolism of glucobrassicin derivatives present in our broccoli sprouts.

The increases on broccoli metabolites were significantly related to the decreases in IL-6 and C-reactive protein levels, what suggests their implication in the modulation of these pro-inflammatory proteins. Studies on cellular models have shown that the mechanism of interaction is common in isothiocyanates and similar to that of endogenous hormones as steroids or vitamins A and D. This type of compounds possesses electrophile groups that interact with nucleophilic moieties of transcription factors, down- or up-regulating their activity [30]; it has been shown that sulforaphane interacts with the redox-sensitive transcription factor Nrf2, to permit its translocation into the nucleus, where it binds to the antioxidant response element (ARE) and activates the synthesis of proteins related to the response to stress, as phase II detoxification enzymes and quinone reductases [31–34]. Besides, SFN inhibits activation of NF- $\kappa$ B, a central transcription factor in inflammation process and the gene expression of proinflammatory mediators [34,35]. This signalling pathway is redox sensitive as depends on the balance between ROS intracellular concentration and GSH levels. Changes on GSH levels by SFN may influence in this anti-inflammatory action. Other authors observed an anti-inflammatory effect by induction of Nrf2-pathway of broccoli sprout extract in human skin [36] and nasal lavage cells [32] in healthy subjects as well as in patients with chronic obstructive pulmonary disease (COPD) [37].

3,3'-DIM has shown to reduce transcriptional activity of NF- $\kappa$ B, what results in lower levels of inflammatory mediators as IL-6, in activated macrophages [38] as well in different models of inflammation in mice [39,40]. It has been pointed out the possible synergistic interaction of both SFN and 3,3'-DIM [7] and the isothiocyanates erucin and sulforaphane are interconvertible [41], so that the anti-inflammatory effects observed with broccoli sprouts intake are likely due to the combined effects of all the hydrolysis products of glucosinolates.

Concerning anthropometric parameters, after 10 week of the daily consumption of broccoli sprouts, weight and body mass index were not altered; however, body fat mass significantly decreased with broccoli intervention. It has been described that the metabolite I3C decreases adipogenesis by suppressing pathways of lipid accumulation mediated by PPAR $\gamma$  [42]; however, we did not detect I3C in 24 h-urine as it is mainly excreted in its metabolite 3,3'-DIM. We did not observe a significant correlation between the increase in 3,3'-DIM and the decrease in fat mass; no further experiments were performed to corroborate an additional hypothesis about the effects on adipogenesis.

A limitation of this study is the lack of a parallel randomised control group which would be ideal to establish causality links between broccoli intakes and the change in biomarker levels. The post-intervention follow up is not an ideal control period as several other factors might have changes in the individuals or the environment. However, the strong changes observed in the

inflammatory markers at the end of the intervention and their recuperation afterwards could be an indication of the beneficial effect of the broccoli that will have to be tested in a proper trial. On the other hand, overweight is frequently associated with other pathologies as hypertension, cardiovascular events, insulin resistance or type 2 diabetes and, due to the complex interactions among them, we limited the study to people with overweight status according to WHO criteria, but without any pathology or clinical disorder. Hence, our result can only be extrapolated to these type of population and not to the overall that could include people with some concomitant pathologies.

## 5. Conclusions

The consumption of broccoli sprouts in a real dietary serving is able to affect IL-6 and C-reactive protein levels in overweight subjects, hence attenuating chronic inflammation. Further research with broccoli sprouts including other biomarkers and mechanistic studies are necessary to elucidate the role of this healthy rich and nutritious food product, but these promising results support the current evidence on the properties of this *Brassica* specie for disease prevention.

## Statement of authorship

López-Chillón MT carried out data analyses and contributed to the interpretation of the findings; Carazo-Díaz MC and Prieto-Merino D performed the statistical analysis; Zafrilla P contributed to data analysis and discussion of the manuscript; Moreno D.A. Principal Investigator and general management of the AGL-2013-46247-P project, contributed with the funding the study, design of experiment, discussion and writing of manuscript. Villaño D contributed to the interpretation of analyses and discussion of the manuscript, statistical management of data and writing of manuscript.

## Conflict of interest

All Co-authors declare no conflicts of interest.

This work was supported by the Spanish Ministry of Economy, Industry and Competitiveness (MINECO) and European Regional Development Fund (ERDF) through Research Project AGL2013-46247-P, and the Grant for Research Groups of Excellence from the Murcia Regional Agency for Science and Technology (Fundación Séneca), Project 19900/GERM/15.

## Acknowledgments

Authors would like to thank the support and collaboration of Aquaporins & Ingredients S.L. specially from their technical department assistance and their facilities for the production of the broccoli sprouts needed in the clinical study according to the controlled certified conditions of the Ecological Agriculture Council of Murcia Region (CAERM) reference ES-ECO-024-MU, to guarantee appropriate standards of quality and safety for consumption.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.clnu.2018.03.006>.

## References

- [1] West LG, Meyer KA, Balch BA, Rossi FJ, Schultz MR, Haas GW. Glucoraphanin and 4-hydroxyglucobrassicin contents in seeds of 59 cultivars of broccoli, raab, kohlrabi, radish, cauliflower, brussels sprouts, kale, and cabbage. *J Agric Food Chem* 2004;52:916–26.
- [2] Baenas N, García-Viguera C, Moreno DA. Elicitation: a tool for enriching the bioactive composition of foods. *Molecules (Basel, Switzerland)* 2014;19:13541–63.
- [3] Dinkova-Kostova AT, Kostov RV. Glucosinolates and isothiocyanates in health and disease. *Trends Mol Med* 2012;18:337–47.
- [4] Wu Q-J, Yang Y, Wang J, Han L-H, Xiang Y-B. Cruciferous vegetable consumption and gastric cancer risk: a meta-analysis of epidemiological studies. *Cancer Sci* 2013;104:1067–73.
- [5] Hu J, Hu Y, Hu Y, Zheng S. Intake of cruciferous vegetables is associated with reduced risk of ovarian cancer: a meta-analysis. *Asia Pac J Clin Nutr* 2015;24:101–9.
- [6] Abdull Razis A, Nicola G, Pagnotta E, Iori R, Ioannides C. 4-Methylsulfonyl-3-butenyl isothiocyanate derived from glucoraphastin is a potent inducer of rat hepatic phase II enzymes and a potential chemopreventive agent. *Arch Toxicol* 2012;86:183–94.
- [7] Jeffery EH, Araya M. Physiological effects of broccoli consumption. *Phytochemistry Rev* 2009;8:283–98.
- [8] Moschen AR, Molnar C, Enrich B, Geiger S, Ebenbichler CF, Tilg H. Adipose and liver expression of interleukin (IL)-1 family members in morbid obesity and effects of weight loss. *Mol Med (Cambridge, Mass.)* 2011;17:840–5.
- [9] Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *J Clin Invest* 2011;121:2111–7.
- [10] Baenas N, Villaño D, García-Viguera C, Moreno DA. Optimizing elicitation and seed priming to enrich broccoli and radish sprouts in glucosinolates. *Food Chem* 2016;204:314–9.
- [11] Houghton CA, Fassett RG, Coombes JS. Sulforaphane: translational research from laboratory bench to clinic. *Nutr Rev* 2013;71:708–26.
- [12] Palermo M, Pellegrini N, Fogliano V. The effect of cooking on the phytochemical content of vegetables. *J Sci Food Agric* 2014;94:1057–70.
- [13] Vermeulen M, Klöpping-Ketelaars IW, van den Berg R, Vaes WH. Bioavailability and kinetics of sulforaphane in humans after consumption of cooked versus raw broccoli. *J Agric Food Chem* 2008;56:10505–9.
- [14] Baenas N, Suárez-Martínez C, García-Viguera C, Moreno DA. Bioavailability and new biomarkers of cruciferous sprouts consumption. *Food Res Int (Ottawa, Ont.)* 2017;100:497–503.
- [15] Domínguez-Perles R, Medina S, Moreno DA, García-Viguera C, Ferreres F, Gil-Izquierdo A. A new ultra-rapid UHPLC/MS/MS method for assessing glucoraphanin and sulforaphane bioavailability in human urine. *Food Chem* 2014;143:132–8.
- [16] Kang YE, Kim JM, Joung KH, Lee JH, You BR, Choi MJ, et al. The roles of adipokines, proinflammatory cytokines, and adipose tissue macrophages in obesity-associated insulin resistance in modest obesity and early metabolic dysfunction. *PLoS One* 2016;11. e0154003–e0154003.
- [17] Jorge ASB, Jorge GC, Paraíso AF, Franco RM, Vieira LJ, Hilzenderger AM, et al. Brown and white adipose tissue expression of IL6, UCP1 and SIRT1 are associated with alterations in clinical, metabolic and anthropometric parameters in obese humans. *Exp Clin Endocrinol Diabetes* 2017;125:163–70.
- [18] Biennu J, Monneret G, Fabien N, Revillard JP. The clinical usefulness of the measurement of cytokines. *Clin Chem Lab Med* 2000;38:267–85.
- [19] Whelton SP, Roy P, Astor BC, Zhang L, Hoogeveen RC, Ballantyne CM, et al. Elevated high-sensitivity C-reactive protein as a risk marker of the attenuated relationship between serum cholesterol and cardiovascular events at older age. *Am J Epidemiol* 2013;178:1076–84.
- [20] Medina S, Domínguez-Perles R, Moreno DA, García-Viguera C, Ferreres F, Gil JJ, Gil-Izquierdo A. The intake of broccoli sprouts modulates the inflammatory and vascular prostanoids but not the oxidative stress-related isoprostanes in healthy humans. *Food Chem* 2015;173:1187–94.
- [21] Riso P, Vendrame S, Del Bo C, Martini D, Martinetti A, Seregini E, et al. Effect of 10-day broccoli consumption on inflammatory status of young healthy smokers. *Int J Food Sci Nutr* 2014;65:106–11.
- [22] Navarro SL, Schwarz Y, Song X, Wang C-Y, Cheng C, Trudo SP, et al. Cruciferous vegetables have variable effects on biomarkers of systemic inflammation in a randomized controlled trial in healthy young adults. *J Nutr* 2014;144:1850–7.
- [23] Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against. *Proc Natl Acad Sci U S A* 1997;94:10367.
- [24] Bones AM, Rossiter JT. The enzymic and chemically induced decomposition of glucosinolates. *Phytochemistry* 2006;67:1053–67.
- [25] Angelino D, Jeffery E. Glucosinolate hydrolysis and bioavailability of resulting isothiocyanates: focus on glucoraphanin. *J Funct Foods* 2014;7:67–76.
- [26] Joseph MA, Moysich KB, Freudenheim JL, Shields PG, Bowman ED, Zhang Y, et al. Cruciferous vegetables, genetic polymorphisms in glutathione S-transferases M1 and T1, and prostate cancer risk. *Nutr Cancer* 2004;50:206–13.
- [27] Higdon JV, Delage B, Williams DE, Dashwood RH. Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res* 2007;55:224–36.
- [28] Sayeed MA, Bracci M, Lazzarini R, Tomasetti M, Amati M, Lucarini G, et al. Use of potential dietary phytochemicals to target miRNA: promising option for breast cancer prevention and treatment? *J Funct Foods* 2017;28:177–93.
- [29] Fujioka N, Ainslie-Waldman CE, Upadhyaya P, Carmella SG, Fritz VA, Rohwer C, et al. Urinary 3,3'-diindolylmethane: a biomarker of glucobrassicin exposure and indole-3-carbinol uptake in humans. *Cancer Epidemiol Biomarkers Prev* 2014;23:282–7.

- [30] Houghton CA, Fassett RG, Coombes JS. Sulforaphane and Other Nutrigenomic Nrf2 activators: can the Clinician's expectation Be matched by the reality? *Oxid Med Cell Longev* 2016;2016: 7857186–7857186.
- [31] Hu C, Eggler AL, Mesecar AD, van Breemen RB. Modification of keap1 cysteine residues by sulforaphane. *Chem Res Toxicol* 2011;24:515–21.
- [32] Riedl MA, Saxon A, Diaz-Sanchez D. Oral sulforaphane increases Phase II antioxidant enzymes in the human upper airway. *Clin Immunol* 2009;130: 244–51.
- [33] Wagner AE, Boesch-Saadatmandi C, Dose J, Schultheiss G, Rimbach G. Anti-inflammatory potential of allyl-isothiocyanate—role of Nrf2, NF-(κ) B and microRNA-155. *J Cell Mol Med* 2012;16:836–43.
- [34] Sturm C, Wagner AE. *Brassica*-derived plant bioactives as modulators of chemopreventive and inflammatory signaling pathways. *Int J Mol Sci* 2017;18.
- [35] Heiss E, Herhaus C, Klimo K, Bartsch H, Gerhäuser C. Nuclear factor kappa B is a molecular target for sulforaphane-mediated anti-inflammatory mechanisms. *J Biol Chem* 2001;276:32008–15.
- [36] Dinkova-Kostova AT, Fahey JW, Wade KL, Jenkins SN, Shapiro TA, Fuchs EJ, et al. Induction of the phase 2 response in mouse and human skin by sulforaphane-containing broccoli sprout extracts. *Cancer Epidemiol Biomarkers Prev: A Publ Am Assoc Cancer Res Cospons Am Soc Prev Oncol* 2007;16:847–51.
- [37] Harvey CJ, Thimmulappa RK, Sethi S, Kong X, Yarmus L, Brown RH, et al. Targeting Nrf2 signaling improves bacterial clearance by alveolar macrophages in patients with COPD and in a mouse model. *Sci Transl Med* 2011;3: 78ra32–78ra32.
- [38] Cho HJ, Seon MR, Lee YM, Kim J, Kim JK, Kim SG, et al. 3,3'-Diindolylmethane suppresses the inflammatory response to lipopolysaccharide in murine macrophages. *J Nutr* 2008;138:17–23.
- [39] Jeon E-J, Davaatseren M, Hwang J-T, Park JH, Hur HJ, Lee AS, et al. Effect of oral administration of 3,3'-diindolylmethane on dextran sodium sulfate-induced acute colitis in mice. *J Agric Food Chem* 2016;64:7702–9.
- [40] Kim YH, Kwon H-S, Kim DH, Shin EK, Kang YH, Park JH, et al. 3,3'-diindolylmethane attenuates colonic inflammation and tumorigenesis in mice. *Inflamm Bowel Dis* 2009;15:1164–73.
- [41] Clarke JD, Hsu A, Riedl K, Bella D, Schwartz SJ, Stevens JF, et al. Bioavailability and inter-conversion of sulforaphane and erucin in human subjects consuming broccoli sprouts or broccoli supplement in a cross-over study design. *Pharmacol Res* 2011;64:456–63.
- [42] Choi Y, Kim Y, Park S, Lee KW, Park T. Indole-3-carbinol prevents diet-induced obesity through modulation of multiple genes related to adipogenesis, thermogenesis or inflammation in the visceral adipose tissue of mice. *J Nutr Biochem* 2012;23:1732–9.