



## Review article

## Effects of Mindfulness-based interventions on physical symptoms in people with multiple sclerosis – a systematic review and meta-analysis

Robert Simpson<sup>a,b,\*</sup>, Sharon Simpson<sup>b</sup>, Nitish Ramparsad<sup>c</sup>, Maggie Lawrence<sup>d</sup>, Jo Booth<sup>d</sup>, Stewart W. Mercer<sup>e</sup><sup>a</sup> Physical Medicine & Rehabilitation, Department of Medicine, Sunnybrook Health Sciences Centre, University of Toronto, Room H390 2075 Bayview Avenue, Toronto, ON M4N 3M5, Canada<sup>b</sup> General Practice and Primary Care, Institute of Health and Wellbeing, University of Glasgow, Glasgow, United Kingdom<sup>c</sup> Robertson Centre for Biostatistics, Institute of Health and Wellbeing, University of Glasgow, Glasgow, United Kingdom<sup>d</sup> Nursing and Community Health, Glasgow Caledonian University, Glasgow, United Kingdom<sup>e</sup> General Practice and Primary Care, Usher Institute, University of Edinburgh, Edinburgh, United Kingdom

## ARTICLE INFO

## Keywords:

Multiple sclerosis  
Fatigue  
Mindfulness  
Systematic review  
Meta-analysis

## ABSTRACT

**Background:** Physical wellbeing is commonly impaired in people with multiple sclerosis (PwMS). This study aims to update our previous systematic review (2014) and conduct a meta-analysis on the efficacy of Mindfulness-based interventions (MBIs) for improving physical symptoms in PwMS.**Methods:** In November 2017 we carried out systematic searches for eligible randomised controlled trials (RCTs) in seven major databases, updating our search in July 2018. We used medical subject headings and key words. Two independent reviewers used pre-defined criteria to screen, data extract, quality appraise, and analyse studies. The Cochrane Collaboration risk of bias tool was used to determine study quality. Physical wellbeing was the main outcome of interest. We used the random effects model for meta-analysis, reporting effect sizes as Standardised Mean Difference (SMD). This study is registered with PROSPERO: CRD42018093171.**Results:** We identified 10 RCTs as eligible for inclusion in the systematic review (including 678 PwMS), whilst seven RCTs (555 PwMS) had data that could be used in our meta-analyses. In general, comorbidity, disability, ethnicity and socio-economic status were poorly reported. MBIs included manualised and tailored interventions, treatment duration 6-9 weeks, delivered face-to-face and online in groups and also individually. For fatigue, against any comparator SMD was 0.24 (0.08 – 0.41),  $I^2 = 0\%$ ; against active comparators only, SMD was 0.10 (-0.14 – 0.34),  $I^2 = 0\%$ . For pain SMD was 0.16 (-0.46 – 0.79),  $I^2 = 77\%$ . Three adverse events occurred across all studies.**Conclusions:** MBIs appear to be an effective treatment for fatigue in PwMS. The optimal MBI in this context remains unclear. Further research into MBI optimisation, cost- and comparative-effectiveness is required.

## 1. Background

Multiple sclerosis (MS) is a complex, poorly understood chronic inflammatory and neurodegenerative condition (Ramagopalan et al., 2010). Common physical symptoms include difficulties with vision, speech, swallow, bowel, bladder and sexual function, chronic pain, spasticity and limited mobility (Ramagopalan et al., 2010). Comorbidity, or the presence of an additional long-term condition besides MS, is common among people with multiple sclerosis (PwMS) (Simpson et al., 2014). Physical comorbidities in MS are associated with more CNS lesions on Magnetic Resonance Imaging (MRI), greater levels

of disability, increased hospitalisations, and higher mortality rates (Marrie, 2017). Furthermore, having additional physical conditions in MS is associated with more stress and worse quality of life (QoL); as the number of additional physical conditions increase, so does the prevalence of mental health impairment (Simpson et al., 2014).

Among physical comorbidities in PwMS, hypertension, hyperlipidaemia and chronic lung disease predominate (Marrie et al., 2015). Specific care guidelines for managing these physical comorbidities in PwMS do not exist (Marrie, 2017). Fatigue and chronic pain are among the commonest symptoms reported by PwMS (Bol et al., 2009; Kratz et al., 2011). The UK National Institute for Care and Clinical Excellence (NICE) recommends

\* Corresponding author at: Physical Medicine & Rehabilitation, Department of Medicine, Sunnybrook Health Sciences Centre, University of Toronto, Room H390 2075 Bayview Avenue, Toronto, ON M4N 3M5, Canada.

E-mail address: [robert.simpson@sunnybrook.ca](mailto:robert.simpson@sunnybrook.ca) (R. Simpson).

<https://doi.org/10.1016/j.msard.2019.101493>

Received 22 January 2019; Received in revised form 28 October 2019; Accepted 31 October 2019

2211-0348/© 2019 Elsevier B.V. All rights reserved.

offering PwMS cognitive behavioural therapy (CBT), aerobic exercise, yoga, or amantadine for fatigue, as well as avoiding stress and treating comorbid anxiety and depression ((2014) NifHaCE 2014). For chronic pain in PwMS, NICE recommends the application of generic treatment approaches ((2014) NifHaCE 2014).

Mindfulness-based interventions (MBIs) fit the UK Medical Research Council criteria for complex interventions (MRC, 2008), with multiple potential active components. Originally introduced in North America in the 1980s as a treatment for people with chronic pain (Kabat-Zinn, 1982), MBIs characteristically include a range of meditation practices, group exercises, psychoeducation and home practices (Kabat-Zinn, 1990; Segal et al., 2012). MBIs have been applied and researched in a range of health conditions and found to be effective treatments for anxiety, stress, recurrent depression and somatisation disorders (Fjorback et al., 2011; Lakhan and Schofield, 2013). In a previous systematic review of the effectiveness of MBIs in PwMS in 2014 (Simpson et al., 2014) we found limited evidence from two randomised controlled trials (RCTs) and a controlled trial to support MBIs as a potential treatment for comorbid fatigue and comorbid pain in the condition, as well as improving standing balance (Simpson et al., 2014). Since 2014, several more RCTs have been published and it is important to determine more definitively whether MBIs are effective treatments for fatigue and pain in PwMS, besides other commonly encountered physical symptoms.

The aim of this review is to conduct a meta-analysis of RCTs testing the efficacy of MBIs in improving physical symptoms in PwMS.

## 2. Methods

### 2.1. Protocol and registration

Our protocol was registered prospectively with the Centre for Reviews and Dissemination, University of York, Prospero ID: CRD42018093171. This body of work also included a meta-analysis of MBI effects on mental wellbeing in PwMS, reported separately (Simpson et al., 2019).

### 2.2. Eligibility for inclusion

We based eligibility on the Study design, Participants, Interventions, Outcomes (SPIO) model (deriving from PICOS) (Richardson et al., 1995). To be eligible for inclusion, studies had to be RCTs, (comparing MBI vs active comparator or care as usual), with no limit placed on sample size. Participants had to be PwMS (of any phenotype), aged 18 years or older. The intervention(s) being tested had to be a recognisable MBI that included core practices of mindful breathing, mindful body awareness, and mindful movement; Mindfulness-based stress reduction (MBSR) and Mindfulness-based cognitive therapy (MBCT) served as reference guides in this regard. Outcomes had to be appropriately validated and report on a definable aspect of physical wellbeing experienced by PwMS (e.g. symptoms such as fatigue, pain, standing balance).

### 2.3. Search strategy

We employed a search strategy from our previous systematic review for use in: Allied and Complementary Medicines Database (AMED), Cochrane Central Register of Controlled Trials, Cumulative Index of Nursing and Allied Health Literature (CINAHL), ExcerptaMedicadatabase (EMBASE), Medical Literature Analysis and Retrieval System Online (MEDLINE), and PsycINFO. As our previous systematic review found the first study in this area was published in 2000, we set our 'years' delimiter to 2000 – 2018. In addition, we also searched ProQuest Dissertations & Theses, reviewed key references from identified studies, searched the grey literature, and approached experts in the field. We carried out our initial search in November 2017 and repeated this in July 2018. Our search strategy as used in MEDLINE is available in Appendix A.

### 2.4. Study selection, storage and screening

We imported search results into COVIDENCE, a data storage package for systematic reviews. Title/abstracts were screened by two reviewers (RS, SS) for potential eligibility using keywords like 'mindfulness' and 'multiple sclerosis'. Selected studies were then assessed against SPIO criteria by two reviewers (JB, RS) to assess ultimate eligibility. A senior, third party reviewer (SM) was available to arbitrate any disagreements.

### 2.5. Data collection/data items

Data from the final list of included studies was extracted guided by CONSORT (Schulz et al., 2010) and TIDieR (Hoffmann et al., 2014) checklist categories (Appendix B).

### 2.6. Quality appraisal

We used the Cochrane Collaboration's tool for assessing risk of bias (RoB) (Higgins et al., 2011) to summarise risk for individual outcomes in selected studies, graded as high, unclear, or low risk. This assessed generation of sequence, concealment of allocation, blinding of participants, outcome assessors and personnel, incomplete outcomes, selective reporting of outcomes, and any other bias. Finally, as outlined by Higgins et al. (2011), an overall RoB within each trial was determined based on the number of individual outcomes falling in to the high, unclear, and low risk categories:

Low = Low RoB for all key domains

Unclear = Low or unclear RoB for all key domains

High = High RoB for one or more key domains

### 2.7. Principal summary measures

The main outcome for this study was impact of MBI on physical symptoms. Main outcome measures were all reported as continuous with mean, standard deviation (SD) values and the number of participants for each treatment group extracted. "Effect size" is reported as the unbiased standardised mean difference (SMD), a positive SMD indicating a finding in support of the intervention having a positive treatment effect (TE). The standardised mean difference was calculated by difference in means between the MBI and the control group at last point of follow-up divided by the pooled last point of follow-up SD. Where effect estimates were reported from adjusted regression models, we extracted these as the SMD with their corresponding SD.

### 2.8. Synthesis of results

Throughout this study we adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Moher et al., 2009) guidance. We used a random-effects meta-analysis, with an inverse variance method for pooling (DerSimonian and Laird, 2015) to determine SMD, as outcome measures were known to vary widely. We report estimates with corresponding 95% confidence intervals (CI) and 'p' values. We used the  $I^2$  statistic to determine variability between studies (Higgins et al., 2003);  $I^2$  representing the percentage of total variability in effect size estimates due to heterogeneity. An  $I^2$  of 0% indicates that all heterogeneity is consistent with sampling error, whilst an  $I^2$  of 100% suggests all variability may be attributable to studies being truly heterogeneous.

To assess for evidence of publication bias, we undertook Funnel plots and Egger's Test for asymmetry (Sterne et al., 2008; Egger et al., 1997).

We carried out all statistical analyses in R version 3.4.0 and using the meta package (Schwarzer, 2007).




Fig. 1. PRISMA flow diagram.

### 3. Results

We identified ten RCTs as eligible for inclusion in the systematic review, with seven studies reporting endpoint data usable in meta-analysis (Fig. 1). We sought additional information from several study authors; one (Cavalera et al., 2018) replied.

#### 3.1. Systematic review

##### 3.1.1. Study characteristics

Three studies took place in Iran (Mahdavi et al., 2016; Nejati et al., 2016; Mosalanejad et al., 2018), three in the UK (Mills and Allen, 2000, Bogosian et al., 2015, Simpson et al., 2017), two in Italy (Cavalera et al., 2018; Carletto et al., 2017), one each in the USA (Senders et al., 2018) and Switzerland (Grossman et al., 2010). Four studies tested a MBI against treatment as usual (Bogosian et al., 2015, Mills and Allen, 2000, Simpson et al., 2017; Grossman et al., 2010), four versus an active comparator (three a psycho-education control Cavalera et al., 2018; Carletto et al., 2017; Senders et al., 2018, one pelvic floor muscle exercises Mosalanejad et al., 2018), and in two the control condition was not clearly specified (Mahdavi et al., 2016; Nejati et al., 2016). Three study sample sizes were based on statistical

power calculations (Cavalera et al., 2018; Carletto et al., 2017; Grossman et al., 2010). Number of study participants ranged from 24 – 150 (median 62). Eight studies reported measuring outcomes at three points in time (baseline, immediately post MBI, and at further follow-up, which varied from 1 month post MBI to 1 year later) (Cavalera et al., 2018; Mosalanejad et al., 2018; Grossman et al., 2010, Simpson et al., 2017, Bogosian et al., 2015, Carletto et al., 2017, Senders et al., 2018, Mills and Allen, 2000), whilst two studies took measures twice, pre and post MBI (Mahdavi et al., 2016; Nejati et al., 2016) (Table 1).

##### 3.1.2. Participant characteristics

There were 678 participants between the 10 RCTs included in the systematic review, versus 555 participants in the seven studies included in the meta-analysis. Participant ethnicity was described in three studies (Bogosian et al., 2015; Simpson et al., 2017; Senders et al., 2018), most were Caucasian. Between all 10 RCTs, the majority of participants were female (76%; n = 517). The extractable mean participant age was 46.0 years (not reported in Mahdavi et al., 2016). One study reported on socioeconomic status (SES) using post-code derived data (Simpson et al., 2017). Three studies described negligible data on employment status of participants (Mills and Allen, 2000; Simpson et al.,

**Table 1**  
Study characteristics.

Study	Country	Study design	Powered(Y/N/ unclear)	Comparator	Sample size (n)	Study attrition (%)	Outcome measures (others)	Data collection
Mills and Allen (2000)	Wales (UK)	Randomised controlled trial	N	Treatment as usual	24	33	Profile of Mood States, Standing balance, Symptom rating questionnaire	oBaseline oPost MBI o3 months post MBI
Grossman et al. (2010)	Switzerland	Randomised controlled trial	Y	Treatment as usual	150	5	Center for Epidemiological Studies Depression, Spielberger Trait Anxiety Inventory, Modified Fatigue Impact Scale, Hamburg Quality of Life Questionnaire in Multiple Sclerosis, Profile of health-related Quality Of Life in Chronic disorders, Goal setting, Neuropsychology assessment	oBaseline oPost MBI o6 months post MBI
Bogossian et al. (2015)	England (UK)	Randomised controlled trial	N	Treatment as usual	40	5	General Health Questionnaire, Hospital Anxiety and Depression Scale, Multiple Sclerosis Impact Scale-29, EuroQol, Fatigue Severity Scale, Numerical Rating Scale (Pain)	oBaseline oPost MBI o3 months post MBI
Mahdavi et al. (2016)	Iran	Randomised controlled trial	N	Indeterminate	24	0	Beck Anxiety Inventory, Beck Depression Inventory, Fatigue Severity Scale, Meta-Worry Questionnaire, Thought Fusion Inventory	oBaseline oPost MBI oPost MBI
Nejati et al. (2016)	Iran	Randomised controlled trial	Unclear	Indeterminate	24	0	Multiple Sclerosis Quality of Life-54, Fatigue Severity Scale	oBaseline oPost MBI oBaseline oPost MBI
Simpson et al. (2017)	Scotland (UK)	Randomised controlled trial	N	Treatment as usual	50	12	Perceived Stress Scale, EuroQol, Multiple Sclerosis Quality of Life Inventory, Mindful Attention Awareness Scale, Self-Compassion Scale-short form, Emotional Lability Questionnaire	o3 months post MBI
Carletto et al. (2017)	Italy	Randomised controlled trial	Y	Psycho-education	90	21	Beck Anxiety Inventory, Beck Depression Inventory, Perceived Stress Scale, Brief Illness Perception Questionnaire, Functional Assessment of Multiple Sclerosis, Fatigue Severity Scale	oBaseline oPost MBI o6 months post MBI
Cavalera et al. (2018)	Italy	Randomised controlled trial	Y	Psycho-education	139	39	Multiple Sclerosis Quality of Life-54, Hospital Anxiety and Depression Scale, Medical Outcomes Sleep Scale, Modified Fatigue Impact Scale	oBaseline oPost MBI o6 months post MBI
Mossalanejad et al. (2018)	Iran	Randomised controlled trial	Unclear	Pelvic floor muscle exercises	75	7	Female Sexual Function Index	oBaseline oPost MBI o1-month post MBI
Senders et al. (2018)	USA	Randomised controlled trial	Y	Psycho-education	62	16	Perceived Stress Scale, Patient-Reported Outcomes Information System, Connor-Davidson Resilience Scale, Paced Auditory Serial Attention Task	oBaseline oMid- intervention oPost MBI o4 months post MBI o8 months post MBI o12 months post- MBI

2017; Carletto et al., 2017). Seven studies reported education status (Cavalera et al., 2018; Mahdavi et al., 2016; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Senders et al., 2018; Grossman et al., 2010), most having school level education as a minimum. The majority (a minimum of 414 or 61%) had a relapsing-remitting phenotype, a minimum of 112 (17%) a secondary progressive phenotype, and a minimum of 27 (4%) had a primary progressive phenotype. Degree of disability was reported in six studies (Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Cavalera et al., 2018, Senders et al., 2018), using the Expanded Disability Status Scale (EDSS) with a range of 2.3 – 6.5. Comorbidity (mental and physical) count was described in one study (Simpson et al., 2017) (mean 2.3, SD 1.7). Four studies (Cavalera et al., 2018; Simpson et al., 2017; Senders et al., 2018; Grossman et al., 2010) described use of psychotropic and/or MS disease modifying drugs (Table 2).

### 3.1.3. Intervention characteristics

MBSR was explicitly used as the MBI in four studies (Cavalera et al., 2018; Simpson et al., 2017; Senders et al., 2018; Grossman et al., 2010) and the loose basis in a further two (Nejati et al., 2016; Carletto et al., 2017), two explicitly used MBCT (Mahdavi et al., 2016; Bogosian et al., 2015), one described the intervention as 'Mindfulness of Movement' (Mills and Allen, 2000), and in the remaining case the foundation for the MBI was unclear (Mosalanjad et al., 2018). Five studies reported on what course materials were provided to those taking part (Mills and Allen, 2000, Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017). An interview was compulsory prior to taking part in three studies (Mahdavi et al., 2016; Nejati et al., 2016; Grossman et al., 2010). Two studies required evidence of impaired mental wellbeing (stress, anxiety) at baseline in order to take part (Bogosian et al., 2015; Senders et al., 2018). Six studies reported on what MBI sessions comprised (Mahdavi et al., 2016; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Mosalanjad et al., 2018, Senders et al., 2018), three provided scant information in this regard (Mills and Allen, 2000; Carletto et al., 2017; Grossman et al., 2010), and in another this information was available in a separate publication, via the study protocol (Cavalera et al., 2018). Home practices were prescribed in six studies (Bogosian et al., 2015; Mills and Allen, 2000; Simpson et al., 2017; Carletto et al., 2017; Senders et al., 2018; Grossman et al., 2010). Teacher characteristics (training/certification/experience) were outlined in seven studies (Cavalera et al., 2018; Mosalanjad et al., 2018; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018), but details were sparse in one (Mosalanjad et al., 2018). MBIs were delivered as groups in nine studies (Cavalera et al., 2018; Mahdavi et al., 2016; Nejati et al., 2016; Mosalanjad et al., 2018; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018), the remaining study delivered a one-to-one MBI (Mills and Allen, 2000). An online platform was used to deliver the MBI in two studies (Cavalera et al., 2018; Bogosian et al., 2015). Four studies reported where the MBI took place (Cavalera et al., 2018; Mosalanjad et al., 2018; Bogosian et al., 2015; Simpson et al., 2017). The majority of studies used eight MBI sessions (Cavalera et al., 2018; Mahdavi et al., 2016; Nejati et al., 2016; Mosalanjad et al., 2018; Bogosian et al., 2015, Simpson et al., 2017; Senders et al., 2018, Carletto et al., 2017), there were nine in one study (Grossman et al., 2010), another used six (Mills and Allen, 2000). Weekly MBI session lengths varied between 1–3 h. There were between five to 25 participants per MBI class across the studies, sessions being administered by 1–2 MBI instructors. The core MBI components were delivered in all studies. However, in six studies the MBI was tailored for PwMS (Cavalera et al., 2018; Bogosian et al., 2015; Carletto et al., 2017; Grossman et al., 2010, Mills and Allen, 2000, Simpson et al., 2017), mostly in advance, but reflexively in one case (Simpson et al., 2017), where mindful movement was simplified to accommodate high levels of

disability. Another study pre-emptively removed mindful movement following stakeholder consultation (Bogosian et al., 2015). Home practice completion and/or session attendance was used to determine treatment adherence in six studies (Mills and Allen, 2000, Cavalera et al., 2018; Bogosian et al., 2015; Simpson et al., 2017; Senders et al., 2018; Grossman et al., 2010). Intervention fidelity was appraised in three studies (Cavalera et al., 2018; Bogosian et al., 2015; Simpson et al., 2017), in one case by an independent observer checking session content against referenced standards (Bogosian et al., 2015). The day retreat, characteristically part of week six in MBSR, was included in three studies (Carletto et al., 2017; Senders et al., 2018; Grossman et al., 2010) (Table 3 outlines intervention characteristics using the Template for Intervention Description and Replication checklist).

### 3.1.4. Outcome characteristics

Seven studies measured the impact of MBI on fatigue (Carletto et al., 2017, Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Senders et al., 2018), three on pain (Bogosian et al., 2015; Simpson et al., 2017; Senders et al., 2018), one on standing balance (Mills and Allen, 2000), one on sleep (Cavalera et al., 2018), and one on female sexual function (Mosalanjad et al., 2018). As all three studies that reported on pain also reported on fatigue, fatigue was thus chosen as the main outcome for our analysis.

Average home practice was reported in three studies (32, 29.2, 32.5 min) (Mills and Allen, 2000; Simpson et al., 2017; Grossman et al., 2010); whilst one study reported median value/ minimum-maximum range (38 min/day; 14 – 80) (Senders et al., 2018). Attrition ranged from 0–39% across the ten studies; those with no attrition were pre-post studies with small sample sizes (Mahdavi et al., 2016; Nejati et al., 2016).

## 3.2. Meta-analysis

### 3.2.1. Effect of MBIs on physical symptom measures

The effect of a MBI on physical symptoms was measured in 10 studies (Cavalera et al., 2018; Mahdavi et al., 2016; Nejati et al., 2016; Mosalanjad et al., 2018; Bogosian et al., 2015; Mills and Allen, 2000; Simpson et al., 2017; Carletto et al., 2017; Senders et al., 2018; Grossman et al., 2010); seven reported endpoint data usable in the meta-analysis (Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018). Seven studies evaluated MBI effect on fatigue (Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Senders et al., 2018, Carletto et al., 2017), where the SMD against any comparator was 0.24 (0.08 – 0.41)  $p < 0.01$ ,  $I^2 = 0\%$  (low heterogeneity) (Fig. 2); against active comparators only the SMD for fatigue was 0.10 (-0.14 – 0.34),  $p = 0.40$ ,  $I^2 = 0\%$  (low heterogeneity) (Fig. 3). Three studies also evaluated MBI effect on pain (besides fatigue) (Bogosian et al., 2015; Simpson et al., 2017; Senders et al., 2018), where the SMD was 0.16 (-0.46 – 0.79),  $p = 0.61$ ,  $I^2 = 77\%$  (substantial heterogeneity) (Fig. 4).

### 3.2.2. Heterogeneity and publication bias

Using the  $I^2$  statistic, heterogeneity was low for fatigue (0%), but substantial for pain (77%). The funnel plot for fatigue identified no evidence of publication bias (Fig. 5). The p-value from Egger's Test of asymmetry from fatigue studies was 0.256.

### 3.2.3. Outcomes by intervention type

Where MBSR was used (four studies Cavalera et al., 2018; Simpson et al., 2017; Senders et al., 2018; Grossman et al., 2010);  $n = 401$ , SMD for fatigue was 0.22 (0.01 – 0.42),  $p = 0.04$ ,  $I^2 = 0\%$ ; for pain (two studies Simpson et al., 2017; Senders et al., 2018) SMD was -0.07 (-0.83 – 0.68),  $p = 0.85$ ,  $I^2 = 74\%$ . Outcomes for MBCT came from

**Table 2**  
Participant characteristics.

Study/ demographic	Mills and Allen (2000)	Grossmanetal. (2010)	Bogosian et al. (2015)	Mahdavi et al. (2016)	Nejadi et al. (2016)	Simpson et al. (2017)	Carletto et al. (2017)	Cavalera et al. (2018)	Mosalanejad- et al. (2018)	Senders et al. (2018)
Ethnicity	Not reported	Not reported	90% British Caucasian	Not reported	Not reported	100% British Caucasian	Not reported	Not reported	Not reported	97% Caucasian
Number of participants (% F)	16 (80%)	150 (80%)	40 (55%)	24 (100%)	24 (46%)	50 (92%)	90 (71%)	139 (65%)	75 (100%)	67 (78%)
Mean age (SD)	49.8 (6.8)	47.3 (10.3)	52.2 (9.1)	NR	32.3 (5.1)	45 (10.9)	44.6 (9.4)	42.7 (8.7)	37.5 (6.5)	52.94 (11.37)
Socio-economic status	Not reported	Not reported	Not reported	Not reported	Not reported	Postcode derived; controlled in analyses	Not reported	Not reported	Not reported	Not reported
Employment status	4 employed (25%)	Not reported	Not reported	Not reported	Not reported	20 employed (40%)	59 employed (65%)	Not reported	Not reported	Not reported
Education status (SD)	Not reported	Mean (SD) 14.1 (1.9) years of education	31 (77.5) college education at least	Completed high school	Completed high school	(56%) university	Not reported	11% elementary school;52% completed high school;38% university	Not reported	60% college education at least
Disease phenotype (%)	Secondary progressive 16 (100%)	Relapsing 123 (82%)	Secondary progressive 23 (57.5%)	Not reported	Not reported	Relapsing 40 (80%)	Relapsing 79 (88%)	Relapsing 131 (93%)	Not reported	Relapsing 41 (67%)
		Secondary progressive 27 (18%)	Primary progressive 17 (42.5%)			Secondary progressive 16 (32%)	Secondary progressive 7 (8%)	Secondary progressive 8 (7%)		Secondary progressive 15 (25%)
						Primary progressive 4 (8%)	Primary progressive 2 (2%)			Primary progressive 4 (6%)
EDSS score	Not reported	Mean (SD) 3.0 (1.1)	Mean (SD) 6.5 (1.5)	Not reported	Not reported	4.4 (1.8)	2.3 (1.7)	Median 3.0	Not reported	Unknown 2 (3%)
Comorbidities	Not reported	Not reported	Not reported	Not reported	Not reported	Mean 2.4 (2.0); Range 0-9	Not reported	1 participant had severe depression on HADS	Excluded if comorbid conditions	4.6 (1.93)
On DMDS (85%)	Not reported	91 (60.1%)	Not reported	Not reported	Not reported	26 (52%)	Not reported	104	Not reported	Not reported
Psychotropic medication (s)	Not reported	34 (55%)	Not reported	Not reported	No	23 (46%)	Not reported	9 (6%)	Not reported	35 (56%)


**Table 3**  
Template for intervention description and replication (TIDieR) checklist.

Study/ checklist item	Mills and Allen, 2000	Grossman et al., 2010	Bogosian et al., 2015	Mahdavi et al., 2016	Nejati et al., 2016	Simpson et al., 2017	Carletto et al., 2017	Cavalera et al., 2018	Mosalanejad et al., 2018	Senders et al., 2018
<b>1. Brief name?</b>	Mindfulness of Movement	Mindfulness-based stress reduction (MBSR)	Mindfulness-based cognitive therapy	Mindfulness-based cognitive therapy	MBSR and Conscious Yoga	Mindfulness-based stress reduction	Modified MBSR – ‘Body Affective Mindfulness’	Mindfulness-based stress reduction	‘Mindfulness’	Mindfulness-based stress reduction
<b>2. Why (stated rationale/theory/ goal)?</b>	Develop moment to moment awareness of breath, posture, movement with compassion	Adaptation of MBSR. Focus on negative thinking, engaging low mood, changing relationship with thoughts, feelings, sensations, no longer avoiding/ reacting to them automatically	Adaptation of MBSR. Focus on negative thinking, engaging low mood, changing relationship with thoughts, feelings, sensations, no longer avoiding/ reacting to them automatically	Adaptation of MBSR. Focus on negative thinking, engaging low mood, changing relationship with thoughts, feelings, sensations, no longer avoiding/ reacting to them automatically	Facilitate the compliance with and adaptation to medical conditions. Pay attention to being present in a non-judgmental manner	Cultivate interested, accepting, non-judgmental attitude to experience, including difficult sensations, emotions, thoughts and behavior	Cultivation of mindful awareness, loving kindness, enrichment of listening, self-compassion, sensorimotor psychotherapy principles ‘window of tolerance’	Cultivate interested, accepting, non-judgmental attitude to experience, including difficult sensations, emotions, thoughts and behavior	Non-judgmental present moment awareness	Cultivate interested, accepting, non-judgmental attitude to experience, including difficult sensations, emotions, thoughts and behavior
<b>3. What-Materials provided to participants?</b>	Written handout, audio and video aids	Headset, webcam, compact discs for home practice	Not reported	Not reported	Leaflets for each session and compact discs for home practice	Course manual, compact discs for home practice <i>Book - Full Catastrophe Living</i>	Not reported	Dedicated website with online multimedia for home practices	Not reported	Not reported
<b>4. What - Procedures –In session?</b>	Had to make a commitment to regular practice	Screened for evidence of distress on General Health Questionnaire	Personal intake interview	Personal intake interview	Personal intake interview	Session content reported in paper –	General description in trial protocol –	General description only -	Session content reported in paper –	Score of at least 1 Item on Perceived Stress Scale
<b>4. What - Procedures –Home practice?</b>	Body awareness, breath awareness, mindful movement, Tui Na self-massage	Raisin exercise, Mindful awareness, body scan, sitting practice, three-minute breathing space, psycho-education, cognitive exercises	Sustained attentional focus on the body and breath, decentered view of thoughts as passing mental events	Body awareness, raise exercise, three-minute breathing, yoga, sitting meditation, psycho-education on stress, mountain meditation	Body awareness, raise exercise, three-minute breathing, yoga, sitting meditation, psycho-education on stress, mountain meditation	Raisin exercise, Mindful breathing, body scan, mindful movement, psycho-education	Emphasis on sensorimotor resources: grounding, self-centring, self-soothing, psycho-education on stress, self-compassion, body scan, breath meditation, walking meditation, yoga exercises	Based on original Mindfulness-based stress reduction protocol	Mindful breathing, body scan, sitting meditation, mountain eating, choice-less awareness, loving kindness, psycho-education	Mindful breathing, body scan, mindful movement, loving kindness, sitting meditation, push-pull exercise, psycho-education on stress
<b>4. What - Procedures –Post-course?</b>	Thirty minutes per day	Ten-twenty minutes per day	Not reported	Not reported	Not reported	Forty-five minutes per day	Forty-five minutes per day	Not reported	Not reported	Forty-five minutes per day
<b>4. What - Procedures –Post-course?</b>	Post course interviews for all participants	Post course interviews for some participants	Not reported	Not reported	Not reported	Post course interviews for some participants	Not reported	Not reported	Not reported	Not reported

(continued on next page)

Table 3 (continued)

Study/ checklist item	Mills and Allen, 2000	Grossman et al., 2010	Bogosian et al., 2015	Mahdavi et al., 2016	Nejati et al., 2016	Simpson et al., 2017	Carletto et al., 2017	Cavalera et al., 2018	Mosalanejad et al., 2018	Senders et al., 2018
<b>5. Who provided?</b>	Not reported	Two experienced (over nine years), certified teachers	Study author: Had completed MBI teacher training	Not reported	Not reported	Two experienced (seven and a half years), certified physician teachers	Trained clinical psychologists, used to working with people with multiple sclerosis	Expert MBSR trainer	Study author	Certified MBSR teacher with sixteen years of experience
<b>6. How - Mode of delivery?</b>	One-to-one, face-to-face	Group, face-to-face, ten-fifteen per group	Group, via Skype, up to five per group	Group, twelve per group	Group, twelve per group	Group, face-to-face, twenty-five per group	Group, number per group not reported	Group, via Skype, average of five per group	Not reported	Group, number per group not reported
<b>7. Where</b>	Unclear	Unclear	Participants' own homes	Unclear	Unclear	NHS Centre for Integrative Care	Unclear	In patients own homes	University hospital out-patient clinic	Not reported
<b>8. When and how much?</b>	Six weekly sessions	Nine weekly two and a half hour sessions Seven-hour practice day at week six	Eight weekly one hour sessions	Eight weekly two hour sessions	Eight weekly two hour sessions	Eight weekly two and a half hour sessions	Eight weekly three hour sessions Seven-hour practice day	Eight weekly sessions (? duration)	Eight weekly ninety minute sessions	Eight weekly two hour sessions Six-hour practice day at week six
<b>9. Tailoring?</b>	Individualised application of core techniques	Exercises did not exceed level of function	Developed with people with multiple sclerosis. MBCT manual adapted for Progressive multiple sclerosis issues Mindful-movement removed	Not reported	Not reported	Developed with people with multiple sclerosis, informed MBSR optimisation for future iteration	Protocol reports tailoring to needs of participants, but not reported in paper	Music meditations and acceptance of multiple sclerosis symptoms introduced	Not reported	Not reported
<b>10. In study modifications?</b>	Not reported	Not reported	Not reported	Not reported	Not reported	Mindful movement simplified	Not reported	Not reported	Not reported	Not reported
<b>11. How well</b>	Average thirty-two minutes home practice per day	Ninety-two percent session attendance; Average twenty-nine-point two minutes home practice per day	Ninety-five percent completed four or more sessions. Home practice not reported	Not reported	Not reported	Sixty percent session attendance; Average thirty-two and a half minutes home practice per day	Not reported	Seventy-nine percent session attendance	Not reported	Eighty-five percent attended six or more sessions. Median home practices thirty-eight minutes per day
<b>12. How well</b>	Not reported	Not reported	Senior clinical psychologist listened to session recordings for every session	Not reported	Not reported	Based on National Institutes of Health (2004)	Not reported	Treatment integrity monitored, but not reported in what way	Not reported	Not reported



TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study

**Fig. 2.** SMD for fatigue vs any comparator. TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study.


a single pilot study (Bogosian et al., 2015) ( $n=40$ ) versus usual care, where effect size for fatigue was 0.29 (-0.18 – 0.76),  $p=0.30$  and the effect size for pain was 0.59 (0.14 – 1.04),  $p<0.05$ . Compared to a psychoeducation control, a study using Body-Affective Mindfulness ( $n=90$ ) (Carletto et al., 2017) had an effect size of 0.19 (-0.22 – 0.60),  $p=0.37$  for effect on fatigue.

**3.3. Study quality**

Study quality was highly variable. Assessment was frequently made challenging by scanty reporting. For unclear reasons, those studies of highest quality (lowest RoB) originated from European countries and the United States. Eight studies outlined random sequence generation (Carletto et al., 2017, Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Mosalanejad et al., 2018, Senders et al., 2018). Five studies were adjudged low risk for allocation concealment (Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018), with the remainder unclear (Cavalera et al., 2018; Mahdavi et al., 2016; Nejati et al., 2016; Mosalanejad et al., 2018; Mills and Allen, 2000). Blinding of assessors was outlined in six studies (Mosalanejad et al., 2018; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018), as was outcome assessor blinding (Mosalanejad et al., 2018; Bogosian et al.,


2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018). Five studies were deemed low risk when assessing reporting of outcomes as incomplete (Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018). One study was assessed as at high risk for selective reporting of outcomes (Mills and Allen, 2000). In terms of overall within trials RoB assessments, five studies were deemed low risk (Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010, Carletto et al., 2017, Senders et al., 2018), two unclear (Mosalanejad et al., 2018, Cavalera et al., 2018), and three as high (Mahdavi et al., 2016; Nejati et al., 2016; Mills and Allen, 2000). Across trials, overall RoB was low for random sequence allocation, unclear for allocation concealment, low for assessor blinding, low for blinding of outcome assessment, unclear for incomplete outcome assessment, low for selective outcome reporting, and low for other sources of bias. Overall, there is an unclear RoB across trials.(Table 4). Appendix C details rationale for RoB assessments.

As all the pain studies were in the low RoB group, Fig. 6 illustrates only the SMD for all trials able to be analysed for fatigue, arranged by RoB categories (low, unclear and high). Low RoB ( $n=5$ ) SMD was 0.29 (0.09 – 0.49);  $I^2=0\%$  (low heterogeneity);  $p<0.01$ . Unclear RoB ( $n=1$ ) SMD was -0.01 (-0.37 – 0.35);  $p=0.95$ . High RoB ( $n=1$ ) SMD was 0.80 (-0.04 – 1.64);  $p=0.06$ . Effect estimates did not vary significantly between RoB allocation in the overall RoB analysis,  $p=0.15$ . The low RoB



TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study

**Fig. 3.** SMD for fatigue vs active comparators. TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study.



TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study

**Fig. 4.** SMD for pain vs any comparator. TE - Treatment effect; seTE - standard error of the TE; SMD - Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study.

studies are most likely to approximate the true effect of an MBI on PwMS who have fatigue, with (generally) larger sample sizes, a higher standard of trial procedures and hence less chance of inadvertent bias.

**3.4. Adverse events**


Two studies reported on adverse events associated with MBI exposure (Simpson et al., 2017; Senders et al., 2018). In one study that used MBSR a participant reported an episode of increased spasticity during mindful body awareness (Senders et al., 2018). In the same study another participant described increased anxiety following the

MBSR retreat (Senders et al., 2018). In another study using MBSR one participant with chronic pain reported increased symptoms following the raisin exercise (Simpson et al., 2017).

**4. Discussion**

**4.1. Summary of main findings**

Ten RCTs that assessed the effects of an MBI on physical symptom outcomes in PwMS were eligible for inclusion in our systematic review; seven of these had data extractable for use in our meta-analysis. Four



**Fig. 5.** Funnel plot for fatigue studies.

**Table 4**  
Risk of bias.

	Mills and Allen (2000)	Grossman et al. (2010)	Bogosian et al. (2015)	Mahdavi et al. (2016)	Nejati et al. (2016)	Simpson et al. (2017)	Carletto et al. (2017)	Cavalera et al. (2018)	Mosalanejad et al. (2018)	Senders et al. (2018)	Across trials – overall RoB
Random sequence generation	Unclear	Low	Low	Unclear	Low	Low	Low	Low	Low	Low	Low
Allocation concealment	Unclear	Low	Low	Unclear	Unclear	Low	Low	Unclear	Unclear	Low	Unclear
Blinding of assessors	Unclear	Low	Low	Unclear	Unclear	Low	Low	Unclear	Low	Low	Low
Blinding of outcome assessment	High	Low	Low	Unclear	Unclear	Low	Low	Unclear	Low	Low	Low
Incomplete outcome data addressed	Unclear	Low	Low	Unclear	Unclear	Low	Low	Unclear	Unclear	Low	Unclear
Selective outcome reporting	High	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low
Other sources of bias	Unclear	Low	Low	High	High	Low	Low	Low	Low	Low	Low
Within trials overall RoB	High	Low	Low	High	High	Low	Low	Unclear	Unclear	Low	Unclear

studies tested an MBI against an active comparator, four tested against treatment as usual, whilst the control condition was unclear in the remaining two studies. Intervention fidelity was reliably assessed in only one study. Sample sizes were frequently small. Follow-ups took place from immediately post-MBI to up to 1 year following course completion.

Six hundred and seventy-eight PwMS were included in these studies. Most (58%) had relapsing phenotypes. Most participants were female; mostly of Caucasian ethnicity. In general, comorbidity and disability levels were poorly reported.

Four studies used MBSR, two were loosely modeled on MBSR; two explicitly used MBCT, one ‘Mindfulness of Movement’, and in one case the basis for the MBI was unclear. Most interventions were provided as groups (n=5-25), delivering core MBI practices in and between sessions. Level of teacher training and experience were not well reported. MBI session attendance +/- home practice (treatment adherence) was described in six studies. Rates of attrition varied considerably (0–39%). Although very few adverse events were described from MBI training, few studies explicitly reported on this outcome.

Five RCTs were categorised as overall low RoB using the Cochrane Collaboration tool, three as high and two as unclear, signifying an overall improvement in study quality since we last assessed this in 2014 (Simpson et al., 2014).

Our meta-analysis indicates that MBIs are modestly effective treatments for fatigue in PwMS, but evidence to support improvements in pain is inconsistent. No MBI is clearly optimal for treating impairment of fatigue in PwMS.

#### 4.2. Comparison with existing literature


In this study we found MBIs moderately effective for improving fatigue (SMD 0.24; 0.08 – 0.41), but inconsistent with regards to effects on pain (SMD 0.16; -0.46 – 0.79) in PwMS.

A 2018 meta-analysis (Phyo et al., 2018) of psychological interventions for treating fatigue in PwMS reported CBT to be moderately effective (SMD 0.32; 0.01 – 0.63) and MBIs to be considerably more effective (SMD 0.62; 0.12 – 1.12), but only included two (Bogosian et al., 2015; Grossman et al., 2010) of the seven (Cavalera et al., 2018; Nejati et al., 2016; Bogosian et al., 2015; Simpson et al., 2017; Grossman et al., 2010; Senders et al., 2018; Carletto et al., 2017) RCTs identified in our current review, likely reflecting an earlier search cut-off date in their study (April 2017).

No previous meta-analysis has assessed the impact of MBI training on pain in PwMS, but in chronic pain populations at large, several meta-analyses have been conducted. A 2014 meta-analysis (Goyal et al., 2014) reported moderate overall treatment effects (Cohen's d) from MBI training (0.33; 0.03 – 0.62), a finding that diminished to a null effect when examining the effect against active comparators. A 2015 meta-analysis (Marikar Bawa et al., 2015) comprising painful musculoskeletal conditions reported small effects (Hedge's g) versus usual care following MBI training on pain intensity (0.16; 0.03 – 0.36; the effect attenuated when compared against active comparators to 0.09; -0.13 – 0.31), moderate effects on perceived pain control (0.58; 0.23 – 0.93), but larger effects on pain acceptance versus usual care (1.58; -0.57 – 3.74). Finally, a 2017 meta-analysis (Hilton et al., 2016) found small overall effects against any comparator, SMD 0.32 (0.09 – 0.54), but included a wide variety of clinical syndromes.

#### 4.3. Strengths of this review

Guided by the PRISMA checklist (Moher et al., 2009), the TIDieR checklist (Hoffmann et al., 2014) and the Cochrane Collaboration tool (Higgins et al., 2011), our multidisciplinary team of experienced reviewers used robust search, appraisal and analysis techniques for extracting and analysing data in this systematic review and meta-analysis.



TE - Treatment effect; seTE - standard error of the TE; SMD – Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study

Fig. 6. Risk of Bias Forest plot for fatigue studies. TE - Treatment effect; seTE - standard error of the TE; SMD – Standardised mean difference; 95%CI - 95% confidence interval; Weight - weight contributed by each study.

4.4. Limitations of this review

Although we assessed quality using a reference standard, the Cochrane Collaboration RoB tool, we did not estimate the strength of any recommendation for use of MBIs in PwMS. Future studies could do so by applying the GRADE criteria (Guyatt et al., 2011).

Meta-analyses of RCTs by design exclude other potentially relevant data, such as that deriving from observational or qualitative research. When considering intervention feasibility, such as acceptability, accessibility and implementability, these alternate study designs can provide important insights into how and why interventions succeed or fail in a given context. However, in this current study, the use of SPIO, the TIDieR checklist and Cochrane Collaboration tool for RoB, means that our evidence synthesis has covered other, related aspects of trial feasibility and execution.

4.5. Strengths and Limitations of the included studies

When considering the strength of evidence for the use of MBIs in PwMS, most studies which assessed impact on fatigue ( $n = 5/7$ ) and all that assessed impact on pain ( $n = 3$ ) were adjudged low RoB. However, despite all studies being RCTs, participant numbers were low ( $n = < 50$ ) in four. Although all MS phenotypes were represented, most participants had relapsing-remitting MS. Furthermore, mean sample age was relatively low (46.0), whilst ethnicity, SES and comorbidity were poorly covered, limiting the generalisability of findings. To complicate matters, several studies tailored their MBIs with minimal/absent prior

justification. Only four compared an MBI against an active comparator condition. Observed effects were mostly small, with a wide range of confidence intervals. Heterogeneity, overall, was low.

Given the well documented high levels of physical comorbidity in PwMS, it is notable that our meta-analysis has only been able to quantify the effects of MBI training on two, albeit common, facets of physical wellbeing, namely fatigue and pain. Other aspects of physical wellbeing were measured in individual studies (e.g. standing balance, sleep and sexual function), where beneficial effects were reported, but meta-analysis was not possible. Future studies could address this evidence gap by measuring the impact of MBI training on other common physical symptoms associated with MS, for example dysarthria, dysphagia, bowel and bladder dysfunction, dynamic balance, in-coordination and spasticity. Although MBSR and MBCT both appear to be effective treatments for fatigue, it is not currently possible to recommend one approach over the other.

4.6. Implications for research

The quality of evidence for MBIs as effective treatments for fatigue in PwMS has improved considerably since our systematic review in 2014. However, adherence to CONSORT (Schulz et al., 2010) reporting was poor in several studies included in the meta-analysis, with three studies assessed overall as high risk and two as unclear according to the Cochrane Collaboration (Higgins et al., 2011) tool. In addition, MBI description was often sparse in detail. Were researchers to adhere more closely to the CONSORT (Schulz et al., 2010) and TIDieR

(Hoffmann et al., 2014) checklists when reporting studies of MBIs for PwMS, the knowledge base in this area could be further enhanced, helping to clarify where further research efforts should focus.

It remains unclear which type of MBI may be best for PwMS with impaired physical wellbeing in general, or fatigue or pain more specifically. Future research could test either MBSR or MBCT against established treatments in this area; by involving people affected with the condition in this endeavor, the co-design, delivery and ongoing development of an optimised MBI course for PwMS could take place (MRC, 2008).

#### 4.7. Implications for clinical practice

MBIs appear to be modestly effective at improving fatigue in PwMS.

## 5. Conclusions

Meta-analytic evidence supports the use of MBIs in PwMS to improve fatigue. Evidence to support the use of MBIs for treating pain in this population is inconsistent. Although the quality of study reporting has become better, room still exists for enhanced reporting in this area.

## Appendix A. MEDLINE search strategy

Search History: OVIDsp - MEDLINE with Full Text  
01 July 2018  
MS and MBSR for MS\_MBSR review 2018/19

- 1 exp Multiple Sclerosis/
- 2 expNeuromyelitisOptica/
- 3 exp Multiple Sclerosis, Chronic Progressive/
- 4 exp Multiple Sclerosis, Relapsing-Remitting/
- 5 "disseminated sclerosis".mp.
- 6 devic.mp.
- 7 "acute disseminated encephalomyelitis".mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 8 encephalomyelitis.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 9 "multiple sclerosis".mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 10 "neuromyelitoptica".mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 11 "optic neuritis".mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 12 "transverse myelitis".mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 13 demyelinat\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 14 myelitis.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 15 ((clinically or radiologically) and isolated syndrome).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 16 (demyelinating and (disease or disorder)).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 17 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16
- 18 exp Mindfulness/
- 19 exp Meditation/
- 20 exp Breathing Exercises/
- 21 (MBSR or MBCT).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 22 relaxation.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 23 (breathing and (exercis\* or techniqu\*)).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 24 vipassana.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]

No clear optimal MBI exists for improving impaired physical wellbeing in PwMS.

## Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

## Funding

This study was funded by the RS McDonald Trust (SC012710)

## Declaration of Competing Interest

We declare no competing interests.

## Acknowledgments

We wish to thank the RS McDonald trust for funding this study.

- 25 yoga.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 26 mindful\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 27 meditat\*.mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 28 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27
- 29 17 and 28
- 30 limit 29 to (english language and humans and yr="2000 -Current")

**Appendix B. Data Extraction Sheet v.1 Simpson et al. (2018)**

---

Bibliographic details	
Authors	
Year	
Country	
Citation	
Title (identifying study as an RCT y/n?)	
Structured abstract	
References identified from reference list	Yes/nolf yes, please provide details:

---



---

Study
Study aims and objectives
Study design
Comparator group
Statistical methods
Power calculation
Inclusion criteria
Exclusion criteria
Stopping criteria
Setting/location where data collected
Trial protocol/ registration
Ethical approval
Funding details y/n?

---



---

Population	Intervention group	Control /comparison group
Sample size		
Recruited from where?		
Age		
Sex		
Socioeconomic status		
Ethnicity		
Marital status		
Living Arrangements		
Educational status		
Employment status		
Disease phenotype		
Use of disease modifying drugs		
Time since diagnosis		
Disability level		
Cognitive impairment		
Comorbid anxiety (% on drug treatment)		
Comorbid depression (% on drug treatment)		
Other comorbidities		

---

---

Intervention
<b>Definition</b> Course content Tailored to population? (If yes, describe) Fidelity to treatment assessed y/n (if so, how) Course completion criteria Mode of delivery (face-to-face, internet etc) Duration & frequency Instructor characteristics No. of participants per group Intervention materials Intervention location Cost to participants Cost effectiveness Transport issues Family/carer involvement Other (specify) Intervention for control group Provide details:

---



---

	Intervention group	Control /comparison group
<b>Feasibility outcomes</b> • Recruitment (to pre-defined target y/n?) • Randomisation • Retention • Adherence (classes attendance/ home-practice completion) • Follow-up (when?) Reasons accounting for attrition reported • Missing data CONSORT flow diagram y/n Adverse events reported y/n? (specify) Standardised outcomes measures (specify) Study-specific outcomes measures (provide details) Sub-group analyses Other outcomes measured (provide details) No. of data collection time points		

---



---

<b>Limitations/conclusions/comments</b>
Limitations noted by the authors Authors' conclusions Reviewer's comments

---



---

Risk of bias assessment (High/unclear/low)
Random sequence generation (selection bias) Allocation concealment (selection bias) Blinding of assessors (performance bias) Blinding of outcome assessment (detection bias) (patient reported outcomes) Incomplete outcome data addressed (attrition bias) Selective outcome reporting (reporting bias) Other sources of bias (i.e. baseline bias)

---

## Appendix C. Cochrane Collaboration Risk of Bias assessment - justifications

**Table C1**

Cochrane Collaboration Risk of Bias – [Mills and Allen \(2000\)](#). "Mindfulness of movement as a coping strategy in multiple sclerosis: a pilot study."

Risk of bias assessment	
Random sequence generation	Unclear – First matched on basis of activities of daily living (ADL) scores, then randomly allocated from that pair only
Allocation concealment	Unclear – Not mentioned in the paper
Blinding of assessors	Unclear – Not mentioned in the paper
Blinding of outcome assessment	High – Paper suggests in discussion that study would be enhanced by including a 'blind or more objective rater'
Incomplete outcome data addressed	Unclear – Some data is reported in methods, but participant data has been omitted on the basis of not being 'complete'. Unclear what this means.
Selective outcome reporting	High – This study is also reported in another paper with different outcome data: <i>Mills N, Allen J, Carey-Morgan S. Does tai chi/qi gong help patients with multiple sclerosis? Journal of Bodywork and Movement Therapies. 2000 Jan 1;4(1):39-48.</i>
Other sources of bias	Unclear – Nothing else of note mentioned in paper

**Table C2**

Cochrane Collaboration Risk of Bias – [Grossman et al. \(2010\)](#). "MS quality of life, depression, and fatigue improve after mindfulness training A randomized trial."

Risk of bias assessment	
Random sequence generation	Low – Baseline assessments prior to randomisation. PI randomised blind, using <a href="http://www.randomizer.org">www.randomizer.org</a> in blocks of 4 - 6
Allocation concealment	Low – PI sent allocation list to coordinator who informed participants in writing of their assignment. This was then re-checked by PI, no deviations found
Blinding of assessors	Low – Investigators blinded to assignment
Blinding of outcome assessment	Low – All PRO measures were entered into a database by blinded personnel
Incomplete outcome data addressed	Low – Consort flow diagram included in report with n randomised, 'n' analysed etc. Intention to Treat (ITT) employed. Missing data imputed by multiple linear regression that adjusted for age, gender, and disease progression
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Well conducted and reported study

**Table C3**

Cochrane Collaboration Risk of Bias – [Bogosian et al. \(2015\)](#). "Distress improves after mindfulness training for progressive MS: A pilot randomised trial."

Risk of bias assessment	
Random sequence generation	Low – Randomisation took place once cohort of 10 participants consented, screened and baseline data collected. Independent unit at KCL Clinical Trials Unit (CTU) handled randomisation, with fixed block sizes of 2
Allocation concealment	Low – As above. Then CTU sent assignment list to PI
Blinding of assessors	Low – Trial assessor blinded to allocation
Blinding of outcome assessment	Low – Statistician, health economist blinded to assignment
Incomplete outcome data addressed	Low – Consort flow diagram included in report with n randomised, 'n' analysed etc. ITT employed. Informative missingness processes explored by sensitivity analysis. Missing baseline variables handled using the missing indicator method
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Well conducted and reported study

**Table C4**

Cochrane Collaboration Risk of Bias – [Mahdavi et al. \(2016\)](#). "The Effectiveness of Mindfulness-Based Cognitive Therapy in Reducing Psychological Symptoms, Meta-Worry and Thought Fusion of Multiple Sclerosis Patients."

Risk of bias assessment	
Random sequence generation	Unclear – Paper only states that participants were selected using a random sampling method
Allocation concealment	Unclear - Paper only states that participants were selected using a random sampling method
Blinding of assessors	Unclear – Not reported in the paper
Blinding of outcome assessment	Unclear – Not reported in the paper
Incomplete outcome data addressed	Unclear – Attrition not reported, nor numbers included in analyses or details regarding missing data. No consort flow diagram.
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	High – No reporting of baseline participant characteristics at all

**Table C5**

Cochrane Collaboration Risk of Bias – [Nejati et al. \(2016\)](#). "The effect of group mindfulness-based stress reduction and consciousness yoga program on quality of life and fatigue severity in patients with MS."

Risk of bias assessment	
Random sequence generation	Low – Each participant's names placed on slip of paper, mixed and drawn randomly
Allocation concealment	Unclear - Paper only states that participants were selected using a random sampling method
Blinding of assessors	Unclear – Not reported in the paper
Blinding of outcome assessment	Unclear – Not reported in the paper
Incomplete outcome data addressed	Unclear – Attrition not reported, nor numbers included in analyses or details regarding missing data
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	High – Paper states study population based on convenience sampling

**Table C6**

Cochrane Collaboration Risk of Bias – [Simpson et al. \(2017\)](#). "Mindfulness-based stress reduction for people with multiple sclerosis—a feasibility randomised controlled trial."

Risk of bias assessment	
Random sequence generation	Low – Post-baseline measures an independent statistician undertook block randomisation and sequence generation
Allocation concealment	Low – Blinded research staff undertook treatment allocation
Blinding of assessors	Low – Research staff were blinded to treatment allocation and participant ID
Blinding of outcome assessment	Low – Anonymous data was collected by a blinded research assistant
Incomplete outcome data addressed	Low – Detailed reporting of missing data, no imputation. Consort flow diagram and details accounting for participant drop-out
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Well conducted and reported study

**Table C7**

Cochrane Collaboration Risk of Bias – [Carletto et al. \(2017\)](#). "The effectiveness of a body-affective mindfulness intervention for multiple sclerosis patients with depressive symptoms: a randomized controlled clinical trial."

Risk of bias assessment	
Random sequence generation	Low – Randomly assigned on 1:1 ratio using a block wise randomisation sequence
Allocation concealment	Low – Sequence determined by an independent researcher blinded to initial assessment. Study co-ordinator communicated assignment to participants
Blinding of assessors	Low – Clinical Psychologists performing assessments were blinded to participant ID
Blinding of outcome assessment	Low – Clinical Psychologists performing assessments were blinded to participant ID
Incomplete outcome data addressed	Low – Both Per Protocol (PP) and ITT performed - ITT explored missing data – data imputation was used for two participants. Consort flow diagram detailing numbers analysed and dropping out. Comparison between completers and dropouts baseline measures and socio-demographics undertaken.
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Well conducted and reported study

**Table C8**

Cochrane Collaboration Risk of Bias – [Cavalera et al. \(2018\)](#). "Online meditation training for people with multiple sclerosis: A randomized controlled trial."

Risk of bias assessment	
Random sequence generation	Low – Participants were randomly assigned 1:1 to MBI and control using <a href="http://www.random.org">www.random.org</a>
Allocation concealment	Unclear – Paper only states that participants were randomly assigned to MBI and control
Blinding of assessors	Unclear – Not reported in the paper
Blinding of outcome assessment	Unclear – Not reported in the paper
Incomplete outcome data addressed	Unclear – Although consort flow diagram included, detailing attrition, reasons accounting for this were insufficiently described. No mention of missing data
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Generally well conducted and reported study

**Table C9**

Cochrane Collaboration Risk of Bias – [Mosallanejad et al. \(2018\)](#). "Investigating the combined effect of pelvic floor muscle exercise and mindfulness on sexual function in women with multiple sclerosis: a randomized controlled trial."

Risk of bias assessment	
Random sequence generation	Low – Block randomisation of six blocks of three patients (ABC, ACB, BAC, BCA, CAB, CBA), then replacement random sampling used to select blocks
Allocation concealment	Unclear – Not discernable from paper
Blinding of assessors	Low – Data collected by third author blinded to group assignment
Blinding of outcome assessment	Low – Data analysed by second author, blinded to group assignment
Incomplete outcome data addressed	Unclear – Consort flow diagram detailing attrition and numbers analysed, but no mention of missing data
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Generally well conducted and reported study

**Table C10**

Cochrane Collaboration Risk of Bias – Senders et al. (2018). "Impact of mindfulness-based stress reduction for people with multiple sclerosis at 8 weeks and 12 months: A randomized clinical trial."

Risk of bias assessment	
Random sequence generation	Low – Statistician generated randomisation scheme stratified by baseline PSS scores with a block size of four (SPSS random number generator)
Allocation concealment	Low – Randomisation scheme maintained by individual external and blinded to study. Allocation concealed from all study staff
Blinding of assessors	Low – Baseline data collected prior to randomisation – PI, statistician and personnel performing data entry were blinded to group assignment
Blinding of outcome assessment	Low – Baseline data collected prior to randomisation – PI, statistician and personnel performing data entry were blinded to group assignment
Incomplete outcome data addressed	Low – Low – Consort flow diagram detailing reasons accounting for attrition and numbers analysed.
Selective outcome reporting	Low – All pre-specified outcomes were reported
Other sources of bias	Low – Well conducted and reported study

## References

- (2014) NifHaCE. Multiple sclerosis: management of multiple sclerosis in primary and secondary care 2014 Available from: <https://www.nice.org.uk/guidance/cg186>.
- Bogossian, A, Chadwick, P, Windgassen, S, et al., 2015. Distress improves after mindfulness training for progressive MS: A pilot randomised trial. *Multiple Sclerosis. J* 1352458515576261.
- Bol, Y, Duits, AA, Hupperts, RM, et al., 2009. The psychology of fatigue in patients with multiple sclerosis: a review. *J. Psychosom. Res.* 66 (1), 3–11.
- Carletto, S, Tesio, V, Borghi, M, et al., 2017. The effectiveness of a body-affective mindfulness intervention for multiple sclerosis patients with depressive symptoms: a randomized controlled clinical trial. *Front. Psychol.* 8, 2083.
- Cavalera, C, Rovaris, M, Mendozzi, L, et al., 2018. Online meditation training for people with multiple sclerosis: A randomized controlled trial. *Multiple Sclerosis. J* 1352458518761187.
- DerSimonian, R, Laird, N., 2015. Meta-analysis in clinical trials revisited. *Contemp. Clin. Trials* 45, 139–145.
- Egger, M, Smith, GD, Schneider, M, et al., 1997. Bias in meta-analysis detected by a simple, graphical test. *Bmj* 315 (7109), 629–634.
- Fjorback, L, Arendt, M, Ørnbøl, E, et al., 2011. Mindfulness-Based Stress Reduction and Mindfulness-Based Cognitive Therapy—a systematic review of randomized controlled trials. *Acta Psychiatr. Scand.* 124 (2), 102–119.
- Goyal, M, Singh, S, Sibinga, EM, et al., 2014. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Int. Med.* 174 (3), 357–368.
- Grossman, P, Kappos, L, D'Souza, M, et al., 2010. MS quality of life, depression, and fatigue improve after mindfulness training: A randomized trial. *Neurology* 75 (13), 1141–1149.
- Guyatt, G, Oxman, AD, Akl, EA, et al., 2011. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J. Clinical Epidemiol.* 64 (4), 383–394.
- Higgins, JP, Altman, DG, Gotzsche, PC, et al., 2011. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *Bmj* 343, d5928.
- Higgins, JP, Thompson, SG, Deeks, JJ, et al., 2003. Measuring inconsistency in meta-analyses. *BMJ: Br. Med. J.* 327 (7414), 557.
- Hilton, L, Hempel, S, Ewing, BA, et al., 2016. Mindfulness meditation for chronic pain: systematic review and meta-analysis. *Annals Behav. Med.* 51 (2), 199–213.
- Hoffmann, TC, Glasziou, PP, Boutron, I, et al., 2014. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *Bmj* 348:g1687.
- Kabat-Zinn, J, 1990. Full Catastrophe Living: the Program of the Stress Reduction Clinic at the University Of Massachusetts Medical Center. Delta, New York.
- Kabat-Zinn, J., 1982. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: Theoretical considerations and preliminary results. *General Hosp. Psych.* 4 (1), 33–47.
- Kratz, AL, Molton, IR, Jensen, MP, et al., 2011. Further evaluation of the motivational model of pain self-management: coping with chronic pain in multiple sclerosis. *Ann. Behav. Med.* 41 (3), 391–400.
- Lakhan, SE, Schofield, KL., 2013. Mindfulness-based therapies in the treatment of somatization disorders: a systematic review and meta-analysis. *PLoS One* 8 (8), e71834.
- Mahdavi, A, Yazdanbakhsh, K, Sharifi, M, 2016. The effectiveness of mindfulness-based cognitive therapy in reducing psychological symptoms. *Meta-Worry Thought Fus.* *Mult. Scleros. Pat.*
- MarikarBawa, F, Mercer, S, Atherton, R, et al., 2015. Does mindfulness improve outcomes in chronic pain patients?: Systematic review and meta-analysis. *Br. J. Gen. Pract.*
- Marrie, RA, Cohen, J, Stuve, O, et al., 2015. A systematic review of the incidence and prevalence of comorbidity in multiple sclerosis: overview. *Multiple Sclerosis. J* 21 (3), 263–281.
- Marrie, RA., 2017. Comorbidity in multiple sclerosis: implications for patient care. *Nature Rev. Neurol.* 13 (6), 375.
- Mills, N, Allen, J., 2000. Mindfulness of movement as a coping strategy in multiple sclerosis: a pilot study. *Gen. Hospital Psych.* 22 (6), 425–431. [https://doi.org/10.1016/S0163-8343\(00\)00100-6](https://doi.org/10.1016/S0163-8343(00)00100-6).
- Moher, D, Liberati, A, Tetzlaff, J, et al., 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann. Int. Med.* 151 (4), 264–269.
- Mosalanejad, F, Afrasiabifar, A, Zolad, M, 2018. Investigating the combined effect of pelvic floor muscle exercise and mindfulness on sexual function in women with multiple sclerosis: a randomized controlled trial. *Clinical Rehabilitation* 0269215518777877.
- MRC, U, 2008. Developing and Evaluating Complex Interventions: New Guidance. Medical Research Council, London.
- Nejati, S, Esfahani, SR, Rahmani, S, et al., 2016. The effect of group mindfulness-based stress reduction and consciousness yoga program on quality of life and fatigue severity in patients with MS. *J. Caring Sci.* 5 (4), 325.
- Phyo, AZZ, Demaneuf, T, De Livera, AM, et al., 2018. The efficacy of Psychological interventions for managing Fatigue in People with multiple Sclerosis: a Systematic Review and meta-analysis. *Front. Neurol.* 9, 149.
- Ramagopalan, SV, Dobson, R, Meier, UC, et al., 2010. Multiple sclerosis: risk factors, prodromes, and potential causal pathways. *Lancet Neurol.* 9 (7), 727–739.
- Richardson, WS, Wilson, MC, Nishikawa, J, et al., 1995. The well-built clinical question: a key to evidence-based decisions. *Acad. J. Club* 123 (3), A12–A13.
- Schulz, KF, Altman, DG, Moher, D, 2010. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMC Med.* 8 (1), 18.
- Schwarzer, G., 2007. Meta: An R package for meta-analysis. *R News* 7 (3), 40–45.
- Segal, ZV, Williams, JMG, Teasdale, JD, 2012. Mindfulness-Based Cognitive Therapy for Depression. Guilford Press.
- Senders, A, Hanes, D, Bourdette, D, et al., 2018. Impact of mindfulness-based stress reduction for people with multiple sclerosis at 8 weeks and 12 months: A randomized clinical trial. *Multiple Sclerosis. J* 1352458518786650.
- Simpson, R, Booth, J, Lawrence, M, et al., 2014. Mindfulness based interventions in multiple sclerosis—a systematic review. *BMC Neurol.* 14 (1), 15.
- Simpson, R, Mair, FS, Mercer, SW, 2017. Mindfulness-based stress reduction for people with multiple sclerosis—a feasibility randomised controlled trial. *BMC Neurol.* 17 (1), 94.
- Simpson, R, Simpson, S, Ramparsad, N, et al., 2019. Mindfulness-based interventions for mental well-being among people with multiple sclerosis: a systematic review and meta-analysis of randomised controlled trials. *J. Neurol., Neurosurg. Psych* jnnp-2018-320165.
- Simpson, RJ, McLean, G, Guthrie, B, et al., 2014. Physical and mental health comorbidity is common in people with multiple sclerosis: nationally representative cross-sectional population database analysis. *BMC Neurol.* 14 (1), 128.
- Sterne, JA, Egger, M, Moher, D, 2008. Addressing reporting biases. *Cochrane Handbook for Systematic Reviews of Interventions: Cochrane Book Series.* pp. 297–333.