



Efficacy of an m-Health Physical Activity and Sleep Health Intervention for Adults: A Randomized Waitlist-Controlled Trial

Beatrice Murawski, MSc,^{1,2} Ronald C. Plotnikoff, PhD,^{1,3} Anna T. Rayward, MD,^{1,2}
Christopher Oldmeadow, PhD,^{2,4,5} Corneel Vandelanotte, PhD,⁶
Wendy J. Brown, PhD,⁷ Mitch J. Duncan, PhD^{1,2}

Introduction: Interventions that improve both physical activity and sleep quality may be more effective in improving overall health. The purpose of the Synergy Study is to test the efficacy of a mobile health combined behavior intervention targeting physical activity and sleep quality.

Study design: Randomized, waitlist-controlled trial.

Setting/participants: This study had an app-based delivery mode, Australia-wide. The participants were 160 adults who reported insufficient physical activity and poor sleep quality in an eligibility survey.

Intervention: The intervention was a mobile app providing educational resources, goal setting, self-monitoring, and feedback strategies. It included 12 weeks of personalized support including weekly reports, tool sheets, and prompts.

Main outcome measures: Outcomes were assessed at baseline, 3 months (primary), and 6 months (secondary endpoint). Self-reported minutes of moderate-to-vigorous intensity physical activity and sleep quality were co-primary outcomes. Resistance training; sitting time; sleep hygiene; sleep timing variability; insomnia severity; daytime sleepiness; quality of life; and depression, anxiety, and stress symptoms were secondary outcomes. Data were collected between June 2017 and February 2018 and analyzed in August 2018.

Results: At 3 months, between-group differences in moderate-to-vigorous intensity physical activity were not statistically significant ($p=0.139$). Significantly more participants in the intervention group engaged in ≥ 2 days/week ($p=0.004$) of resistance training. The intervention group reported better overall sleep quality ($p=0.009$), subjective sleep quality ($p=0.017$), sleep onset latency ($p=0.013$), waketime variability ($p=0.018$), sleep hygiene ($p=0.027$), insomnia severity ($p=0.002$), and lower stress symptoms ($p=0.032$) relative to waitlist controls. At 6 months, group differences were maintained for sleep hygiene ($p=0.048$), insomnia severity ($p=0.002$), and stress symptoms ($p=0.006$). Differences were observed for bedtime variability ($p=0.023$), sleepiness ($p<0.001$), daytime dysfunction ($p=0.039$), and anxiety symptoms ($p=0.003$) at 6 months, but not 3 months.

From the ¹Priority Research Centre for Physical Activity and Nutrition, University of Newcastle, Callaghan, New South Wales, Australia; ²Faculty of Health and Medicine, School of Medicine and Public Health, University of Newcastle, Callaghan, New South Wales, Australia; ³Faculty of Education and Arts, School of Education, University of Newcastle, Callaghan, New South Wales, Australia; ⁴Faculty of Health, Center for Clinical Epidemiology and Biostatistics, Callaghan, New South Wales, Australia; ⁵Clinical Research Design and Statistics Unit, Hunter Medical Research Institute, New Lambton, New South Wales, Australia; ⁶Physical Activity Research Group, Appleton Institute, Central Queensland University,

Rockhampton, Queensland, Australia; and ⁷Centre for Research on Exercise, Physical Activity and Health, School of Human Movement and Nutrition Sciences, University of Queensland, St. Lucia, Queensland, Australia

Address correspondence to: Mitch J. Duncan, PhD, ATC Building Level 3, The University of Newcastle, University Drive, Callaghan, New South Wales 2308, Australia. E-mail: mitch.duncan@newcastle.edu.au.

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2019.05.009>

Conclusions: This remotely delivered intervention did not produce statistically significant between-group differences in minutes of moderate-to-vigorous intensity physical activity. Significant short-term differences in resistance training and short- and medium-term differences in sleep health in favor of the intervention were observed.

Trial registration: This study is registered at anzctr.org.au ACTRN12617000376347.

Am J Prev Med 2019;57(4):503–514. © 2019 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

INTRODUCTION

Large proportions of the adult population are insufficiently active¹ and also report poor sleep health.^{2–5} Accruing <150 minutes/week of physical activity (PA) is considered insufficient,⁶ and poor sleep health is characterized by the presence of one or multiple complaints relating to the duration, quality, or timing of sleep and daytime functioning.⁷ The two behaviors are separately associated with increased chronic disease risk (e.g., cardiovascular disease or Type 2 diabetes) and considerable economic burden.^{8–14}

Given the high prevalence and associated burden of insufficient PA and poor sleep health in adults,¹⁵ wide-reaching interventions for PA and sleep health are needed. Mobile health (m-Health) interventions have the capacity to deliver accessible, scalable, and cost effective interventions,¹⁶ and are known to improve PA¹⁷ and reduce the severity of clinical sleep complaints.¹⁸ A meta-analysis of sleep interventions administered to adults without clinical sleep complaints reported that interventions were effective (Hedge's $g=0.54$), yet few were delivered using m-Health.¹⁹ Behavior change interventions, which implement evidence-based strategies that conceptually align with theoretic frameworks, are thought to be more effective than those not informed by theory.²⁰ This is particularly important when multiple behaviors are combined in a single intervention.²¹ Further, given that both PA and sleep are influenced by individual and environmental factors, it is useful for interventions to be guided by a theory that acknowledges this relationship.

Insufficient PA and poor sleep tend to co-occur,¹⁵ and there is evidence that PA and sleep share a bi-directional relationship.^{22–24} Consequently, interventions targeting PA and sleep concurrently may yield larger improvements in both behaviors and produce greater health benefits than single-behavior interventions.²⁵ However, it appears that no previous m-Health studies have addressed PA and sleep health simultaneously.^{26,27}

This study aimed to test the efficacy of a novel m-Health intervention to improve PA and sleep health in a randomized waitlist-controlled trial.

METHODS

Prospective registration occurred through the Australian New Zealand Clinical Trials Registry (ACTRN12617000376347). The Human Research Ethics Committee, University of Newcastle, Australia (H-2016-0181) granted ethical approval. Data were collected between June 2017 and February 2018. Trial design, methods, and measures are detailed elsewhere.²⁸

Study Population

In June–August 2017, social media (Facebook) advertisements, lifestyle magazine editorials, and e-Newsletters invited interested individuals to take part. Participants were eligible if they were aged 18–55 years, lived in Australia, reported being insufficiently physically active (<90 minutes/week), and rated their sleep quality as *fairly bad* or *very bad* during screening. Exclusion criteria were a BMI <18.5 or >35, recent pregnancy or childbirth (<12 months), any contraindications for being more physically active or changing sleep behaviors, diagnosed sleep disorders (e.g., chronic insomnia or sleep apnea), hypnotics use, shift work, frequent jetlag-inducing travel, current use of an app/tracker to self-monitor PA or sleep, and no access to an Internet-enabled device (smartphone or tablet). Following completion of baseline assessments, participants ($n=160$) were randomly allocated to the intervention or a waitlist-control group (1:1 ratio). The concealed allocation sequence (numbered opaque envelopes) was generated according to recommended methods for permuted randomization using blocks of 4 and 8.²⁹

Measures

The intervention group ($n=80$) received access to the Balanced app,^{28,30} which provided a platform for personalized goal setting, daily logging with dynamic feedback, and comprehensive educational content for PA (i.e., moderate-to-vigorous intensity PA [MVPA], daily steps, and resistance training [RT]) and for sleep (i.e., bedtimes/waketimes, sleep quality, and sleep hygiene practices). Participants received a series of tool sheets (printed materials via mail, followed up with an electronic copy once a month) that was e-mailed to participants providing guidance on how to set goals, develop action plans, and manage stress. During the 3-month intervention period, participants were e-mailed individualized weekly summary reports based on progress in relation to goals (for both behaviors), as well as weekly SMS including educational content and separate SMS prompts to re-engage with self-monitoring, if necessary (i.e., self-monitoring on <4 days in the last 7 days). The behavior change strategies are described in the context of the theoretic framework (i.e., Social Cognitive Theory) in [Appendix Table 1](#), available online. Following randomization,

participants were mailed a printed handbook including hard copies of the tool sheets and guidance specific to the initial stages of app installation, usage, and troubleshooting, and a pedometer (Yamax SW200) to monitor steps. Waitlist participants ($n=80$) were offered full access to the intervention after the 6-month assessment.

Sociodemographic and behavioral variables were assessed at baseline. The primary endpoint occurred at 3 months and the secondary endpoint at 6 months. Two co-primary and 21 secondary outcomes were assessed at all 3 time points (baseline, 3 months, and 6 months). All data were collected via online survey between June 2017 and February 2018. One of the two co-primary outcomes was weekly minutes of MVPA. This was assessed using the Active Australia Questionnaire,³¹ which is reliable and sensitive to change.^{32,33} Sleep quality was the other co-primary outcome and was assessed using the Pittsburgh Sleep Quality Index (PSQI), which has also shown good reliability and sensitivity to change in intervention studies.^{34,35} Secondary outcomes included the 7 PSQI component scores, RT (frequency and duration/week),³⁶ minutes/day of sitting (Workforce Sitting Questionnaire),³⁷ sleep hygiene practices (Sleep Hygiene Index),³⁸ sleep timing variability (Sleep Timing Questionnaire),³⁹ insomnia symptom severity (Insomnia Severity Index),⁴⁰ daytime sleepiness (Epworth Sleepiness Scale),⁴¹ health-related quality of life (RAND-12),⁴² the energy/fatigue subscale of RAND-36,⁴³ as well as depression, anxiety, and stress symptoms (DASS-21).⁴⁴ To measure engagement, the self-monitoring data participants logged in the Balanced app (defined as logged data on a given day for any of the following: active time, steps, RT, sleep duration, sleep quality, sleep hygiene) were exported to calculate the average number of days for which data were logged and the time to nonusage attrition (defined as ≥ 14 consecutive days of nonengagement at any given point within a person's 84-day intervention period), as previously used.⁴⁵ Participant satisfaction with the app was assessed using the System Usability Scale (scores ranged from 0 to 100, with higher scores indicating greater satisfaction) at 3 months only.⁴⁶ The outcomes assessed and instruments used in this study are described elsewhere.²⁸

Assuming an α of 0.025 (adjusting for use of two co-primary outcomes), power of 0.80, moderate effects at the 3-month primary endpoint (Cohen's d [d]=0.45, mean=88 minutes, SD=194 minutes for PA; and $d=0.65$, mean=1.55, SD=2.41 for sleep quality⁴⁷), and a pre-post correlation of 0.60 (between baseline and 3 months), a total of 60 participants per group were required for PA and 35 for sleep quality. Thus, the larger sample was used for this study. To account for dropout, the sample size was inflated by 25% (calculated as $60/[1-0.25]$), resulting in 80 participants needed per group.⁴⁸

Statistical Analysis

Differences in sample characteristics (e.g., age, gender, baseline levels of PA, and sleep quality) between completers and noncompleters (lost to follow-up) were examined using t -tests (continuous data) and chi-squared tests (categorical data).

Between-group differences at 3 months and 6 months were estimated using generalized linear mixed models, except for PSQI component scores (mixed effects ordered logistic regression). Owing to positive skewness in the data, both RT outcomes were analyzed as dichotomized outcomes (with RT frequency dichotomized as

<2 days/week or ≥ 2 days/week, as per guidelines,⁶ and RT duration dichotomized as <10 minutes/week or ≥ 10 minutes/week). All models included fixed effects for group and time, group $\times$ time interaction, the baseline value of the outcome, and a random intercept for individuals. White–Huber SEs were used if departures of homoscedasticity or normality were observed.⁴⁹ Residual diagnostics informed the specification of family and link functions (Table 2). Models were interpreted using α levels of 0.025 for co-primary outcomes and 0.05 for secondary outcomes.

The impact of missing data was assessed using sensitivity analyses. Missing data were imputed using chained equations and predicted mean matching. Twenty data sets were imputed, using baseline values of the outcome, predictors of missingness, and any variables that predicted a given outcome. The models specified for complete-case analyses were repeated using pooled estimates derived from imputed data sets and coefficients compared for consistency, with little deviation from complete-case analyses indicating robustness of findings.

Secondary analyses using generalized linear mixed models with a binomial logit link function were conducted to examine the proportion of participants meeting guidelines for PA (≥ 150 minutes of MVPA combined with RT on ≥ 2 days/week)⁶ and reporting good sleep (PSQI total score <5).³⁴ Data were analyzed in August 2018 using Stata, version 14.2.

RESULTS

The flow of participants throughout the trial is shown in Figure 1. The baseline sample included 128 women and 32 men, most of whom were middle-aged and overweight. Most were married or in a relationship, highly educated, employed in a professional occupation, and reported having 1 or more chronic conditions. Sociodemographic, health, and behavioral characteristics are provided in Table 1.

Groups did not differ in the proportion of withdrawals ($p=0.181$). Participants who were lost to follow-up reported more severe depression symptoms ($p=0.035$) and lower mental health ($p=0.012$). Complete data (primary outcomes) from 125 participants were available at the 3-month primary endpoint, which corresponds to an overall retention of 78%. Participant retention at the 6-month follow-up was 56%. Dropout rates (defined as formal withdrawal from the trial) were 9% in both groups. Reasons for withdrawal are listed in Figure 1.

Throughout the 84-day intervention period, participants (intervention group only) logged data for at least 1 of the 2 behaviors on an average of 38.2 (SD=30.09) days. Ten percent of participants did not log any data during this period. Nonusage attrition occurred for 89% of participants. The average number of days to nonusage attrition was 32 (SD=25) days. The average number of days on which data were logged and the proportion of participants logging no data did not differ between PA (36 days and 12.5%, respectively) and sleep (37 days and 10%, respectively). Intervention group participants




Figure 1. Participant flow chart.

^an for complete-case analyses at 3 months.

^bn for complete-case analyses at 6 months.

PA, physical activity.

(n=58; assessed at 3 months) reported good usability and acceptance, consistent with a mean system usability score of 70.8 (SD=19.71).⁴⁶

At 3 months, the estimated between-group difference in MVPA was 109 minutes in favor of the intervention group, which was not statistically significant (Table 2), corresponding to a small effect size ($p=0.139$, $d=0.24$). At 6 months, this difference was reduced to 5 minutes ($p=0.952$, $d=0.01$).

The groups showed significant differences in the relative odds of engaging in at least 2 days of RT per week (OR=20.56, 95% CI=2.69, 157.46, $p=0.004$) and ≥ 10 minutes of RT/week (OR=6.71, 95% CI=1.52, 29.65, $p=0.012$), favoring the intervention at 3 months, but these were not maintained at 6 months. Differences in

average sitting time were not statistically significant at either time point (Table 2).

The between-group difference in average sleep quality (PSQI total score) at 3 months was -1.3 points, with medium-sized effect estimates showing significantly better sleep quality in the intervention group ($p=0.009$, $d=0.48$). This difference was slightly attenuated at 6 months and no longer statistically significant ($p=0.040$, $d=0.46$). The intervention group was more likely to report improved subjective sleep quality ($p=0.017$) and sleep onset latency ($p=0.013$) (Table 3). Small-to-medium effect sizes in favor of the intervention were found at 3 months for waketime variability ($p=0.018$, $d=0.40$), sleep hygiene ($p=0.027$, $d=0.40$), and insomnia severity ($p=0.002$, $d=0.56$). At 6 months, significant

Table 1. Baseline Sociodemographic, Health, and Behavioral Characteristics of Study Participants

Characteristics	IG (n=80)	WLG (n=80)
Age, years, mean (SD)	41.1 (9.84)	41.9 (10.07)
Gender, n (%)		
Male	14 (17.50)	18 (22.50)
Female	66 (82.50)	62 (77.50)
BMI, mean (SD)	28.7 (4.64)	27.2 (4.01)
Weight status		
Normal weight (≤ 25.0)	23 (28.75)	28 (35.00)
Overweight (25.1–30)	23 (28.75)	33 (41.25)
Obese (> 30.0)	34 (42.50)	19 (23.75)
Marital status, n (%)		
Single	22 (27.50)	20 (25.00)
Married/de facto	48 (60.00)	45 (56.25)
Divorced/separated	10 (12.50)	12 (15.00)
Widowed/not stated	2 (2.50)	1 (1.25)
Ethnicity, n (%)		
Caucasian	75 (93.75)	71 (88.75)
Asian	3 (3.75)	7 (8.75)
Not stated	2 (2.50)	2 (2.50)
Area of residence ^a		
Major city	56 (70.00)	52 (65.00)
Regional or remote	24 (30.00)	28 (35.00)
Education in years, mean (SD)	16.3 (2.71)	15.8 (2.87)
Occupation, n (%)		
Professional	51 (63.75)	49 (61.25)
White-collar	16 (20.00)	9 (11.25)
Blue-collar	0 (0.00)	4 (5.00)
Not working ^b	13 (16.25)	17 (21.25)
Hours of work, n (%)		
Daytime only	67 (83.75)	62 (77.50)
Other (including not working)	13 (16.25)	18 (22.50)
Annual income, n (%)		
<\$30,000	13 (16.25)	20 (25.00)
\$30,001–\$50,000	13 (16.25)	9 (11.25)
\$50,001–\$70,000	24 (30.00)	17 (21.25)
\$70,001–\$100,000	14 (17.50)	16 (20.00)
$\geq$ \$100,001	9 (11.25)	13 (16.25)
Not stated	7 (8.75)	5 (6.25)
Chronic disease status, n (%)		
None	26 (32.50)	28 (35.00)
1	22 (27.50)	19 (23.75)
≥ 2	32 (40.00)	33 (41.25)
Alcohol consumption, n (%) ^c		
Never	9 (12.00)	16 (21.62)
Monthly or less	23 (30.67)	13 (17.57)
2–4 times/month	18 (24.00)	19 (25.68)
≥ 2 times/week	25 (33.34)	26 (35.13)
Caffeine consumption, n (%) ^c		
2–4 times/month or less	20 (26.66)	13 (17.58)
≥ 2 times/week	55 (73.34)	61 (82.42)

(continued on next page)

Table 1. Baseline Sociodemographic, Health, and Behavioral Characteristics of Study Participants (*continued*)

Characteristics	IG (n=80)	WLG (n=80)
Smoking, n (%) ^c		
Yes	6 (8.00)	5 (6.76)
No	69 (92.00)	69 (93.24)
Physical activity, mean (SD)		
MVPA minutes/week	164.0 (165.45)	191.3 (244.12)
Resistance training frequency, n (%)		
<2 days/week	72 (90.00)	76 (95.00)
≥2 days/week	8 (10.00)	4 (5.00)
Resistance training duration, n (%)		
<10 minutes/week	65 (81.25)	73 (91.25)
≥10 minutes/week	15 (18.75)	7 (8.75)
Sitting minutes, mean (SD)	661.4 (197.71)	671.9 (180.43)
Sleep quality, mean (SD)		
PSQI total score ^d	9.2 (3.07)	9.2 (2.86)
Subjective quality	2.0 (0.56)	1.9 (0.56)
Sleep onset latency	1.8 (1.01)	1.7 (1.06)
Sleep duration	1.0 (0.91)	1.2 (0.96)
Sleep efficiency	1.1 (1.03)	1.1 (1.03)
Sleep disturbances	1.6 (0.51)	1.5 (0.55)
Sleep medication	0.2 (0.53)	0.3 (0.66)
Daytime dysfunction	1.6 (0.67)	1.7 (0.73)
Timing variability ^e		
Bedtime	3.9 (1.96)	3.5 (1.85)
Waketime	2.7 (1.51)	2.4 (1.25)
Sleep hygiene ^f	32.3 (6.72)	32.4 (6.63)
Insomnia severity ^g	12.4 (4.23)	12.7 (3.82)
Daytime sleepiness ^h	8.9 (4.68)	7.9 (4.42)
Symptom severity		
Depression ⁱ	11.3 (7.87)	12.6 (8.84)
Anxiety ^j	6.9 (5.94)	7.1 (6.83)
Stress ^k	15.3 (6.02)	15.4 (7.46)
Quality of life ^l		
Mental health	47.5 (4.96)	47.44 (5.13)
Physical health	44.6 (7.74)	44.19 (8.00)
Energy/fatigue	54.0 (10.74)	55.19 (9.63)

^aArea of remoteness was determined via residential postcode using the Accessibility and Remoteness Index of Australia (ARIA).

^bNot working included participants who were on home duties, retired, or a student.

^cFor the lifestyle items, only 149/160 participants provided valid data.

^dScores range from 0 to 21 with scores >5 indicating poor quality sleep.

^eScores range from 1 to 11 where lower scores indicate less variability in bed- or waketimes.

^fScores range from 13 to 65 (lower scores indicate better sleep hygiene).

^gScores range from 0 to 7 (no clinically significant insomnia), 8–14 (subthreshold insomnia), 15–21 (moderate clinical insomnia), 22–28 (severe clinical insomnia).

^hScores range from 0 to 24 where higher scores indicate higher levels of daytime sleepiness.

ⁱScores for depression symptoms range from 0 to 9 (normal), 10–13 (mild), 14–20 (moderate), 21–27 (severe), 29+ (extremely severe).

^jScores for anxiety symptoms range from 0 to 7 (normal), 8–9 (mild), 10–14 (moderate), 15–19 (severe), 20+ (extremely severe).

^kScores stress symptoms range from 0 to 14 (normal), 15–18 (mild), 19–25 (moderate), 26–33 (severe), 34+ (extremely severe).

^lHigher scores for mental and physical health and the energy/fatigue subscale indicate better quality of life.

IG, intervention group; MVPA, moderate-to-vigorous intensity physical activity; PSQI, Pittsburgh Sleep Quality Index; WLG, waitlist-control group.

Table 2. Marginalized Mean Estimates and Results From Tests of Between-Group Differences for Continuous Outcomes Using Complete Cases

Outcomes	3 months ^a				6 months ^b			
	IG, mean (SD)	WLG, mean (SD)	p-value	d	IG, mean (SD)	WLG, mean (SD)	p-value	d
Co-primary								
MVPA minutes/week ^{c,d}	428.4 (523.41)	319.7 (378.23)	0.139	0.24	405.3 (491.45)	400.2 (497.80)	0.952	0.01
Sleep quality (PSQI) ^e	6.7 (3.04)	8.0 (2.34)	0.009	0.48	6.3 (2.98)	7.5 (2.57)	0.040	0.46
Secondary								
Sitting minutes ^e	612.3 (160.91)	653.7 (202.12)	0.205	0.22	579.7 (187.83)	581.7 (197.27)	0.960	0.01
Bedtime variability ^c	3.6 (1.70)	4.1 (2.10)	0.171	0.24	3.4 (1.46)	4.2 (1.92)	0.023	0.47
Waketime variability ^c	2.5 (1.84)	3.0 (1.95)	0.018	0.40	2.6 (1.06)	3.0 (1.92)	0.236	0.22
Sleep hygiene ^e	30.0 (4.27)	31.6 (3.98)	0.027	0.40	30.6 (4.33)	32.8 (6.02)	0.048	0.42
Insomnia severity ^e	9.3 (3.80)	11.3 (3.50)	0.002	0.56	8.5 (4.23)	11.4 (4.11)	0.002	0.69
Daytime sleepiness	7.1 (3.44)	8.0 (3.31)	0.103	0.29	5.7 (2.92)	8.4 (3.98)	<0.001	0.74
Depression symptoms ^{c,d}	10.6 (7.62)	12.6 (7.97)	0.120	0.26	10.9 (8.01)	13.3 (9.49)	0.190	0.27
Anxiety symptoms ^d	6.4 (3.65)	7.5 (5.04)	0.148	0.25	5.9 (3.54)	8.9 (4.70)	0.003	0.57
Stress symptoms ^d	13.6 (4.20)	15.4 (4.97)	0.032	0.38	13.0 (5.75)	16.3 (5.24)	0.006	0.62
Mental health ^e	44.4 (6.81)	44.9 (7.12)	0.689	0.07	47.6 (5.53)	45.0 (8.12)	0.071	0.37
Physical health ^e	47.6 (5.21)	46.8 (5.54)	0.400	0.15	46.5 (4.95)	47.3 (5.14)	0.467	0.16
Energy/fatigue ^e	52.4 (8.29)	53.4 (11.08)	0.541	0.11	54.7 (9.90)	51.1 (11.49)	0.118	0.33

Note: Boldface indicates statistical significance at $p < 0.025$ for co-primary outcomes and $p < 0.05$ for secondary outcomes.

Analyzed using a GLMM with gaussian distribution and log link function; all GLMM were fitted using the robust variance estimator.

Cohen's d (the magnitude of effects is interpreted as small [0.2], medium [0.5], or large [0.80]).

^aAt 3 months, 125 observations ($n=59$ in IG; $n=66$ in WLG) were available for analyses of MVPA, sleep quality, mental health, physical health and energy/fatigue, and 124 observations ($n=59$ in IG; $n=65$ in WLG) were available for analyses of all other outcomes.

^bAt 6 months, 89 observations ($n=35$ in IG; $n=54$ in WLG) were available for analyses of MVPA, sleep quality, mental health, physical health and energy/fatigue, and 88 observations ($n=34$ in IG; $n=53$ in WLG) were available for analyses of all other outcomes.

^cAnalyzed using a GLMM with gamma distribution and log link function.

^dA small positive constant (+1) was added to minutes of MVPA and the DASS-21 scores for the purpose of data analysis and these outcomes are reported with this constant included.

^eAnalyzed using a GLMM with gaussian distribution and identity link function.

DASS, Depression Anxiety and Stress Scales; GLMM, generalized linear mixed model; IG, intervention group; MVPA, moderate-to-vigorous intensity physical activity; PSQI, Pittsburgh Sleep Quality Index; WLG, waitlist-control group.

differences in favor of the intervention group were maintained for sleep hygiene ($p=0.048$, $d=0.42$) and insomnia severity ($p=0.002$, $d=0.69$), with an increase in the magnitude of differences for insomnia severity. Additional significant differences at 6 months, which were not statistically significant at 3 months, were observed for bedtime variability ($p=0.023$, $d=0.47$), daytime sleepiness ($p<0.001$, $d=0.74$), and daytime dysfunction (OR=0.28, 95% CI=0.08, 0.94, $p=0.039$) (Tables 2 and 3).

No significant differences were observed for mental or physical health-related quality of life, energy/fatigue levels, or for depression symptoms, at either time point (all $p>0.05$). The intervention group reported significantly lower stress symptom severity relative to waitlist controls at 3 months ($p=0.032$, $d=0.38$) with an additional increase in magnitude at 6 months ($p=0.006$, $d=0.62$). Further, differences in anxiety symptoms at 6 months were statistically significant, in favor of the intervention group ($p=0.003$, $d=0.57$).

Results from analyses using imputed data are provided in Appendix Tables 2 and 3, available online, showing robustness of findings at 3 months for all

outcomes, except for stress symptoms, which was no longer statistically significant ($p=0.078$). Group differences in the co-primary outcome of sleep quality (PSQI total score) were statistically significant at 6 months based on imputed data ($p=0.015$). Differences in bedtime variability, sleep hygiene, and anxiety measured at 6 months, which were statistically significant based on complete-case analysis, no longer reached statistical significance (all $p>0.05$).

Secondary analyses using complete cases showed, at 3 months, participants in the intervention group were significantly more likely to meet aerobic exercise and RT guidelines,⁶ relative to participants in the waitlist-control group (OR=16.32, 95% CI=2.24, 119.00, $p=0.004$). This difference was not maintained at 6 months (OR=1.05, 95% CI=0.08, 13.13, $p=0.970$) (Appendix Figure 1A, available online). The proportion of participants reporting good sleep (Appendix Figure 1B, available online) was significantly higher in the intervention group relative to the control group at 3 months (OR=13.13, 95% CI=2.94, 58.64, $p=0.001$), but not at 6 months (OR=4.47, 95% CI=0.96, 20.79, $p=0.056$). Results from analyses

Table 3. ORs, 95% CIs, and Results From Tests of Between-Group Differences for Categorical Outcomes Using Complete Cases

Variable	3 months ^a			6 months ^b		
	OR	SE (95% CI)	p-value	OR	SE (95% CI)	p-value
Resistance training on ≥ 2 days/week ^c	20.56	21.36 (2.69, 157.46)	0.004	1.05	1.35 (0.08, 13.13)	0.970
Resistance training for ≥ 10 minutes/week ^c	6.71	5.09 (1.52, 29.65)	0.012	1.84	1.72 (0.30, 11.43)	0.511
Subjective sleep quality ^{d,e}	0.36	0.15 (0.16, 0.84)	0.017	0.89	0.50 (0.29, 2.68)	0.832
Sleep onset latency ^{d,e}	0.27	0.14 (0.10, 0.76)	0.013	0.48	0.32 (0.13, 1.74)	0.263
Sleep duration ^{d,e}	0.49	0.25 (0.18, 1.32)	0.158	0.45	0.29 (0.13, 1.56)	0.208
Sleep efficiency ^{d,e}	0.52	0.23 (0.21, 1.25)	0.107	0.64	0.33 (0.23, 1.77)	0.392
Sleep disturbances ^{d,e}	0.34	0.22 (0.10, 1.18)	0.089	0.25	0.20 (0.05, 1.22)	0.086
Sleep medication use ^{d,e}	0.47	0.42 (0.08, 2.72)	0.402	0.07	0.10 (0.00, 1.19)	0.066
Daytime dysfunction ^{d,e}	0.80	0.41 (0.29, 2.19)	0.665	0.28	0.17 (0.08, 0.94)	0.039

Note: Boldface indicates statistical significance at $p < 0.05$.

^aAt 3 months, 125 observations ($n=59$ in IG; $n=66$ in WLG) were available for analyses.

^bAt 6 months, 89 observations ($n=35$ in IG; $n=54$ in WLG) were available for analyses.

^cORs for resistance training frequency and duration were calculated using GLMMs with binomial distribution, logit link function and robust variance estimator.

^dEstimates for the seven PSQI composites represent proportional ORs, robust SEs, and 95% CIs based on mixed effects ordered logistic regression that tested between-group differences in the likelihood of shifting to another level of the variable (lower PSQI composites indicate better sleep quality, thus OR < 1 indicates the intervention group was less likely to move up a level or report worse outcomes for the composites).

^eA small positive constant (+1) was added to PSQI component scores for the purpose of data analysis.

GLMM, generalized linear mixed model; IG, intervention group; PSQI, Pittsburgh Sleep Quality Index; WLG, waitlist-control group.

using imputed data were consistent with these results, except the proportion of participants reporting good sleep quality at 6 months was significantly higher in the intervention group relative to the control group at 6 months (OR=4.05, 95% CI=1.11, 14.75, $p=0.034$).

DISCUSSION

The Synergy Study was a combined behavior m-Health intervention that improved sleep quality and a range of secondary outcomes, including RT, sleep time variability, sleep hygiene, subjective sleep quality, sleep onset latency, insomnia severity, and symptoms of stress and anxiety. Moreover, participants were more likely to meet guidelines for aerobic PA and RT and report good sleep quality after the 3-month intervention.

Although group differences in MVPA were not statistically significant at either endpoint, several promising changes were observed. At 3 months, the intervention group engaged in an additional 109 minutes/week of MVPA, relative to waitlist control, which is encouraging given that even small improvements in MVPA are beneficial for long-term health.⁵⁰ The effect size of this non-significant difference ($d=0.24$) was consistent with previous meta-analyses of m-Health PA interventions.⁴⁸ The lack of statistically significant between-group differences in MVPA may have been because of the large variation in activity at all time points and the increased activity reported in the control group, which is commonly observed.⁵¹ Secondary analyses showed a significantly greater proportion of intervention group

participants (37.3%) met PA guidelines (≥ 150 minutes of MVPA and RT on ≥ 2 days/week). These improvements may be attributable to the detailed strategies provided for both MVPA and RT and thus supports their use in multibehavior interventions. This finding is relevant from a public health perspective, given that most adults do not engage in RT,⁵² and only approximately 15% meet guidelines for both MVPA and RT, although this would confer significant reductions for morbidity and mortality.^{53–55} However, this outcome should be interpreted with caution, as the analyses of meeting PA guidelines were exploratory. Also, although both minutes of MVPA and frequency of RT increased, the higher proportion of participants meeting PA guidelines at 3 months appeared to be largely driven by an increased frequency of RT.

A unique contribution of this study is that it assessed sleep quality at 3 and 6 months, which is longer than the intervention periods and follow-up intervals of many sleep interventions.⁵⁶ The magnitude of the between-group difference in favor of the intervention group at 3 months is consistent with findings from trials in subclinical population groups.¹⁹ Although sleep quality continued to improve at 6 months, the difference between groups was no longer statistically significant for complete-case analysis but was for analysis based on multiple imputation (at $\alpha=0.025$). Given that the observed effect sizes at 3 months ($d=0.48$) and 6 months ($d=0.46$) were consistent, this may have been because of improvements in both groups and the lack of power at the 6-month time point. The intervention was designed for a

population with poor sleep quality but without a diagnosed sleep disorder. Baseline values of the PSQI, Insomnia Severity Index, and the Epworth Sleepiness Index suggest that most participants did have poor sleep quality but did not have a clinical sleep disorder (e.g., above clinical threshold for insomnia or sleep apnea). A smaller margin for improvement relative to that typically seen in clinical population groups was a function of studying a subclinical group with lower baseline symptom severity. Accordingly, the shift in scores observed in the Synergy Study was deemed satisfactory, especially given that the effect size ($d=0.48$) associated with between-group differences in sleep quality was comparable to that reported in a systematic review of Internet-delivered cognitive behavioral therapy for insomnia ($d=0.49$).⁵⁶ This is important considering the high percentage of the population with poor sleep quality, but without a clinical sleep disorder and limited access to practitioner-based treatment.^{57,58} PSQI scores <5 indicate remission of sleep problems,⁵⁹ and despite the magnitude of improvement observed in the intervention group, average scores remained >5 , which is consistent with most studies in subclinical populations.¹⁹ However, the intervention group had 32% more participants reporting good sleep quality (PSQI scores <5) (Appendix Figure 1B, available online) indicating the intervention may have considerable public health utility. The hypothesized synergistic relationship between PA and sleep was not examined in this study. However, changes in MVPA and sleep quality were lower in magnitude than anticipated. It is possible the magnitude of change in PA was not large enough to leverage larger increases in sleep quality and vice versa, or the study duration was too short to detect this.

The intervention group further exhibited significantly better sleep hygiene practices and improvements in subjective sleep quality and sleep onset latency. More-pronounced improvements in sleep onset latency for the intervention group were likely a result of adherence to and improvements in sleep hygiene practices, which were targeted specifically in the Synergy Study. Moreover, PA is associated with reduced sleep onset latency,⁶⁰ and the large amount of additional MVPA reported in the intervention group (adjusted group difference=109 minutes, $d=0.24$), albeit not statistically significant, combined with the greater increase in RT may have contributed to improvements in sleep health indicators (i.e., sleep onset latency), which is consistent with the literature.^{61,62} These improvements, combined with those seen for insomnia severity, which capture clinically relevant characteristics of poor sleep health, support the overall finding that the intervention was efficacious in improving sleep.

The lack of group differences in health-related quality of life might indicate that changes in these parameters take longer than 3–6 months to manifest and may be of small magnitude in a population with nonacute conditions.⁶³ Long-term follow-up assessments therefore are warranted. The Synergy Study provided detailed stress management resources but none specifically for depression or anxiety, which may explain the significantly greater improvements in stress symptoms observed in the intervention group. Given that high stress levels are associated with engagement in unhealthy behaviors and poorer sleep quality,⁶⁴ facilitating stress management to reduce symptoms is important for multiple behavior interventions, particularly those targeting sleep.

App usability ratings were fair, time to nonusage attrition was 32 days, and 89% of participants suffered nonusage attrition. Although there are few app usage data available from multiple behavior interventions, the proportion of participants making at least one entry (90%) is similar to that observed in single-behavior m-Health programs.⁶⁵ However, there is no evidence that defines the minimum amount of app usage needed for behavior change to occur and whether continuous usage differs from intermittent usage with regard to the magnitude of behavior change it confers. Time to nonusage attrition in the current study (32 days) appears to indicate moderate usage in comparison with other studies for which time to nonusage attrition ranged from 1.5 weeks to 25 weeks.⁶⁶ These results suggest that targeting 2 behaviors simultaneously does not adversely impact app usage rates. Despite fair participant ratings for app usability and time to nonusage attrition, almost all participants in the intervention group still suffered nonusage attrition during the intervention period. Several devices exist that allow automated self-monitoring of PA and sleep.⁶⁷ However, manual data entry was used for pragmatic reasons (e.g., cost) and this may have contributed to nonusage attrition, although it is unknown which method (manual or automated) is optimal for use in behavior change interventions. Moreover, it is possible that some participants reached personal goals relating to PA and sleep sooner than others or lost motivation over time. Furthermore, participants may have engaged in PA and sleep hygiene practice more frequently than indicated by the completed logs, possibly because of not feeling any need to keep track, or because of time restrictions. This is in line with findings from a study indicating that participants only log approximately 60% of their objectively measured daily activity.²⁴

To the authors' knowledge, this was the first trial of its kind to target both PA and sleep with an m-Health intervention using an RCT design. Trial strengths included its potential for wide reach, given the remote delivery,

which made it accessible for those living outside metropolitan areas, as well as the personalized approach, which reinforced personally meaningful goals.

Limitations

Several factors reduced the power to detect significant between-group differences in MVPA. Despite requiring that participants report <90 minutes/week to be eligible, 41.25% reported doing >150 minutes at baseline. Discrepancies between the levels of PA participants reported at the eligibility screening and those reported at baseline may have been because of the different methods of assessment used (e.g., eligibility survey: single item versus baseline survey: multiple items). This may indicate that using a single-item measure has limited usefulness as a screening tool for interventions.^{68,69} Self-report measures of PA are subject to recall bias, and there was large variation in minutes of PA at baseline and follow-up in both groups. In addition, the waitlist-control group reported substantial increases in activity, which reduced the difference between groups. Given that participants volunteered to take part, this could have been because of high levels of readiness to make changes to PA and sleep behaviors upon enrollment in the study, as seen in previous trials.^{51,70} However, participants' readiness to change behavior was not assessed in this study. Moreover, the study was powered based on effect sizes that assumed a synergistic effect between PA and sleep and this may have been overestimated. The use of objective measures (e.g., accelerometry and polysomnography) however, was not feasible in this trial. Moreover, the self-report sleep measure (PSQI) was likely able to better capture the restorative effects of improved sleep, which is not possible using objective measures.⁷¹ A number of secondary outcomes were examined, and this may have increased the risk of Type 1 errors. The classification of RT duration may have been somewhat arbitrary, and it is unknown if a minimum duration of 10 minutes confers a health benefit. Finally, it is possible that access to an Internet-enabled device as an eligibility criterion has reduced the representativeness of the sample. However, the associated risk of bias may be minimal, given the widespread ownership of smartphones in Australia.⁷²

CONCLUSIONS

This remotely delivered intervention produced short-term improvements in RT and short- and medium-term improvements in sleep health. Some of the group differences seen at 3 months were not sustained at the 6-month follow-up. The capacity of m-Health interventions to foster long-term engagement and sustainable changes in behavior remains to be determined, but

m-Health interventions may be useful at least in the short-term to promote a combination of healthy habits.

ACKNOWLEDGMENTS

This research was in part supported by a Future Leader Fellowship (ID 100029) and Vanguard Grant (ID 100629) from the National Heart Foundation of Australia, awarded to Mitch J. Duncan. Mitch J. Duncan is supported by a Career Development Fellowship from the National Health and Medical Research Council (APP1141606). Corneal Vandelanotte is supported by a Future Leader Fellowship (ID 100427) from the National Heart Foundation of Australia.

No financial disclosures were reported by the authors of this paper.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2019.05.009>.

REFERENCES

1. WHO. Global strategy on diet, physical activity and health. Physical inactivity: A global public health problem. www.who.int/dietphysicalactivity/factsheet_inactivity/en/. Accessed April 1, 2016.
2. Hoyos C, Glozier N, Marshall NS. Recent evidence on worldwide trends on sleep duration. *Curr Sleep Med Rep*. 2015;1(4):195–204. <https://doi.org/10.1007/s40675-015-0024-x>.
3. Buman MP, Phillips BA, Youngstedt SD, Kline CE, Hirshkowitz M. Does nighttime exercise really disturb sleep? Results from the 2013 National Sleep Foundation Sleep in America Poll. *Sleep Med*. 2014;15(7):755–761. <https://doi.org/10.1016/j.sleep.2014.01.008>.
4. Duncan MJ, Kline CE, Rebar AL, Vandelanotte C, Short CE. Greater bed- and wake-time variability is associated with less healthy lifestyle behaviors: a cross-sectional study. *Z Gesundh Wiss*. 2016;24(1):31–40. <https://doi.org/10.1007/s10389-015-0693-4>.
5. Adams RJ, Appleton SL, Taylor AW, et al. Sleep health of Australian adults in 2016: results of the 2016 Sleep Health Foundation national survey. *Sleep Health*. 2017;3(1):35–42. <https://doi.org/10.1016/j.sleh.2016.11.005>.
6. The Department of Health. Australia's Physical Activity and Sedentary Behaviour Guidelines for Adults (18–64 years). www.webcitation.org/6pseaziKS. Published 2014. Accessed February 15, 2017.
7. Buysse DJ. Sleep health: can we define it? Does it matter? *Sleep*. 2014;37(1):9–17. <https://doi.org/10.5665/sleep.3298>.
8. Shiroma EJ, Lee IM. Physical activity and cardiovascular health: lessons learned from epidemiological studies across age, gender, and race/ethnicity. *Circulation*. 2010;122(7):743–752. <https://doi.org/10.1161/circulationaha.109.914721>.
9. Aune D, Norat T, Leitzmann M, Tonstad S, Vatten LJ. Physical activity and the risk of type 2 diabetes: a systematic review and dose–response meta-analysis. *Eur J Epidemiol*. 2015;30(7):529–542. <https://doi.org/10.1007/s10654-015-0056-z>.
10. Shan Z, Ma H, Xie M, et al. Sleep duration and risk of type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care*. 2015;38(3):529–537. <https://doi.org/10.2337/dc14-2073>.
11. Hoevenaer-Blom MP, Spijkerman AM, Kromhout D, van den Berg JF, Verschuren WM. Sleep duration and sleep quality in relation to 12-year cardiovascular disease incidence: the MORGEN study. *Sleep*. 2011;34(11):1487–1492. <https://doi.org/10.5665/sleep.1382>.

12. Deloitte Access Economics. Asleep on the Job: Costs of Inadequate Sleep in Australia. Sleep Health Foundation. www.sleephealthfoundation.org.au/news/special-reports/asleep-on-the-job-costs-of-inadequate-sleep-in-australia.html. Accessed January 4, 2018.
13. Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet*. 2016;388(10051):1311–1324. [https://doi.org/10.1016/s0140-6736\(16\)30383-x](https://doi.org/10.1016/s0140-6736(16)30383-x).
14. Kruk J. Health and economic costs of physical inactivity. *Asian Pac J Cancer Prev*. 2014;15(18):7499–7503. <https://doi.org/10.7314/apjcp.2014.15.18.7499>.
15. Rayward AT, Duncan MJ, Brown WJ, Plotnikoff RC, Burton NW. A cross-sectional cluster analysis of the combined association of physical activity and sleep with sociodemographic and health characteristics in mid-aged and older adults. *Maturitas*. 2017;102:56–61. <https://doi.org/10.1016/j.maturitas.2017.05.013>.
16. Price M, Yuen EK, Goetter EM, et al. mHealth: a mechanism to deliver more accessible, more effective mental health care. *Clin Psychol Psychother*. 2014;21(5):427–436. <https://doi.org/10.1002/cpp.1855>.
17. Muller AM, Alley S, Schoeppe S, Vandelanotte C. The effectiveness of e- & mHealth interventions to promote physical activity and healthy diets in developing countries: a systematic review. *Int J Behav Nutr Phys Act*. 2016;13:109. <https://doi.org/10.1186/s12966-016-0434-2>.
18. Ye YY, Chen NK, Chen J, et al. Internet-based cognitive-behavioural therapy for insomnia (ICBT-i): a meta-analysis of randomised controlled trials. *BMJ Open*. 2016;6(11):e010707. <https://doi.org/10.1136/bmjopen-2015-010707>.
19. Murawski B, Wade L, Plotnikoff RC, Lubans DR, Duncan MJ. A systematic review and meta-analysis of cognitive and behavioral interventions to improve sleep health in adults without sleep disorders. *Sleep Med Rev*. 2018;40:160–169. <https://doi.org/10.1016/j.smrv.2017.12.003>.
20. McEwan D, Beauchamp MR, Kouvousis C, et al. Examining the active ingredients of physical activity interventions underpinned by theory versus no stated theory: a meta-analysis. *Health Psychol Rev*. 2018;13(1):1–17. <https://doi.org/10.1080/17437199.2018.1547120>.
21. McSharry J, Olander EK, French DP. Do single and multiple behavior change interventions contain different behavior change techniques? A comparison of interventions targeting physical activity in obese populations. *Health Psychol*. 2015;34(9):960–965. <https://doi.org/10.1037/hea0000185>.
22. Kline CE. The bidirectional relationship between exercise and sleep: implications for exercise adherence and sleep improvement. *Am J Lifestyle Med*. 2014;8(6):375–379. <https://doi.org/10.1177/1559827614544437>.
23. Dolezal BA, Neufeld EV, Boland DM, Martin JL, Cooper CB. Interrelationship between sleep and exercise: a systematic review. *Adv Prev Med*. 2017;2017:1364387. <https://doi.org/10.1155/2017/1364387>.
24. Buman MP, Epstein DR, Gutierrez M, et al. BeWell24: development and process evaluation of a smartphone “app” to improve sleep, sedentary, and active behaviors in U.S. Veterans with increased metabolic risk. *Transl Behav Med*. 2016;6(3):438–448. <https://doi.org/10.1007/s13142-015-0359-3>.
25. Pepin V, Boucetta S, Chan-Thim E, et al. The role of sleep and physical activity on the risk for cardiovascular disease. *Curr Cardiovasc Risk Rep*. 2014;8(12):1–10. <https://doi.org/10.1007/s12170-014-0413-6>.
26. James E, Freund M, Booth A, et al. Comparative efficacy of simultaneous versus sequential multiple health behavior change interventions among adults: a systematic review of randomised trials. *Prev Med*. 2016;89:211–223. <https://doi.org/10.1016/j.ypmed.2016.06.012>.
27. Meader N, King K, Wright K, et al. Multiple risk behavior interventions: meta-analyses of RCTs. *Am J Prev Med*. 2017;53(1):e19–e30. <https://doi.org/10.1016/j.amepre.2017.01.032>.
28. Murawski B, Plotnikoff RC, Rayward AT, et al. Randomised controlled trial using a theory-based m-health intervention to improve physical activity and sleep health in adults: the Synergy Study protocol. *BMJ Open*. 2018;8(2):e018997. <https://doi.org/10.1136/bmjopen-2017-018997>.
29. Doig GS, Simpson F. Randomization and allocation concealment: a practical guide for researchers. *J Crit Care*. 2005;20(2):187–191. <https://doi.org/10.1016/j.jcrc.2005.04.005>.
30. Duncan MJ, Vandelanotte C, Trost SG, et al. Balanced: a randomised trial examining the efficacy of two self-monitoring methods for an app-based multi-behaviour intervention to improve physical activity, sitting and sleep in adults. *BMC Public Health*. 2016;16:670. <https://doi.org/10.1186/s12889-016-3256-x>.
31. Australian Institute of Health and Welfare. *The Active Australia Survey: A Guide and Manual for Implementation, Analysis and Reporting*. Canberra: AIHW; 2003.
32. Reeves MM, Marshall AL, Owen N, Winkler EA, Eakin EG. Measuring physical activity change in broad-reach intervention trials. *J Phys Act Health*. 2010;7(2):194–202. <https://doi.org/10.1123/jpah.7.2.194>.
33. Brown W, Burton N, Marshall A, Miller Y. Reliability and validity of a modified self-administered version of the Active Australia physical 2 activity survey in a sample of mid-age women. *Aust N Z J Public Health*. 2008;32(6):535–541. <https://doi.org/10.1111/j.1753-6405.2008.00305.x>.
34. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4).
35. Backhaus J, Junghanns K, Broocks A, Riemann D, Hohagen F. Test–retest reliability and validity of the Pittsburgh Sleep Quality Index in primary insomnia. *J Psychosom Res*. 2002;53(3):737–740. [https://doi.org/10.1016/s0022-3999\(02\)00330-6](https://doi.org/10.1016/s0022-3999(02)00330-6).
36. Bampton EA, Johnson ST, Vallance JK. Profiles of resistance training behavior and sedentary time among older adults: associations with health-related quality of life and psychosocial health. *Prev Med Rep*. 2015;2:773–776. <https://doi.org/10.1016/j.pmedr.2015.08.017>.
37. Chau JY, van der Ploeg HP, Dunn S, Kurko J, Bauman AE. A tool for measuring workers’ sitting time by domain: the Workforce Sitting Questionnaire. *Br J Sports Med*. 2011;45(15):1216–1222. <https://doi.org/10.1136/bjsports-2011-090214>.
38. Mastin DF, Bryson J, Corwyn R. Assessment of sleep hygiene using the Sleep Hygiene Index. *J Behav Med*. 2006;29(3):223–227. <https://doi.org/10.1007/s10865-006-9047-6>.
39. Monk TH, Buysse DJ, Kennedy KS, et al. Measuring sleep habits without using a diary: the sleep timing questionnaire. *Sleep*. 2003;26(2):208–212. <https://doi.org/10.1093/sleep/26.2.208>.
40. Morin CM, Belleville G, Belanger L, Ivers H. The Insomnia Severity Index: psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep*. 2011;34(5):601–608. <https://doi.org/10.1093/sleep/34.5.601>.
41. Johns MW. A new method for measuring daytime sleepiness: the Epworth Sleepiness Scale. *Sleep*. 1991;14(6):540–545. <https://doi.org/10.1093/sleep/14.6.540>.
42. Ware J Jr., Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996;34(3):220–233. <https://doi.org/10.1097/00005650-199603000-00003>.
43. Hays RD, Morales LS. The RAND-36 measure of health-related quality of life. *Ann Med*. 2001;33(5):350–357. <https://doi.org/10.3109/07853890109002089>.
44. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol*. 2005;44(2):227–239. <https://doi.org/10.1348/014466505x29657>.
45. Kolt GS, Rosenkranz RR, Vandelanotte C, et al. Using Web 2.0 applications to promote health-related physical activity: findings from the WALK 2.0 randomised controlled trial. *Br J Sports Med*. 2017;51(19):1433–1440. <https://doi.org/10.1136/bjsports-2016-096890>.

46. SUS BJ. A quick and dirty usability scale. In: Jordan PW, Thomas B, Weerdmeester BA, McClelland AL, eds. *Usability Evaluation in Industry*. Vol. 189. London: Taylor & Francis; 1996:194.
47. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Mahwah, NJ: L. Erlbaum Associates; 1988.
48. Davies CA, Spence JC, Vandelanotte C, Caperchione CM, Mummery WK. Meta-analysis of internet-delivered interventions to increase physical activity levels. *Int J Behav Nutr Phys Act*. 2012;9:52. <https://doi.org/10.1186/1479-5868-9-52>.
49. White H. A heteroskedasticity-consistent covariance-matrix estimator and a direct test for heteroskedasticity. *Econometrica*. 1980;48(4):817–838. <https://doi.org/10.2307/1912934>.
50. Warburton DE, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *CMAJ*. 2006;174(6):801–809. <https://doi.org/10.1503/cmaj.051351>.
51. Waters L, Reeves M, Fjeldsoe B, Eakin E. Control group improvements in physical activity intervention trials and possible explanatory factors: a systematic review. *J Phys Act Health*. 2012;9(6):884–895. <https://doi.org/10.1123/jpah.9.6.884>.
52. Loustalot F, Carlson SA, Kruger J, Buchner DM, Fulton JE. Muscle-strengthening activities and participation among adults in the United States. *Res Q Exerc Sport*. 2013;84(1):30–38. <https://doi.org/10.1080/02701367.2013.762289>.
53. Bennie JA, Pedisic Z, van Uffelen JGZ, et al. The descriptive epidemiology of total physical activity, muscle-strengthening exercises and sedentary behaviour among Australian adults - results from the National Nutrition and Physical Activity Survey. *BMC Public Health*. 2016;16:73. <https://doi.org/10.1186/s12889-016-2736-3>.
54. Ihalainen JK, Schumann M, Eklund D, et al. Combined aerobic and resistance training decreases inflammation markers in healthy men. *Scand J Med Sci Sports*. 2018;28(1):40–47. <https://doi.org/10.1111/sms.12906>.
55. Dankel SJ, Loenneke JP, Loprinzi PD. Participation in muscle-strengthening activities as an alternative method for the prevention of multimorbidity. *Prev Med*. 2015;81:54–57. <https://doi.org/10.1016/j.ypmed.2015.08.002>.
56. Zachariae R, Lyby MS, Ritterband LM, O'Toole MS. Efficacy of internet-delivered cognitive-behavioral therapy for insomnia: a systematic review and meta-analysis of randomized controlled trials. *Sleep Med Rev*. 2016;30:1–10. <https://doi.org/10.1016/j.smrv.2015.10.004>.
57. Cunningham D, Junge MF, Fernando AT. Insomnia: prevalence, consequences and effective treatment. *Med J Aust*. 2013;199(8):S36–S40. <https://doi.org/10.5694/mja13.10718>.
58. Espie CA. “Stepped care”: a health technology solution for delivering cognitive behavioral therapy as a first line insomnia treatment. *Sleep*. 2009;32(12):1549–1558. <https://doi.org/10.1093/sleep/32.12.1549>.
59. Buysse DJ, Germain A, Moul DE, et al. Efficacy of brief behavioral treatment for chronic insomnia in older adults. *Arch Intern Med*. 2011;171(10):887–895. <https://doi.org/10.1001/archinternmed.2010.535>.
60. 2018 Physical Activity Guidelines Advisory Committee. *Physical Activity Guidelines Advisory Committee Scientific Report*. Washington, DC: HHS; 2018.
61. Kovacevic A, Mavros Y, Heisz JJ, Fiatarone Singh MA. The effect of resistance exercise on sleep: a systematic review of randomized controlled trials. *Sleep Med Rev*. 2018;39:52–68. <https://doi.org/10.1016/j.smrv.2017.07.002>.
62. Loprinzi PD, Loenneke JP. Engagement in muscular strengthening activities is associated with better sleep. *Prev Med Rep*. 2015;2:927–929. <https://doi.org/10.1016/j.pmedr.2015.10.013>.
63. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med Care*. 2003;41(5):582–592. <https://doi.org/10.1097/01.mlr.0000062554.74615.4c>.
64. Ng DM, Jeffery RW. Relationships between perceived stress and health behaviors in a sample of working adults. *Health Psychol*. 2003;22(6):638–642. <https://doi.org/10.1037/0278-6133.22.6.638>.
65. Guertler D, Vandelanotte C, Kirwan M, Duncan MJ. Engagement and nonusage attrition with a free physical activity promotion program: the case of 10,000 Steps Australia. *J Med Internet Res*. 2015;17(7):e176. <https://doi.org/10.2196/jmir.4339>.
66. Vandelanotte C, Duncan MJ, Kolt GS, et al. More real-world trials are needed to establish if web-based physical activity interventions are effective. *Br J Sports Med*. In press. Online July 3, 2018. <https://doi.org/10.1136/bjsports-2018-099437>.
67. Duncan M, Murawski B, Short CE, et al. Activity trackers implement different behavior change techniques for activity, sleep, and sedentary behaviors. *Interact J Med Res*. 2017;6(2):e13. <https://doi.org/10.2196/ijmr.6685>.
68. Caperchione CM, Duncan MJ, Rosenkranz RR, et al. Recruitment, screening, and baseline participant characteristics in the WALK 2.0 study: a randomized controlled trial using web 2.0 applications to promote physical activity. *Contemp Clin Trials Commun*. 2016;2:25–33. <https://doi.org/10.1016/j.conctc.2015.12.004>.
69. Vandelanotte C, Reeves MM, Brug J, De Bourdeaudhuij I. A randomized trial of sequential and simultaneous multiple behavior change interventions for physical activity and fat intake. *Prev Med*. 2008;46(3):232–237. <https://doi.org/10.1016/j.ypmed.2007.07.008>.
70. Blanton S, Morris DM, Prettyman MG, et al. Lessons learned in participant recruitment and retention: the EXCITE trial. *Phys Ther*. 2006;86(11):1520–1533. <https://doi.org/10.2522/ptj.20060091>.
71. Rogers LQ, Courneya KS, Oster RA, et al. Physical activity and sleep quality in breast cancer survivors: a randomized trial. *Med Sci Sports Exerc*. 2017;49(10):2009–2015. <https://doi.org/10.1249/MSS.0000000000001327>.
72. Australian Communications and Media Authority. *Access to Internet broadband and mobile phones family households*. Media and Communications in Australian Families series. 2008;3. www.acma.gov.au/-/media/Research-and-Analysis/Report/pdf/no3_access_to_internet_broadband_and_mobile_phones_family_households-pdf.pdf?la=en. Accessed May 9, 2019.