



Review article

Enhanced recovery after hepatectomy: A systematic review

Pauline Rouxel, Helene Beloeil *



Department of Anaesthesiology and Critical Care Medicine, CHU de Rennes, Université de Rennes, Inserm NumCan, CIC 1414, 35000 Rennes, France

ARTICLE INFO

Article history:

Received 13 February 2018
 Received in revised form 12 April 2018
 Accepted 2 May 2018
 Available online 26 May 2018

Keywords:

ERP
 Hepatectomy
 Liver
 Fast-track

ABSTRACT

Background: Hepatectomy is a surgery with high postoperative complication rates. Enhanced recovery after surgery (ERAS) clinical pathways in liver surgery have been studied and may become a standard of care. However, few specific recommendations have been published so far.

Objective: The aim of this study was to assess the efficacy and safety of the enhanced recovery program in liver surgery.

Methods: Randomised controlled trials (RCTs) comparing ERAS group with traditional care published between 2007 and 2017 were included in this review. The outcomes were length of stay (LoS), complications, mortality and readmission rate for all liver surgeries except transplantation.

Results: Five hundred and twenty-four patients randomised in 4 RCTs were analysed. Two hundred and fifty-four patients were in ERAS group and 270 patients in traditional care (TC) group. Two studies compared cares in laparoscopic surgery and 2 in open surgery. Postoperative LoS was significantly lower in the ERAS group whereas readmission and mortality rate were similar. ERAS group had also significant lower complication rate in 2 studies of the 4. The complication rate in the 2 other studies was similar.

Conclusion: ERAS protocols in liver surgery appeared to be safe and effective. Recent recommendations from the ERAS group in liver surgery are the only ones published so far. Other studies evaluating ERAS components in liver surgery and recommendations from scientific societies are needed to spread this clinical care pathway.

© 2018 Société française d'anesthésie et de réanimation (Sfar). Published by Elsevier Masson SAS. All rights reserved.

Contents

1. Introduction	30
2. Methods	30
2.1. Search strategy and study selection	30
2.2. Inclusion and exclusion criteria	30
2.3. Data extraction	30
2.4. Quality assessment	30
3. Results	30
3.1. Literature selection	30
3.2. Articles characteristics	30
3.3. Outcomes	30
3.3.1. Complications	30
3.3.2. Length of stay	31
3.3.3. Quality of life	31
3.3.4. Readmission rate	33
3.3.5. Postoperative mortality	33
4. Discussion	33
5. Conclusion	34
References	34

* Corresponding author. Pôle Anesthésie-réanimation, CHU de Rennes, Université de Rennes, Inserm NumCan, CIC 1414, 2, avenue Henri-le-Guilloux, 35033 Rennes cedex, France.

E-mail address: helene.beloeil@chu-rennes.fr (H. Beloeil).

1. Introduction

Fast-track surgery was introduced in 1997 by Kehlet [1] and has been widely applied since. The primary goal of this clinical pathway was to improve postoperative patient outcomes and decrease hospitalisation costs. In fact, enhanced recovery or fast-track programs have been shown to be safe, effective and mostly beneficial for the patient. Indeed, Wu et al. [2] reported lower complication rate in ERAS clinical pathways. Colorectal surgery has been the most studied in ERAS programs. Thereby, recent recommendations for enhanced recovery in colorectal surgery have been established by the French Society of abdominal Surgery (SFCD) and the French Society of Anaesthesia and Intensive Care Medicine (SFAR) [3]. The following items are associated with a benefit preoperative counseling and feeding, carbohydrate intake before surgery, immunonutrition before colorectal cancer surgery, goal-directed intraoperative fluid management laparoscopic procedure, multi-modal analgesia, early ambulation and feeding and minimal drain policy. Nowadays, an increasing number of centres tries to apply this clinical pathway to others surgeries like cardiovascular [4] and orthopaedic [5] surgeries. Items of ERAS in colorectal surgery have been applied to hepatic surgery. However, liver surgery has its own complications and high mortality-morbidity rate (15% to 48%) [6,7]. Therefore, it is not clear if ERAS program in liver surgery is safe and effective.

The aim of this review was to assess the safety and efficacy of ERAS protocols versus traditional care by comparing length of stay in hospital (LoS), complications, mortality and readmission rates for patients after liver surgery. We hypothesised that ERAS protocols are relevant and thus can be used on a daily basis.

2. Methods

We performed a systematic review of literature according to Preferred Reporting Items for Systematic reviews and Meta-Analyses guidelines (PRISMA).

2.1. Search strategy and study selection

We searched the Cochrane Central Register of Controlled Trials, Medline and completed clinical trials in Clinicaltrials.gov from January 2007 to November 2017. And we restricted the search to English language. The search terms were (fast-track or fast track or enhanced recovery or ERAS OR rehabilitation) and (hepatectomy or hepatectomies or liver resection or liver resections or hepatic resection or hepatic resections or liver surgery or liver surgeries) not transplantation. We ensure that terms associated individually could give the same result. Relevance was established first on the title then the abstract and on full-texts if information was not found in the abstract. All relevant articles were exported to Zotero[®] (reference management Software).

2.2. Inclusion and exclusion criteria

The criteria for eligibility were (1) full-text randomised clinical trials (RCTs) which: (2) compared traditional care versus enhanced recovery (3) in liver surgery (laparoscopic or open) and (4) considered at least one outcome among: length of stay (LoS), morbidity, mortality and readmission rates. We did not reach study authors to identify additional studies. Exclusion criterion were (1) prospective or retrospective cohort study or case-control study, (2) studies considering transplantation surgery, (3) studies not comparing ERAS care versus traditional care, (4) studies consisted of unpublished data with only the abstract. Guidelines, reviews and meta-analysis were considered apart.

2.3. Data extraction

Year of study publication, authors, title, study characteristics, type of surgery, number of patients and their characteristics (age, ASA score, gender), ERAS protocol/items, length of stay, readmission, morbidity and mortality rates, primary and secondary outcomes and methodological quality (JADAD score) were extracted in a spread sheet. Two members reviewed data of relevant studies.

2.4. Quality assessment

Quality assessment was performed using the JADAD composite scale (1–5), which attributes 1 point for blinding, appropriate method of blinding, randomisation, appropriate method of randomisation and description of outputs and withdrawals while attribute –1 point if blinding is inappropriate or if randomisation is inappropriate.

3. Results

3.1. Literature selection

Initial search found 388 articles. There was no duplicate. After reviewing titles and abstracts, 323 articles did not concern liver surgery or enhanced recovery. Of the 65 articles left, 34 were not RCTs, 23 were review and meta-analyses, 1 was recommendations/guidelines and 3 were RCTs with no traditional care controlled group. Finally, we reviewed 4 articles. The PRISMA diagram is shown (Fig. 1).

3.2. Articles characteristics

The characteristics of the included studies are presented in Table 1 and Table 2. A total of 524 patients were included in 4 RCTs: 254 in ERAS group and 270 in traditional care group. In the study by Liang et al. [8], it is important to notice that a part of the study was not prospective. Patients were ASA I, II or III. Within the 4 studies, 2 studied laparoscopic surgery and the 2 other studies concerned open surgery. Procedures were conducted mostly for colorectal liver metastases, hepatocellular carcinoma (HCC) or benign diseases. In the study by Jones et al. [9], the 2 groups were not comparable. The ERAS group had more patients with colorectal metastases. Therefore, they received more preoperative neoadjuvant chemotherapy versus traditional care group ($P = 0.021$). In the other studies [8,10,11], groups were similar. ERAS components were described for each study and are presented in Table 3. All studies used a similar approach in perioperative management and described discharge criteria by evaluating analgesia, tolerance for solid food and willingness to go home. Of note, “good pain control” was not defined in studies.

3.3. Outcomes

All results are presented in Table 4.

3.3.1. Complications

All studies graded complications according to the Dindo-Clavien classification scored grade I to V [12]. Liang et al. [8] and Ni et al. [11] found that the morbidity rate, considering all complications (grade I to V), was significantly lower in the ERAS group whereas He et al. [10] and Jones et al. [9] did not. This difference may be explained by the sample size. Indeed, number of patient range from single to double (from $n = 86$ in He [10] and $n = 91$ in Jones [9], up to $n = 187$ in Liang [8] and $n = 160$ in Ni [11]).




Fig. 1. PRISMA flow chart.

Moreover, the complication rate according to the Dindo–Clavien classification was the primary outcome of only one study (Ni et al. [11]). An estimation of the sample size was done before inclusion to perform a power of 80% for 2-tailed log-rank test. They needed to include 160 patients, 80 in each group, to report 18% difference between the ERAS group and the TC group in the overall complication rates. Finally, they randomised 80 patients in each group and found a significant difference between ERAS group and TC group (30% vs 46.3%, $P = 0.03$). Among all complications, we can notice that significant differences were reported on grade I Dindo–Clavien classification while number of grade II to V complications were similar. In the study by Liang et al. [8], deep vein thrombosis was significantly lower in ERAS than traditional care group (0 vs 6 event, $P = 0.03$) whereas in Ni et al., it was nausea and vomiting (7 vs 18 event, $P = 0.02$). In the study by Jones et al. [9], grade complication details were not reported. Even if overall complication rate did not reach the statistical significance in this study (17% for ERAS vs 31% for traditional, $P = 0.126$), we can notice that liver

surgery-specific complication rates were similar (15% ERAS vs 11% traditional, $P = 0.612$) and total number of general complications was lower in ERAS group (4 vs 20, $P = 0.009$).

3.3.2. Length of stay

All studies reported a significant reduction of postoperative length of stay. The definition of LoS was different in the study by Jones et al. [9]. Indeed, LoS included days after eventual readmissions whereas the other studies had the same definition of postoperative LoS: from the day of surgery until the day of discharge. Days of hospitalisation before the surgery were not included in the LoS. We noticed that Jones et al. [9] included ASA II patients whereas others studies included patients ASA I and II. Patients were also older than in other studies and had open surgery. But despite all, they had the shortest LoS of all studies (in ERAS or TC). Furthermore, length of stay was the primary outcome of the study as well as in He et al. [10] and Liang et al. [8] studies. None of the 3 studies performed a sample size calculation before inclusions.

3.3.3. Quality of life

Three studies investigated the quality of life after surgery. He et al. [10] primary outcome was a composite including quality of life (QoL) applying the EQ-5D test and length of hospital stay. They found that the QoL after surgery was under the baseline measure in both groups. However, one month after surgery, QoL was considerably improved in the ERAS group (median AUC 36.9 versus 35.2, $P = 0.04$). In the study by Jones et al. [9], QoL was a secondary outcome. They used the same test (EQ-5D) and found similar results. In the study by Ni et al. [11], one of the secondary outcome was general comfort questionnaire measures (GCQ). In that study,

Table 1

JADAD composite scale score for each study. Points attribution: 1 point for blinding, appropriate method of blinding, randomisation, appropriate method of randomisation and description of outputs and withdrawals while attribute -1 point if blinding was inappropriate or if randomisation was inappropriate.

Author	Randomisation	Double blinding	Withdrawals and dropouts	JADAD SCORE
He, F. [10]	+2	0	+1	3
Jones, C. [9]	+2	0	+1	3
Liang, X. [8]	+1	0	0	1
Ni, C.Y. [11]	+2	0	+1	3

Table 2
 Characteristics of the studies: ASA: American Society of Anesthesiologists; ERAS: enhanced recovery after surgery; TC: traditional care; HCC: hepatocellular carcinoma.

Results of the studies									
Author	Year	Type of cases	Nb. of patients (n)		Study endpoints	Complication & morbidity	Postoperative length of stay (days) Median (interquartile range)	Readmission (n)	Mortality (n)
			ERAS	TC		ERAS/TC	ERAS/TC	ERAS/TC	ERAS/TC
He, F. [10]	2015	RCT	48	38	Primary: quality of life; LoS Secondary: % of readmission; mortality; functional recovery; complication; hospital costs; conversions; blood loss	7 (15%)/6 (16%) P=0.88	6 (4–8)/10 (7–15) P=0.04	1/1 P=1	0/0 P=NA
Jones, C. [9]	2013	RCT	46	45	Primary: LoS Secondary: Pain score; morbidity; mortality (30 days follow-up); volume of IV fluids; time to return of: bowel sounds, flatus, full diet, bowel opening and mobilisation	17%/31% P=0.126	4 (3–5)/7 (6–8) P<0.001	2/0 P=0.153	1/1 P=0.987
Liang, X. [8]	2016	RCT	80	107	Primary: LoS; hospitalization cost Secondary: resumption of oral intake; pain score; readmissions; complications	18 (22%)/47 (44%) P=0.002	6±2.6/9.9±5.9 P<0.001	3/5 P=0.6	0/0 P=NA
Ni, C.Y. [11]	2013	RCT	80	80	Primary: complication rate Secondary: C-reactive protein level; LoS; liver functional status on POD 5; general comfort questionnaire measures	24 (30%)/37 (46%) P=0.03	6.9±2.8/8.0±3.7 P=0.018	NC	NC

Table 3
 Summary of ERAS components.

	He, F. 2015 [10]	Jones, C. 2013 [9]	Liang, X. 2016 [8]	Ni, C.Y. 2013 [11]
Preoperative				
Preoperative education or exercise	a	a	a	a
Avoid bowel preparation	a	a	a	a
Avoid preanesthetic medications	NA	a	NA	a
Carbohydrate drinks up to 2 h before surgery	a	a	a	a
Minimal preoperative fasting (2 h)	a	NA	NA	NA
Perioperative				
Antibiotic prophylaxis	a	a	a	NA
Epidural analgesia	a	a	NA	a
Short-acting i.v. anesthetic agent	NA	a	a	NA
Prevention of hypothermia	a	a	a	NA
Avoidance of excessive i.v. fluids	NA	a	a	a
No routine drainage of the peritoneal cavity	a	a	a	a
No nasogastric tube or early removal after surgery	a	a	a	a
Postoperative				
Not sent routinely to the ICU	NA	NA	NA	a
Early removal of urinary catheter (within the 1st 24 h)	a	No, 1st 48 h	a	a
Early oral intake	a 4 h	a	a 6 h	POD 2 a
Early mobilisation	a	a	a	a
Early discontinuation of i.v. fluids	a	a	a	NA
Early restoration of normal diet	Liquid diet before 12 h	a	Semiliquid POD 2 Normal POD 3	POD 3 a
Glucose control	a	a	NA	NA
Nausea and vomiting prophylaxis	a	a	NA	NA
Multimodal analgesia	Thoracic epidural	Epidural+\$(1 g paracetamol + tramadol 50–100 mg)/6 h	Trocar incision analgesia + PCIA + 40 mg ParecoxibNa/12h ± 50 mg tramadol	Epidural or fentanyl transdermal patch
Discharge criteria				
Normal or decreasing serum bilirubin	NA	a	NA	a
Good pain control with oral analgesia only	a	a	a	a
Tolerance of solid food	a	a	a	a
No i.v. fluids	NA	a	a	a
Mobile independently or at the preoperative level	a	a	NA	a
Willingness to go home	a	a	a	a
Normal body temperature	a	NA	a	NA
No incision infection	a	NA	NA	NA

ERAS: Enhanced Recovery After Surgery, NA: Not Available, i.v.: intra venous, ICU: Intensive Care Unit, PCIA: Patient-Controlled Intravenous Analgesia.

^a Component present in the study.

Table 4
Results of the studies.

Author	Year	Type of cases	Nb. of patients (n)		Study endpoints	Complications		Postoperative length of stay (days) Median (interquartile range)	Readmission (n)	Mortality (n)
			ERAS	TC		Grade I	Grade II to V			
He, F. [10]	2015	RCT	48	38	Primary: quality of life; LoS Secondary: % of readmission; mortality; functional recovery; complication; hospital costs; conversions; blood loss	4 (8%)/2 (5%) P=NC	3 (6%)/4 (10.5%) P=NC	6 (4–8)/10 (7–15) P=0.04	1/1 P=1	0/0 P=NA
Jones, C. [9]	2013	RCT	46	45	Primary: LoS Secondary: Pain score; morbidity; mortality (30 days follow-up); volume of IV fluids; time to return of: bowel sounds, flatus, full diet, bowel opening and mobilisation	2 (4%)/4 (9%) P=NC	12 (26%)/24 (53%) P=NC	4 (3–5)/7 (6–8) P<0.001	2/0 P=0.153	1/1 P=0.987
Liang, X. [8]	2016	RCT	80	107	Primary: LoS; hospitalisation cost Secondary: resumption of oral intake; pain score; readmissions; complications	5 (6%)/14 (13%) P=NC	13 (16%)/33 (31%) P=0.02	6 ± 2.6/9.9 ± 5.9 P<0.001	3/5 P=0.6	0/0 P=NA
Ni, C.Y. [11]	2013	RCT	80	80	Primary: complication rate Secondary: C-reactive protein level; LoS; liver functional status on POD 5; general comfort questionnaire measures	10 (12.5%)/20 (25%) P=0.04	14 (17.5%)/17 (21%) P=0.55	6.9 ± 2.8/8.0 ± 3.7 P=0.018	NC	NC

the mean GCQ measures was higher in the ERAS group than in the TC group (101.2 ± 13 vs 93.4 ± 21.4 , $P < 0.01$).

3.3.4. Readmission rate

None of the studies reported a significant difference in terms of readmission. Reasons for readmission were: urinary tract infection for traditional care group, postoperative bile leakage and abdominal collection ($n = 2$) for ERAS group. Liang et al. did not precise reasons for the 8 readmissions. The Ni et al. study [11] did not consider readmission as an outcome.

3.3.5. Postoperative mortality

None of the studies found a significant difference in mortality. There were only 2 deaths in Jones et al., which were the consequence of postoperative small liver syndrome due to extensive preoperative neoadjuvant chemotherapy. In Ni et al. study [11], mortality was not reported.

4. Discussion

The aim of this review was to assess the safety and efficacy of ERAS protocols in liver surgery. With 4 studies representing 524 patients, we found that LoS was in favor of ERAS group while readmission and mortality rate were not different than TC. Two studies [8,11] found lower complication rate in ERAS group while the 2 others [9,10] did not report any significant difference between the 2 groups. Thus, ERAS seems to be safe and efficient for open and laparoscopic liver surgery. Furthermore, He et al. [10] analysed quality of life after surgery and established that patients in ERAS group had a higher score at 1 month than traditional care

group (median area under the curve respectively: 36.9 versus 35.2, $P = 0.04$). Several reviews and meta-analysis are available on this subject. Two meta-analysis [13,14] reported a reduced LoS in the ERAS group despite the heterogeneity of the studies ($I^2 = 77\%$ and $I^2 = 89\%$). In these meta-analyses, the complication rate was lower in ERAS group ($P < 0.0001$, $I^2 = 0\%$ and $P < 0.0001$, $I^2 = 0\%$). Furthermore, as shown by Liang et al. [8] and Ni et al. [11], the meta-analysis by Li et al. [13] found a significant difference not only in grade I complications (OR 0.55, 95% CI: 0.31–0.98, $P = 0.04$, $I^2 = 0\%$) but also in grade II to V complications (OR 0.49, 95% CI: 0.32–0.76, $P = 0.001$, $I^2 = 0\%$). No statistical difference was shown for readmission rate (OR 1.16, 95% CI: 0.38–3.54, $P = 0.08$). All 4 selected studies reported a reduction in LoS despite applying different ERAS components. Indeed, analgesia regimens were all multimodal but included different techniques (from epidural to morphine PCA or fentanyl transdermal patch). No clear differences in ERAS components can explain the differences in complication rate when comparing the 4 studies. Jones et al. applied the most ERAS components when compared with the 3 other studies and did not observe any difference in complication rate. He et al. [10] reported lower hospital costs in the ERAS group (7742 ± 1200 ¥ vs. 9470 ± 1540 ¥, $P = 0.03$). Liang et al. also compared hospitalisation costs: \$ 6871 ± 2571 in the ERAS group and \$ 7948 ± 3630 in the control group ($P = 0.020$). On the opposite, Joliat et al. [15] did not find any significant difference in total hospital cost (€ 38,726 for the ERAS group and € 42,356 (26,898) for the pre-ERAS group, $P = 0.467$). A recent meta-analysis [16] found that hospitalization costs were lower in ERAS group (SMD = -0.31, 95% CI = -0.47 to -0.14, $P = 0.0002$, fixed-effects model: $P = 0.53$, $I^2 = 0\%$). The heterogeneity of the results and the comparison of costs in different countries/heath system do

not allow us to draw any conclusion. More studies with identical evaluation criterion are needed to evaluate the total hospitalization cost for patient in an ERAS clinical pathway.

Our study has limitations. The quality assessment left us with only 4 studies to analyse, one of them (Liang et al. [8]) being just partly prospective. Furthermore, patient eligible standards, treatment programmes, number of ERAS elements used and discharge criteria were quite different between studies. Moreover, inclusion of cirrhotic patients in the selected studies was heterogeneous. This is important as cirrhotic patients can develop more complications. In Liang et al. study [8], there was a non-statistically significant trend towards more cirrhotic patients in the conventional group and this could partly explain the higher rate of complications or longer hospital stay. It should also be noted that 3 of the 4 selected articles included Chinese patients introducing potential bias of selection. It is difficult to know the degree of adherence to ERAS elements that was applied by the care team. Finally, nowadays it seems difficult to design a RCT comparing ERAS with a group including no ERAS components. Jones et al. [9] minimise that bias by using an independent blinded clinician for the evaluation of discharge criteria.

In order to homogenise ERAS items in liver surgery, recommendations have been published by Melloul et al. [17] in 2016. Regarding the evidence of the literature about ERAS in liver surgery, they strongly recommended, among others: preoperative counselling and education of patients, no preoperative fasting more than 6 h, antibiotic prophylaxis, minimal invasive approach, no nasogastric tube routinely, maintenance of normothermia during liver resection, eat normal food at postoperative day 1, no routine thoracic epidural analgesia (wound infusion catheter or intrathecal opiates can be good alternatives combined with multimodal analgesia), multimodal approach to postoperative nausea, vomiting and analgesia and target guiding fluid. These components are the same as these recommended in colorectal surgery. Nevertheless, some essential items for colorectal surgery like laparoscopic surgery, early mobilisation, early removal of urinary catheter and no routine drainage of the peritoneal cavity are not actually reported as advisable for liver surgery. The ERAS group recommendations are the only ones published so far. Recommendations from scientific society are needed.

5. Conclusion

ERAS in hepatic surgery is feasible, safe, efficient and worthwhile for patients. Thereby, recommendations have been established for liver surgery. Nevertheless, additional studies of quality applying strong grade recommendations and evaluating the weak ones are essential to optimise ERAS clinical pathway.

Funding

Funding was solely institutional.

Disclosure of interest

The authors declare that they have no competing interest.

References

- [1] Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth* 1997;78:606–17.
- [2] Wu S-J, Xiong X-Z, Lu J, et al. Fast-track programs for liver surgery: a meta-analysis. *J Gastrointest Surg* 2015;19:1640–52.
- [3] *Rehabilitation rapide apres une chirurgie colorectale, programmée*. SFAR–Société Française d'Anesthésie et de Réanimation.; 2015 [Accessed 22 November 2017]<http://sfar.org/rehabilitation-rapide-apres-une-chirurgie-colorectale-programmee/>.
- [4] Muehling B, Schelzig H, Steffen P, et al. A prospective randomized trial comparing traditional and fast-track patient care in elective open infrarenal aneurysm repair. *World J Surg* 2009;33:577–85.
- [5] Jones EL, Wainwright TW, Foster JD, et al. A systematic review of patient reported outcomes and patient experience in enhanced recovery after orthopaedic surgery. *Ann R Coll Surg Engl* 2014;96:89–94.
- [6] Benzoni E, Molaro R, Cedolini C, et al. Liver resection for HCC: analysis of causes and risk factors linked to postoperative complications. *Hepatogastroenterology* 2007;54:186–9.
- [7] Karanjia ND, Lordan JT, Fawcett WJ, et al. Survival and recurrence after neoadjuvant chemotherapy and liver resection for colorectal metastases: a ten year study. *Eur J Surg Oncol* 2009;35:838–43.
- [8] Liang X, Ying H, Wang H, et al. Enhanced recovery program versus traditional care in laparoscopic hepatectomy. *Medicine (Baltimore)* 2016;95:e2835.
- [9] Jones C, Kelliher L, Dickinson M, et al. Randomized clinical trial on enhanced recovery versus standard care following open liver resection. *Br J Surg* 2013;100:1015–24.
- [10] He F, Lin X, Xie F, et al. The effect of enhanced recovery program for patients undergoing partial laparoscopic hepatectomy of liver cancer. *Clin Transl Oncol* 2015;17:694–701.
- [11] Ni CY, Yang Y, Chang YQ, et al. Fast-track surgery improves postoperative recovery in patients undergoing partial hepatectomy for primary liver cancer: a prospective randomized controlled trial. *Eur J Surg Oncol* 2013;39:542–7.
- [12] Dindo D, Demartines N, Clavien P-A. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–13.
- [13] Li L, Chen J, Liu Z, et al. Enhanced recovery program versus traditional care after hepatectomy: a meta-analysis. *Medicine (Baltimore)* 2017;96:e8052.
- [14] Zhao Y, Qin H, Wu Y, et al. Enhanced recovery after surgery program reduces length of hospital stay and complications in liver resection: a PRISMA-compliant systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2017;96:e7628.
- [15] Joliat G-R, Labgaa I, Hübner M, et al. Cost-benefit analysis of the implementation of an enhanced recovery program in liver surgery. *World J Surg* 2016;40:2441–50.
- [16] Wang C, Zheng G, Zhang W, et al. Enhanced recovery after surgery programs for liver resection: a meta-analysis. *J Gastrointest Surg* 2017;21:472–86.
- [17] Melloul E, Hübner M, Scott M, et al. Guidelines for perioperative care for liver surgery: enhanced recovery after surgery (ERAS) society recommendations. *World J Surg* 2016;40:2425–40.