



ELSEVIER

## ERRATUM

# Erratum to “Arthroplasty for glenohumeral arthritis in shoulders with a previous Bristow or Latarjet procedure” [J Shoulder Elbow Surg 2018;27:1607-1613]



Laurent B. Willemot, MD, Bassem T. Elhassan, MD, John W. Sperling, MD, MBA, Robert H. Cofield, MD, Joaquin Sánchez-Sotelo, MD, PhD

Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN, USA

In the abovementioned article earlier/incorrect versions of **Tables I** and **II** were presented for publication. The numbers reported in the text are correct. The correct tables are available here:

**Table I** Features of the three groups of shoulder arthroplasty included in the study

	HA (5)	TSA (14)	RTSA (11)	Total (30)
M:F	2:3	11:3	4:7	17:13
Age (years)	34.2 (SD 8.7)	51.2 (SD 11.8) ( $P < .001$ )	57.7 (SD 8.7) ( $P < .001$ )	50.7 (SD 12.8)
Coracoid transfer	Bristow 6	Bristow 14	Bristow 8 Latarjet 3	Bristow 27 Latarjet 3
Interval BL - SA (years)	12.7 (SD 9.0)	22.5 (SD 8.8)	19.8 (SD 18.1)	19.9 (SD 13.1)
Clinical FU (years)	4.3 (SD 1.7)	6.4 (SD 6.4)	2.8 (SD 1.2)	4.9 (SD 4.8)
Rx FU (years)	4.6 (SD 2.1)	5.9 (SD 7.2)	2.2 (SD 1.1)	4.1 (SD 5.1)
Glenoid bone grafting		3 (21.4%)	4 (36.3%)	7 (23.3%)

HA, hemiarthroplasty; TSA, total shoulder arthroplasty; RTSA, reverse total shoulder arthroplasty; M, male; F, female; BL, Bristow-Latarjet; SA, shoulder arthroplasty; FU, follow-up; SD, standard deviation.

**Table II** Outcome, complications and reoperations

	HA (5)	TSA (14)	RTSA (11)	Total (30)
Complications	4 (80.0%)	7 (50.0%)	2 (18.18%) vs. HA ( $P = .02$ ) vs. TSA ( $P = NS$ )	13 (43.3%)
Revisions	3 (60.0%)	6 (42.9%)	0 (0.0%) vs. HA ( $P = .0058$ ) vs. TSA ( $P = .015$ )	9 (30.0%)
Δ Pain score (preop=>postop)	-1.5 ( $P = .039$ )	-3.3 ( $P < .001$ )	-4.2 ( $P < .001$ )	-3.4 ( $P < .001$ )
Δ Elevation (degrees)	33.8 (SE 18.0) (n.s.)	35.5 (SE 11.0) ( $P = .009$ )	57.7 (SE 10.7) ( $P = .003$ )	44.6 (SE 7.1) ( $P < .001$ )
Δ External rotation (degrees)	13.8 (SE 11.4) (n.s.)	26.4 (SE 8.3) ( $P = .010$ )	18.6 (SE 7.4) ( $P = .038$ )	21.2 (SE 4.9) ( $P < .001$ )
Δ Internal rotation	0.8 (SE 0.8) (n.s.)	0.4 (SE 0.4) (n.s.)	0.3 (SE 0.6) (n.s.)	0.4 (SE 0.3) n.s.
Satisfaction	Satisfied 2 Dissatisfied 3	Excellent 2 Satisfied 5 Dissatisfied 7	Excellent 5 Satisfied 5 Dissatisfied 1	Excellent 7 Satisfied 12 Dissatisfied 11

HA, hemiarthroplasty; TSA, total shoulder arthroplasty; RTSA, reverse total shoulder arthroplasty; M, male; F, female; BL, Bristow-Latarjet; SA, shoulder arthroplasty; FU, follow-up; SE, standard error; NS, not significant.

DOI of original article: <https://doi.org/10.1016/j.jse.2018.02.062>