



Carbohydrate oxidation and glucose utilisation under hyperglycaemia in aged and young males during exercise at the same relative exercise intensity

James J. Malone¹ · Minoo Bassami² · Sarah C. Waldron³ · Iain T. Campbell³ · Andrew Hulton⁴ · Dominic Doran⁵ · Don P. MacLaren⁵

Received: 26 June 2018 / Accepted: 11 October 2018 / Published online: 23 October 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Purpose The purpose of the present study was to investigate the age-related carbohydrate oxidation and glucose utilisation rate response during exercise at the same relative intensity under hyperglycaemia in aged and young males.

Methods 16 endurance-trained aged ($n = 8$; 69.1 ± 5.2 year) and young ($n = 8$; 22.4 ± 2.9 year) males were studied during 40 min of cycling exercise ($60\% \dot{V}O_{2\max}$) under both hyperglycaemic and euglycaemic (control) conditions. Venous blood samples were collected at baseline, post-infusion, mid- and post-exercise. Carbohydrate and fat oxidation rates were determined at both 15 and 35 min during exercise, and glucose utilisation rates were calculated.

Results The aged group displayed significantly lower rates of carbohydrate oxidation during exercise during maintained hyperglycemia (15 min = 2.3 ± 0.4 vs. 1.6 ± 0.5 g min^{-1} ; 35 min = 2.3 ± 0.5 vs. 1.5 ± 0.5 g min^{-1}) and control (15 min = 2.2 ± 0.4 vs. 1.6 ± 0.7 g min^{-1} ; 35 min = 1.9 ± 0.7 vs. 1.3 ± 0.7 g min^{-1}) conditions ($P = 0.01$). The rate of glucose utilisation during exercise was also significantly reduced (85.76 ± 23.95 vs. 56.67 ± 15.09 $\mu\text{M kg}^{-1} \text{min}^{-1}$). There were no differences between age groups for anthropometric measures, fat oxidation, insulin, glucose, NEFA, glycerol and lactate ($P > 0.05$) although hyperglycemia resulted in elevated glucose and insulin, and attenuated fat metabolite levels.

Conclusion Our findings highlight that ageing results in a reduction in carbohydrate oxidation and utilisation rates during exercise at the same relative exercise intensity.

Keywords Carbohydrate oxidation · Fat oxidation · Glucose clamp · Insulin · Ageing · Endurance

Communicated by Guido Ferretti.

Iain T. Campbell: Currently retired from University Hospital of South Manchester.

✉ James J. Malone
malonej2@hope.ac.uk

¹ School of Health Sciences, Liverpool Hope University, Taggart Avenue, Liverpool L16 9JD, UK

² Faculty of Physical Education and Sports Sciences, Allameh Tabataba'i University, Tehran, Iran

³ Department of Anaesthesia, University Hospital of South Manchester, Manchester, UK

⁴ School of Biosciences and Medicine, University of Surrey, Guildford, UK

⁵ Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

Abbreviations

AMPK	Adenosine 5'-monophosphate-activated protein kinase
ANOVA	Analysis of variance
DXA	Dual-energy X-ray absorptiometry
ELISA	Enzyme-linked immunosorbent assay
FPG	Fasting plasma glucose
GLUT-4	Insulin-sensitive glucose transporter
HOMA2	Homeostasis model assessment
HOMA2-%B	β -Cell function
HOMA2-%S	Insulin sensitivity
HOMA2-IR	Insulin resistance
HR _{max}	Maximal heart rate
INF	Glucose infusion rate
NEFA	Non-esterified fatty acids
$\dot{V}CO_2$	Volume of carbon dioxide
$\dot{V}O_2$	Volume of oxygen
$\dot{V}O_{2\max}$	Maximal oxygen uptake
RER	Respiratory exchange ratio

SC Space correction
UC Correction for urinary loss of glucose

and glucose utilisation rates in comparison to trained aged males at the same relative exercise intensity.

Introduction

Ageing is associated with a decline in physical activity which may lead to alterations in carbohydrate metabolism, particularly in terms of glucose utilisation (DiPietro et al. 2006) and oxidation rates (Sial et al. 1996). Skeletal muscle is the major site relating to insulin-mediated glucose disposal, with the level of oxidative capacity suggested to relate to the effectiveness of insulin action (Short et al. 2003). The rate of glucose uptake by muscle depends on both exercise and circulating levels of insulin concentration (Kern et al. 1990). Previously untrained aged adults display a reduced carbohydrate oxidative capacity compared to young untrained adults at the same relative intensity (Meredith et al. 1989; Coggan et al. 1992b). This is particularly problematic as a lack of exercise training in aged adults may lead to reduced insulin sensitivity and ultimately insulin resistance (Lee et al. 2015; Colberg et al. 2016).

Despite the reported reductions in carbohydrate oxidation due to ageing at the same relative intensity, evidence suggests that long-term endurance training can attenuate the effects of ageing (Coggan et al. 1992a; Dube et al. 2008). Comparisons between trained and sedentary aged adults have revealed an improvement in insulin sensitivity and glucose disposal within trained aged adults (Pratley et al. 1995; Lanza et al. 2008). Training intervention studies have also found improvements in insulin-mediated glucose disposal and physical fitness parameters in previously sedentary aged adults (Cox et al. 1999; Evans et al. 2005; DiPietro et al. 2006; Bloem and Chang 2008). Thus, it may be possible to offset some of the adverse physiological effects of ageing through regular endurance exercise. Bassami et al. (2007) previously found that a relative exercise intensity of 60% $\dot{V}O_{2max}$ produced peak fat oxidation rates in aged adults. Subsequently, this would allow for the measurement of fat oxidation and the relative contribution of carbohydrate oxidation during exercise.

Previous work investigating carbohydrate metabolism in trained aged adults compared to young has utilised the hyperinsulinemic clamp technique at rest (Dubé et al. 2016). However, no study to date has investigated the response under conditions of hyperglycaemia to quantify insulin resistance/sensitivity alongside oxidation rates during exercise. In this study, we investigated the age-related carbohydrate oxidation and glucose utilisation rate response during a steady-state exercise protocol at the same relative exercise intensity under maintained hyperglycaemia. We hypothesised that trained young males would demonstrate higher rates of carbohydrate oxidation

Methods

Participants

Sixteen trained males who regularly participated in endurance exercise (running = 6 and cycling = 10, > 3 times per week) were recruited for the study and separated into the aged ($n = 8$) and young ($n = 8$) group categories (Table 1). Participants were recruited from local running and cycling clubs through online advertisements and follow-up via email/phone. The participants were medically screened by a qualified physician prior to study commencement. Following the initial screening and testing process, four aged participants declined to take any further part in the study (i.e. 12 recruited, with 8 aged participants completing the study). All participants were non-smokers, free from any metabolic conditions (e.g. diabetes, cardiovascular disease and musculoskeletal issues) and not on any medications known to influence glucose/fat metabolism. Informed consent was obtained from all individual participants and ethics approval was granted by the Liverpool John Moores University ethics committee prior to beginning the study.

Table 1 Participant anthropometric and physical characteristics according to age group (mean \pm SD)

Variable	Young	Aged
Age (years)	22.4 \pm 2.9	69.1 \pm 5.2*
Training frequency (sessions per week)	4 \pm 1	4 \pm 1
Training duration (total per week – mins)	258 \pm 30	251 \pm 31
Height (m)	1.8 \pm 0.1	1.7 \pm 0.1*
Body mass (kg)	78.6 \pm 4.3	76.3 \pm 10
Body mass index (kg m ⁻²)	24.8 \pm 1.7	25.9 \pm 2.4
Body fat (%)	18.9 \pm 3.9	20.8 \pm 4.3
Lean body mass (kg)	59.5 \pm 4	57.7 \pm 7
$\dot{V}O_{2max}$ (ml kg ⁻¹ min ⁻¹)	55.8 \pm 5.1	41.1 \pm 12.2*
HR _{max} (beats min ⁻¹)	190.5 \pm 8.7	160 \pm 10.3*
HOMA2-%B	63.6 \pm 22.4	125.3 \pm 78.2*
HOMA2-%S	230.6 \pm 102.0	146.0 \pm 95.2
HOMA2-IR	0.5 \pm 0.3	1.0 \pm 0.5

$\dot{V}O_{2max}$ maximal oxygen uptake, HR_{max} maximal heart rate, HOMA homeostatic model assessment, %B β -cell function, %S insulin sensitivity, IR insulin resistance

*Significant difference ($P < 0.05$) between age groups

Experimental design

This study was a single-blind, repeated measures design that took place over a 3-week period. The study consisted of an initial anthropometric and physiological assessment with protocol familiarisation, followed by two steady-state exercise protocols during which glucose or saline infusion took place in a sequential order. The glucose infusion trial took place first because the rate of saline infusion needed to be equivalent to that of glucose infusion (MacLaren et al. 1999). However, the participant was not informed of which infusion condition they were undertaking. Each visit to the laboratory was separated by a 7-day period to ensure sufficient recovery between the trials. Figure 1 outlines the protocols and sampling time points of the experimental protocol. Participants were required to refrain from drinking alcohol and strenuous exercise 24 h before trials. Furthermore, participants were required to fast for 12 h prior to commencement, with all trials carried out in the morning (08:00).

Protocols

Anthropometric and physiological assessment

On arrival at the laboratory, participants were assessed for height and weight using a calibrated stadiometer and digital scale, respectively (Seca, GmbH & Co., Germany). Following this initial assessment, fat and lean body mass were determined using dual-energy X-ray absorptiometry (DXA) (Hologic QDR Series Discovery A, Bedford, MA, USA). Prior to scanning, the DXA unit was calibrated using criterion phantom devices provided by the manufacturer to minimise scanning errors. Participants followed standard protocols of food and fluid intake prior to each scan. The participants were placed with their hands in a pronated flat position, with the legs secured with straps to avoid overlap within the lower limbs in minimal clothing (i.e. t-shirt and

shorts). The same technician analysed all scans using manual analysis to determine total body fat and lean body mass.

To determine maximal oxygen uptake ($\dot{V}O_{2\max}$), each participant completed an incremental maximal cycling test on a Monark cycle ergometer (874E, Vansbro, Sweden). The test began at 90 W and was increased by 30 W every 3 min until volitional exhaustion. Expired gas was collected for the final 60 s of each 3-min stage using the Douglas bag method (Douglas 1911). A mouthpiece connected to a two-way valve was used for the collection of gas samples (Cranlea & Company, Birmingham, England), which were subsequently analysed for oxygen and carbon dioxide concentrations (Servomex 5200S, Crowborough, UK). $\dot{V}O_{2\max}$ was confirmed using established physiological criteria from the British Association of Sport and Exercise Science (BASES), and included oxygen uptake reaching a plateau with increasing work rate, a heart rate close to age-predicted maximal values, and a rating of perceived exertion (RPE) of 20. Following determination of individual $\dot{V}O_{2\max}$ values, work rates equivalent to 60% $\dot{V}O_{2\max}$ were calculated by interpolating the relationship of $\dot{V}O_{2\max}$ and work rate (W).

Steady-state exercise infusion protocol

On arrival to the laboratory, participants were instructed to void urine before lying on a medical bed in a supine position for insertion of a 16-gauge IV cannula into the antecubital vein of the right hand under local anaesthetic for infusion of glucose (Infusion Pump, Colleague, Baxter Healthcare, IL, USA). The left hand was placed in a hotbox to arterialise the blood and a 20-gauge cannula was inserted retrogradely into the dorsal vein of the left hand for sampling arterialised venous blood. The use of a heated superficial hand vein as a replacement for an artery has been validated for measurement of glucose kinetics in humans (Abumrad et al. 1981). A baseline blood sample (20 ml) was taken after 20-min rest, after which a priming infusion of 20% dextrose was initiated




Fig. 1 Schematic representation of the experimental protocol. *BS* blood sample, *RES* respiratory gas collection, *U* urine sample

into the right vein for 30 min to increase the blood glucose concentration to 10 mM in accordance to the method by DeFronzo et al. (1979). During this period, blood glucose concentration was measured (Hemocue AB, Angelholm, Sweden) and the infusion rate adjusted every 5 min based on the negative feedback principle. At the end of the 30-min prime infusion of glucose, a further blood sample (20 ml) was taken.

Thereafter, participants cycled at 60% $\dot{V}O_{2\max}$ for 40 min on a Monark cycle ergometer (874E, Vansbro, Sweden) following a 5-min warm-up period at 90 W. Dextrose was continually infused to maintain blood glucose concentration at 10 mM, with glucose concentration measured and infusion rate adjusted every 5 min accordingly. Participants were required to maintain a constant cycle cadence of between 60 and 70 rpm on the cycle ergometer during the exercise period. Two more venous blood samples (20 ml) were taken during exercise (20 min) and immediately after exercise (40 min). Indirect calorimetry was performed during exercise for the measurement of oxygen volume ($\dot{V}O_2$), carbon dioxide volume ($\dot{V}O_{2\max}$) and respiratory exchange ratio (RER). Samples were collected at 15 and 35 min using the Douglas bag method previously detailed and whole-body substrate oxidation rates (carbohydrate and fat) were determined using the stoichiometric equations of Frayn (1983).

On completion of the exercise protocol, participants lay down on a medical bed whilst the glucose infusion rate was gradually slowed. Participants were provided with exogenous carbohydrate sources (e.g. sandwiches, chocolate and sports drinks) to prevent rebound hypoglycaemia. During this recovery period, blood glucose was continually measured every 5 min until glucose levels were stable. A urine sample was collected immediately following the recovery period to measure any spill over of glucose.

In the saline control trial, participants were infused with a 0.9% saline solution instead of receiving the dextrose solution. The exercise protocol, indirect calorimetry and blood sample collection time points were identical to the previous dextrose infusion trial.

Blood sample analysis

Venous blood samples were withdrawn using a sterile plastic syringe and gauge needle, which were treated using different anticoagulants specific to each variable. Blood samples were centrifuged (Sigma 3-18K, Osterode am Harz, Germany) for 15 min (speed = 3950 rpm, RCF = 3000, temperature = 4 °C). Plasma was removed and stored in an ultralow-temperature freezer (Thermoforma, Ohio, USA) at – 80 °C until further analysis. Plasma samples were defrosted and analysed for non-esterified fatty acids (NEFA), glycerol, glucose and lactate concentrations using a fully automated bench-top clinical chemistry analyser (Daytona RX, Randox Laboratories

Ltd, Crumlin, UK). Plasma samples were also analysed for insulin concentration using a solid-phase enzyme-linked immunosorbent assay (ELISA) kit (Demeditec Diagnostics GmbH, Germany) based on the sandwich principle. For the lactate analysis, only four of the eight participants in the elderly group were successfully analysed for lactate concentrations.

HOMA scores

Insulin resistance (HOMA2-IR), insulin sensitivity (HOMA2-%S) and β -cell function (HOMA2-%B) in fasting state were determined using a homeostasis model assessment (HOMA2). These values were calculated using a computer programme based on the equations of Levy et al. (1998). The programme is based on a corrected non-linear model that has been calibrated in line with current insulin assays. The calculations are made from fasting plasma insulin (FPI, measured in $\mu\text{IU ml}^{-1}$) and fasting plasma glucose (FPG, measured in mmol l^{-1}) amounts.

Rate of glucose utilisation

The rate of glucose utilisation was calculated using a series of formula described by DeFronzo et al. (1979). The utilisation rate is based on the computation of an M value, which is a measure of glucose tolerance. The M value is calculated for 5-min intervals throughout the hyperglycaemic clamp according to the equation:

$$M = \text{INF} - \text{UC} - \text{SC},$$

where INF is the glucose infusion rate, UC is the correction for urinary loss of glucose and SC is the space correction, with all values computed in dimensions of $\mu\text{M kg}^{-1} \text{min}^{-1}$. The M value is calculated from the mean of the 5-min intervals from 0 to 20 min and 20 to 40 min of the glucose infusion trial.

Statistical analysis

A three-way mixed design ANOVA for trial (glucose and saline), age (young and aged) and time (15 and 35 min) was employed to examine the differences in the mean values of fat and carbohydrate oxidation measured during exercise. In addition, blood parameters (glucose, NEFA, glycerol, lactate and insulin) measured at – 30, 0, 20 and 40 min (time) across two trials (glucose and saline) and two age groups (young and aged) were analysed using a three-way mixed design ANOVA. Glucose utilisation rates were analysed across two time points (0–20 and 20–40 min) and age group (young and aged) using a two-way mixed design ANOVA. Where the ANOVA revealed a significant effect, post hoc tests were completed using the Bonferroni correction

method. Anthropometric and training data, HOMA2-IR, HOMA2-%B and HOMA2-%S were compared between age groups (young and aged) using an independent *t* test or non-parametric equivalent (Mann–Whitney test) after checking for normality. HR_{max} was the only parameter that was found to violate the assumption of normality and, therefore, analysed using the non-parametric equivalent test. Alpha significance level was set at 0.05 for all analyses. All data were analysed using Statistical Package for Social Sciences (version 24.0, SPSS Inc., Chicago, IL, USA). Assumptions for all statistical analyses were explored according to the methods of Field (2009). Values are reported as mean \pm SD.

Results

Anthropometric and physiological assessments

The aged group displayed lower $\dot{V}O_{2max}$, HR_{max} and higher HOMA2-%B scores in comparison to the young group ($P < 0.05$) (Table 1). No differences were observed for body mass, body mass index, body fat %, lean body mass, HOMA2-%S and HOMA2-IR ($P > 0.05$).

Oxidation rates

Carbohydrate oxidation

The young group displayed significantly higher overall rates of carbohydrate oxidation compared to the aged group during both maintained hyperglycaemia (15 min = 2.3 ± 0.4 vs. 1.6 ± 0.5 g min^{-1} ; 35 min = 2.3 ± 0.5 vs. 1.5 ± 0.5 g min^{-1}) and control (15 min = 2.2 ± 0.4 vs. 1.6 ± 0.7 g min^{-1} ; 35 min = 1.9 ± 0.7 vs. 1.3 ± 0.7 g min^{-1}) conditions ($P = 0.01$; Fig. 2a). Both groups displayed significantly higher carbohydrate oxidation rates at 15 min compared to 35 min during exercise ($P = 0.04$). There was a significant reduction in oxidation rates during the saline control trial at 35 min in both groups compared to the glucose infusion trial ($P = 0.01$).

Fat oxidation

Fat oxidation rates were significantly higher at 35 min compared to 15 min during control conditions for the young group (0.6 ± 0.3 vs. 0.4 ± 0.2 g min^{-1} , $P = 0.04$) and aged group (0.4 ± 0.4 vs. 0.3 ± 0.2 g min^{-1} , $P = 0.04$) (Fig. 2b). However, there were no time differences found for both groups during maintained hyperglycaemia ($P < 0.05$). In addition, there was no significant difference found in fat oxidation rates between young and aged groups ($P < 0.05$).

Blood analysis

Glucose

There were no significant differences found between age groups across both trials ($P < 0.05$) (Fig. 3a). Glucose infusion resulted in a significantly elevated glucose levels compared to the control trial in both young (9.3–10.1 vs. 4.6–4.7 mmol l^{-1} , respectively) and aged (10.4–10.7 vs. 4.7–5.3 mmol l^{-1} , respectively) groups ($P = 0.01$).

Insulin

Despite displaying higher overall values in the young compared to aged group during maintained hyperglycaemia (13.6 ± 11.0 and 8.9 ± 8.9 $\mu IU ml^{-1}$), this did not reach statistical significance ($P = 0.08$) (Fig. 3b). There were significantly higher values found during maintained hyperglycaemia compared to control in both groups when comparing baseline (–30 min) and post-exercise (40 min) values ($P = 0.01$). No difference was found both within and between groups during exercise (i.e. 0–40 min) for either trial ($P < 0.05$).

NEFA

There was a significant increase observed during control compared to maintained hyperglycaemia in both age groups ($P = 0.01$) (Fig. 3c). Specifically, differences were found during exercise (20 min; $P = 0.04$) and post-exercise (40 min; $P = 0.01$) compared with baseline (–30 min). The aged group displayed higher NEFA concentrations during exercise within the control trial (0.29–0.50 vs. 0.57–0.78 mmol l^{-1}). However, these differences were determined as non-significant ($P = 0.07$). No differences were observed between age groups during maintained hyperglycaemia ($P < 0.05$).

Glycerol

The aged group displayed higher concentrations of glycerol during both trials; however, there was no statistical significance found between age groups ($P = 0.67$) (Fig. 3d). The control trial resulted in higher concentrations for both groups, with differences found at –30 vs. 40 min ($P = 0.01$), 0 vs. 40 min ($P = 0.01$) and 20 vs. 40 min ($P = 0.01$).

Lactate

Blood lactate concentrations were similar between both age groups across both trials ($P = 0.94$) (Fig. 3e). Values were found to be significantly elevated immediately post-exercise compared to both baseline (–30 min; $P = 0.01$) and post-infusion (0 min; $P = 0.01$) across both trials. There was also




Fig. 2 Oxidation rates during glucose and saline infusion trials for young and aged groups. **a** Carbohydrate and **b** fat. *G* glucose, *S* saline. *Significant difference for age; #significant difference for time (15 vs. 35 min); §significant difference for trial (saline vs. glucose)

a significantly higher lactate production during maintained hyperglycaemia compared to control conditions ($P=0.02$).

and 20–40 min (54.4 vs. 63.7%) compared to the young group, respectively (Fig. 4b).

Glucose utilisation rates

Significantly higher overall utilisation rates were observed in the young compared to aged group at both 0–20 min (85.76 ± 23.95 and $56.67 \pm 15.09 \mu\text{M kg}^{-1} \text{min}^{-1}$, respectively) and at 20–40 min (104.87 ± 17.79 and $57.60 \pm 29.30 \mu\text{M kg}^{-1} \text{min}^{-1}$, respectively) ($P=0.01$) (Fig. 4a). However, there were no significant differences found within groups at 0–20 min and 20–40 min following glucose infusion ($P<0.05$). When glucose utilisation rates were expressed relative to total carbohydrate oxidation, the aged group displayed lower percentages at both 0–20 min (47.8 vs. 52.1%)

Discussion

The present study examined the age-related carbohydrate and fat oxidation response and glucose utilisation rates during a steady-state relative exercise protocol under maintained hyperglycaemia and control conditions. We hypothesized that trained young males would demonstrate higher rates of carbohydrate oxidation and glucose utilisation in comparison to trained aged males. Our findings were in agreement with this hypothesis, with the trained aged group displaying lower rates of carbohydrate oxidation despite similar rates of fat oxidation. In addition, glucose utilisation rates




Fig. 3 Blood analysis measures at baseline (−30 min), post-infusion (0 min), during exercise (20 min) and post-exercise (40 min) during glucose and saline infusion trials for young and aged groups. **a** Glu-

ucose; **b** insulin; **c** NEFA; **d** glycerol; **e** lactate concentrations. #Significant difference for time (compared to baseline); \$significant difference for trial (saline vs. glucose)


were significantly reduced in trained aged adults throughout the exercise protocol. The trained aged adults displayed a reduced insulin response to hyperglycaemia, although this was not deemed statistically different to trained young adults. There were no age-related differences in NEFA, glycerol and lactate concentrations between both groups.

During moderate exercise, aged adults displayed a reduction in carbohydrate oxidation despite similar fat oxidation rates to younger adults. This is in agreement with previous studies which observed reduced carbohydrate oxidation using indirect calorimetry in aged athletes (Sial et al. 1996; Dubé et al. 2016). Dubé et al. (2016) found aged athletes are able to maintain similar levels of fat oxidation at moderate intensity exercise in comparison to their young counterparts. However, when the intensity of exercise increased, the level

of carbohydrate oxidation was blunted in aged adults. This may be explained by the possibly reduced glycogen storage capacity of trained aged adults (Dubé et al. 2016). Sial et al. (1996) reported reduced fat oxidation rates despite similar carbohydrate oxidation rates as observed in the present study. One possible explanation for such discrepancies may be due to the lower training status of the aged adults used in the study ($\dot{V}O_{2max} = 41.1$ vs. 31.4 ml kg⁻¹ min⁻¹). Training intervention studies have also reported an improvement in fat oxidation rates in previously sedentary aged adults (Sial et al. 1998). Therefore, it would appear that despite regular endurance training protecting the ability to oxidise fat, there is an aged-related decline in carbohydrate oxidation.

Having stated the above, it must be borne in mind that the aged athletes exercised at a lower absolute exercise intensity

Fig. 4 a Glucose utilisation rates during exercise for young and aged during the glucose infusion trial. **b** Glucose utilisation rates relative to overall carbohydrate oxidation rates during exercise for young and aged groups. *Significant difference between age groups ($P < 0.05$)



than the young athletes (i.e. 85 W vs. 127 W; a 33% lower exercise intensity). The difference in carbohydrate oxidation between the groups, therefore, could be associated with the fact that the 33% lower intensity undertaken by the aged group is reflected by a 30% reduction in carbohydrate oxidation. So, most of the reduced carbohydrate oxidation may be associated with lower absolute exercise intensity. However, other factors need to be considered since not all the reduced carbohydrate oxidation is due to exercise intensity. Sial et al. (1996) previously compared both relative and absolute exercise intensities in aged and young groups' 60 min of cycle ergometer exercise under euglycaemic conditions. The authors found the aged group displayed 35% higher carbohydrate oxidation rates at the same absolute intensity but 40% lower when compared at the same relative intensity as the young group. This would agree with our present findings and thus the absolute intensity at which aged adults' exercises may play a key role in how they handle and oxidise

carbohydrate under both hyperglycaemic and euglycaemic conditions.

Ageing has previously been associated with a decrease in glucose tolerance (Scheen 1997; Meneilly and Tessier 2001) and increased peripheral resistance to the action of insulin (Broughton and Taylor 1991; Ryan 2000). The HOMA2 data revealed that trained aged adults had significantly higher HOMA2-%B values compared to the trained young group. Despite a trend towards lower HOMA2-%S and higher HOMA2-IR values in the trained aged group, this was deemed to not be statistically different to the young group. Similar findings were evident for the insulin response during exercise, with a trend towards lower values in the aged group but were not deemed statistically significant. As the glucose utilisation rates were significantly lower in the aged group, this would suggest that there may be evidence of reduced insulin sensitivity within the aged group. One possible cause may be due to lower levels of GLUT-4 in

skeletal muscle, which has previously been associated with adults demonstrating reduced insulin sensitivity (Dela et al. 1994). However, training intervention studies have found that aged adults maintain the ability to increase concentrations of GLUT-4 in skeletal muscle (Cox et al. 1999; Kim et al. 2004; Biensø et al. 2015). Although speculative, as muscle biopsies were not taken in the present study, it may be pertinent that aged adults can increase their GLUT-4 concentrations but are still lower than their younger counterparts which limit the oxidation and disposal of exogenous carbohydrate sources during exercise.

Another possible mechanism for the reduced uptake of glucose in aged adults may relate to skeletal muscle function involving adenosine 5'-monophosphate-activated protein kinase (AMPK). AMPK has previously been described as a key “master switch” in metabolism regarding the regulation of fuel transport for oxidation (Vigelsø et al. 2016). Whilst AMPK plays a major role in the regulation of intracellular fatty acid oxidation, it has previously been reported that it may also regulate insulin sensitivity via stimulating GLUT-4 expression (Jessen et al. 2003). Previous research has reported ageing-related insulin resistance associated with impaired AMPK- α activity (Qiang et al. 2007; Morris et al. 2010). It has been suggested that regular exercise may increase the recruitment of the AMPK signalling system and thus reduce the level of glucose intolerance in aged populations (Winder and Hardie 1999). In the present study, glucose utilisation rates expressed as % of total carbohydrate oxidation were lower in the aged group across the exercise protocol. This would imply that ageing results in a reduction in the ability to utilise exogenous carbohydrate intake, possibly due to the limited activity of AMPK. However, further research is required investigating the AMPK response within endurance-trained aged adult humans.

Previous research has reported an increase in fat mass and decrease in lean body mass with advancing age in adults (Vermeulen et al. 1999). The present study revealed no difference between aged and young adults in terms of lean body mass, body fat % and body mass index when matched for exercise type and frequency. Longitudinal training data support this notion, with significant reductions in fat mass reported in previously sedentary aged adults (Evans et al. 2005). Despite similar anthropometric characteristics between age groups, the aged adults displayed significantly reduced fitness levels compared to young adults (e.g. $\dot{V}O_{2\max} = 41.1$ vs. 55.8 ml kg⁻¹ min⁻¹, respectively). This is in agreement with previous research comparing trained adults across different age groups (Sial et al. 1996; Dubé et al. 2016). The decline in fitness levels is despite an apparent ability for aged adults to still evidence sufficient peripheral adaptation to exercise interventions (Coggan et al. 1992a; Sial et al. 1998). Dubé et al. (2016) suggested that this reduction in peak fitness is down to central factors rather

than peripheral due to similar skeletal muscle capillarization and mitochondrial oxidative capacity between trained aged and young adults. Despite these differences between age groups, previous research would suggest that regular endurance exercise enhances overall fitness and metabolic capacity relative to sedentary adults in aged adults (Amati et al. 2011).

There were no significant differences observed between age groups for blood metabolic markers during exercise. Both lactate and glycerol values increased during exercise in both groups compared to baseline values. Hagberg et al. (1988) previously reported no difference in blood lactate values between trained older and younger runners during 1 h of treadmill running at 70% $\dot{V}O_{2\max}$. It has also been found that following a 16-week endurance training intervention on older adults, there were no significant differences in glycerol kinetics during exercise compared to younger adults (Sial et al. 1998). These findings would suggest that aged adults regularly engaged in endurance training maintain the ability to regulate glycerol metabolism during exercise. Conversely, Sial et al. (1996) reported that aged adults displayed reduced glycerol and free fatty acid kinetics in comparison to a young control group matched for anthropometric variables (e.g. lean body mass). However, the aged subjects used in this study were of a lower fitness level compared to the present study ($\dot{V}O_{2\max} = 31.4$ vs. 41.1 ml kg⁻¹ min⁻¹). Therefore, this suggests that differences between age groups may relate to the level of fitness attained by aged adults later in life.

There are several limitations that must be noted relating to the present study. First, the type and intensity of the exercise protocol used in the present study (i.e. 60% of $\dot{V}O_{2\max}$) was chosen as it has been previously shown to produce peak fat oxidation levels in aged adults (Bassami et al. 2007). This does not account for individual variation and the difference with the younger group. However, controlling the intensity in this way allows direct comparison of metabolic variables across age groups. The exercise protocol was only limited to 40 min to allow sufficient time for changes in metabolism during exercise. It must be noted that longer duration protocols have been used previously when investigating the metabolic response during exercise (MacLaren et al. 1999). Therefore, the findings from this study are limited to type and duration of protocol utilised. In addition, we chose to compare the two age groups relative to their own individual $\dot{V}O_{2\max}$ values (i.e. at 60% $\dot{V}O_{2\max}$). Future work should investigate the same relative and absolute exercise intensities to tease out the ageing effects and the intensity of effort effects. The sample size of this study was relatively small per group and it must be acknowledged that further work with larger sample sizes would further enhance this research area in the future.

In conclusion, the results from the present study demonstrate that trained aged adults display lower rates of

carbohydrate oxidation despite similar rates of fat oxidation when exercising at the same relative intensity. The effects of exercise intensity per se probably accounts for the major difference reported here. Likewise, glucose utilisation rates were significantly reduced in trained aged adults during steady-state exercise. Trained aged adults displayed a reduced insulin response to hyperglycaemia during exercise although this was not deemed statistically significant. There were no age-related differences in NEFA, glycerol and lactate concentrations between both aged groups. As trained aged adults appear to have a reduced ability to oxidise and utilise exogenous carbohydrate during exercise, it would be suggested that the use of carbohydrate supplements (e.g. energy gels and drinks) must be taken with caution. Overuse of such supplements may lead to an increase in fat mass and subsequently lead to aged-related health issues in the future.

Author contributions JJM, AH and MB carried out the majority of the study data collection. SCW and ITC carried out the hyperglycaemic and saline clamp technique on all participants and provided medical supervision throughout the study. DPM and DD conceived and developed the original study idea and supervised the project. JJM led the write up of the study manuscript, with input from all authors. All authors read and approved the manuscript.

References

- Abumrad N, Rabin D, Diamond M, Lacy W (1981) Use of a heated superficial hand vein as an alternative site for the measurement of amino acid concentrations for the study of glucose and alanine kinetics in man. *Metabolism* 30:936–940. [https://doi.org/10.1016/0026-0495\(81\)90074-3](https://doi.org/10.1016/0026-0495(81)90074-3)
- Amati F, Dube JJ, Alvarez-Carnero E et al (2011) Skeletal muscle triglycerides, diacylglycerols, and ceramides in insulin resistance: another paradox in endurance-trained athletes? *Diabetes* 60:2588–2597. <https://doi.org/10.2337/db10-1221>
- Bassami M, Ahmadizad S, Doran D, MacLaren DPM (2007) Effects of exercise intensity and duration on fat metabolism in trained and untrained older males. *Eur J Appl Physiol* 101:525–532. <https://doi.org/10.1007/s00421-007-0523-7>
- Biensø RS, Olesen J, Gliemann L et al (2015) Effects of exercise training on regulation of skeletal muscle glucose metabolism in elderly men. *J Gerontol Ser A* 70:866–872. <https://doi.org/10.1093/geronol/glv012>
- Bloem CJ, Chang AM (2008) Short-term exercise improves beta-cell function and insulin resistance in older people with impaired glucose tolerance. *J Clin Endocrinol Metab* 93:387–392. <https://doi.org/10.1210/jc.2007-1734>
- Broughton D, Taylor R (1991) Deterioration of glucose tolerance with age: the role of insulin resistance. *Age Ageing* 20:221–225
- Coggan A, Spina R, King D et al (1992a) Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. *J Appl Physiol* 72:1780–1786
- Coggan AR, Spina RJ, King DS et al (1992b) Histochemical and enzymatic comparison of the gastrocnemius muscle of young and elderly men and women. *J Gerontol Biol Sci* 47:B71–B76. <https://doi.org/10.1093/GERONJ/47.3.B71>
- Colberg SR, Sigal RJ, Yardley JE et al (2016) Physical activity/exercise and diabetes: a position statement of the american diabetes association. *Diabetes Care* 39:2065–2079. <https://doi.org/10.2337/dc16-1728>
- Cox JH, Cortright RN, Dohm GL, Houmard JA (1999) Effect of aging on response to exercise training in humans: skeletal muscle GLUT-4 and insulin sensitivity. *J Appl Physiol* 86:2019–2025
- Defronzo R, Tobin J, Andres R (1979) Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol Endocrinol Metab* 237:E214–E223
- Dela F, Ploug T, Handberg A et al (1994) Physical training increases muscle GLUT4 protein and mRNA in patients with NIDDM. *Diabetes* 43:862–865
- DiPietro L, Dziura J, Yeckel CW, Neufer PD (2006) Exercise and improved insulin sensitivity in older women: evidence of the enduring benefits of higher intensity training. *J Appl Physiol* 100:142–149. <https://doi.org/10.1152/jappphysiol.00474.2005>
- Douglas G (1911) A method for determining the total respiratory exchange in man. *J Physiol* 42:17–18
- Dube JJ, Amati F, Stefanovic-Racic M et al (2008) Exercise-induced alterations in intramyocellular lipids and insulin resistance: the athlete's paradox revisited. *AJP Endocrinol Metab* 294:E882–E888. <https://doi.org/10.1152/ajpendo.00769.2007>
- Dubé J, Broskey N, Despines A et al (2016) Muscle characteristics and substrate energetics in lifelong endurance athletes. *Med Sci Sports Exerc* 48:472–480. <https://doi.org/10.1007/128>
- Evans EM, Racette SB, Peterson LR et al (2005) Aerobic power and insulin action improve in response to endurance exercise training in healthy 77–87 year olds. *J Appl Physiol* 98:40–45. <https://doi.org/10.1152/jappphysiol.00928.2004>
- Frayn KN (1983) Calculation of substrate oxidation rates in vivo from gaseous exchange. *J Appl Physiol* 55:628–634. <https://doi.org/10.1152/jappphysiol.1983.55.2.628>
- Hagberg JM, Seals DR, Yerg JE et al (1988) Metabolic responses to exercise in young and older athletes and sedentary men. *J Appl Physiol* 65:900–908
- Jessen N, Pold R, Buhl ES et al (2003) Effects of AICAR and exercise on insulin-stimulated glucose uptake, signaling, and GLUT-4 content in rat muscles. *J Appl Physiol* 94:1373–1379. <https://doi.org/10.1152/jappphysiol.00250.2002>
- Kern M, Wells J, Stephens J, Elton C (1990) Insulin responsiveness in skeletal muscle is determined by glucose transporter (Glut4) protein level. *Biochem J* 270:397–400
- Kim HJ, Lee JS, Kim CK (2004) Effect of exercise training on muscle glucose transporter 4 protein and intramuscular lipid content in elderly men with impaired glucose tolerance. *Eur J Appl Physiol* 93:353–358. <https://doi.org/10.1007/s00421-004-1214-2>
- Lanza IR, Short DK, Short KR et al (2008) Endurance exercise as a countermeasure for aging. *Diabetes* 57:2933–2942. <https://doi.org/10.2337/db08-0349>
- Lee SW, Youm Y, Lee WJ et al (2015) Appendicular skeletal muscle mass and insulin resistance in an elderly Korean population: the Korean social life, health and aging project-health examination cohort. *Diabetes Metab J* 39:37–45. <https://doi.org/10.4093/dmj.2015.39.1.37>
- Levy J, Matthews D, Hermans M (1998) Correct homeostasis model assessment (HOMA) evaluation uses the computer program. *Diabetes Care* 21:2191–2192
- MacLaren DP, Reilly T, Campbell IT, Hopkin C (1999) Hormonal and metabolic responses to maintained hyperglycemia during prolonged exercise. *J Appl Physiol* 87:124–131
- Meneilly G, Tessier D (2001) Diabetes in elderly adults. *J Gerontol A Biol Sci Med Sci* 56:M5–M13
- Meredith CN, Frontera WR, Fisher EC et al (1989) Peripheral effects of endurance training in young and old subjects. *J Appl Physiol* 66:2844–2849

- Morris E, Rivas D, Fielding R (2010) Increased intramuscular triglycerides are associated with increased AMPK alpha1 and cleaved SREBP1 in aged skeletal muscle. *FASEB J* 24:lb677–lb677
- Pratley RE, Hagberg JM, Rogus EM, Goldberg AP (1995) Enhanced insulin sensitivity and lower waist-to-hip ratio in master athletes. *Am J Physiol* 268:E484–E490
- Qiang W, Weiqiang K, Qing Z et al (2007) Aging impairs insulin-stimulated glucose uptake in rat skeletal muscle via suppressing AMPKalpha. *Exp Mol Med* 39:535–543. <https://doi.org/10.1038/emm.2007.59>
- Ryan A (2000) Insulin resistance with aging: effects of diet and exercise. *Sport Med* 30:327–346
- Scheen A (1997) Non-insulin-dependent diabetes mellitus in the elderly. *Baillieres Clin Endocrinol Metab* 11:389–406
- Short KR, Vittone JL, Bigelow ML et al (2003) Impact of aerobic exercise training on age-related changes in insulin sensitivity and muscle oxidative capacity. *Diabetes* 52:1888–1896. <https://doi.org/10.2337/diabetes.52.8.1888>
- Sial S, Coggan AR, Carroll R et al (1996) Fat and carbohydrate metabolism during exercise in elderly and young subjects. *Am J Physiol Endocrinol Metab* 271:E983–E989
- Sial S, Coggan AR, Hickner RC, Klein S (1998) Training-induced alterations in fat and carbohydrate metabolism during exercise in elderly subjects. *Am J Physiol Endocrinol Metab* 274:E785–E790
- Vermeulen A, Goemaere S, Kaufman JM (1999) Testosterone, body composition and aging. *J Endocrinol Invest* 22:110–116
- Vigelsø A, Gram M, Dybbøe R et al (2016) The effect of age and unilateral leg immobilization for 2 weeks on substrate utilization during moderate-intensity exercise in human skeletal muscle. *J Physiol* 594:2339–2358. <https://doi.org/10.1113/JP271712>
- Winder WW, Hardie DG (1999) AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes. *Am J Physiol* 277:E1–E10