



Adult vaccination for pneumococcal disease: a comparison of the national guidelines in Europe

C. Bonnavé¹ · D. Mertens¹ · W. Peetermans² · K. Cobbaert³ · B. Ghesquiere³ · M. Deschodt^{4,5} · J. Flamaing^{1,4}

Received: 22 October 2018 / Accepted: 10 January 2019 / Published online: 18 February 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Pneumococcal disease constitutes a major global health problem. Adults aged over 50 years and younger adults with specific chronic health conditions are at risk for invasive pneumococcal disease, associated with substantial morbidity and mortality. In Europe, two vaccine types are used in adults for pneumococcal immunization: pneumococcal polysaccharide vaccine (PPV23) and pneumococcal conjugate vaccine (PCV13). To provide an overview and to compare the national guidelines for pneumococcal immunization for adults in Europe. In November 2016, national guidelines on pneumococcal vaccination for adults of 31 European countries were obtained by Google search, the website of European Centre for Disease Prevention and Control, and contacting public health officials. In our analysis, we distinguished between age-based and risk-based guidelines. In October 2017, we used the same method to retrieve guideline updates. We observed great variability regarding age, risk groups, vaccine type, and use of boosters. In age-based guidelines, vaccination is mostly recommended in adults aged over 65 years using PPV23. Boosters are generally not recommended. An upper age limit for vaccination is reported in three countries. In the immunocompromised population, vaccination with both vaccines and administration of a booster is mostly recommended. In the population with chronic health conditions, there is more heterogeneity according vaccine type, sequence, and administration of boosters. Asplenia is the only comorbidity for which all countries recommend vaccination. The great variability in European pneumococcal vaccination guidelines warrants European unification of the guidelines for better control of pneumococcal disease.

Keywords Pneumococcal vaccination · Adults · Europe · Recommendations · PCV13 · PPV23

C. Bonnavé and D. Mertens shared first authorship.

M. Deschodt and J. Flamaing shared last authorship.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10096-019-03485-3>) contains supplementary material, which is available to authorized users.

✉ J. Flamaing
Johan.flamaing@uzleuven.be

¹ Department of Geriatrics, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium

² Department of Internal Medicine, University Hospitals Leuven, Leuven, Belgium

³ Department of Geriatrics, AZ Delta Roeselare, Roeselare, Belgium

⁴ Department of Chronic Diseases, Metabolism and Aging, KU Leuven, Leuven, Belgium

⁵ Department of Public Health, University of Basel, Basel, Switzerland

Introduction

Pneumococcal infections constitute a major global health problem. It leads to a number of diseases like otitis media, pneumonia, meningitis, and bacteremia [1]. The most frequent manifestation of pneumococcal disease in adults is pneumonia, where *Streptococcus pneumoniae* is responsible for up to 60% of the cases of community-acquired pneumonia [2].

In the USA, data of the Centers for Disease Control and Prevention showed 17 cases of invasive pneumococcal disease (IPD) per 100,000 persons in all age groups [3]. In England and Wales, an incidence of IPD of 7 per 100,000 persons in all age groups is reported and an incidence of 21 per 100,000 in persons aged 65 years and older [2].

Risk groups for developing pneumococcal infections and complications are children aged < 2 years, adults aged 50 years and older, and persons with specific comorbidities [1]. Case fatality rate of IPD may reach 15–20% in adults and even 30–40% in elderly [4, 5].

The pneumococcus is a common human commensal in the nasopharynx, with asymptomatic carriage reported from 5 to 93% in all age groups. Carriage diminishes greatly with age: whereas up to 60% of the school-aged children may be colonized, only 5–10% of adults and < 5% of nursing home residents are [2, 3, 6, 7]. *S. pneumoniae* is classified by the presence of capsular polysaccharides, with at least 93 different serotypes identified. Up to 60% of the IPD infections (in Belgium even up to 83%) are caused by serotypes covered by PPV23 [5, 8].

Currently, two different pneumococcal vaccines are approved for use in adults: the polysaccharide vaccine containing 23 different capsular polysaccharides (PPV23) and the conjugate vaccine containing 13 serotypes of *S. pneumoniae* (PCV13) [9]. The PPV23 triggers a T cell-independent immune response with serotype-specific antibody formation. Although the PPV23 offers an extended serotype coverage, much of the antibody response after vaccination is transitory, resulting in a decline in protection after 2–4 years. Also, immunogenicity is probably lower among young children, elderly, and immunocompromised persons [10, 11]. PCV13 triggers a T cell dependent immune response, with formation of both serotype-specific antibodies and B memory cells, resulting in a better immune response [10]. As PCV13 stimulates the formation of mucosal antibodies and thus prevents acquisition of nasal carriage of vaccine types, a reduced spread by children of the serotypes included in PCV13 was observed, thereby protecting unvaccinated individuals in all age groups (“herd protection effect”) [3].

As the global burden of pneumococcal disease is high, most European countries have implemented national vaccination guidelines. However, the immunization guidelines vary greatly among countries in Europe in terms of age and risk groups for vaccination [1]. The aim of this review is to provide an overview and compare the national guidelines in Europe for pneumococcal immunization in adults.

Methods

Search strategy

A twofold search strategy was conducted in November 2016 to identify the national guidelines on pneumococcal vaccination for adults aged ≥ 18 years in 31 European countries. Immunization schedules for patients with hematological stem cell transplantation were excluded, as this was outside the scope of our review. Firstly, we used the Vaccine Schedule on the website of the European Centre for Disease Prevention and Control (ECDC; www.ecdc.europa.eu) and searched the reference websites provided for each country to obtain their official immunization guidelines. If no vaccination guidelines could be retrieved for a country by

using this method, we performed a Google search (“pneumococcal vaccination guidelines adult [Name of the country],” translated in the official language of this country by using Google Translate). In case both search strategies failed, we contacted the public health officials of this country (as was the case for Bulgaria, Cyprus, Croatia and Estonia). In October 2017, we used the same method to retrieve guideline updates.

Data extraction

To translate the vaccination guidelines from their original language, Google Translate or interpreters were used.

Data were extracted in Word and Excel tables to synthesize information regarding study characteristics, vaccine type, sequence of vaccination, and use of boosters (supplementary Tables 1–3 and Tables 1 and 2). In our analysis, we distinguished between age-based and risk-based guidelines. In the risk-based guidelines, we beheld three different risk groups: patients with specific chronic health conditions, an immunocompromised group, and patients with other predisposing risk factors for pneumococcal disease. The risk factor-based guidelines concern all adults aged 18 years and older, unless otherwise specified in supplementary Table 3.

Data synthesis and analysis

For the age-based guidelines and the different risk factors, we calculated the absolute numbers and percentages of vaccine and booster recommendation and vaccine type used (Tables 1 and 2). This data extraction was performed by one researcher (CB) and verified by a second researcher (DM).

Results

Identification of national vaccination guidelines

We were able to identify guidelines for adult pneumococcal immunization for 28 of the 31 reviewed European countries (90%). No guidelines were found for Cyprus, Latvia, and Romania. In all of the 28 countries, risk-based guidelines were available, while age-based guidelines were available in 27 countries (96%). No age-based guidelines were found for Croatia. More than 90% of these countries (93%; $n = 26$) updated their guidelines in 2014 or later.

In both age- and risk-based guidelines, a great variability was observed regarding to the age group eligible for vaccination and the type of recommended vaccine (see supplementary Tables 2–3 and Tables 1 and 2).

Table 1 Resume age-based guidelines

Age-based guidelines (healthy population)— <i>n</i> = 27	Number	Percent	Countries
Recommended starting age of vaccination			
≥ 50 years	5	18.5	Austria [12], Estonia [13], Hungary [14], Lithuania [15, 16], Poland [17]
≥ 59 years	1	3.7	Slovakia [18]
≥ 60 years	5	18.5	Bulgaria [19], Germany [20], Iceland [21], the Netherlands [22, 23], Spain [24]
≥ 65 years	13	48.1	Belgium [25], Czech Republic [26], Denmark [27], Finland [28], Greece [29], Ireland [30], Italy [31], Luxembourg [32], Malta [33], Norway [34], Slovenia [18], Sweden [35], the UK [36]
Vaccination not recommended	3	11.1	France [37], Switzerland [38], Portugal [39]
Upper limit of age for vaccination reported:			
	3	11.1	Belgium [25], Luxembourg [32], the Netherlands [22, 23]
Vaccine type recommended in primary vaccination			
PCV13	1	3.7	Estonia [13]
PPV23	11	40.7	Bulgaria [19], Germany [20], Iceland [21], Ireland [30], Malta [33], the Netherlands [22, 23], Norway [34], Poland [17], Spain [24], Sweden [35], the UK [36]
PCV13 and PPV23	7	25.9	Austria [12], Belgium [25], Czech Republic [26], Greece [29], Hungary [14], Luxembourg [27]
Several options possible	4	14.8	Denmark [28], Finland [15], Lithuania [16, 18], Slovenia [40]
Recommended but vaccine type not specified	1	3.7	Slovakia [25]
Sequence and recommended interval if both vaccines recommended (primary vaccination)			
PCV13 first	9	100	
- Interval between the two vaccines ≥ 8 weeks	5	55.6	Belgium [27], Denmark [14], Hungary [32], Luxembourg [31], Italy [12]
- Interval between the two vaccines ≥ 1 year	3	33.3	Austria [28], Finland [29], Greece [26]
- Interval not specified	1	11.1	Czech Republic [26]
PPV23 first	0	0	
Administration of booster			
Booster recommended	5	18.5	Czech Republic [26], Denmark [27], Estonia [13], the Netherlands [22, 23], Norway [34]
No booster recommended	12	44.4	Austria [12], Belgium [25], France [37], Germany [20], Iceland [21], Italy [31], Switzerland [38], Luxembourg [32], Malta [33], Portugal [39], Sweden [35], the UK [36]
Booster to consider	5	18.5	Bulgaria [19], Ireland [30], Lithuania [15, 16], Slovenia [18], Spain [24]
Not reported	5	18.5	Finland [28], Greece [29], Hungary [14], Poland [17], Slovakia [40]
Type of vaccine used for booster			
PCV13	10	100	Bulgaria [19], Czech Republic [26], Denmark [27], Estonia [13], Ireland [30], Lithuania [15, 16], the Netherlands [22, 23], Norway [34], Slovenia [18], Spain [47]
PPV23	0	0	

n number of countries, *PPV23* pneumococcal polysaccharide vaccine, *PCV13* pneumococcal conjugate vaccine. Sources for vaccine recommendations per country: see references 20 to 49

Age-based guidelines

France, Switzerland and Portugal do not recommend systematical vaccination in healthy adults. The remaining European countries with age-based guidelines (*n* = 24; 89%)

advise vaccinating healthy adults with a minimum age of 50 years (*n* = 5; 18.5%), 59 years (*n* = 1; 4%), 60 years (*n* = 5; 18.5%), or 65 years (*n* = 13; 48%). An upper age limit for vaccination was mentioned in Belgium and Luxembourg (both 85 years) and in the Netherlands (75 years).

Table 2 Resume risk factor-based guidelines

	Vaccination recommended		Vaccination to consider		Vaccine type								Booster							
	n	%	n	%	PCV13		PPV23		PCV13 + PPV23		Several options		Recommended		No booster		To consider		Not reported	
					n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Chronic disease																				
Chronic kidney disease	25	89.3	2	7.1	1	3.7	7	25.9	14	51.9	5	18.5	10	37	7	25.9	4	14.8	6	22.2
Chronic lung disease	25	89.3	2	7.1	1	3.7	8	29.6	12	44.4	6	22.2	8	29.6	9	33.3	4	14.8	6	22.2
Chronic heart disease	24	85.7	2	7.1	1	3.8	7	26.9	12	46.2	6	23.1	7	26.9	9	34.6	4	15.4	6	23.1
Chronic liver disease	22	78.6	2	7.1	1	4.2	7	29.2	11	45.8	5	20.8	7	29.2	8	33.3	4	16.7	5	20.8
Diabetes mellitus	21	75	2	7.1	1	4.3	8	34.8	9	39.1	5	21.7	6	26.1	8	34.8	4	17.4	5	21.7
Immunocompromising conditions																				
Asplenia	28	100	0	0	1	3.6	4	14.3	18	64.3	5	17.9	16	57.1	3	10.7	2	7.1	7	25
Immunodeficiency	23	82.1	2	7.1	1	4	2	8	16	64	6	24	10	40	5	20	3	12	7	28
HIV	19	67.9	1	3.6	1	5	4	20	12	60	3	15	7	35	6	30	3	15	4	20
Hematological malignancy	18	64.3	1	3.6	1	5.3	3	15.8	11	57.9	4	21.1	7	36.8	6	31.6	3	15.8	3	15.8
Immunosuppressive therapy	18	64.3	2	7.1	1	5	3	15	12	60	4	20	7	35	7	35	2	10	4	20
Solid organ transplant	17	60.7	1	3.6	1	5.6	2	11.1	11	61.1	4	22.2	8	44.4	5	27.8	3	16.7	2	11.1
Parenchymatous malignancy	9	32.1	2	7.1	0	0	1	9.1	8	72.7	2	18.2	5	45.5	3	27.3	2	18.2	1	9.1
Other risk factors for pneumococcal disease																				
CSF leak	25	89.3	0	0	1	4	5	16	16	64	4	16	10	40	7	28	3	12	5	20
Cochlear implant	23	82.1	1	3.6	1	4.2	4	16.7	15	62.5	4	16.7	10	41.7	7	29.2	2	8.3	5	20.8
Alcoholism	12	42.9	2	7.1	0	0	5	35.7	6	42.9	3	21.4	2	14.3	5	35.7	3	21.4	4	28.6
Smoking	10	35.7	1	3.6	0	0	4	36.4	6	54.5	1	9.1	3	27.3	4	36.4	2	18.2	2	18.2
Occupational risk	6	21.4	1	3.6	0	0	3	42.9	3	42.9	1	14.3	2	28.6	2	28.6	2	28.6	1	14.3
Institutionalized patients	5	17.9	0	0	0	0	2	40	0	0	3	60	1	20	0	0	1	20	3	60
Neuromuscular disease	5	17.9	0	0	0	0	3	60	1	20	1	20	1	20	4	80	0	0	0	0
Medical history of IPD	3	10.7	2	7.1	0	0	0	0	4	80	1	20	3	60	1	20	1	20	0	0
BMI > 40 kg/m ²	2	7.1	0	0	1	50	1	50	0	0	0	0	0	2	100	0	0	0	0	0
Celiac disease	0	0	2	7.1	0	0	0	0	1	50	1	50	0	0	1	50	1	50	0	0
Homeless patients	0	0	2	7.1	0	0	0	0	1	50	1	50	0	0	1	50	1	50	0	0
Drug addicts	0	0	2	7.1	0	0	0	0	1	50	1	50	0	0	1	50	1	50	0	0

BMI body mass index, CSF cerebrospinal fluid, HIV human immunodeficiency virus, IPD invasive pneumococcal disease, n number of countries, PPV23 pneumococcal polysaccharide vaccine, PCV13 pneumococcal conjugate vaccine

For primary vaccination of the healthy adult, four out of ten countries recommends the use of PPV23 only ($n = 11$; 40.7%). One country (Estonia) recommends the use of PCV13 only, while seven other countries (26%) recommended both PCV13 and PPV23 for primary vaccination. In four countries (15%), the type of vaccine recommended depends on the preference of the treating physician (Finland, Denmark, Lithuania, Slovenia). In Slovakia, the recommended vaccine type is not specified.

If both vaccines are recommended for primary vaccination, which is the case in nine countries, PCV13 is always administered first. The recommended interval between the two

vaccinations (PCV13 and PPV23) differed between ≥ 8 weeks in Belgium, Denmark, Hungary, Luxembourg, and Italy and ≥ 1 year in Finland, Greece, and Austria. Slovakia recommends primary vaccination but did not specify the recommended vaccine type.

Twelve countries (44.4%) do not advise a booster in healthy adults. Five national guidelines recommend a booster (18.5%), and five countries consider a booster under certain conditions (e.g. patients aged > 65 years, who received their first vaccine at least 5 years ago when < 65 years). In five guidelines, administration of boosters was not reported (18.5%). If a booster is given, all countries use PPV23.

Risk-based guidelines

The *chronic disease group* includes vaccination guidelines for chronic kidney ($n = 25$; 89%), lung ($n = 25$, 89%), heart ($n = 24$, 86%), and liver diseases ($n = 22$, 79%) and diabetes mellitus ($n = 21$, 75%). Mostly PCV13 and PPV23 or only PPV23 are recommended (see Table 2). PCV13 only is rarely used. A booster is recommended in 26 to 37% of the countries (depending on the underlying health condition), is considered in about 15% of the countries, and is not recommended in 26 to 35% of the countries, respectively. In one fifth of the countries, no booster guidelines were reported.

In the immunocompromised population, vaccination is recommended in the majority of the countries. Asplenia is the only condition for which all countries advise vaccination. Immunization of patients with immunodeficiency is recommended in 23 countries. About 65% of the countries recommend vaccination for HIV patients, hematologic malignancies, patients taking immunosuppressive drugs, and patients after solid organ transplant. Thirty-two percent of these countries recommend pneumococcal immunization in patients with parenchymatous malignancies. The majority of the countries that recommend vaccinations in these immunocompromised patients (58 to 72% depending on the underlying condition) would vaccinate with PCV13 and PPV23. Ten to 20% of the countries recommend immunization with PPV23 only. Again, PCV13 only is rarely recommended. A booster is recommended in 40% of the countries that report risk-based guidelines (depending on the underlying condition). A booster is recommended for patients with asplenia in six out of ten countries. In up to 25% of the risk-based guidelines, no booster guidelines are reported.

The third category includes *various risk factors* for pneumococcal disease. For patients with cerebrospinal fluid leak or a cochlear implant, most countries (89% and 82% respectively) recommend vaccination, mostly with PCV13 and PPV23 (64% and 62.5% respectively). Four out of ten countries advise a booster in this risk group. About 40% of the countries recommend vaccination for smokers and alcoholics, and two out of ten would vaccinate patients with occupational risk (such as welders), institutionalized patients, and patients with neuromuscular disease. In these five previous risk groups, half of the countries vaccinate with PPV23 only, and roughly a quarter recommends administration of a booster. For patients with previous invasive pneumococcal disease or pneumococcal pneumonia, vaccination is only recommended in three countries. Sweden and Switzerland are the only countries that would vaccinate patients with BMI > 40 kg/m². Norway and the Netherlands would consider pneumococcal immunization for patients with celiac disease, homeless patient groups, and drug addicts.

Discussion

In this study, comparing the European guidelines for pneumococcal vaccination of adult patients at elevated risk for pneumococcal infection, we observed a great variability in vaccine recommendation despite increasing evidence of pneumococcal vaccine being effective in older adults.

Although no concluding evidence about the efficacy of PPV23 was reported in a Cochrane review (probably due to a lack of statistical power), the benefit of PPV23 was more pronounced when limiting the analysis to trials that assessed the incidence of vaccine-serotype disease [41]. Also, two recent meta-analyses showed PPV23 vaccine efficacy [9, 42]. Falkenhorst et al. found a significant vaccine efficacy of 73% for IPD and of 64% for pneumococcal pneumonia in patients aged ≥ 60 years [9]. Kraicer-Melamed et al. described a 50% vaccine efficacy for IPD in patients aged ≥ 50 years [42]. The authors state that the use of recent and high quality data led to this results, contrary to the Cochrane review.

Concerning PCV13, the most important study is the CAPITA trial (2015). This placebo-controlled study analyzed the vaccine efficacy of PCV13 in 85,000 patients aged ≥ 65 years. A vaccine efficacy of 45.6%, 45%, and 75% in the prevention of respectively vaccine-type community acquired pneumonia, non-bacteremic and non-invasive community-acquired pneumonia, and invasive pneumococcal disease was demonstrated. Efficacy persisted throughout the trial (mean follow-up 3.97 years) suggesting that PCV13 should be part of the vaccination program in older adults [43]. However, the results of the CAPITA trial should be interpreted with caution as patients with immunodeficiency were excluded and as it took place before implementation of childhood pneumococcal immunization in the Netherlands. The latter thereby ignores the herd effect, and the study possibly overestimates the efficacy of PCV13 [9].

The CAPITA trial also examined the long-term immunogenicity of PCV13 in pneumococcal vaccine-naïve older adults, demonstrating high antibody titers during the first 2 years following vaccination after a single dose of PCV13. In adults aged ≥ 80 years, weaker responses were observed, but antibody levels remained above baseline [14].

Cost-effectiveness analysis is often issued by health authorities to substantiate inclusion of pneumococcal vaccination in the adult vaccination program with (partial) reimbursement of the costs. In 2012, the Infection Prevention Institute in Poland performed a cost-effectiveness analysis on the implementation of a public vaccination program with PPV23 only in the elderly aged ≥ 65 years. When taking into account the medical and economic burden of pneumococcal disease of that time, presuming a PPV23-vaccine efficacy of 64% for IPD in elderly and assuming a 50% reimbursement of the PPV23 vaccine from their national health fund, they concluded that the use of PPV23 in healthy elderly, would be highly cost-effective [4].

A cost-effectiveness analysis in the UK (2015) stated that adding PCV13 to the existing vaccination program with PPV23 in healthy elderly would substantially raise costs of immunization programs and would not be cost-effective [44].

Despite some disadvantages of PPV23 (lack of development of an immune memory, reduced immunogenicity in the elderly), and some conflicting results on efficacy of this vaccine as mentioned above, the more recent systematic reviews and this cost-effectiveness analysis provide arguments for a vaccination strategy in the healthy elderly that includes PPV23.

Furthermore, as there is to our knowledge no study comparing the efficacy of PCV13 and PPV23 in elderly patients, and considering the economic aspect and weaknesses of the CAPITA trial mentioned above, updating vaccination guidelines and recommending the use of both PCV13 and PPV23 for the healthy elderly (as done in the USA by the Advisory Committee on Immunization Practices (ACIP) in 2014) should be done cautiously [45]. More studies on pneumococcal vaccination of the healthy elderly are needed, including a new efficacy trial for PCV13 that takes into account the herd protection effect.

As vaccine-induced immunity wanes after a few years, considering the lack of development of an immune memory, a booster vaccine can be considered. The efficacy of boosters however is uncertain, mainly because a state of hyporesponsiveness or immune tolerance is described following revaccination with PPV23. This especially seems relevant if a booster is administered within 5 years after primary vaccination. Musher et al. demonstrated that persons who received PPV23 within 1 year almost did not respond to revaccination, and those who had received PPV23 within 1 to 5 years responded less well than those who had no previous or more remote vaccination [46, 47]. Another systematic review, comparing the effectiveness, immunogenicity, and safety of revaccination with PPV23 in persons aged ≥ 50 years, demonstrated a reduced antibody response during the first months after revaccination (thereafter, immunogenicity of primary and repeated vaccination was comparable). There were no major safety concerns regarding revaccination [48]. All of the studies mentioned above, however, are immunogenicity studies that evaluate antibody responses after immunization, therefore lacking clinical endpoints or data about clinical effectiveness. As the available evidence concerning revaccination with PPV23 is not unambiguous, neither are the guidelines concerning the use of boosters in healthy elderly.

The strength of this article lies in the comprehensive overview of 31 European countries and the USA, which is to our knowledge the largest comprehensive overview up to present. A limitation of the study includes the use of Google Translate or interpreters and thus the lack of a formal forward-backward translation. Although this could have led to incorrect translations or interpretations, the immunization guidelines were very straightforward, often written in bullet points.

Conclusion

Pneumococcal vaccination has greatly reduced incidence of vaccine-type pneumococcal morbidity and mortality. There is great variability in European pneumococcal vaccination guidelines for adults, regarding age, risk groups, and vaccine type. Most countries did not have clear recommendations about the administration of boosters. For ease of implementation, European unification of the immunization guidelines is needed. This remains difficult despite increasing evidence on pneumococcal vaccine effectiveness in elderly.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval Not applicable.

Informed consent Not applicable

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Castiglia P (2014) Recommendations for pneumococcal immunization outside routine childhood immunization programs in Western Europe. *Adv Ther* 31(10):1011–1044
2. Chalmers JD, Campling J, Dicker A, Woodhead M, Madhava H (2016) A systematic review of the burden of vaccine preventable pneumococcal disease in UK adults. *BMC Pulm Med* 16(1):77
3. Yildirim I, Shea KM, Pelton SI (2015) Pneumococcal disease in the era of pneumococcal conjugate vaccine. *Infect Dis Clin N Am* 29(4):679–697
4. Grzesiowski P, Aguiar-Ibanez R, Kobryn A, Durand L, Puig PE (2012) Cost-effectiveness of polysaccharide pneumococcal vaccination in people aged 65 and above in Poland. *Human Vaccines Immunotherapeutics* 8(10):1382–1394
5. Verhaegen J, Flamaing J, De Backer W, Delaere B, Van Herck K, Surmont F et al (2014) Epidemiology and outcome of invasive pneumococcal disease among adults in Belgium, 2009–2011. *Euro Surveill* 19(31):14–22
6. Chidiac C (2012) Pneumococcal infections and adult with risk factors. *Medecine et maladies infectieuses* 42(10):517–524
7. Flamaing J, Peetermans WE, Vandeven J, Verhaegen J (2010) Pneumococcal colonization in older persons in a nonoutbreak setting. *J Am Geriatr Soc* 58. United States. p. 396–8
8. Mirsaiedi M, Ebrahimi G, Allen MB, Aliberti S (2014) Pneumococcal vaccine and patients with pulmonary diseases. *Am J Med* 127(9):886 e1–886 e8
9. Falkenhorst G, Remschmidt C, Harder T, Hummers-Pradier E, Wichmann O, Bogdan C (2017) Effectiveness of the 23-valent pneumococcal polysaccharide vaccine (PPV23) against pneumococcal disease in the elderly: systematic review and meta-analysis. *PLoS One* 12(1):e0169368

10. Papadatou I, Spoulou V (2016) Pneumococcal vaccination in high-risk individuals: are we doing it right? *Clin Vaccine Immunol* 23(5):388–395
11. van Deursen AMM, van Houten MA, Webber C, Patton M, Scott DA, Patterson S et al (2017) Immunogenicity of the 13-valent pneumococcal conjugate vaccine in older adults with and without comorbidities in the Community-Acquired Pneumonia Immunization Trial in Adults (CAPiTA). *Clin Infect Dis* 65(5):787–795
12. Rendi-Wagner P, Paulke-Korinek M. Impfplan Österreich 2017. Wien: Bundesministerium für Gesundheit und Frauen (BMGF); 2017
13. Estonian Society for Infectious Diseases (2014) Immuniseerimiskava väliselt vaksineeritavate isikute sihtrühmad ja nendele näidustatud vaktsiinid. Estonia
14. Mészner Z, Jelenik Z, Kulcsár A. Felnőttkori oltások ütemezésének ajánlása. Budapest: Nemzeti Egészségfejlesztési Intézet támogatásával; 2017
15. Suaugusiųjų skiepijimo rekomendacijos. Lithuania; 2013
16. A A. Suaugusiųjų rizikos grupių skiepijimo lietuvoje rekomendacijos. Vilnius 2015. p. 20
17. Posobkiewicz M. W sprawie Programu Szczepień Ochronnych na rok 2017. Poland: Ministra Zdrowia; 2017
18. zdravje NIzj. Priporočila za cepljenje odraslih in otrok starih 5 let in vec proti pnevmokoknum okuzbam. Slovenia; 2016
19. Sanofi Pasteur Europe. PNEUMO 23, solution injectable, Vaccin pneumococque polysaccharidique, Information de l'utilisateur. Lyon: Sanofi Pasteur; 2017
20. Robert Koch Institut (2017) Empfehlungen der Ständigen Impfkommission (STIKO) am Robert Koch-Institut - 2017/2018. Ständige Impfkommission (STIKO, Berlin
21. Directorate of Health. Leiðbeiningar um bólusetningu gegn pneumókókkum. Reykjavík; 2014
22. Nederland G (2018) Vaccinatie van ouderen tegen pneumokokken. Nederland: Gezondheidsraad Nederland
23. voor In-R, Milieu Ve (2017) LCI-richtlijn pneumokokkose—invasieve pneumokokkenziekte. Nederland
24. publico S. Vacunacion en adultos. Spain: Ministerio de sanidad y consumo; 2004
25. Hoge Gezondheidsraad. Advies Vaccinatie tegen Pneumococcen (2014). Brussel: Federale Overheidsdienst Volksgezondheid, Veiligheid van de Voedselketen en Leefmilieu; 2014
26. Česká vakcinologická společnost. Očkovací kalendář pro dospělé 2015. Czech Republic; 2015
27. Statens Serum Institut. Suggestions for pneumococcal vaccination outside the childhood vaccination program, Tularaemia (rabbit fever) after thick bites in Denmark. Denmark: Epi-News; 2014
28. Terveysten ja Hyvinvoinnin Laitos (2017) Kansallisen rokotusohjelman ulkopuoliset pneumokokkirokotukset. Helsinki
29. ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ ΓΕΝΙΚΗ Δ/ΝΣΗ ΔΗΜΟΣΙΑΣ ΥΓΕΙΑΣ & ΥΠΗΡΕΣΙΩΝ ΥΓΕΙΑΣ. Πρόγραμμα Εμβολιασμών Ενηλίκων 2017. Athens; 2017
30. Immunisation Guidelines for Ireland. Dublin: National Immunisation Advisory Committee of the Royal College of Physicians of Ireland; 2015
31. Direzione Generale della Prevenzione Sanitaria del Ministero della Salute. Piano Nazionale Prevenzione Vaccinale (PNPV) 2017-2019. Rome; 2017
32. Staub T, Biver A, Berthet F, Arendt V, Faber J, Federspiel C, et al. Vaccination contre le pneumocoque pour les populations à risque ou âgées de 65 ans et plus—Recommandations du conseil supérieur des maladies infectieuses. Luxembourg: Conseil supérieur des maladies infectieuses; 2015
33. Gauci C, Farrugia Sant'Angelo V, Attard Montalto S, Barbara C, Caruana P, Piscopo T, et al. (2011) The official guide to immunisations. Malta: Advisory Committee on immunisation policy
34. Norway F (2015) Pneumokokkvaksinosjon—Pneumokokkonjugatvaksine inngar i barnevaksinasjonsprogrammet. Norway
35. Sweden F (2016) Rekommendationer om pneumokokkvaccination till riskgrupper. Sweden
36. England PH (2017) The green book—immunisation against infectious disease—chapter 25—Pneumococcal. United Kingdom
37. Ministère des Affaires Sociales et de la Santé. Calendrier des vaccinations et recommandations vaccinales 2017. Paris: Direction générale de la Santé; 2017
38. (CFV) OfdlSpeCfplv. Plan de vaccination suisse 2017. Suisse: Confederation Suisse; 2017. Report No.: 2016-OEG-01
39. Programa Nacional de Vacinação—PNV 2017. Portugal: Direção de Serviços de Prevenção da Doença e Promoção da Saúde. Médicos e Enfermeiros do Sistema de Saúde; 2017
40. Republiky ÚVZS. Povinné a odporúčané očkovanie podľa dosiahnutého veku a rizika nákazy v Slovenskej republike na rok 2016. Slovakia: Ministerstva zdravotníctva; 2016
41. Moberley S, Holden J, Tatham DP, Andrews RM (2013) Vaccines for preventing pneumococcal infection in adults. *Cochrane Database Syst Rev*. (1):Cd000422
42. Kraicer-Melamed H, O'Donnell S, Quach C (2016) The effectiveness of pneumococcal polysaccharide vaccine 23 (PPV23) in the general population of 50 years of age and older: a systematic review and meta-analysis. *Vaccine* 34(13):1540–1550
43. Bonten MJ, Huijts SM, Bolkenbaas M, Webber C, Patterson S, Gault S et al (2015) Polysaccharide conjugate vaccine against pneumococcal pneumonia in adults. *N Engl J Med* 372(12):1114–1125
44. van Hoek AJ, Miller E (2016) Cost-effectiveness of vaccinating immunocompetent ≥ 65 year olds with the 13-valent pneumococcal conjugate vaccine in England. *PLoS One* 11(2):e0149540
45. Tomczyk S, Bennett NM, Stoecker C, Gierke R, Moore MR, Whitney CG et al (2014) Use of 13-valent pneumococcal conjugate vaccine and 23-valent pneumococcal polysaccharide vaccine among adults aged ≥ 65 years: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Morb Mortal Wkly Rep* 63(37):822–825
46. Musher DM, Rueda AM, Nahm MH, Graviss EA, Rodriguez-Barradas MC (2008) Initial and subsequent response to pneumococcal polysaccharide and protein-conjugate vaccines administered sequentially to adults who have recovered from pneumococcal pneumonia. *J Infect Dis* 198(7):1019–1027
47. Musher DM, Sampath R, Rodriguez-Barradas MC (2011) The potential role for protein-conjugate pneumococcal vaccine in adults: what is the supporting evidence? *Clin Infect Dis* 52(5):633–640
48. Remschmidt C, Harder T, Wichmann O, Bogdan C, Falkenhörst G (2016) Effectiveness, immunogenicity and safety of 23-valent pneumococcal polysaccharide vaccine revaccinations in the elderly: a systematic review. *BMC Infect Dis* 16(1):711