



Evaluation of Vascular Event Risk while on Long-term Anti-retroviral Suppressive Therapy [EVERLAST]: Protocol for a prospective observational study

Fred Stephen Sarfo^{a,b,*}, Michelle Nichols^c, Mulugeta Gebregziabher^c, Raelle Tagge^c, Shadrack Osei Asibey^b, Carolyn Jenkins^c, Bruce Ovbiagele^d

^a Kwame Nkrumah University of Science & Technology, Kumasi, Ghana

^b Komfo Anokye Teaching Hospital, Kumasi, Ghana

^c Medical University of South Carolina, Charleston, USA

^d Department of Neurology, University of California, San Francisco, USA

ARTICLE INFO

Keywords:

HIV
Cardiovascular risk
Stroke
Neurocognitive dysfunction
Hypertension
Sub-Saharan Africa
Protocol

ABSTRACT

Background & objective: Cardiovascular disease (CVD) risk among the HIV population is high due to a combination of accelerated atherosclerosis from the pro-inflammatory milieu created by chronic HIV infection and the potentially adverse metabolic side effects from cART (combination antiretroviral therapy) medications. Although sub-Saharan Africa (SSA) bears 70% of the global burden of HIV disease, there is a relative paucity of studies comprehensively assessing CVD risk among people living with HIV on the continent. The overarching objective of the Evaluation of Vascular Event Risk while on Long-term Anti-retroviral Suppressive Therapy (EVERLAST) Study is to characterize the burden of CVD among HIV patients on ART in Ghana, and explore factors influencing it.

Methods: The EVERLAST study incorporates prospective CVD risk assessments and a convergent mixed methods approach. This prospective study will evaluate CVD risk by measuring Carotid Intimal Media Thickness (CIMT) and presence of traditional medical and lifestyle vascular risk among 240 Ghanaian HIV patients on anti-retroviral therapy compared with age- and sex-matched HIV uninfected ($n = 240$) and HIV positive ART naïve controls ($n = 240$). A contextual qualitative analysis will also be conducted to determine attitudes/perceptions of various key local stakeholders about CVD risk among HIV patients. The primary outcome measure will be CIMT measured cross-sectionally and prospectively among the three groups. A host of secondary outcome variables including CVD risk factors, CVD risk equations, HIV associated neurocognitive dysfunction and psychological well-being will also be assessed.

Conclusion: EVERLAST will provide crucial insights into the unique contributions of ART exposure and environmental factors such as lifestyle, traditional beliefs, and socio-economic indicators to CVD risk among HIV patients in a resource-limited setting. Ultimately, findings from our study will be utilized to develop interventions that will be tested in a randomized controlled trial to provide evidence to guide CVD risk management in SSA.

1. Introduction & rationale

1.1. Background

Of the approximately 36.9 million people living globally with Human Immunodeficiency Virus (HIV) at the end of 2017, 70% resided in Sub-Saharan Africa (SSA) [1]. Fortunately, the introduction of

potent, combination antiretroviral therapy (cART) has resulted in significant declines in mortality and morbidity due to HIV infection [2,3]. By mid-2015, 15.8 million people living with HIV were receiving cART resulting in a 35% decline in new HIV infections, a 24% decline in AIDS-related deaths and 7.8 million lives saved. However, while the initiation of cART has led to a global decline in deaths from AIDS related opportunistic infections and malignancies, there has been a

* Corresponding author at: Division of Neurology, Department of Medicine, Kwame Nkrumah University of Science & Technology, Private Mail Bag, Kumasi, Ghana.

E-mail address: stephensarfo78@gmail.com (F.S. Sarfo).

<https://doi.org/10.1016/j.ensci.2019.100189>

Received 10 September 2018; Received in revised form 23 March 2019; Accepted 29 March 2019

Available online 05 April 2019

2405-6502/ © 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

corresponding surge in deaths from cardiovascular disease (CVD) and non-AIDS related cancers [4–10,11–13].

In North American cohorts, nearly 50% of deaths are non-AIDS related with 7–19% of all deaths attributed to CVD [14,15–18]. Persistent immune activation from HIV infection, even in the presence of suppressed viremia on cART, has been associated with accelerated atherosclerosis among patients who are HIV-infected [6,19,20], reflecting the increasing rates of myocardial infarction and stroke in this population [4–9]. Several cross-sectional studies among mainly white populations have assessed the CVD risk profiles among HIV-infected patients using cardiovascular prognostication scores, such as the Framingham Risk Score (FRS), the Prospective Cardiovascular Münster Study (PROCAM) scores, the Systematic Coronary Risk Evaluation (SCORE) equation and the Pooled Cohort Atherosclerotic CVD risk (ASCVD) equation; these were developed, proposed, and validated for older, high-income, non-HIV-infected populations [21–27]. Based on results of a large multicenter HIV cohort study conducted in Europe and North America-The Data Collection on Adverse Effects on Anti-HIV Drugs Cohort-DAD, the investigators have proposed the first so-called DAD (Data Collection on Adverse Events of Anti-HIV Drugs) equation to estimate CVD risk among HIV patients which takes into account exposure to protease inhibitor and abacavir [28]. There is, however, no consensus on which cardiovascular prognostication equation to use to assess risk among HIV patients in routine clinic settings and utility of these risk scores for HIV infected patients in resource-limited settings is unknown.

It is now estimated that nearly 80% of all deaths from non-communicable diseases occur in Low and Middle Income Countries (LMICs) where deaths from non-communicable diseases (NCDs) are second only to the disease burden of HIV/AIDS [29,30]. This is largely attributable to enhancements in socio-economic status of these populations leading to the adoption of Western diets and lifestyles with an associated burgeoning epidemic of key CVD risk factors such as hypertension, diabetes mellitus, and dyslipidemia. As a result, CVD event rates are soaring [31]. These untoward secular trends in many LMICs further heighten an existing predisposition to CVD risk among the HIV population due to a combination of accelerated atherosclerosis from the pro-inflammatory milieu created by chronic HIV infection and the potentially adverse metabolic side effects from cART medications [11–13].

A study reported significantly more CVD events among HIV patients on cART in SSA compared with those treated in the USA with events rate of 8.4 (2.4–18.4) vs. 5.0 (2.7–9.2) per 1000 person-years [32]. Up to 24% of Ugandan HIV patients receiving cART vs 14% of those who were ART naïve had evidence of asymptomatic systemic atherosclerosis based on sonographic measurements of carotid intimal media thickness (CIMT) [33]. Furthermore, CIMT values among a predominantly female South African HIV+ population were considerably elevated and were associated with established CVD risk factors such as hypertension, high Body Mass Index (BMI), diabetes mellitus, total and low-density lipoprotein cholesterol, estimated glomerular filtration rate, metabolic syndrome and Framingham Heart Risk score rather than HIV-related factors [34]. In both of these studies, non-HIV infected controls were not also evaluated thereby greatly limiting the validity of relative risk calculations of CVD observed. These findings, however, herald the emergence of CVD risk as an important public health issue within the HIV population in SSA. However, there is a surprising paucity of data on studies evaluating interventions that could mitigate the burden of cardiovascular comorbidity and events among HIV patients on ART in SSA where the vast majority of HIV patients receiving ART reside. Prompt and accurate detection, as well as appropriate management of CVD risk factors in this population, is urgent given poor vascular outcomes in LMICs [35,36].

1.2. Objectives and underlying hypothesis

The overarching objective of the Evaluation of Vascular Event Risk

while on Long-term Anti-retroviral Suppressive Therapy (EVERLAST) Study is to characterize the burden of CVD risk among HIV patients on ART in Ghana, and explore factors influencing it. To accomplish this, we will assess prevalence and predictors of CVD risk factors among 240 HIV patients on ART compared with 480 age and sex-matched controls (HIV+ ART-naïve, $n = 240$ and HIV uninfected, $n = 240$) cross-sectionally and also prospectively over the course of 12 months. The following are the specific aims and their underlying hypotheses of the EVERLAST Study:

- To compare the prevalence of CVD among HIV patients on ART for > 1 year vs. matched controls.
 - o The primary endpoint will be CIMT burden.
 - o We will also secondarily assess the contributions of demographic factors (including age, sex, socio-economic, and urban/rural status), traditional CVD risk factors (hypertension, diabetes, dyslipidemia, smoking), and HIV-related factors (CD4 counts, viral load, ART regimens), to the prevalence of CVD among HIV patients on ART.
 - o We hypothesize that CVD prevalence will be higher in HIV patients on ART than age-matched ART naïve HIV patients and HIV uninfected controls in SSA. CVD risk will be higher in older individuals, longer exposure to ART, higher socioeconomic strata, men, and elevated risk factor prognostication score (e.g. Framingham Risk Score, DAD risk score).
- To obtain information from HIV patients, caregivers, providers, hospital administrators, and community leaders about barriers and facilitators of adherence to goals for CVD risk reduction, using focus groups, key informant interviews, and surveys. From the input and guidance of the various groups, we will develop an integrated protocol-driven risk factor patient self-management intervention, aimed at increasing patient self-efficacy and intrinsic motivation and comprising systematic involvement of enlightened, engaged providers. Within SSA, our hypothesis is that it is feasible to conduct useful mixed methodological research that incorporates local barriers/facilitators aimed at crafting a CVD risk reduction intervention for HIV patients.
- To prospectively assess over 12 months of follow-up, changes in CIMT, incidence of hypertension and diabetes mellitus, CVD events such as stroke, heart failure, neurocognitive dysfunction, quality of life, psychosocial stress, anxiety and stigma. Additionally, we will assess all cause mortality during follow-up.

2. Methods

2.1. Design

EVERLAST incorporates both cross-sectional and prospective CVD risk assessments and a convergent mixed methods approach. The cross-sectional and prospective study will evaluate CVD risk by measuring CIMT and presence of traditional medical and lifestyle vascular risk among 240 Ghanaian HIV patients on antiretroviral therapy compared with age- and sex-matched HIV uninfected ($n = 240$) and HIV positive ART naïve controls ($n = 240$). A contextual qualitative analysis will also be conducted to determine attitudes/perceptions of various key local stakeholders about CVD risk among HIV patients.

2.2. Setting and study populations

Kumasi, a city in Ghana, is among the country's largest metropolitan areas with a population of > 2 million, and is home to Komfo Anokye Teaching Hospital (KATH), a tertiary referral center and teaching hospital with 1000 beds [46]. KATH is the second largest tertiary referral center in Ghana and has one of the largest HIV clinics in-country. The HIV clinic was set up in late 2003 and to date has enrolled 11,500 patients, out of which nearly 6000 are currently on antiretroviral

therapy. The clinic has 4 HIV specialists, 10 HIV nurses, 6 pharmacists, and 15 public health practitioners who run a 5-day clinic/week where approximately 200 patients are seen weekly. The center has facilities for CD4 count and viral load monitoring with the support of the National AIDS Commission. First-line therapy according to National guidelines is zidovudine or tenofovir + lamivudine + efavirenz and second line therapy is protease-inhibitor based with ritonavir-boosted *Lopinavir* plus 2 NRTI backbone of either *Abacavir*/zidovudine/tenofovir. The Kumasi site has been involved in several prospective studies on co-infections in HIV, as well as randomized controlled trials on cART with international collaborators. [37–45]. In addition, KATH has Neurologists and Cardiologists whose expertise will contribute in the assessment of Neurological and Cardiovascular disorders in EVERLAST.

2.3. Eligibility criteria

2.3.1. Inclusion criteria for HIV positive individuals on ART

1. Above the age of 30 years; male or female
2. Receiving uninterrupted antiretroviral therapy for the at least 1 year
3. Legally competent

2.3.2. Exclusion criteria

1. Failure to meet all inclusion criteria
2. Severe cognitive impairment/dementia

2.3.3. Inclusion criteria for HIV positive not receiving ART

1. Above the age of 30 years; male or female
2. Has not had previous exposure to antiretroviral therapy
3. Legally competent

2.3.4. Exclusion criteria

1. Failure to meet all inclusion criteria
2. Severe cognitive impairment/dementia

2.3.5. Inclusion criteria for HIV uninfected subjects

1. Above the age of 30 years; male or female
2. Sero-negative for HIV antibodies on screening
3. Legally competent

2.3.6. Exclusion criteria

1. Failure to meet all inclusion criteria
2. Severe cognitive impairment/dementia

2.4. Recruitment and study evaluations

Two hundred and forty (240) HIV infected adult patients on Antiretroviral therapy for > 1 year and 240 age-and sex-matched HIV infected but ART-naïve patients enrolled for care at the HIV clinic at KATH will be recruited. Two hundred and forty (240) age-and sex matched HIV uninfected individuals will be recruited as controls from the community.

A standardized data collection form will be developed to collect information on the following:

- socio-demographic variables (age, gender, ethnicity, educational status, income, marital status);
- HIV disease characteristics (current CD4 cell count, HIV-1 viral load where available, past and current history of ART, duration on ART), and hepatitis B and C co-infection;
- known risk factors for CVD (hypertension, age, sex, cigarette

smoking, family history of premature CVD). Hypertension will be based on blood pressure levels and will also be classified in the traditional categories of awareness, treatment and control.

- details of dietary habits (for estimation of intake of fruits, vegetables, fried foods, meat, fish, added salts, etc.), tobacco use, exposure to environmental tobacco, alcohol use, as well as physical activity, sleep patterns, substance abuse (e.g. cocaine, amphetamine), which will be assessed at enrollment.
- anthropometric variables such as waist circumference, abdominal circumference, body mass index to categorize patients into underweight, ideal, overweight and obesity, which will also be collected.
- lipid profile, and glucose as well as HBA1C levels, which will be obtained from blood samples.

2.5. Study outcomes

The primary endpoint is CIMT. CIMT was chosen because it is a reliable surrogate marker for CVD events [47–51]. It is an intermediate phenotype for early atherosclerosis and can be measured relatively simply and non-invasively on B-mode ultrasound. For these reasons, it has been increasingly used as an endpoint in clinical trials. Study participants will undergo carotid Doppler ultrasonic evaluation for evidence of clinical or sub-clinical carotid artery disease- a validated surrogate marker of atherosclerosis [57]. To achieve reliable ultrasonic measurements of the common carotid artery intimal media thickness, a standardized protocol and strict quality control procedures will be followed by two experienced sonographers who will be blinded to the participant's group status and risk factor levels to ensure achievement of unbiased results [58,59]. CIMT will be measured at 1 cm portions of the distal left and right common carotid artery far walls with a linear transducer (transducer frequency of 7.5 MHz) with axial resolution of 0.10 mm, and calculated automatically over 3 cardiac cycles following the Mannheim consensus [59]. The average thickness of the left and right carotid arteries will be used as the outcome measure.

Definitions for CVD risk factors, events and CVD risk prognostic scores to be assessed as secondary outcomes are shown in Table 1.

2.6. Assessments for neurocognitive dysfunction & psychological well-being

Cognitive status of study participants will be assessed using Montreal Cognitive Assessment (MOCA) scale [60] and the Modified HIV Dementia Scale [61]. The hospital anxiety and depression scale [62] will be employed in the three groups to assess presence of depression or anxiety. Also the 9-item Patient Health Questionnaire [63] will be used to assess the quality of health of the study participants. Scores on the HIV Dementia scale will be compared between the three groups and specific neurocognitive domains will be compared. We will report on the prevalence of HIV-associated neurocognitive dysfunction (HAND) and assess associations between HAND and a range of factors including demographic (age, gender, educational status, etc), HIV-specific factors (viral load, CD4 counts, ART treatment exposure, opportunistic infections, ART regimen) and cardiovascular risk factors (hypertension, diabetes mellitus, dyslipidemia, and CIMT).

2.7. Prospective follow-up evaluations

Study participants will be followed for 12 months at which point Doppler ultrasound will be repeated to assess rate of change. Also blood pressure measurements will be performed at months 3, 6, 9 and 12 at clinic visits with the patient seated and well rested for 3 blood pressure measurements using an automated blood pressure monitor. Serum biochemical panels including lipid profile, serum creatinine and HBA1c will be repeated at month 12. Neurocognitive function assessments will be performed again at month 12 to assess and compare the trajectory of cognitive function among the three groups. For ethical reasons, HIV positive individuals who at baseline are ART naïve, will be started on

Table 1
Definitions of CVD variables to be collected.

Variable	Brief description
Arterial hypertension	Systolic BP > 140 mmHg and/or diastolic BP > 90 mmHg or (> 135/85 mm Hg in Diabetes patients) or taking antihypertensive drugs.
Diabetes mellitus	At least two fasting plasma glucose levels > 7.0 mmol/l, random blood glucose > 11.1 mmol/l, HBA1C > 6.5% or if patients had a history of diabetes treatment.
Dyslipidemia	Total cholesterol > 6.2 mmol/l, HDL-cholesterol < 1.03 mmol/l, or fasting triglycerides > 2.26 mmol/l.
Metabolic syndrome	According to the updated NCEP criteria: Triglyceride of at least 1.7 mmol/l, HDL-C of 1.0 mmol/l or less in men and of 1.3 mmol/l or less in women, hypertension (SBP > 130 mmHg or DBP > 85 mmHg), or use of anti-hypertensive medications and a glucose level > 5.6 mmol/l or diagnosis of diabetes mellitus [52].
Cardiovascular disease	History of stroke, carotid artery disease (symptomatic e.g., TIA or stroke) or > 50% stenosis on Doppler ultrasonography, myocardial infarction, stable/unstable angina, peripheral artery disease.
Family history of premature cardiovascular event	CV event occurring in a male first-degree relative < 55 years or female first-degree relative < 65 years.
DAD equation	DAD equation includes age, sex, systolic BP, serum cholesterol; HDL-cholesterol; diabetes, smoking status, family history of CVD, current use of Abacavir, indinavir, or lopinavir; and the number of years on indinavir or lopinavir. 5-year risk of coronary heart disease is classified as low (< 1%), moderate (1–5%), high (5–10%) or very high (> 10%) [28]
Framingham equation	Variables included in the FRS equation include age, sex, systolic BP, antihypertensive therapy (yes or no), serum total cholesterol and HDL-cholesterol values, current smoking status (yes or no). Participants with low or moderate 10-year risk will be re-evaluated based on the presence of aggravating factors: family history of CVD, metabolic syndrome, serum creatinine ≥ 1.5 mg/dl, hsCRP > 3.0 mg/dl or albuminuria > 30 μ g/mg. Patients presenting at least one of these aggravating factors will be re-classified into high-risk category [52,53].
PROCAM equation	Variables included in the PROCAM equation include age, sex, known diabetes or fasting blood glucose level > 120 mg/dl, systolic BP, serum TG, LDL, and HDL-cholesterol levels, current nicotine consumption and family history of CVD. The 10-year risk of acute coronary events is classified as low (< 10%), moderate (10–20%) or high (> 20%) [54].
SCORE equation	Variables included in the SCORE equation include sex, age, smoking, systolic BP, and either total cholesterol or the total/HDL cholesterol. Persons are considered low risk (< 1%), moderate risk (1%–5%) and high risk ($\geq 5\%$) [55].
Pooled cohort atherosclerotic CVD risk (ASCVD)	Variables included in the ASCVD equation include age, sex, race, total cholesterol, HDL-cholesterol, systolic BP, antihypertensive drug use, diabetes, and smoking status [56].

BP = blood pressure; HDL = high density lipoprotein; LDL = Low density Lipoprotein; TIA = Transient Ischemic Attack; NCEP = National Cholesterol Education Program; FRS = Framingham Risk Score; hsCRP = highly sensitive C-reactive protein.

ART during prospective follow-up in accordance with local guidelines.

2.9. Statistical analysis plan

2.8. Sample size justification

The study design is a cross-sectional study where a total of 720 age and sex-matched subjects, 240 each from three HIV and ART groups (HIV + on ART, HIV + not on ART, HIV uninfected) are selected from the population to assess their CVD status including symptomatic and sub-clinical carotid artery disease, cerebrovascular events, and coronary heart diseases. Previous studies have shown that the effect size comparing CVD rates among these groups range from a rate ratio of 1.5 (comparing HIV + ART to HIV) to 2.0 (comparing HIV + ART to no-HIV) [64]. A study in Uganda showed that the rate of subclinical atherosclerosis in the ART-naïve group was 14% while that in the HIV + ART group was 24% (RR = 1.5) [33]. Assuming 5% type I error, a sample size of 240 in each group will allow detecting an effect size of odds ratio 1.5 with a power of 80% using a likelihood ratio test. To compare three groups using a likelihood ratio chi-square test ($\alpha = 0.025$), a sample size of 720 (240 per group) will provide an 80% power to detect an effect size of 0.10 (Cohen's effect size based on $\sqrt{\text{chi square}/N}$) using a 2 degrees of freedom Chi-Square to study the associations' between HIV status (HIV + on ART, HIV + not on ART, HIV uninfected) and CVD risk accounting for missing data and additional covariates.

Prior to addressing each hypothesis, univariate descriptive statistics and frequency distributions will be calculated as appropriate for all biological variables (including gender, age) comparing individuals by HIV status (no-HIV, HIV +, HIV + ART). Briefly, box plots will be used to examine the relative distribution of variables stratified by HIV status. Non-parametric and equivalent parametric statistics will be utilized to compare groups. Appropriate regression models (linear regression for continuous outcomes such as carotid-media thickness and logistic regression for binary outcomes such as stroke) will be used to estimate the association between the covariates and each of the outcomes. When building models for each specific aim, the first stage of the model-building algorithm will involve testing if the individual covariates are correlated with the main outcome variables (CVD status indicators). A liberal $\alpha = 0.20$ will be used for these unadjusted analyses [65]. Once the initial pool of candidate predictors has been identified, regression models consisting of multiple covariates will be fitted to identify potential confounders and effect modifiers. To achieve unbiased and robust results, optimal combinations of predictors, including interaction effects, will be identified and used for further analysis based upon whether or not they are a confounder, by whether they did not improve the model fit, or increased the standard error of the parameter estimate of the primary covariates. Finally, each model will be rigorously assessed for collinearity and goodness of fit using residual

analysis. Model diagnostics will be performed using tools (in SAS or STATA) that detect outliers and influential data points. We will use diagnostic measures such as residual deviance, the hat matrix diagonal and residual chi-squared deviance and the difference between chi-square goodness-of-fit when an observation is deleted [62]. Plots of these against predicted values will be used to investigate the influence of each data point on the model. We will handle missing data using several techniques including multiple imputation and propensity score methods [66,67].

2.9.1. Qualitative studies to assess knowledge, attitudes and practices of patients and healthcare providers on CVD among HIV patients

Guided by a social ecological model [68], community engaged research [69], and NIH best practices for mixed methods research [70], Ghanaian researchers and Medical University of South Carolina partners will conduct a mixed methods study to answer the following questions: **a)** What are the individual, interpersonal, health system, and community factors (with an emphasis on barriers and facilitators) influencing CVD risk reductions among Ghanaian HIV patients on ART?; **b)** What are recommended intervening strategies to develop a successful intervention research study that focuses on reducing barriers and enhancing facilitators for CVD risk reduction in Ghanaian HIV patients on ART?

2.10. Procedures

The team will review available data related to HIV patients on ART and CVD risk factors and based on quantitative findings, develop Focus Group and semi-structured Key Informant Interview (KII) guides to further answer the research questions and refine intervention design. Interview guides will embrace a social ecological perspective (i.e., patient, family/caregiver, health systems, and community factors) and will focus on Steps 1 and 2 of the 6 SQuID (Steps in Quality Intervention Development) [71]. We will conduct 7 focus groups ($n = 45\text{--}60$ participants) followed by 10–12 KIIs. The focus groups will last 60–90 min and the KIIs will last approximately 45–60 min and will occur at a location convenient to participants. A trained moderator will conduct the sessions with another member of the investigative team taking field notes during the session. Written consent will be obtained from participants prior to the start of any session. The purposive sample (based on demographics of the HIV patients prescribed ART) for the focus groups and KIIs (based on the clinical care providers (physicians, nurses, pharmacists, hospital administrators) will be recruited through both purposive and snowball sampling techniques. We will purposively sample and stratify the focus groups to include: two focus groups with patients with HIV; two focus groups with patients with HIV and diagnosed CVD; one focus group with family members, friends, caregivers of persons with HIV and diagnosed CVD; and two focus groups with community leaders. A focus group guide will ensure a systematic and consistent approach on obtaining information from each focus group to identify: (1) beliefs about the concept of Cardiovascular risk in HIV disease; (2) existing self-management strategies and experiences; (3) literacy levels on hypertension, diabetes mellitus, dyslipidemia lifestyle and therapeutic management as well as the consequences of undiagnosed and untreated CVD risk factors; and (4) community level factors that may influence CVD risk reductions among HIV patients on ART. The KIIs will be conducted with the representative clinical and administrative staff identified from each of the groups above. KIIs will provide insight into health-system level factors important for optimization of interventions to improve clinical outcomes among this population.

2.11. Data management and analyses

All focus groups and KIIs will be digitally recorded and transcribed verbatim for qualitative analysis. All data will be uploaded and stored

on REDCap, a secure data storage system. Transcripts of the audio-recordings and field notes from the de-identified KIIs and focus groups will be used for data analysis. The team will review transcriptions with actual recordings to check for accuracy and authenticity. The transcripts will be imported into the text analysis software (i.e. NVIVO) [72] for data management, and thematic analysis by the qualitative methodologist. We will employ the ‘Framework Analysis’ approach [73], which includes five key stages: familiarization, identifying a thematic framework, indexing, charting, mapping, and interpretation to meet the processes of Steps 1–2 and partially address Step 3 of the 6 SQuID [71]. The investigative team will share findings through a process known as member checking [74] with at least three focus group and KII participants to ensure credibility of findings. Findings from the quantitative component (Aim 1) will be integrated with findings from the qualitative component (Aim 2) as part of the convergent mixed methodological approach [70]. Using a convergent approach will allow investigators to more thoroughly understand the clinical needs of this population and to tailor intervention refinements to optimize clinical outcomes.

3. Discussion

EVERLAST is premised on systematically assessing the prevalence and predictors of CVD risk among a cohort of HIV-patients receiving ART in an urban setting in Ghana. The inclusion of two control groups - HIV positive ART-naïve subjects and HIV uninfected subjects, will allow for a rigorous and unbiased evaluation of the relative risk of both ART exposure and presence of HIV to CVD risk among HIV patients ART in SSA. A convergent mixed methods research approach will be employed in EVERLAST to identify the locally-relevant and contextually-appropriate barriers and facilitators of adherence to goals of CVD risk reduction among the HIV population that may not be captured with survey methodology alone. The objective is to develop an intervention that incorporates currently unknown community member input into intervention that can be tested in a future study. If the developed intervention proves to be ultimately effective, it could serve as a scalable strategy for managing CVD risk among HIV patients in SSA. By comprehensively assessing the nature of CVD burden among HIV patients in SSA and employing a mixed-methods approach to develop an intervention to address it, the Evaluation of Vascular Event Risk while on Long-term Anti-retroviral Suppressive Therapy (EVERLAST) Study will provide crucial insights into the unique contributions of ART exposure and environmental factors such as lifestyle, traditional beliefs and socio-economic indicators to CVD risk among HIV patients in a resource-limited setting. A 12-month follow-up is planned to allow us assess the progression of carotid intimal media thickness in the three groups, assess the incidence of vascular risk factors such as hypertension, dyslipidemia and diabetes mellitus and determine the trajectory neurocognitive dysfunction in relation to vascular risk factor burden in HIV.

To our knowledge, this will be the first study to: **1)** comprehensively assess the synergy between HIV positivity, antiretroviral drug exposure, lifestyle patterns, and CVD in SSA; **2)** explore the knowledge, attitudes and practices of HIV patients, caregivers, healthcare providers, system administrators, and public health officials towards CVD risk and its management among patients with HIV populations; **3)** develop an intervention for optimal management of CVD in HIV patients in SSA to be rigorously evaluated in randomized trial fashion in a future study. This will help inform policy makers on the need to avert the catastrophic outcomes of cardiovascular diseases observed among HIV patients in particular and indeed among the general population in Ghana and across SSA [75–91].

Anticipated challenges and proposed solutions include **1)** Subject Accrual: Assessment visits will be scheduled, whenever possible, on days subjects already have scheduled clinic visits or plan to visit the city. **2)** Meeting sample size: There is a high HIV patient load of

approximately 100 cases per week. At a recruitment rate of 15 cases per week, recruitment of 480 HIV patients seems a realistic objective. Moreover, Controls will be recruited through our community outreach programs. During these weekly outreach sessions up to 100 individuals from the community participate in screening programs for CVD prevention so recruiting 240 HIV uninfected subjects is attainable. **3) Missing Data:** Given the one-time nature of the cross-sectional clinical assessments and focus group participation, we anticipate little missing data.

Status of study

Follow-up of study participants on-going.

Conflicts of interest

None.

Funding

Grant R21 TW010479-01 from the National Institute of Health.

References

- [1] AIDS by Numbers, UNAIDS, (2015).
- [2] A. Mocroft, B. Ledergerber, C. Katlama, O. Kirk, P. Reiss, A. d'Arminio Monteforte, et al., Decline in the AIDS and death rates in the EuroSIDA study: an observational study, *Lancet* 362 (9377) (2003) 22–29.
- [3] T.C. Quinn, HIV epidemiology and effects of antiviral therapy on long-term consequences, *AIDS* 22 (Suppl. 3) (2008) S7–S12.
- [4] J.S. Currier, J.D. Lundgren, A. Carr, D. Klein, C.A. Sabin, P.E. Sax, et al., Epidemiological evidence for cardiovascular disease in HIV-infected patients and relationship to highly active antiretroviral therapy, *Circulation* 118 (2008) e29–e35.
- [5] Group DADS, N. Friis-Moller, P. Reiss, C.A. Sabin, R. Weber, A. Monteforte, et al., Class of antiretroviral drugs and the risk of myocardial infarction, *N. Engl. J. Med.* 356 (2007) 1723–1735.
- [6] V.A. Triant, HIV infection and coronary heart disease: an intersection of epidemics, *J. Infect. Dis.* 205 (Suppl. 3) (2012) S355–S361.
- [7] M.S. Freiberg, C.C. Chang, L.H. Kuller, M. Skanderson, E. Lowy, K.L. Kraemer, et al., HIV infection and risk of acute myocardial infarction, *JAMA Int. Med.* 173 (2013) 614–622.
- [8] L.D. Rasmussen, F.N. Engsig, H. Christensen, J. Gerstoft, G. Kronborg, C. Pedersen, et al., Risk of cerebrovascular events in persons with and without HIV: a Danish nationwide population-based cohort study, *AIDS* 25 (2011) 1673–1746.
- [9] F.C. Chow, S. Regan, S. Feske, J.B. Meigs, S.K. Grinspoon, V.A. Triant, Comparison of ischemic stroke incidence in HIV-infected and non-HIV infected patients in a US health care system, *J. Acquir. Immune Defic. Syndr.* 60 (2012) 351–358.
- [10] B. Ovbiagele, A. Nath, Increasing incidence of ischemic stroke in patients with HIV infection, *Neurology* 76 (5) (2011) 444–450.
- [11] G. Ortiz, S. Koch, J.G. Romano, A.M. Forteza, A.A. Rabinstein, Mechanisms of ischemic stroke in HIV-infected patients, *Neurology* 68 (16) (2007) 1257–1261.
- [12] M. Depairon, S. Chessex, P. Sudre, N. Rodondi, N. Doser, J.P. Chave, et al., Premature atherosclerosis in HIV-infected individuals-focus on protease inhibitor therapy, *AIDS* 15 (3) (2001) 329–334.
- [13] C. Jerico, H. Knobel, N. Calvo, M.L. Sorli, A. Guelar, J.L. Gimeno-Bayon, et al., Sub-clinical carotid atherosclerosis in HIV-infected patients: role of combination antiretroviral therapy, *Stroke* 37 (3) (2006) 812–817.
- [14] UNAIDS, Epidemiological Slides – How AIDS Changed Everything Report; 2015, (2015).
- [15] C. Lawden, T. May, E. Rosenthal, C. Burty, F. Bonnet, D. Costagliola, et al., Changes in causes of death among adults infected by HIV between 2000 and 2005: the “Mortalite 2000 and 2005” surveys (ANRS EN19 and Mortavic), *J. Acquir. Immune Defic. Syndr.* 48 (5) (2008) 590–598.
- [16] F.J. Palella Jr., R.K. Baker, A.C. Moorman, J.S. Chmiel, K.C. Wood, J.T. Brooks, et al., Mortality in the highly active antiretroviral therapy era: changing causes of death and disease in the HIV outpatient study, *J. Acquir. Defic. Syndr.* 43 (1) (2006) 27–34.
- [17] Data Collection on Adverse Events of Anti-HIV drugs (D:A:D) Study Group, C. Smith, C.A. Sabin, J.D. Lundgren, R. Thiebaut, R. Weber, et al., Factors associated with specific causes of death amongst HIV-positive individuals in the D:A:D Study, *AIDS* 24 (10) (2010) 1537–1548.
- [18] A.J. Rodger, R. Lodwick, M. Schechter, S. Deeks, J. Amin, R. Gilson, et al., Mortality in well controlled HIV in the continuous antiretroviral therapy arms of the SMART and ESPRIT trials compared with the general population, *AIDS* 27 (6) (2013) 973–979.
- [19] D.A. Duprez, J. Neuhaus, L.H. Kuller, R. Tracy, W. Belloso, S. De Wit, et al., Inflammation, coagulation and cardiovascular disease in HIV-infected individuals, *PLoS ONE* 7 (9) (2012) e44454.
- [20] R.C. Kaplan, A.L. Landay, H.N. Hodis, S.J. Gange, P.J. Norris, M. Young, et al., Pmc3400505; potential cardiovascular disease risk markers among HIV-infected women initiating antiretroviral treatment, *J. Acquir. Immune Defic. Syndr.* 60 (2012) 359–368.
- [21] C. Knobel, C. Jerico, M. Montero, M.L. Sorli, M. Velat, A. Guelar, et al., Global cardiovascular risk in patients with HIV infection: concordance and differences in estimates according to three risk equations (Framingham, SCORE, and PROCAM), *AIDS Patient Care STDs* 21 (7) (2007) 452–457.
- [22] G.E. Vrentzos, J.A. Papadakis, E.S. Ganotakis, K.I. Paraskevas, I.F. Gazi, N. Tzanakis, et al., Predicting coronary heart disease risk using the Framingham and PROCAM equations in dyslipidemic patients without overt vascular disease, *Int. J. Clin. Pract.* 61 (10) (2007) 1643–1653.
- [23] M.M. Moreira Guimaraes, D. Bartolomeu Greco, A.H. Ingles Garcés, A.R. de Oliveira Jr., R. Bastos Foscolo, et al., Coronary heart disease risk assessment in HIV-infected patients: a comparison of Framingham, PROCAM, and SCORE risk assessment functions, *Int. J. Clin. Pract.* 64 (6) (2010) 739–745.
- [24] N. Edwards-Jackson, S.J. Kerr, H.V. Tieu, J. Ananworanich, S. Hammer, K. Ruxruntham, et al., Cardiovascular risk assessment in persons with HIV infection in the developing world: comparing three risk equations in a cohort of HIV-infected Thais, *HIV Med.* 12 (8) (2011) 510–515.
- [25] R.C. Kaplan, L.A. Kingsley, A.R. Sharrett, X. Li, J. Lazar, P.C. Tien, et al., Ten-year predicted coronary heart disease in HIV-infected men and women, *Clin. Infect. Dis.* 45 (8) (2007) 1074–1081.
- [26] E.M. Lima, D.M. Gualandro, P.C. Yu, C. Giuliano Ide, A.C. Marques, D. Calderaro, et al., Cardiovascular prevention in HIV patients: results from a successful intervention program, *Atherosclerosis* 204 (1) (2009) 229–232.
- [27] J. Begovac, G. Dragovic, K. Viskovic, J. Kusic, M.P. Mihanovic, D. Lukas, et al., Comparison of four international cardiovascular disease prediction models and the prevalence of eligibility for lipid lowering therapy in HIV infected patients on antiretroviral therapy, *Croat. Med. J.* 56 (2015) 14–23.
- [28] N. Friis-Moller, R. Thiebaut, P. Reiss, R. Weber, A.D. Monforte, S. De Wit, et al., Predicting the risk of cardiovascular disease in HIV-infected patients: the data collection on adverse effects of anti-HIV drugs study, *Eur. J. Cardiovasc. Prev. Rehabil.* 17 (2010) 491–501.
- [29] <http://www.who.int/about/structure/organigram/nmh/en/>.
- [30] Global burden of diseases, http://www.who.int/healthinfo/global_burden_disease/gbd/en/, (2010).
- [31] G.A. Mensah, G.A. Roth, U.K.A. Sampson, A.E. Moran, V.L. Feigin, M.H. Forouzanfar, et al., Mortality from cardiovascular diseases in sub-Saharan Africa, 1990–2013: a systematic analysis of data from the Global burden of disease study 2013, *Cardiovasc. J. Afr.* 26 (2015) S6–S10.
- [32] C.W. Wester, J.R. Koethe, B.E. Shepherd, S.E. Stinnette, P.F. Rebeiro, A.M. Kipp, et al., Non-AIDS-defining events among HIV-1-infected adults receiving combination antiretroviral therapy in resource-replete versus resource-limited urban setting, *AIDS* 25 (2011) 1471–1479.
- [33] I. Sinabulya, J. Kayima, C. Longenecker, M. Luwedde, F. Semitala, A. Kambugu, et al., Subclinical carotid atherosclerosis among HIV-infected adults attending HIV/AIDS care at two large ambulatory HIV clinics in Uganda, *PLoS ONE* 9 (2) (2014) e89537.
- [34] A.F. Schoffelen, E. de Groot, H.A. Tempelman, F.L. Visseren, A.I. Hoepelman, R.E. Barth, Carotid intimal media thickness in mainly female HIV-infected subjects in rural South Africa: association with cardiovascular but not HIV-related factors, *Clin. Infect. Dis.* 61 (10) (2015) 1606–1614.
- [35] V.L. Feigin, C.M. Lawes, D.A. Bennett, S.L. Barker-Collo, V. Parag, Worldwide stroke incidence and early case fatality in 56 population-based studies: a systematic review, *Lancet Neurol.* 8 (4) (2009) 355–369.
- [36] W.N. Kernan, B. Ovbiagele, H.R. Black, D.M. Bravata, M.I. Chimowitz, M.D. Ezekowitz, et al., Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association, *Stroke* 45 (7) (2014 Jul) 2160–2236.
- [37] F.S. Sarfo, M.A. Sarfo, D.R. Chadwick, Incidence and risk factors for neuropsychiatric events among Ghanaian HIV patients on long-term non-nucleoside reverse transcriptase inhibitor-based therapy, *eNeuroSci* 3 (2016) 21–25.
- [38] David Chadwick, F.S. Sarfo, Elaine Kirk, Victoria Parris, George Bedu-Addo, Dorcas Owusu, et al., Tenofovir is associated with increased tubular proteinuria and asymptomatic renal tubular dysfunction in Ghana, *BMC Nephrol.* 16 (2015) 195.
- [39] F.S. Sarfo, K.A. Eberhardt, A. Dompheh, E.O. Kuffour, M. Soltan, M. Schachschneider, J.F. Drexler, et al., *Helicobacter pylori* infection is associated with higher CD4 T cell counts and lower HIV-1 viral loads in ART-naïve HIV-positive patients in Ghana, *PLoS ONE* 10 (11) (2015 Nov 24) e0143388, <https://doi.org/10.1371/journal.pone.0143388>.
- [40] F.S. Sarfo, M.A. Sarfo, B. Norman, R. Phillips, G. Bedu-Addo, D. Chadwick, Risk of deaths, AIDS-defining and non-AIDS defining events among Ghanaians on long-term antiretroviral therapy, *PLoS ONE* 9 (10) (2014) e111400.
- [41] F.S. Sarfo, M.A. Sarfo, B. Norman, R. Phillips, D. Chadwick, Incidence and determinants of nevirapine and efavirenz-related skin rashes in West Africans: Nevirapine's epitaph? *PLoS ONE* 9 (4) (2014 Apr 11) e94854.
- [42] F.S. Sarfo, A. Kassim, R. Phillips, A.M. Geretti, D.R. Chadwick, Long-term responses to first-line antiretroviral therapy in HIV and hepatitis B co-infection in Ghana, *J. Inf. Secur.* 69 (5) (2014) 481–489.
- [43] F.S. Sarfo, M.A. Sarfo, A. Kasim, R. Phillips, M. Booth, D. Chadwick, Long-term effectiveness of first-line non-nucleoside reverse transcriptase inhibitor (NNRTI)-based antiretroviral therapy in Ghana, *J. Antimicrob. Chemother.* 69 (1) (2014) 254–261.

- [44] B. Ovbiagele, 'Al pie de la letra': crafting a report card for elderly Spanish-only-speaking patients with stroke. *Stroke*, J. Cereb. Circ. 41 (2010) 771–777.
- [45] A.W. Dromerick, M.C. Gibbons, D.F. Edwards, D.E. Farr, M.L. Giannetti, B. Sánchez, ... C.S. Kidwell, Preventing recurrence of thromboembolic events through co-ordinated treatment in the District of Columbia, *Int. J. Stroke* 6 (5) (2011) 454–460.
- [46] Index Mundi with data from CIA World Factbook Data, Ghana Demographic Profile, Retrieved from, 2014. http://www.indexmundi.com/ghana/demographics_profile.html.
- [47] M.A. Espeland, D.H. O'Leary, J.G. Terry, T. Morgan, G. Evans, H. Mudra, Carotid intima-media thickness as a surrogate for cardiovascular disease events in trials of HMG-CoA reductase inhibitors, *Curr. Control. Trials Cardiovasc. Med.* 6 (2005) 3.
- [48] D.H. O'Leary, J.F. Polak, R.A. Kronmal, T.A. Manolio, G.L. Burke, S.K. Wolfson Jr., for the Cardiovascular Health Study Collaborative Research Group. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults, *N. Engl. J. Med.* 340 (1999) 14–22.
- [49] M.W. Lorenz, H.S. Markus, M.L. Bots, M. Rosvall, M. Sitzer, Prediction of clinical cardiovascular events with carotid intima-media thickness: a systematic review and meta-analysis, *Circulation* 115 (2007) 459–467.
- [50] K. Kitagawa, H. Hougaku, H. Yamagami, H. Hashimoto, T. Itoh, Y. Shimizu, et al., Carotid intima-media thickness and risk of cardiovascular events in high-risk patients: Results of the Osaka Follow-Up Study for Carotid Atherosclerosis 2 (OSACA2 Study), *Cerebrovasc. Dis.* 24 (2007) 35–42.
- [51] P. Greenland, J.S. Alpert, G.A. Beller, E.J. Benjamin, M.J. Budoff, Z.A. Fayad, et al., 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, *Circulation* 122 (2010) e584–e636.
- [52] J.I. Cleeman, Executive summary of the third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adults treatment panel III), *JAMA* 285 (19) (2001) 2486–2497.
- [53] K.M. Anderson, P.W.F. Wilson, P.M. Odell, W.B. Kannel, An updated coronary risk profile. A statement for health professionals, *Circulation* 83 (1) (1991) 356–362.
- [54] G. Assmann, P. Cullen, H. Schulte, Simple scoring scheme for calculating the risk of acute coronary events on the 10-year follow-up of the Prospective Cardiovascular Munster (PROCAM) study, *Circulation* 105 (3) (2002) 310–315.
- [55] R.M. Conroy, K. Pyörälä, A.P. Fitzgerald, S. Sans, A. Menotti, G. De Backer, et al., Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project, *Eur. Heart J.* 24 (2003) 987–1003.
- [56] N.J. Stone, J. Robinson, A.H. Lichtenstein, C.N. Merz, C.B. Blum, R.H. Eckel, et al., ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American college of cardiology/American heart association task force on practice guidelines, *Circulation* 129 (25 Suppl 2) (2013) S1–S45.
- [57] L.E. Chambless, G. Heiss, A.R. Folsom, W. Rosamond, M. Szklo, A.R. Sharrett, et al., Association of coronary heart disease incidence with carotid arterial thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987–1993, *Am. J. Epidemiol.* 146 (6) (1997) 483–494.
- [58] E. De Groot, G.K. Hovingh, A. Wiegman, P. Duriez, A.J. Smit, J.C. Fruchart, et al., Measurement of arterial wall thickness as a surrogate marker for atherosclerosis, *Circulation* 109 (23 Suppl 1) (2004) III33–III38.
- [59] P.J. Touboul, M.G. Hennerici, S. Meairs, H. Adams, P. Amarenco, M. Desvarieux, et al., Mannheim intima-media thickness consensus, *Cerebro. Vasc. Dis.* 18 (4) (2004) 346–349.
- [60] Z.S. Nasreddine, N.A. Phillips, V. Bedirian, et al., The Montreal Cognitive Assessment, MOCA: a brief screening tool for mild cognitive impairment, *J. Am. Geriatr. Soc.* 53 (4) (2005) 695–699.
- [61] C. Power, O.A. Selnes, J.A. Grim, J.C. McArthur, HIV Dementia Scale: a rapid screening test, *J. Acquir. Immune Defic. Syndr. Hum. Retrovirol.* 8 (3) (1995) 273–278.
- [62] A.S. Zigmond, R.P. Snaith, The hospital anxiety and depression scale, *Acta Psychiatr. Scand.* 67 (6) (1983) 361–370.
- [63] K. Kroenke, R.L. Spitzer, J.B. Williams, The PHQ-9: validity of a brief depression severity measure, *J. Gen. Intern. Med.* 16 (9) (2001 Sep) 606–613.
- [64] F.M. Islam, J. Wu, J. Jansson, D.P. Wilson, Relative risk of cardiovascular disease among people living with HIV: a systematic review and meta-analysis, *HIV Med.* 13 (2012) 453–468.
- [65] D.W. Hosmer, S. Lemeshow, *Applied Logistic Regression*, 2nd ed., John Wiley & Sons, 2000.
- [66] A. Agresti, *Categorical Data Analysis*, John Wiley & Sons, 2002.
- [67] R.J.A. Little, D.B. Rubin, *Statistical Analysis with Missing Data*, John Wiley & Sons Inc, New York, NY, 2002.
- [68] O.R. Young, F. Berkhout, G.C. Gallopin, M.A. Janssen, E. Ostrom, S. van der Leeuw, The globalization of socio-ecological systems: an agenda for scientific research, *Glob. Environ. Chang.* 16 (3) (2006) 304–316.
- [69] M. Minkler, N. Wallerstein, *Community Based Participatory Research for Health*, John Wiley & Sons, San Francisco, 2003, pp. 135–154.
- [70] J.W. Creswell, V.L. Plano Clark, *Designing and Conducting Mixed Methods Research*, 3rd ed., Sage Publications, Washington, D.C., 2018.
- [71] D. Wright, E. Wimbush, K. Jepson, et al., Six steps in quality intervention development (6 SQuID), *J. Epidemiol. Community Health* 16 (2015) 2015, <https://doi.org/10.1136/jech-2015-205952> (published online first: November).
- [72] NVivo Qualitative Data Analysis Software, QSR International Pty Ltd. Version 10, (2012).
- [73] A. Srivastava, S.B. Thomson, Framework analysis: a qualitative methodology for applied policy research, *Joag* 4 (2) (2009) 72–79.
- [74] B.J. Breitmayer, L. Ayres, K.A. Knafl, Triangulation in qualitative research: evaluation of completeness and confirmation purposes. *Image, J. Nurs. Scholarsh.* 25 (3) (1993) 237–243 [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. Fall.
- [75] F.S. Sarfo, Dyslipidemia among African adults, *Lancet Glob. Health* 6 (9) (2018) e940–e941.
- [76] F.S. Sarfo, B. Ovbiagele, M. Gebregziabher, K. Wahab, R. Akinyemi, A. Akpalu, et al., Stroke among young West Africans: evidence from the SIREN (Stroke Investigative Research and Educational Network) Large multisite case-control study, *Stroke* 49 (5) (2018) 1116–1122.
- [77] F.S. Sarfo, O. Opare-Sem, M. Agyei, J. Akassi, D. Owusu, M. Owolabi, et al., Risk factors for stroke occurrence in a low HIV endemic West African country: a case-control study, *J. Neurol. Sci.* 395 (2018) 8–16.
- [78] F.S. Sarfo, L.M. Mobula, J. Plange-Rhule, D. Ansong, D. Ofori-Adjei, Incident stroke among Ghanaians with hypertension and diabetes: a multicenter, prospective cohort study, *J. Neurol. Sci.* 395 (2018) 17–24.
- [79] F.S. Sarfo, L.M. Mobula, G. Burnham, D. Ansong, J. Plange-Rhule, O. Sarfo-Kantanka, et al., Factors associated with uncontrolled blood pressure among Ghanaians: evidence from a multicenter hospital-based study, *PLoS ONE* 13 (3) (2018) e0193494.
- [80] L.M. Mobula, F.S. Sarfo, K.A. Carson, G. Burnham, L. Arthur, D. Ansong, et al., Predictors of glycemic control in type-2 diabetes mellitus: evidence from a multicenter study in Ghana, *Tran. Metab. Syndr. Res.* 1 (2018) 1–8.
- [81] F.S. Sarfo, L.M. Mobula, O. Sarfo-Kantanka, S. Adamu, J. Plange-Rhule, D. Ansong, et al., Estimated glomerular filtration rate predicts incident stroke among Ghanaians with diabetes and hypertension, *J. Neurol. Sci.* 396 (2018) 140–147.
- [82] L.T. Appiah, F.S. Sarfo, C. Agyemang, H.O. Tweneboah, N.A.B.A. Appiah, G. Bedu-Addo, et al., Current trends in admissions and outcomes of cardiac diseases in Ghana, *Clin. Cardiol.* 40 (10) (2017) 783–788.
- [83] O. Sarfo-Kantanka, F.S. Sarfo, E. Oparebea Ansah, B. Eghan, N.K. Ayisi-Boateng, E. Acheamfour-Akokuah, Secular trends in admissions and mortality rates from diabetes mellitus in the central belt of Ghana: a 31-year review, *PLoS ONE* 11 (11) (2016) e0165905.
- [84] M.O. Owolabi, F. Sarfo, R. Akinyemi, M. Gebregziabher, K. Wahab, A. Akpalu, et al., Dominant modifiable risk factors for stroke in Ghana and Nigeria (SIREN): a case-control study, *Lancet Glob. Health* 6 (4) (2018) e436–e446.
- [85] F.S. Sarfo, J. Akassi, G. Kyem, S. Adamu, D. Awuah, O.S. Kantanka, et al., Long-term outcomes of stroke in a Ghanaian outpatient clinic, *J. Stroke Cerebrovasc. Dis.* 27 (4) (2018) 1090–1099.
- [86] F.S. Sarfo, J. Akassi, D. Awuah, S. Adamu, C. Nkyi, M. Owolabi, et al., Trends in stroke admission and mortality rates from 1983 to 2013 in central Ghana, *J. Neurol. Sci.* 357 (1, 2) (2015) 240–245.
- [87] M.O. Owolabi, S. Akarolo-Anthonny, R. Akinyemi, D. Arnett, M. Gebregziabher, C. Jenkins, et al., The burden of stroke in Africa: a glance at the present and a glimpse into the future, *Cardiovasc. J. Afr.* 26 (2 Suppl 1) (2015) S27–S38.
- [88] F.S. Sarfo, G. Kyem, B. Ovbiagele, J. Akassi, O. Sarfo-Kantanka, M. Agyei, et al., One-year rates and determinants of poststroke systolic blood pressure control among Ghanaians, *J. Stroke Cerebrovasc. Dis.* 26 (1) (2017) 78–86.
- [89] F.S. Sarfo, J. Akassi, N.K. Antwi, V. Obese, S. Adamu, A. Akpalu, et al., Highly prevalent hyperuricaemia is associated with adverse clinical outcomes among Ghanaian stroke patients: an observational prospective study, *Ghana. Med. J.* 49 (3) (2015) 165–172.
- [90] F.S. Sarfo, J.W. Acheampong, L.T. Appiah, E. Oparebea, A. Akpalu, G. Bedu-Addo, The profile of risk factors and in-patient outcomes of stroke in Kumasi, Ghana, *Ghana. Med. J.* 48 (3) (2014) 127–134.
- [91] M. Owolabi, F. Sarfo, V.J. Howard, M.R. Irvin, M. Gebregziabher, R. Akinyemi, et al., Stroke in indigenous Africans, African Americans, and European Americans: interplay of racial and geographic factors, *Stroke* 48 (5) (2017) 1169–1175.