




Official Journal of the European Paediatric Neurology Society



Original article

Executive function and general intellectual functioning in dyskinetic cerebral palsy: Comparison with spastic cerebral palsy and typically developing controls



Olga Laporta-Hoyos ^{a,b}, Júlia Ballester-Plané ^{a,b}, David Leiva ^c,
Teresa Ribas ^a, Júlia Miralbell ^a, Clara Torroja-Nualart ^c,
Maria Eugenia Russi ^d, Esther Toro-Tamargo ^e, Mar Meléndez-Plumed ^e,
Francisca Gimeno ^f, Alfons Macaya ^g, Roser Pueyo ^{a,b,*}

^a Institut de Neurociències, Departament de Psicologia Clínica i Psicobiologia, Universitat de Barcelona, Passeig Vall d'Hebron 171, 08035, Barcelona, Spain

^b Institut de Recerca Sant Joan de Déu, Santa Rosa 39-57, 08950, Esplugues de Llobregat, Spain

^c Departament de Psicologia Social i Psicologia Quantitativa, Universitat de Barcelona, Passeig Vall d'Hebron 171, 08035, Barcelona, Spain

^d Servei de Neurologia, Hospital Universitari Sant Joan de Déu, Passeig Sant Joan de Déu 2, 08950, Esplugues de Llobregat, Spain

^e Servei de Rehabilitació i Medicina Física, Hospital Universitari Vall d'Hebron, Passeig Vall d'Hebron 119-129, 08035, Barcelona, Spain

^f Serveis de Salut i Rehabilitació, Associació de la Paràlisi Cerebral (ASPACE), Camí Tres Pins 31-35, 08038, Barcelona, Spain

^g Grup de Recerca en Neurologia Pediàtrica, Vall d'Hebron Institut de Recerca, Hospital Universitari Vall d'Hebron, Passeig Vall d'Hebron 119-129, 08035, Barcelona, Spain

ARTICLE INFO

Article history:

Received 15 December 2018

Received in revised form

14 May 2019

Accepted 14 May 2019

Keywords:

Dyskinetic cerebral palsy

Spastic cerebral palsy

Executive function

Intellectual functioning

Goal setting

ABSTRACT

Aim: To comprehensively describe intellectual and executive functioning (EF) in people with dyskinetic cerebral palsy (DCP), by comparing their performance with that of: 1) age- and sex-matched typically developing controls (TDC); and 2) participants with spastic cerebral palsy (SCP) matched for age, sex, term/preterm and gross motor function classification system (GMFCS).

Method: This cross-sectional study was conducted by the University of Barcelona in collaboration with five institutions. Participants were people with DCP ($n = 52$; 24 females, median age 20.5 y: 5mo, interquartile range [IQR] = 13.75 y: 7mo; GMFCS I–V). As comparison groups, participants with SCP ($n = 20$; 10 females, median age = 20.5 y: 5.5mo, IQR = 13.75 y 9mo; GMFCS I–V) and TDC ($n = 52$; 24 females, median age = 20 y: 4mo, IQR = 12 y 7mo) were included. Intelligence and EF were assessed using common tests in all participants.

* Corresponding author. Departament de Psicologia Clínica i Psicobiologia Universitat de Barcelona, Passeig de la Vall d'Hebron 171, 08035, Barcelona, Spain. Fax: +34 93 402 15 84.

E-mail address: rpueyo@ub.edu (R. Pueyo).

<https://doi.org/10.1016/j.ejpn.2019.05.010>

1090-3798/© 2019 European Paediatric Neurology Society. Published by Elsevier Ltd. All rights reserved.

Results: Both CP groups had lower intelligence than TDC and performed poorer in almost all EF tasks. Intelligence was higher in DCP than SCP ($z = -2.51$, $p = 0.01$). Participants with DCP also performed significantly better in goal-setting tasks ($z = 2.27$, $p = 0.03$) and information processing ($z = -2.54$, $p = 0.01$) than those with SCP.

Conclusion: People with DCP present lower general intellectual functioning and poorer EF across multiple domains than typically developing controls. People with DCP have higher general intellectual functioning and better EF than people with SCP when levels of motor severity are similar.

© 2019 European Paediatric Neurology Society. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Although cerebral palsy (CP) is primarily a disorder of movement and posture, it often involves disturbances of communication and cognition¹ which may have an impact on quality of life.² Dyskinetic cerebral palsy (DCP) is the second largest CP group, with high rates among children born at term and among children with normal birth weight. It is agreed that DCP presents poorer gross motor function than other CP types³ but there is less scientific evidence of the presence of poorer cognitive abilities in comparison with other CP types, especially when levels of gross motor severity are equivalent between groups.^{3–10} Dyskinetic cerebral palsy is characterized by involuntary, uncontrolled, recurring and occasionally stereotyped movements. Primitive reflex patterns predominate, muscle tone is varying, and dyskinesia is differentiated into dystonia and choreoathetosis.^{11,12}

Some studies have reported that people with DCP present higher rates of cognitive difficulties. Specifically, a study including 474 participants with DCP (16% walked without aids, 24% with aids and 59% were confined to a wheelchair) and 4746 participants with bilateral spastic cerebral palsy (SCP) (36% walked unaided, 42% unable to walk) found learning disability to be more frequent in children with DCP (52%) than in children with bilateral SCP (33%).³ Similarly, another study reported that people with DCP have a lower intelligence quotient (IQ) than people with SCP.⁶ Verbal IQ was also found to be significantly higher among children with spastic diplegia ($n = 49$) and hemiplegia ($n = 35$), and lower in those with DCP ($n = 11$).⁷ Neither of those studies reported the Gross Motor Function Classification System (GMFCS) level separately for participants with DCP and other CP types. Only Himmelmann et al.³ reported gross motor severity by CP type. While 59% of participants with DCP needed wheelchair, only 2% of participants with SCP needed it. Interestingly, Pueyo et al.⁴ did not find differences between bilateral DCP, mixed CP, and SCP in nonverbal reasoning using the same measure in all participants. Strauss et al.⁵ found that in a sample of people with tetraplegic CP with severe motor dysfunction (defined as tetraplegia with no functional hand use and inability to crawl, creep, scoot, or walk), 95% of those with SCP had an IQ < 50 and only 2% had no intellectual disability. Among the participants with DCP, only 40% ($n = 8$) had an IQ < 50, and 20%

($n = 4$) had an IQ ≥ 70 or higher. However, the authors reported in the study that “the reliability of the cognitive assessments [was] untested”. Indeed, another study⁸ found that the percentage of learning disability was higher (100%) in people with spastic tetraplegia (GMFCS IV $n = 2$, V $n = 21$) than in DCP (60%; GMFCS I $n = 2$, II $n = 4$, III $n = 5$, IV $n = 15$, V $n = 26$). Moreover, a recent study suggested that “cognitive quotient” in tetraplegic SCP may be significantly lower than other CP types, including DCP.⁹ Finally, Dalvand et al. found that the lowest IQ level was more frequent in spastic tetraplegia than in DCP. Specifically, in this study hypotonic, spastic tetraplegic, and hemiplegic but not dyskinetic participants had the highest odds to assign higher ratings in impaired IQ.¹⁰ Overall, these studies might suggest that when people with DCP are comparable to other CP types in terms of gross motor severity, they might not present poorer cognitive performance. Some aspects of these studies, however, preclude drawing firm conclusions. For example, some studies have small samples of participants with DCP, gross motor severity by CP type is not always reported and groups are not comparable in terms of gestational age and gross motor severity in most studies. Moreover, the different CP classifications and the different terms used between some studies make it difficult to compare results. Table 1 shows further details about the above-mentioned studies.

Interestingly, a systematic review focused on cognition in childhood dystonia has been recently published.⁹ Although this work is not specifically focused on secondary dystonia due to CP, it indicates that people with DCP often have mild deficits in memory, impairments of visuospatial functions, information processing speed, and social cognition. This review stresses that available data is very limited and that there is a strong need for case–control studies assessing cognition and using standardized neuropsychological tests, with particular emphasis on attention and executive functioning skills among others. In this review it is further encouraged to assess control groups consisting of patients with other movement disorders.¹³

Executive function (EF) is necessary for the successful completion of everyday, novel, goal-directed activities and has been associated with quality of life in DCP.¹⁴ Most studies of EF in CP focus on SCP or unilateral CP, and have shown that people with CP perform significantly worse than typically developing children¹⁵ in all EF domains described by Anderson.¹⁶ Briefly, these domains include attentional control

Table 1 – Studies comparing/reporting percentages of general intellectual functioning in dyskinetic and spastic cerebral palsy.

	n	Age range	Motor assessment/Motor severity by CP type	CP type (n)	IQ assessment	Results
Stadskleiv et al., 2018 ⁹	70	5 y:1 m–17 y:7 m	GMFCS/- (GMFCS range of the whole sample I–V)	DCP (8) SCP tetraplegia (9) SCP hemiplegia (35) SCP diplegia (18)	Cognitive quotient measured by one of the following options: 1) Cognitive Ability Quotient: Verbal comprehension Test for Reception of grammar Vocabulary score (British Picture Vocabulary Scale and/or Receptive Vocabulary) Perceptual reasoning WPPSI-III/WISC-IV (substituting Block Design with Picture Completion for children with severe fine motor impairments) 2) Developmental Quotient BSID-III/Receptive Vocabulary WPPSI, WISC-R	Significantly lower cognitive quotient in the tetraplegic SCP group DCP 75% IQ ≥ 85 SCP tetraplegia 33% IQ ≥ 85 SCP hemiplegia 86% IQ ≥ 85 SCP diplegia 78% IQ ≥ 85 Mixed CP 76% IQ ≥ 85
Dalvand et al., 2012 ¹⁰	662	3 y–14 y	GMFCS/-	DCP (53) SCP tetraplegia (218) SCP hemiplegia (57) SCP diplegia (223) Mixed CP (63) Ataxic CP (18) Hypotonic (30)	WPPSI, WISC-R	DCP 26% IQ ≥ 85; 9% IQ 71–84; 19% IQ 50–70; 45% IQ < 50 SCP tetraplegia 8% IQ ≥ 85; 5% IQ 71–84; 10% IQ 50–70; 77% IQ < 50 SCP hemiplegia 25% IQ ≥ 85; 21% IQ 71–84; 28% IQ 50–70; 26% IQ < 50 SCP diplegia 24% IQ ≥ 85; 18% IQ 71–84; 22% IQ 50–70; 35% IQ < 50 Mixed CP 33% IQ ≥ 85; 21% IQ 71–84; 13% IQ 50–70; 33% IQ < 50

Sigurdardottir et al., 2011 ⁷	111	4 y:6 y	GMFCS/- (GMFCS range of the whole sample I–V)	DCP (11) SCP tetraplegia (10) SCP hemiplegia (35) SCP diplegia (49) Ataxic CP (6)	n = 29 BSID-II n = 4 Reynell Zinkin Developmental Scales n = 9 Columbia Mental Maturity Scale n = 4 The Leiter International Performance Scale	Verbal IQ was lower in DCP than SCP
Himmelmann et al., 2009 ³	5220	Years of birth 1991–1996	Severe: needing a wheelchair Moderate: ambulation with aids Mild: ambulation without aids/ DCP: 16% mild, 24% moderate, 59% severe SCP bilateral: 36% mild, 42% severe	DCP (474) SCP bilateral (4746)	Measure not specified Categorized into IQ above/ below 50	DCP 52% IQ<50 SCP bilateral 33% IQ<50
Sigurdardottir et al., 2008 ⁶	118	4 y–6y: 6 m	GMFCS/- (GMFCS range of the whole sample I–V)	DCP (14) SCP tetraplegia (28) SCP hemiplegia (31) SCP diplegia (45)	50% WPPSI 26% WPPSI in combination with the other developmental scales 12% BSID-II 10% BSID-II in combination with other developmental scales n = 4 Reynell Zinkin Scales n = 5 Columbia Mental Maturity Scale n = 1 Leiter International Performance Scale-Revised in combination with TONI-2	Median IQ was lower in DCP than SCP DCP 29% IQ/DQ ≥ 85; 14% IQ/DQ 70 to 84; 21% IQ/DQ 50 to 69; 36% IQ/DQ <50 SCP tetraplegia 50% IQ/DQ ≥85; 14% IQ/DQ 70 to 84; 21% IQ/DQ 50 to 69; 14% IQ/DQ <50 SCP hemiplegia 61% IQ/DQ ≥85; 19% IQ/DQ 70 to 84; 13% IQ/DQ 50 to 69; 6% IQ/DQ <50 SCP diplegia 53% IQ/DQ ≥85; 18% IQ/DQ 70 to 84; 22% IQ/DQ 50 to 69; 7% IQ/DQ <50
Himmelmann et al., 2006 ⁸	70	4 y–8y	GMFCS/DCP (GMFCS I n = 2, II n = 4, III n = 5, IV n = 15, V n = 26) SCP (GMFCS IV n = 2, V n = 21)	DCP (50) SCP tetraplegia (20)	WPPSI-R, WISC-III, Griffith scales or estimated from clinical observation	DCP 40% IQ > 70; 18% IQ 50-70; 42% IQ<50 SCP tetraplegia 100% IQ<50
Strauss et al., 2005 ⁵	446	–	Severe: tetraplegia with no functional hand use and inability to crawl, creep, scoot, or walk/All participants with severe motor dysfunction	DCP (20) SCP (426)	Measure not specified	DCP 20% IQ ≥ 70; 40% IQ<25 (profound mental retardation) ⁺ SCP 2% IQ ≥ 70; 95% IQ<50 severe mental retardation (of which 75% profound IQ<25)

(continued on next page)

Table 1 – (continued)

n	Age range	Motor assessment/Motor severity by CP type	CP type (n)	IQ assessment	Results
19	16 y–38 y	-/All bilateral	DCP tetraplegia (2) DCP diplegia (2) DCP triplegia (2) SCP tetraplegia (5) SCP diplegia (6) SCP triplegia (2)	RCPM	No statistically significant differences between DCP and SCP

-. data not provided; +: The term used is the same that authors use; CP: cerebral palsy; BSID, Bayley Scales of Infant Development; DCP: dyskinetic cerebral palsy; DQ: developmental quotient; GMFCS: Gross Motor Function Classification System; ICD10: International Statistical Classification of Diseases and Related Health Problems 10th Revision; IQ: intelligence quotient; m: months; RCPM: Raven's coloured progressive matrices; SCP: spastic cerebral palsy; TONI-2: Test of Nonverbal Intelligence, 2nd Edition; WPPSI: Wechsler Preschool and Primary Scale of Intelligence; WISC: Wechsler Intelligence Scale for Children; y: years.

(capacity to selectively attend to specific stimuli, to focus attention for a prolonged period, and impulse control), cognitive flexibility (ability to shift between response sets, learn from mistakes, devise alternative strategies, divide attention, apply working memory and process multiple sources of information simultaneously), goal setting (ability to initiate an activity and devise a plan to complete it) and information processing (fluency, efficiency and speed of output).¹⁶ Although it has been proposed that DCP may be characterized by an executive dysfunction,⁴ this hypothesis has still to be tested. The hypothesis is consistent with the lesions of the basal ganglia and thalamic systems which may impair focused attention and EF¹⁷ and are frequently described in people with DCP.¹⁸ Executive functions depend on the integrity of the entire brain but they are mainly mediated by the frontal lobes and their connections with posterior and subcortical brain regions.^{15,16} Specifically, some circuits in the basal ganglia originate in the prefrontal and limbic regions of the cortex which are known to be involved in the EF.¹⁵ If we focus on the differences between CP types, one study⁴ found that EF was the only function in which mean performance was poorer in DCP participants than in participants with SCP; however, the differences were not statistically significant, the sample was small, not all domains of EF were assessed, and other variables that may influence EF such as gestational age and gross motor severity were not controlled. When aiming to identify the association between spasticity/dyskinesia and cognition, influential variables other than CP type, such as gross motor severity and prematurity, should be taken into account, since they may prompt additional cognitive impairments.¹⁹

The International Classification of Functioning, Disability and Health considers intellectual functions to be an essential Core Set for people with CP,²⁰ and intellectual functions play an important role in communication,²¹ employment²² and quality of life.¹⁴ Nevertheless, few studies so far have focused on cognitive functions in DCP or have analysed the EF profile in depth, and the results they present are partly conflicting. Most of them have used different measures for assessing cognition in the participants, and the results vary depending on whether the participants with DCP are compared with participants with severe or moderate forms of SCP. No study to date has compared cognitive functioning between a relatively large sample of people with DCP and a group of participants with SCP who are similar in terms of gross motor severity and prematurity. To better understand the cognitive correlates of different CP types, neuropsychological studies able to differentiate between subtypes of the condition are required.²³ The results would help to guide the design of more appropriate interventions and follow-up programs focused on DCP.

Thus, in the present study we aim to map general intellectual functioning and EF in people with DCP by comparing their performance with: 1) typically developing controls (TDC) matched for age and sex, and 2) participants with SCP matched for age, sex, gestational age and gross motor severity. Following on from previous studies, our primary hypothesis was that general intellectual functioning and EF would be poorer in participants with CP compared with TDC. Our second hypothesis was that, with similar gross motor

severity in the two CP groups, general intellectual functioning would be higher in participants with DCP than in those with SCP. This hypothesis is in accordance with the studies mentioned above which seem to preliminarily indicate that when people with DCP are comparable to other CP types in terms of gross motor severity, their cognitive performance tends to be as good as that of the other CP types or better. Furthermore, taking into account the brain lesions described in DCP and the results of the only previous study to analyse EF differences between CP types,⁴ we expected performance on EF to be poorer in participants with DCP.

2. Method

2.1. Participants

This study recruited participants with DCP and a comparison group with SCP from the main hospitals in Barcelona that monitor people with CP (the Pediatric Neurology Department and Rehabilitation and Physical Medicine Department at the Hospital Universitari Vall d'Hebron and the Neurology Department at the Hospital Sant Joan de Déu), other institutions attending people with CP (Health Services and Rehabilitation Services of the ASPACE Cerebral Palsy Association, Centro Ocupacional Sinia, and Nadís). Some of them were participants in a previous study⁴ who were contacted and invited to participate in the current study. Cases were also compared with a group of TDC composed of relatives/friends of the participants with CP and people recruited through advertisements.

The inclusion criteria for participants with clinical diagnosis of CP with predominant dyskinetic features were (1) age older than 6 years and (2) ability to understand instructions, as evaluated by the Spanish Grammar Screening Test (receptive part).²⁴ Exclusion criteria were (1) presence of severe visual or auditory disability and (2) lack of an intelligible yes/no response. In case of signs of sensory impairment, only participants in whom the deficit was corrected or/and the sensory impairment did not prevent evaluation were included.

After the recruitment of the participants with DCP, participants who had a clinical diagnosis of SCP and met the inclusion and exclusion criteria described above were recruited and matched by sex (male/female) and age with participants with DCP (Supplemental material 1). Since the influence of age on cognitive performance is stronger in younger people, the age matching criterion was more flexible with older participants: <20 years old \pm 2 years; 20 < 30 years old \pm 4 years; \geq 30 years old \pm 8 years. In order to control for variables other than the CP type, additional inclusion criteria were implemented to ensure that each participant of the SCP group matched a participant with DCP with regards to term vs preterm (\geq 37 weeks, <37 weeks) and as ambulant vs non-ambulant (GMFCS I-II-III vs IV-V). That is, the prematurity and ambulation level of participants with SCP included was driven by the characteristics of participants with DCP.

Physicians from the institutions mentioned above informed their patients with DCP or SCP who met the inclusion criteria, or their careers, about the possibility of

participating in this project. The diagnosis of CP type was based on the neurologist's clinical assessment. All physicians agreed to define DCP as the CP type characterized by abnormal patterns of posture and/or movement accompanied by involuntary, uncontrolled, recurring, and occasionally stereotyped movements. Impaired muscle tone regulation, movement control, and coordination may comprise dystonic and choreoathetotic patterns.¹¹ Only participants who clearly presented predominant dyskinetic symptoms were included in the DCP group. Participants who presented spastic and dyskinetic symptoms to the same extent were not considered eligible. Participants were further contacted by phone to double-check inclusion/exclusion criteria, to explain the participation procedure, and to take part in the study; the recruitment and data collection period was from 2012 to 2016.

Typically developing people without brain pathology were matched one-to-one by age and sex with participants with DCP. Controls were ineligible if they had been born preterm, were diagnosed with a neurological or psychiatric disorder, or were illicit substance consumers.

All procedures performed in the study complied with the ethical standards of the 1964 Helsinki declaration. Ethical approval was obtained by the University of Barcelona's (CBUB) Institutional Ethics Committee, Institutional Review Board (IRB 00003099, assurance number: FWA00004225; <http://www.ub.edu/recerca/comissiobioetica.htm>) and the Hospital Universitari Vall d'Hebron. Written informed consent was obtained from all participants or their careers.

2.2. Neuropsychological assessment

Tests were carefully chosen to allow most of the participants to answer in an autonomous way and to permit, when possible, the use of assistive technology for communication. Participants were encouraged to use the response technique best suited to their degree of disability and the communication devices they normally used. See Supplemental material 2 for test details and adaptations used.

The Raven's Coloured Progressive Matrices (RCPM) test was used to measure general intellectual functioning and the four EF¹⁶ domains were assessed as follows:

Attentional control

- Inhibition and sustained attention components were assessed using an adapted version of the Stop Signal Task (SST) of the Cambridge Neuropsychological Test Automated Battery (CANTAB).²⁵
- Selective verbal attention was assessed using the digit forward span from either the Wechsler Intelligence Scale for Children (WISC-IV)²⁶ or the Wechsler Adult Intelligence Scale (WAIS-III).²⁷
- Selective visual attention was assessed using the spatial forward span from the Wechsler Nonverbal Scale of Ability (WNV).²⁸

Cognitive flexibility

- Feedback utilization was assessed using the 64-item computerized version of the Wisconsin Card Sorting Test (WCST).²⁹

- Verbal working memory was assessed using the digit backward span from either WISC-IV²⁶ or WAIS-III.²⁷
- Visual working memory was assessed using the backward spatial span from the WNV.²⁸

Goal setting ability was evaluated by means of the Stockings of Cambridge (SOC) test from the CANTAB.²⁵ Three scores were used: problems solved in minimum moves, mean moves in items of two moves and mean moves in items of five moves.

Information processing was assessed only in verbal participants, using a lexical verbal fluency task.

2.3. Background measures

Gross motor function was determined based on the GMFCS³⁰ while manual ability was measured by the Manual Ability Classification System (MACS).³¹ The CFCS (Communication Function Classification System) was used to categorize communication.³² Gestational age and epilepsy status were recorded from parent interviews and medical records.

2.4. Statistical analysis

Statistical analyses were performed using R version 3.3.1 (<https://www.r-project.org/>). Descriptive statistics were calculated, and a global test based on Kruskal–Wallis and Chi-squared test was used to check differences between groups in matching and descriptive variables. Medians and interquartile ranges (IQR) were used due to the presence of skewness in the empirical distribution of some numerical variables.

Matched pairs were not included in the analyses as blocks because only one measure was recorded for each pair and, therefore, the block effect could not be separated from error. As a result, global tests based on Kruskal–Wallis (H statistic) were performed to analyse differences in cognitive performance between groups. Pairwise contrasts, based on the Mann–Whitney U test, were performed in cases in which the global test yielded a significant result. The Hodges–Lehmann estimator, as well as a 95% bootstrap-percentile confidence interval, was estimated in order to report the differences between groups. The p values were corrected for multiple comparisons (for global and posteriori contrasts) and we used an alpha level of 0.05 for all statistical tests. As for the reported effect size, probability of superiority was used. Given its relationship with Cohen's d under certain conditions, probabilities greater than 0.56, 0.64, and 0.71 (or lower than 0.44, 0.36, and 0.29, depending on the order in which groups are compared) could be regarded respectively as small, medium, and large effects. Further details about the statistical analysis are available in [Supplemental material 3](#).

3. Results

3.1. Participants

The final sample comprised 52 participants with DCP with age range 6–62 and median age 20.5 y: 5mo, IQR = 13.75 y:

7mo without visual/auditory abnormalities precluding neuropsychological assessment, able to understand instructions and, at least, able to answer yes/no. Five of these 52 participants had sensorineural hearing loss (two mild, one moderate and two severe hearing loss – the last three using hearing aids), 17 subjects with corrected refractive errors (myopia, astigmatism and hypermetropia), four subjects with strabismus (two corrected and two non-corrected), two who showed a slight delay in signal transmission in visual evoked potentials but without pathological values and one with decreased visual acuity in one eye. The remaining cases did not report any hearing or visual problem. A comparison group of 52 age- and sex-matched TDC with an age range 7–59 and age median age 20 y: 4mo (IQR = 12 y: 7mo) were recruited. The final sample for the group with SCP comprised 20 participants (10 females), age range 7–65 and median age = 20.5 y: 5.5mo (IQR = 13.75 y: 9mo). The recruitment process is described in [Fig. 1](#). Due to absence or unintelligible speech and motor severity or comprehension difficulties, some participants were unable to complete some of the cognitive tests used. These missing values are indicated in the footnote of [Table 3](#).

Given the difficulty of finding participants with SCP who met the matching criteria and due to the completion of the research project, a one-to-one matching was not possible. The range of age and GMFCS of participants were similar in both CP groups. For detailed information on the matching, see [Supplemental material 1](#). There were no significant differences in age and sex between the three groups, as regards the rates of term, preterm and very preterm participants ($\chi^2 = 2.67$, $p = 0.26$) or GMFCS between CP groups. Demographic and clinical data for the three groups are shown in [Table 2](#).

3.2. General intellectual functioning differences between the three groups

Both CP groups had significantly lower scores than TDC ([Table 3](#), [Fig. 2](#)), and participants with DCP had significantly higher scores than participants with SCP. The differences were large for DCP and SCP vs TDC and medium for DCP vs SCP.

3.3. Executive function differences between the three groups

Attentional control: Inhibition/sustained attention measured by the SST did not differ significantly between the three groups. Both selective verbal and visual attention were significantly poorer in DCP and SCP than TDC; there were no differences between CP groups.

Cognitive flexibility: Feedback utilization as well as verbal and visual working memory were significantly poorer in DCP and SCP than TDC. There were no significant differences between CP groups in any cognitive flexibility domain.

Goal setting: The mean number of moves in problems that can be solved in two moves did not differ significantly between the three groups. The mean number of moves in problems that can be solved in five moves was significantly lower in TDC and DCP than in SCP. There were no significant differences between DCP and TDC. Both CP groups had significantly worse global scores (problems solved in the




Fig. 1 – Flowchart showing recruitment process for dyskinetic and spastic cerebral palsy participants. DCP, dyskinetic cerebral palsy; SCP, spastic cerebral palsy.

Table 2 – Participants' characteristics.

Matching variables	DCP	SCP	TDC	Differences: Statistic (p-value)*
Age	20.5: 5 (13.75: 7)	20.5: 5.5 (13.75: 9)	20: 4 (12: 7)	H = 0.01 (p = 0.99)
Md years: months (IQR years: months)				
Sex	24/28	10/10	24/28	$\chi^2 = 0.02$ (p = 0.99)
n (female/male)				
Gestational age	4/6/42	4/1/15	–	$\chi^2 = 2.67$ (p = 0.26)
n (<32 weeks/32–36 weeks/≥37 weeks)				
Gestational age participants born preterm n/median weeks (IQR weeks)	10/34 (6.5)	5/28 (4.5)	–	z = 1.48 (p = 0.12)
Gross motor function (GMFCS) (n)	I (15) II (8) III (6) IV (11) V (12)	I (5) II (3) III (4) IV (5) V (3)	–	z = 0.14 (p = 0.89)
Other participants' characteristics				
Motor distribution	44/8	16/4		
n (bilateral/unilateral)				
Manual ability (MACS) (n)	I (5) II (10) III (17) IV (10) V (10)	I (1) II (5) III (6) IV (4) V (4)	–	z = 0.16 (p = 0.88)
Communication (CFCS) (n)	I (17) II (23) III (6) IV (6)	I (6) II (10) III (2) IV (2)	–	z = 0 (p = 1)
Epilepsy status ^b	30/16/6	11/4/5	–	$\chi^2 = 0.93$ (p = 0.63)
n (no epilepsy/active/resolved)				
Aetiology ^a				
HIE	22	6	–	$\chi^2 = 7.47$ (0.11)
Intra-cranial haemorrhage, infarction or hydrocephalus	9	7	–	
Infection	2	2	–	
Kernicterus	2	0	–	
Unclassifiable	17	5	–	
Main communication system ^c n (Speech/Speech ^g gestures, communication board and other tools/gaze, facial expressions and gestures/communication board/alphabetic communication board)	37/5/3/5/2	17/1/2/0/0	–	

CFCS, Communication function classification system; DCP, dyskinetic cerebral palsy; GMFCS, Gross motor function classification system; HIE, hypoxic-ischemic encephalopathy; IQR, interquartile range; MACS, Manual ability classification system; SCP, spastic cerebral palsy; TDC, typically developing controls; –, not applicable; χ^2 Pearson's Chi-squared test; z, transformed score from Mann–Whitney U test.

^a Aetiology classification was based on clinical criteria complemented by the information from available neuroimaging, and HIE criteria were based on Himmelmann et al.⁴⁰

^b The International League Against Epilepsy criteria were used to determine epilepsy status; *The level of significance was set at a p-value < 0.05 and p values were corrected for multiple comparisons (for global and posteriori contrasts); –, not applicable (note that all typically developing controls were born at term).

^c Here there are presented the main communication system used in their daily life but most of participants used other complementary systems depending on the context or nature of the content they want to communicate.

minimum number of moves) than TDC. There were no significant differences between CP groups in problems solved in the minimum number of moves.

Information processing: Both CP groups showed significantly poorer verbal fluency than TDC. Verbal fluency was significantly higher in DCP than in SCP.

Interestingly, in the variables in which both CP groups performed worse than TDC but did not differ from each other, mean scores were always higher (though not significantly) in participants with DCP (Table 3, Fig. 2).

4. Discussion

The present study contributes to the characterization of cognitive impairments in DCP. To our knowledge, it is the first study to compare general intellectual functioning and performance in all EF domains in a relatively large sample of people with DCP against samples of TDC and people with SCP who are similar in terms of age, sex, gestational age and gross motor severity.

Table 3 – Descriptive statistics and comparisons of cognitive scores between participants with dyskinetic cerebral palsy, typically developing controls and participants with spastic cerebral palsy.

	n/Md (IQR)			General contrasts H (<i>p</i> value)*	Posteriori contrasts	z (adjusted <i>p</i> value)	HL estimator (95% CI)	Effect size (probability of superiority)
	DCP	TDC	SCP					
General intellectual functioning								
	52/27.5 (11)	52/35 (2)	20/24.5 (9.75)	55.49 (<i>p</i> < 0.01)	TDC - DCP	5.96 (<i>p</i> < 0.01)	7 (3;–9)	0.79 ^d
					SCP - DCP	–2.51 (<i>p</i> = 0.01)	–5 (–9;–1)	0.29 ^e
					SCP - TDC	–6.22 (<i>p</i> < 0.01)	–11 (–15;–9)	0.02 ^d
Executive functioning								
Attentional control								
Inhibition and sustained attention ^a	49/283 (17)	52/283 (13.25)	18/291.5 (18.75)	2.02 (<i>p</i> = 0.36)	TDC - DCP	–		
					SCP - DCP	–		
					SCP - TDC			
Selective verbal attention ^a	47/5 (2)	52/6 (2)	19/5 (2)	24.76 (<i>p</i> < 0.01)	TDC - DCP	4.61 (<i>p</i> < 0.01)	1 (1;2)	0.68 ^e
					SCP - DCP	0.00 (<i>p</i> = 1)	0 (–1;1)	0.38 ^f
					SCP - TDC	–3.36 (<i>p</i> < 0.01)	–1 (–2;–1)	0.16 ^d
Selective visual attention ^a	47/5 (2)	52/6 (1)	19/4 (1)	46.21 (<i>p</i> < 0.01)	TDC - DCP	5.79 (<i>p</i> < 0.00)	2 (1;2)	0.68 ^e
					SCP - DCP	–1.64 (<i>p</i> = 0.10)	–1 (–1;0)	0.38 ^f
					SCP - TDC	–5.29 (<i>p</i> < 0.00)	–2 (–3;–2)	0.16 ^d
Cognitive flexibility								
Feedback utilization ^{†,a}	50/9.5 (8)	52/7.5 (4)	18/10 (16.25)	7.44 (<i>p</i> = 0.03)	TDC - DCP	–2.21 (<i>p</i> = 0.04)	–2 (–4;0)	0.34 ^e
					SCP - DCP	0.88 (<i>p</i> = 0.38)	1 (–2;6)	0.55 ^f
					SCP - TDC	2.19 (<i>p</i> = 0.04)	3 (0;9)	0.64 ^f
Verbal working memory ^{a,c}	47/4 (2)	52/5 (2)	19/3 (1)	37.06 (<i>p</i> < 0.01)	TDC - DCP	4.83 (<i>p</i> < 0.01)	2 (1;2)	0.71 ^e
					SCP - DCP	0.00 (<i>p</i> = 0.20)	0 (–1;0)	0.28 ^e
					SCP - TDC	–5.24 (<i>p</i> < 0.01)	–2 (–3;–1)	0.03 ^d
Visual working memory ^{a,c}	47/5 (2.5)	52/6 (0.25)	18/4 (1)	50.55 (<i>p</i> < 0.01)	TDC - DCP	5.85 (<i>p</i> < 0.01)	2 (1;2)	0.75 ^d
					SCP - DCP	–1.54 (<i>p</i> = 0.12)	–1 (–1;0)	0.28 ^d
					SCP - TDC	–5.96 (<i>p</i> < 0.01)	–2 (–3;–2)	0.02 ^d
Goal setting								
Mean moves in tests of two moves ^{i,c}	48/2 (0)	52/2 (0)	17/2 (0)	5.57 (<i>p</i> < 0.07)	TDC - DCP	–		
					SCP - DCP	–		
					SCP - TDC	–		
Mean moves in tests of five moves ^{i,c}	46/6.62 (2.44)	52/6.5 (1.75)	17/7.75 (1.75)	13.79 (<i>p</i> < 0.01)	TDC - DCP	–1.80 (<i>p</i> = 0.07)	–0.50 (–1;0)	0.36 ^f
					SCP - DCP	2.27 (<i>p</i> = 0.03)	1.12 (0.25;1.75)	0.67 ^e
					SCP - TDC	3.68 (<0.01)	1.50 (0.75;2.25)	0.77 ^d

(continued on next page)

Table 3 – (continued)

	n/Md (IQR)		General contrasts H (p value)*	Posteriori contrasts	z (adjusted p value)	HL estimator (95% CI)	Effect size (probability of superiority)
	DCP	TDC					
Problems solved in minimum moves ^a	48/8 (3)	52/9 (2.25)	24.69 (p < 0.01)	TDC - DCP SCP - DCP SCP - TDC	3.48 (p < 0.01) -1.91 (p = 0.06) -4.55 (p < 0.01)	2 (1;2) -1 (-3;0) -3 (-4;-2)	0.64 ^f 0.29 ^e 0.10 ^d
Information processing Lexical verbal fluency ^b	37/30 (23)	52/39.5 (14.75)	36.21 (p < 0.01)	TDC - DCP SCP - DCP SCP - TDC	4.32 (<0.01) -2.54 (p = 0.01) -5.15 (p < 0.01)	13 (7;19) -11 (-20;-2) -25 (-31;-18)	0.76 ^d 0.27 ^d 0.09 ^d

*The level of significance was set at a p-value < 0.05 and p values were corrected for multiple comparisons (for global and posteriori contrasts); † Higher scores indicate worse performance; -, not applicable; CI, Bootstrapped percentile confidence interval based on 10,000 bootstrap samples; DCP, dyskinetic cerebral palsy; H, Kruskal-Wallis statistic; HL, Hodges-Lehmann estimator; IQR, interquartile range; Md, median; SCP, spastic cerebral palsy; TDC, typically developing controls; z, transformed score from Mann-Whitney U-test. Reasons for missing data (handled with pairwise deletion):
^a absence or inintelligible speech and motor severity; ^b absence or inintelligible speech; ^c No comprehension of test instructions; Effect size: ^d , large effect size; ^e , medium effect size; ^f , small effect size.

As expected, the comparison between TDC and CP groups showed that people with CP had lower general intellectual functioning and poorer executive function across all the domains described by Anderson.¹⁶ Differences between CP and TDC in all EF domains have previously been reported in unilateral CP.¹⁵ In the present study, performance did not differ significantly between TDC and CP groups for inhibition/sustained attention and for the easiest items of goal setting. The lack of differences between CP and TDC on the easiest items of goal setting shows that CP performance may be similar that of TDC in very basic planning tasks. The lack of differences in inhibition/sustained attention is possibly due to the fact that SST may overestimate the performance of some of the CP participants on “stop trials” (25% of the total score used), as motor slowness precludes inhibition errors.

Our results corroborate those of previous studies which have indicated that learning or intellectual disability is more common in spastic tetraplegia than in DCP.^{5,8,9} Although our results seem to be at odds with those of Sigurdardottir et al.,⁶ when those authors took into account participants with DCP who could not be assessed by the same test as the rest of the sample (and were therefore assessed by different measures) the highest proportion of children with intellectual impairment was found in the group with spastic tetraplegia. In fact, the authors themselves concluded that cognitive skills might be masked by limitations of motor control.⁶ Our results, are also at odds with those of other studies.^{3,4,7} The differences with regard to two of these studies^{4,7} may be due to the small size of the sample of participants with DCP. Himmelmann et al.,³ however, studied a large sample of DCP participants; the differences between our study and theirs might be due to the fact that they did not control for GMFCS and prematurity, and that general intellectual functioning was reported dichotomously using an IQ cut-off point of 50. It has been suggested that the more severe the motor impairments, the higher the percentage of cognitive impairment³ and so DCP has often been associated with poorer cognitive outcomes. However, our results show that, at similar levels of gross motor severity, dyskinetic forms may present higher general intellectual functioning than spastic forms. Accordingly, some studies suggest that intellectual disability, rather than the degree of motor involvement, is a predictor of verbal comprehension abilities.^{21,33,34} Overall, these findings seem to support the hypothesis that people with DCP, even if they often present poorer gross motor functioning than people with SCP, do not necessarily present poorer cognitive functioning. It is worthy of note that similar mismatches have previously been found between motor impairment and communication abilities in DCP.³⁵

Participants with DCP performed significantly better for goal setting and information processing than participants with SCP. Better performance in goal setting was observed for the most difficult items; that is, both CP groups performed similarly on relatively easy items, but people with DCP performed significantly better when more complex planning is required. In fact, their performance on the most difficult items was close to the performance of TDC, since no significant differences were found. Given that information processing was assessed with a verbal fluency task, the conclusions about the better performance observed in DCP than SCP must be




Fig. 2 – Boxplots showing raw scores (y axis) by groups (x axis) of intellectual functioning, goal setting (mean moves in tests of five moves) and information processing. Only boxplots showing significant differences between participants with dyskinetic and spastic cerebral palsy in Table 3 are presented. * $p < 0.05$; ** $p < 0.01$; † Higher scores indicate poorer performance in this task; DCP, dyskinetic cerebral palsy; IQR, interquartile range; Md, median; SCP, spastic cerebral palsy; TDC, typically developing controls.

limited to verbal participants. This fact may have biased information processing scoring toward a better performance in both CP groups. Information processing is inherently temporal, so clinical testing of this process is affected by a participant's ability to form a response. In the authors' opinion, comparing performance against people with similar motor difficulties as we have done by comparing DCP and SCP groups is a promising way to control this limitation. Further studies should focus on developing tasks that control this effect at the maximum level possible but, presently, this is a limitation difficult to overcome when researching performance in this kind of tasks in people with CP.

Overall, our results show that people with DCP perform worse than TDC and better than SCP in both general intellectual functioning and EF indicating a general tendency, rather than a specific dysexecutive deficit. The performance of people with DCP tends to be closer to TDC on EF than on general intellectual functioning. At this point it has to be noted that the measure used to estimate general intellectual functioning (Raven's coloured progressive matrices) is a matrix reasoning task. Although the Raven's progressive matrices is regarded as a quintessential measure of fluid intelligence,³⁶ it aims to measure the ability to deduct relationships³⁷; other aspects considered in psychometric tests of cognitive abilities such as verbal ability or crystallized intelligence are not considered. Moreover, it has been suggested that the Raven's coloured progressive matrices gives comprehensive information on cognitive performance and it is sensitive to the structural state of the brain in DCP.³⁸ The executive function results comparing DCP and SCP groups do not confirm DCP as a CP type specifically characterized by an executive dysfunction. Further research including neuropsychological assessment of other cognitive domains is needed to conclusively reject the hypothesis of a dysexecutive deficit in DCP.

The strengths of this study include the recruitment of a large sample of participants with DCP taking into account that this CP type is relatively rare; a wide assessment using reliable and common measures in all participants and the lack of differences in age, sex, GMFCS and prematurity between groups. The limitations include the absence of measurement of dyskinesia and speech production using a quantitative scale, the wide age range of the sample and the lack of a wider description of the functioning and disability of the participants according the International Classification of Functioning, Disability and Health. Moreover, neuro-ophthalmological disorders are among the main symptoms in CP³⁹ and their interaction with manual ability may have an effect on cognitive performance. To control this effect, the execution time was not considered when scoring performance, but SST and verbal fluency are influenced by execution time. In this regard, it is interesting that there were no significant differences in manual ability between DCP and SCP groups ($z = 0.16$, $p = 0.88$). Additionally, considering neuro-imaging as a further matching variable between groups would provide meaningful information and further strength the results of our study. Finally, participants with SCP included in the present study are not representative of the entire SCP population and the sample is small. Caution is therefore required when interpreting the results referring to this group. This is due to the fact that participants with SCP were included *a posteriori*, based on prematurity, ambulation, age and sex characteristics of participants with DCP.

In conclusion, the present study identified difficulties across multiple executive function domains compared with TDC and a better cognitive functioning in people with DCP than SCP. Although performance observed in our sample cannot be generalized to the broad population of DCP (we have included only participants with enough comprehension, an

intelligible yes/no response and without severe visual or auditory disabilities), the study of specific cognitive functions is only possible in subsamples of DCP such as these. The results are clinically relevant as they suggest that cognitive functions may have been underestimated and masked by gross motor severity in people with DCP. The present study highlights the importance of properly assessing general and specific cognitive functions in CP, even in the most severe cases. Our study may help to broaden the understanding of the clinical consequences of dyskinesia for cognitive function and, by extension, the interaction between cognitive function, muscle tone, and specific brain damage in early childhood. Once again, the results indicate that observations made in SCP cannot be generalized to dyskinetic forms. A comprehensive understanding of cognitive functioning in each CP type would contribute to improving the accuracy of prognosis and also to the design of educational approaches. Although this study increases our current knowledge about DCP, further studies are needed. Harmonizing nomenclature and cognitive measures between studies is also important, as this would allow researchers to compare results, perform meta-analyses and multicentre studies, as well as to pool data for the sake of achieving larger sample sizes.

What this paper adds

This is the first study to compare general intellectual functioning and executive function in a relatively large sample of people with dyskinetic cerebral palsy (DCP) against samples of typically developing controls (TDC) and people with spastic cerebral palsy (SCP) who are similar in terms of age, sex, being term/preterm and gross motor severity. The present study contributes, therefore, to the characterization of cognitive impairments in DCP. Participants with DCP present poorer general intellectual functioning and executive function than TDC in terms of attentional control, cognitive flexibility, goal setting and information processing. People with DCP display stronger general intellectual functioning and executive function than those with SCP. The results suggest that cognitive functions may have been underestimated and masked by gross motor severity in people with DCP. Observations made in SCP cannot be generalized to dyskinetic forms. It is important to properly assess general and specific cognitive functions in people with cerebral palsy (CP) even in the most severe cases.

Funding

This study was supported by the Ministerio de Ciencia e Innovación (grant code PSI2011/24386), by the Ministerio de Economía y Competitividad (PSI2016-75979-R AEI/FEDER, UE) and by Generalitat de Catalunya (2014SGR98 and 2017SGR748). Olga Laporta received a research grant from Ministerio de Educación, Cultura y Deporte of the government of Spain (grant code FPU13/06435).

Conflict of interest

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Acknowledgments

The authors thank all participants and their families for their collaboration. We also thank the Vigatans and ASDI centers for their collaboration during the enrolment process, as well as the UTAC service of the Faculty of Psychology at the University of Barcelona.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejpn.2019.05.010>.

REFERENCES

- Novak I, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012;130(5):e1285–312. <https://doi.org/10.1542/peds.2012-0924>.
- Colver A, Rapp M, Eisemann N, Ehlinger V, Thyen U, Dickinson HO, et al. Self-reported quality of life of adolescents with cerebral palsy: a cross-sectional and longitudinal analysis. *Lancet* 2015;385(9969):705–16. [https://doi.org/10.1016/S0140-6736\(14\)61229-0](https://doi.org/10.1016/S0140-6736(14)61229-0).
- Himmelman K, McManus V, Hagberg G, Uvebrant P, Krageloh-Mann I, Cans C. Dyskinetic cerebral palsy in Europe: trends in prevalence and severity. *Arch Dis Child* 2009;94(12):921–6. <https://doi.org/10.1136/adc.2008.144014>.
- Pueyo R, Junque C, Vendrell P. Neuropsychologic differences between bilateral dyskinetic and spastic cerebral palsy. *J Child Neurol* 2003;18(12):845–50. <https://doi.org/10.1177/088307380301801204>.
- Strauss D, Rosenbloom L, Day S, Shavelle R. Cognitive function in tetraplegic cerebral palsy with severe motor dysfunction. *Dev Med Child Neurol* 2005;47(8):562. <https://doi.org/10.1017/S0012162205001106>.
- Sigurdardottir S, Eiriksdottir A, Gunnarsdottir E, Meintema M, Arnadottir U, Vik T. Cognitive profile in young Icelandic children with cerebral palsy. *Dev Med Child Neurol* 2008;50(5):357–62. <https://doi.org/10.1111/j.1469-8749.2008.02046.x>.
- Sigurdardottir S, Vik T. Speech, expressive language, and verbal cognition of preschool children with cerebral palsy in Iceland. *Dev Med Child Neurol* 2011;53(1):74–80. <https://doi.org/10.1111/j.1469-8749.2010.03790.x>.
- Himmelman K, Beckung E, Hagberg G, Uvebrant P. Gross and fine motor function and accompanying impairments in cerebral palsy. *Dev Med Child Neurol* 2006;48(6):417–23. <https://doi.org/10.1017/S0012162206000922>.
- Stadskleiv K, Jahnsen R, Andersen GL, von Tetzchner S. Neuropsychological profiles of children with cerebral palsy. *Dev Neurorehabil* 2018;21(2):108–20. <https://doi.org/10.1080/17518423.2017.1282054>.

10. Dalvand H, Dehghan L, Hadian MR, Feizy A, Hosseini SA. Relationship between gross motor and intellectual function in children with cerebral palsy: a cross-sectional study. *Arch Phys Med Rehabil* 2012;**93**(3):480–4. <https://doi.org/10.1016/j.apmr.2011.10.019>.
11. Christine C, Dolk H, Platt MJ, Colver A, Prasauskiene A, Krägeloh-Mann I, et al. Recommendations from the SCPE collaborative group for defining and classifying cerebral palsy. *Dev Med Child Neurol* 2007;**109**:35–8.
12. Graham HK, Rosenbaum P, Paneth N, Dan B, Lin J-P, Damiano DL, et al. Cerebral palsy. *Nat Rev Dis Prim* 2016:16005. <https://doi.org/10.1038/nrdp.2016.5>.
13. Coenen MA, Eggink H, Tijssen MA, Spikman JM. Cognition in childhood dystonia: a systematic review. *Dev Med Child Neurol* 2017. <https://doi.org/10.1111/dmcn.13632>.
14. Laporta-Hoyos O, Ballester-Plané J, Póo P, Macaya A, Meléndez-Plumed M, Vázquez E, et al. Proxy-reported quality of life in adolescents and adults with dyskinetic cerebral palsy is associated with executive functions and cortical thickness. *Qual Life Res* 2017;**26**(5):1209–22. <https://doi.org/10.1007/s11136-016-1433-0>.
15. Bodimeade HL, Whittingham K, Lloyd O, Boyd RN. Executive function in children and adolescents with unilateral cerebral palsy. *Dev Med Child Neurol* 2013;**55**(10):926–33. <https://doi.org/10.1111/dmcn.12195>.
16. Anderson PJ. Towards a developmental model of executive function. In: Anderson V, Jacobs R, Anderson PJ, editors. *Executive functions and the frontal lobes: a lifespan perspective*. Philadelphia, PA, US: Taylor & Francis; 2008. p. 3–23.
17. Bottcher L, Flachs EM, Uldall P. Attentional and executive impairments in children with spastic cerebral palsy. *Dev Med Child Neurol* 2010;**52**(2):2010. <https://doi.org/10.1111/j.1469-8749.2009.03533.x>.
18. Himmelmann K, Ahlin K, Jacobsson B, Cans C, Thorsen P. Risk factors for cerebral palsy in children born at term. *Acta Obstet Gynecol Scand* 2011;**90**(10):1070–81. <https://doi.org/10.1111/j.1600-0412.2011.01217.x>.
19. Anderson P, Doyle LW. Victorian Infant Collaborative Study Group. Neurobehavioral outcomes of school-age children born extremely low birth weight or very preterm in the 1990s. *JAMA* 2003;**289**(24):3264–72. <https://doi.org/10.1001/jama.289.24.3264>.
20. Schiari V, Selb M, Cieza A, O'Donnell M. International classification of functioning, disability and Health Core sets for children and youth with cerebral palsy: a consensus meeting. *Dev Med Child Neurol* 2015;**57**(2):149–58. <https://doi.org/10.1111/dmcn.12551>.
21. Vos RC, Dallmeijer AJ, Verhoef M, Van Schie PEM, Voorman JM, Wiegerink DJHG, et al. Developmental trajectories of receptive and expressive communication in children and young adults with cerebral palsy. *Dev Med Child Neurol* 2014;**56**(10):951–9. <https://doi.org/10.1111/dmcn.12473>.
22. Murphy KP, Molnar GE, Lankasky K. Employment and social issues in adults with cerebral palsy. *Arch Phys Med Rehabil* 2000;**81**(6):807–11. <https://doi.org/10.1053/apmr.2000.6798>.
23. Fennell EB, Dikel TN. Cognitive and neuropsychological functioning in children with cerebral palsy. *J Child Neurol* 2001;**16**(1):58–63. <https://doi.org/10.1177/088307380101600110>.
24. Toronto AS. *Screening test of Spanish grammar*. Evenston, IL: Northwestern University Press; 1973.
25. Cambridge Cognition. *Cambridge neuropsychological test automated battery*. Cambridge: Cambridge Cognition; 1999.
26. Wechsler D. *Wechsler intelligence scale for children-fourth edition (WISC-IV)*. San Antonio, TX: The Psychological Corporation; 2003.
27. Wechsler D. *Wechsler adult intelligence scale-third edition (WAIS-III)*. San Antonio, TX: The Psychological Corporation; 1997.
28. Wechsler D, Naglieri JA. *Wechsler nonverbal scale of ability (WNV)*. Pearson; 2006.
29. Kongs SK, Thompson LL, Iverson GL, Heaton RK. *Wisconsin card sortin test-64 card version*. Lutz, FL: Psychological Assessment Resources; 2000.
30. Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH. Content validity of the expanded and revised gross motor function classification system. *Dev Med Child Neurol* 2008;**50**(10):744–50. <https://doi.org/10.1111/j.1469-8749.2008.03089.x>.
31. Eliasson A-C, Krumlinde-Sundholm L, Rösblad B, Beckung E, Arner M, Ohrvall A-M, et al. The Manual Ability Classification System (MACS) for children with cerebral palsy: scale development and evidence of validity and reliability. *Dev Med Child Neurol* 2006;**48**(7):549–54. <https://doi.org/10.1017/S0012162206001162>.
32. Hidecker MJC, Paneth N, Rosenbaum PL, Kent RD, Lillie J, Eulenberg JB, et al. Developing and validating the communication function classification system for individuals with cerebral palsy. *Dev Med Child Neurol* 2011;**53**(8):704–10. <https://doi.org/10.1111/j.1469-8749.2011.03996.x>.
33. Mei C, Reilly S, Reddihough D, Mensah F, Pennington L, Morgan A. Language outcomes of children with cerebral palsy aged 5 years and 6 years: a population-based study. *Dev Med Child Neurol* 2016;**58**(6):605–11. <https://doi.org/10.1111/dmcn.12957>.
34. Geytenbeek JMM, Heim MMJ, Vermeulen RJ, Oostrom KJ. Assessing comprehension of spoken language in nonspeaking children with cerebral palsy: application of a newly developed computer-based instrument. *Augmentative Altern Commun* 2010;**26**(2):97–107. <https://doi.org/10.3109/07434618.2010.482445>.
35. Monbaliu E, de Cock P, Ortibus E, Heyrman L, Klingels K, Feys H. Clinical patterns of dystonia and choreoathetosis in participants with dyskinetic cerebral palsy. *Dev Med Child Neurol* 2015. <https://doi.org/10.1111/dmcn.12846>. n/a-n/a.
36. Strauss E, Spreen O, Sherman EMS. *A Compendium of neuropsychological tests: administration, norms, and commentary, 3*. Oxford: Oxford University Press; 2006.
37. Raven JC, Court JH, Seisdedos Cubero N. *Raven matrices progresivas escalas: CPM color, SPM general, APM superior*. Madrid: TEA; 2001.
38. Ballester-Plané J, Laporta-Hoyos O, Macaya A, Póo P, Meléndez-plumed M, Vázquez É, et al. Measuring intellectual ability in cerebral palsy: the comparison of three tests and their neuroimaging correlates. *Res Dev Disabil* 2016;**56**:83–98. <https://doi.org/10.1016/j.ridd.2016.04.009>.
39. Fazzi E, Signorini SG, La R, Chiara P, Walter B, Galli J, et al. Neuro-ophthalmological disorders in cerebral palsy: ophthalmological, oculomotor, and visual aspects. *Dev Med Child Neurol* 2012. <https://doi.org/10.1111/j.1469-8749.2012.04324.x>.
40. Himmelmann K, Hagberg G, Uvebrant P. The changing panorama of cerebral palsy in Sweden. X. Prevalence and origin in the birth-year period 1999 – 2002. *Acta Paediatr* 2010;**1**:1337–43. <https://doi.org/10.1111/j.1651-2227.2010.01819.x>.