



Endocrine

Presented at the Academic Surgical Congress 2019

Extent of surgery for low-risk thyroid cancer in the elderly: Equipoise in survival but not in short-term outcomes



Alan Zambeli-Ljepović, BS^a, Frances Wang, MS^a, Michaela A. Dinan, PhD^a,
Terry Hyslop, PhD^a, Michael T. Stang, MD^a, Sanziana A. Roman, MD^b,
Julie A. Sosa, MD, MA^b, Randall P. Scheri, MD^{a,*}

^a Department of Surgery, Duke University Medical Center, Durham, NC^b Department of Surgery, University of California at San Francisco, San Francisco, CA

ARTICLE INFO

Article history:

Accepted 31 May 2019

Available online 6 July 2019

ABSTRACT

Background: Total thyroidectomy is more common than lobectomy for low-risk papillary thyroid cancer, despite equipoise in survival. Because postoperative morbidity increases with age, we aimed to investigate how the extent of thyroidectomy affects short-term outcomes among older patients.

Methods: Using the Surveillance, Epidemiology, and End Results–Medicare database, we identified patients aged ≥ 66 years who were treated between 1996 and 2011 for papillary thyroid cancer with tumors ≤ 2 cm in diameter. We used multivariable logistic regression to evaluate the effect of extent of surgery on complications, emergency-department visits, and unplanned readmissions.

Results: Among 3,341 selected patients, 77.3% were female, mean age was 72.9 years, and tumors averaged 0.8 cm in diameter. A total of 67.6% of patients underwent total thyroidectomy, and 32.4% underwent lobectomy. Total thyroidectomy was associated with complications (odds ratio = 1.99) and readmissions (odds ratio = 1.59; both $P < 0.01$). Complications were higher in female patients (odds ratio = 1.34), black patients (versus white patients, odds ratio = 1.65), and those with ≥ 2 comorbidities (vs 0, odds ratio = 1.43; all $P < 0.01$). Black patients and those with ≥ 2 comorbidities had more emergency-department visits (odds ratio = 1.50 and 1.92, respectively) and readmissions (odds ratio = 2.19 and 2.29, respectively; all $P < 0.01$).

Conclusion: Total thyroidectomy for older adults with low-risk papillary thyroid cancer may lead to potentially avoidable complications and readmissions, particularly for black and female patients. In many cases, lobectomy may be a safer and less costly alternative.

© 2019 Elsevier Inc. All rights reserved.

Introduction

Thyroid cancer has the fastest-increasing incidence of any cancer worldwide.¹ Over the past 30 years, the incidence of thyroid cancer in the United States has tripled, affecting 64,300 people in 2016.² Eighty-five percent of new cases are diagnosed as papillary thyroid cancer (PTC), for which the 10-year survival rate exceeds 95%. The American Thyroid Association categorizes PTC as low risk when tumors are ≤ 4 cm in diameter, do not feature aggressive histology (eg, tall-cell, insular, or diffuse sclerosing variants), and are not associated with extrathyroidal invasion or metastasis.³ Before 2015, the

American Thyroid Association universally recommended total thyroidectomy, except in cases of subcentimeter tumors (papillary thyroid microcarcinoma), for which thyroid lobectomy was the recommended alternative.^{4–6} The current (as of 2015) guidelines consider either procedure to be appropriate for all low-risk tumors.³

Despite this equipoise in operative recommendations for low-risk PTC, total thyroidectomy is still performed more commonly, even for subcentimeter tumors.⁷ Extent of surgery does not impact survival,⁸ but total thyroidectomy is associated with having nearly double the complication rate of thyroid lobectomy.⁹ Older patients are particularly vulnerable: those aged >70 years have higher complication rates, longer lengths of stay, and greater hospital charges after any type of thyroid surgery.¹⁰ As the Centers for Medicare and Medicaid Services (CMS) expands penalties for postsurgical complications and readmissions, extent of thyroidectomy may have financial implications as well.¹¹

Presented at the 14th Annual Academic Surgical Congress, Houston, TX.

* Reprint requests: Randall P. Scheri, MD, Department of Surgery, Duke University Medical Center, 463 Seeley G. Mudd Bldg., Box 3513, Durham, NC 27710.

E-mail address: r.scheri@duke.edu (R.P. Scheri).<https://doi.org/10.1016/j.surg.2019.05.035>

0039-6060/© 2019 Elsevier Inc. All rights reserved.

Our aim was to use the Surveillance, Epidemiology, and End Results (SEER)-Medicare database to investigate the association between extent of surgery and complications, emergency-department (ED) visits, and readmissions among patients aged ≥ 66 years with low-risk PTC with tumors ≤ 2 cm in diameter. This size threshold selects a majority of tumors for which current guidelines permit either total thyroidectomy or lobectomy (≤ 4 cm), while excluding larger tumors (2–4 cm) that many clinicians would not be comfortable treating with lobectomy. We hypothesized that total thyroidectomy is independently associated with complications, ED visits, and readmissions.

Methods

Data source

Data were collected from the SEER-Medicare linked database. SEER captures population-level demographic, clinical, and survival data for approximately 28% of all cancer patients in the United States.¹² Medicare insures 97% of those aged ≥ 65 years in the United States and collects data on health care services provided throughout a person's Medicare eligibility. Medicare Part D files, available to us for the years 2008 to 2012 ($n = 1,224$, 36.6% of the entire cohort), provide information on filled prescriptions. The biennially linked database has been used to study factors associated with cancer care, including the following: sociodemographic characteristics, physician and hospital characteristics, diagnostics, surgery, chemotherapy, radiation, comorbid conditions, complications, screening, relapse, and costs.

Cohort selection

The 17 SEER registries with uninterrupted data from 1997 to 2011 were used to select all patients aged ≥ 66 years with a primary diagnosis of PTC (International Classification of Diseases-O-3 codes 8050, 8060, 8140, 8260, 8340, 8341, 8342, 8343, and 8344), including classic and follicular variants and excluding aggressive variants (eg, tall-cell, insular, or diffuse sclerosing), who underwent thyroid lobectomy (Current Procedural Terminology codes 60210, 60212, and 60220) or total thyroidectomy (Current Procedural Terminology codes 60240 and 60271) ≤ 180 days after diagnosis. To select the subset of low-risk PTC (T1N0M0) that most clinicians would feel could be treated with lobectomy, we excluded patients with the following: tumors > 2 cm in diameter, cancer in situ, extrathyroidal extension, nodal or distant metastases, or a PTC diagnosis at autopsy.

We restricted the minimum patient age to 66 years to ensure continuous Medicare Part A and Part B coverage for at least 1 year before diagnosis; we further required coverage for 2 years after diagnosis (or until death). We excluded patients with a missing month of diagnosis, a second malignant diagnosis ≤ 1 year within the diagnosis of PTC, or procedure codes suggesting multiple thyroid surgeries (eg, code for completion thyroidectomy, multiple codes for lobectomy). This increased the accuracy of dates and reduced the potential confounding effects of coexisting diagnoses and multiple surgical procedures.

To maximize the completeness of Medicare claims data, we only included patients whose PTC diagnosis in SEER could be matched with a synchronous, primary diagnosis of PTC on an inpatient, outpatient, or carrier-based Medicare claim.

Independent variable and outcomes

The independent variable, extent of surgery, was dichotomized into total thyroidectomy or lobectomy, based on the thyroid

surgical procedure coded within 180 days after diagnosis. Medicare modifier codes were used to verify that the claims of interest were associated with thyroid surgery.

The primary outcomes were postoperative complications and unplanned encounters. Complications were identified using International Classification of Diseases-9 diagnosis codes from postoperative encounters. General surgical complications (eg, pneumonia, cystitis, wound infection) and endocrine-specific complications (eg, hypoparathyroidism, hoarseness [a sign of recurrent laryngeal nerve injury]) were recorded during the first 30 postoperative days and are henceforth termed ≤ 30 -day complications. To estimate rates of permanent injury to neck structures, endocrine complications also were examined between 6 and 18 months after surgery and termed ≥ 6 -month complications.¹³ Rates of filled prescriptions for calcitriol were used as the most specific measure of postoperative hypoparathyroidism.

Unplanned encounters comprised presentations to the ED not resulting in admission, henceforth termed ED visits, and postoperative hospital admissions for any cause except radioactive iodine (RAI) administration, henceforth termed readmissions. Exclusion of patients with more than one coded surgery ensured that readmissions did not include hospitalization for completion thyroidectomy. Given recent data demonstrating that up to 48% of readmissions occur after the traditional postoperative window of 30 days, these encounters were examined over 90 postoperative days.^{14–16}

Other variables

Primary outcomes were correlated with clinical variables, including comorbidity, tumor size, histology, and (multi)focality. Comorbidity was assessed using the 2014 National Cancer Institute Comorbidity Index, which is based on the Charlson Comorbidity Index.¹⁷ Demographic variables included patient age, sex, race, and local census tract characteristics, such as population density and income quartile, which were obtained from SEER data. Vital status and follow-up times were acquired from the SEER Patient Entitlement and Diagnosis Summary File.

Statistical analysis

Baseline patient characteristics are presented as frequencies and proportions for categorical variables and means with standard deviations (SDs) or medians with interquartile ranges for continuous variables. Comparisons between patients undergoing total thyroidectomy versus lobectomy were made using χ^2 tests for categorical variables and Student's *t* tests for continuous variables.

Multivariable logistic regression was used to estimate odds ratios (ORs) and 95% confidence intervals associated with extent of surgery. The model for postsurgical complications and encounters was adjusted for patient age, sex, race, comorbidity index, RAI administration, tumor size and focality, and other clinicopathologic disease characteristics.

All *P* values are 2 sided and set at a significance level of 0.05. All statistical analyses were performed using SAS software (version 9.4; SAS Institute, Cary, NC). This study was approved by the Duke University Health System Institutional Review Board.

Results

Demographic, clinical, and surgical characteristics

A total of 3,341 patients with PTC with tumors ≤ 2 cm met eligibility criteria (Fig 1). The mean age at diagnosis was 72.9




Fig 1. Patient selection diagram. *Defined as a primary diagnosis of PTC on an inpatient, outpatient, or carrier-based Medicare claim within 3 months of the SEER-reported diagnosis. †SEER records from 1997 to 2011 were searched for the incident diagnosis of PTC; Medicare claims spanning 1996 to 2013 were used to verify diagnosis-related procedures and to determine insurance status.

Table 1
Characteristics of patients aged ≥66 years who underwent surgery between 1996 and 2011 for PTC with tumors ≤2 cm

	Entire cohort	Extent of surgery		P value
		Lobectomy	Total thyroidectomy	
N	3,341	1,081	2,260	
Mean age (y) (SD)	72.9 (5.5)	73.2 (5.6)	72.81 (5.5)	.05
Female sex	2,583 (77.3%)	828 (76.6%)	1,755 (77.7%)	.49
Race				.20
White	2,901 (86.8%)	930 (86.0%)	1,971 (87.2%)	
Black	188 (5.6%)	72 (6.7%)	116 (5.1%)	
Other	252 (7.5%)	79 (7.3%)	173 (7.7%)	
Charlson Comorbidity Index				.82
1	850 (25.4%)	270 (25.0%)	580 (25.7%)	
≥2	600 (18.0%)	200 (18.5%)	400 (17.7%)	
Tumor size				<.01
Mean (SD) (cm)	0.8 (0.6)	0.6 (0.5)	0.8 (0.6)	
≤1 (cm)	2,396 (71.7%)	906 (83.8%)	1,490 (65.9%)	
1.1–2 (cm)	945 (28.3%)	175 (16.2%)	770 (34.1%)	
Histology				.12
Classic papillary	2,170 (65.0%)	682 (63.1%)	1,488 (65.8%)	
Follicular variant	1,171 (35.0%)	399 (36.9%)	772 (34.2%)	
Multifocal disease, 2004–2011*	805 (31.7%)	138 (18.5%)	667 (37.2%)	<.01

Percentages are based on column totals and may not add to 100% owing to rounding. P values were computed by t test and χ^2 tests for continuous and categorical variables, respectively.

* Focality was not recorded before 2004. Percentages are based on 2,681 patients (75.9%) who had complete focality data.

Table IIFrequency of postoperative complications and encounters in patients aged ≥ 66 years with PTC with tumors ≤ 2 cm between 1996 and 2011

	Entire cohort	Extent of surgery		P value
		Lobectomy	Total thyroidectomy	
N	3,341	1,081	2,260	
Surgical setting (LOS in days)				<.01
Outpatient	1,032 (30.9%)	408 (37.7%)	624 (27.6%)	
Inpatient (1)	1,371 (41.0%)	425 (39.3%)	946 (41.9%)	
Inpatient (≥ 2)	938 (28.1%)	248 (22.9%)	690 (30.5%)	
≤ 30 -day complications	682 (20.4%)	146 (13.5%)	536 (23.7%)	<.01
General complications	224 (6.7%)	64 (5.9%)	160 (7.1%)	.21
Hoarseness	88 (2.6%)	19 (1.8%)	69 (3.1%)	.03
Vocal cord paralysis	36 (1.1%)	13 (1.2%)	23 (1.0%)	.63
Hypoparathyroidism	127 (3.8%)	22 (2.0%)	105 (4.6%)	<.01
Taking calcitriol*	107 (8.7%)	<11 (<3.4%)†	>96 (>10.7%)†	<.01
≥ 6 -month endocrine complications	386 (11.6%)	117 (10.8%)	269 (11.9%)	.36
Hoarseness	171 (5.1%)	46 (4.3%)	125 (5.5%)	.12
Vocal cord paralysis	43 (1.3%)	<11 (<1.0%)†	>32 (>1.4%)†	.11
Hypoparathyroidism	86 (2.6%)	26 (2.4%)	60 (2.7%)	.67
Taking calcitriol*	51 (4.2%)	14 (4.3%)	37 (4.1%)	.86
90-day postoperative encounters				
ED visits	379 (11.3%)	111 (10.3%)	268 (11.9%)	.18
Readmissions	494 (14.8%)	105 (9.7%)	389 (17.2%)	<.01

Percentages are based on column totals and may not add to 100% due to rounding. An outpatient surgical setting is defined as discharge on the same calendar day as the surgery; overnight observation stays are categorized as inpatient (LOS = 1 day). General complications include cardiovascular, pulmonary, renal, infectious, and wound-related etiologies. P values were computed by Wald test.

LOS, length of stay; ED, emergency department.

* Analysis limited to patients with Medicare part D (N = 1,224, 36.6% of the entire cohort).

† True values suppressed owing to small cell size (N < 11), per data use agreement.

years (SD 5.5 years; Table I). Most patients were white (86.8%) and female (77.3%). Mean tumor size was 0.8 cm (SD 0.6 cm). Histologically, 65.0% of tumors were classic PTC; the remainder represented the follicular variant. From 2004 to 2011, the years for which focality was recorded, 805 (31.7%) tumors were multifocal.

Most patients (67.6%) underwent total thyroidectomy, and 32.4% underwent lobectomy. Lobectomy was more likely than total thyroidectomy to be performed in the outpatient setting (37.7% vs 27.6%, $P < 0.01$; Table II). Longer length of stay (≥ 2 days) was more likely with total thyroidectomy than lobectomy (42.2% vs 36.9%, $P < 0.01$). Survival at 1 year was 98.0%.

Short-term (≤ 30 days) complications

A total of 682 patients (20.4%) experienced complications in the first 30 postoperative days. Complications were more common after total thyroidectomy compared with lobectomy (23.7% vs 13.5%, $P < 0.01$). Specifically, total thyroidectomy was followed by higher rates of hoarseness (3.1% vs 1.8%, $P = 0.03$), hypocalcemia (12.7% vs 4.9%, $P < 0.01$), hypoparathyroidism (4.6% vs 2.0%, $P < 0.01$), and proportion of patients filling calcitriol prescriptions (>10.7% vs <3.4%, $P < 0.01$). There was no significant difference in rates of general complications (eg, pneumonia, wound infection) between total thyroidectomy and lobectomy (7.1% vs 5.9%, $P = 0.21$).

On multivariable analysis (Fig 2), ≤ 30 -day complications were independently associated with total thyroidectomy (OR = 1.99), a comorbidity index ≥ 2 (OR = 1.43), female sex (OR = 1.34), and black compared with white race (OR = 1.65; all $P < 0.01$). Sex-based differences may be explained, in part, by differential rates of hypocalcemia (female versus male patients: 11.2% vs 6.9%, $P < 0.01$). Potentially owing to small sample sizes, race-based differences could not be attributed to any one particular complication.




Fig 2. Multivariable regression analysis of postoperative complications and readmissions for patients aged ≥ 66 years with T1N0M0 PTC between 1996 and 2011 (N = 3,307). The ≤ 30 -day period includes both general complications (eg, pneumonia, cystitis, wound infection) and endocrine complications (eg, hypoparathyroidism, hoarseness, vocal cord paralysis). The ≥ 6 -month period only includes endocrine complications. Aside from variables shown, analysis adjusts for the following: age, RAI administration, tumor size, (multi)focality, income, residence, year of diagnosis, and National Cancer Institute designation of treatment center. These variables did not have clinically significant effects on outcomes. *Sample size is reduced in multivariable analysis due to exclusion of 34 patients (1%) with missing values.

Long-term (≥ 6 months) complications

A total of 386 patients (11.6%) had endocrine complications recorded ≥ 6 months after surgery (Table II). Complication rate was not significantly different between total thyroidectomy (11.9%) and lobectomy (10.8%, $P = 0.36$), although total thyroidectomy more often resulted in ≥ 2 complications (1.5% vs <0.9%, respectively, $P < 0.01$). Overall, 5.1% of patients had hoarseness, 4.5% had

hypocalcemia, 2.6% had hypoparathyroidism, and notably, 4.2% of patients were still filling prescriptions for calcitriol at ≥ 6 months after surgery. After multivariable adjustment (Fig 2), only female sex remained associated with ≥ 6 -month complications (OR = 1.37, $P = 0.03$).

Postoperative encounters

During the first 90 postoperative days, 379 (11.3%) patients presented to the ED but did not require admission (Table II). The median number of days between surgery and ED visit was 24. The most common reasons for these ED visits were general (non-endocrine) surgical complications (32.2%), followed by cardiac (29.3%) and endocrine (26.9%) complications. ED visit rates did not vary significantly by extent of surgery (total thyroidectomy versus lobectomy: 11.9% vs 10.3%, $P = 0.18$).

After excluding admissions for RAI administration and completion thyroidectomy, 494 (14.8%) patients were readmitted ≤ 90 days after surgery. The median time to readmission was 41 days. Most readmissions (57.7%) were for endocrine complications; readmissions were more common after total thyroidectomy compared with lobectomy (17.2% vs 9.7%, respectively, $P < 0.01$). After multivariable adjustment (Fig 2), readmissions remained associated with total thyroidectomy (OR = 1.59), comorbidity index ≥ 2 (vs 0, OR = 2.29), and black race (versus white, OR = 2.19; all $P < 0.01$).

Discussion

This nationally representative study is the first to show that rates of both complications and unanticipated readmissions are higher after total thyroidectomy compared with lobectomy for elderly patients with low-risk PTC. Female sex and black race are independently associated with complications and readmissions. In the absence of evidence that extent of surgery affects survival in low-risk PTC, our findings suggest that lobectomy may often be the safer surgical choice.

Higher ≤ 30 -day endocrine complication rates after total thyroidectomy underscore the increased risk of hypoparathyroidism and recurrent laryngeal nerve injury associated with a bilateral procedure. Our findings are consistent with a 2014 nationwide analysis of outcomes after 62,722 thyroid surgeries.⁹ In that study, Hauch et al demonstrated that complication rates nearly doubled after total thyroidectomy compared with lobectomy (20.4% vs 10.8%), even in the hands of high-volume surgeons. We corroborate that these differences are largely driven by higher rates of hypocalcemia following total thyroidectomy. Although the multivariable model used by Hauch et al adjusted for diagnosis (Graves' disease versus benign versus malignant), it did not distinguish among cancer types or risk strata, likely increasing the proportion of complicated cases (eg, larger or locally invasive tumors) in the total thyroidectomy group. Our data suggest that even for small, intrathyroidal tumors, total thyroidectomy compared with lobectomy is associated with doubled odds of complications and should be used judiciously.

Unlike ≤ 30 -day complications, ≥ 6 -month complication rates do not appear to differ by extent of surgery. This stands in contrast to prior published work. A meta-analysis examining longer-term complications of thyroid surgery found that, relative to lobectomy, total thyroidectomy resulted in 1.85 \times and 3.17 \times higher rates of permanent recurrent laryngeal nerve and parathyroid injuries, respectively.¹⁸ As in the study by Hauch et al, the total thyroidectomy group in this meta-analysis included larger and more aggressive tumors, which would increase posttotal thyroidectomy complication rates. In our cohort, fewer complicated total

thyroidectomy cases likely resulted in fewer events at ≥ 6 months and therefore less power to detect a difference in rates of permanent injury between total thyroidectomy and lobectomy.

Differences in complication rates may also have been biased toward the null by the presence of hypoparathyroidism among some patients in the lobectomy group. Most likely, these patients had previously (ie, at a time not captured in this study) undergone lobectomy. The procedure coded as "lobectomy" during our observation period would, in truth, have been a completion thyroidectomy, a more plausible cause of hypoparathyroidism. Alternatively, others have suggested that, even with lobectomy, exploration of the contralateral lobe, ischemia of ≥ 2 parathyroid glands, or postoperative hematoma formation may be sufficient to cause hypoparathyroidism.^{19,20}

Readmission rates, another measure of postoperative morbidity, were also significantly higher for patients undergoing total thyroidectomy compared with those undergoing lobectomy in this cohort. Previously published readmissions data vary, as do methodologies. Collier et al also compared 90-day readmission rates by extent of surgery.²¹ Their study of 59,427 patients found an overall readmission rate (17.6%) similar to the one we observed (14.8%) but a 32% lower adjusted risk of readmission after total thyroidectomy compared with lobectomy. Importantly, Collier et al did not analyze unplanned readmissions separately: 98.9% of readmissions after lobectomy for thyroid cancer were for completion thyroidectomy or administration of RAI, which were planned treatments that exaggerated the postlobectomy readmission rate. In contrast, and more in line with our findings, Iannuzzi et al examined 30-day unplanned readmissions after 34,046 thyroid surgeries, demonstrating a significantly lower readmission rate after lobectomy (2.1%) compared with after total thyroidectomy (2.8%).²² Although their cohort was not stratified by diagnosis, more complicated thyroidectomy procedures (total thyroidectomy with lymph node dissection, substernal thyroidectomy) were treated separately, increasing the likelihood that their lobectomy and total thyroidectomy groups were similar to ours. Lower average age (54 years) and a shorter readmission windows likely contributed to lower overall readmission rates relative to our cohort.

Increased readmission rates after total thyroidectomy compared with lobectomy have three major implications. First, most readmissions were prompted by endocrine complications, emphasizing that even the increased risk of short-term complications with total thyroidectomy warrants careful consideration. In fact, unplanned readmissions may portend poor survival outcomes. In a cohort that included all stages of thyroid cancer, Tuggle et al found that unplanned hospitalizations were significantly associated with death at 1 year, compared with nonhospitalization.²³ A low mortality rate in our cohort precluded a similar analysis. Second, half of all readmissions occurred after postoperative day 41; this supports recent findings that events during the first 30 postoperative days do not fully capture surgical outcomes.^{14–16} Instead, a 60- or 90-day postoperative period may better define a complete "episode of care" if CMS bundled payments expand to include thyroid surgery. Third, as the CMS Hospital Readmission Prevention Program increases penalties for readmissions after a broader range of surgeries, avoiding total thyroidectomy when lobectomy is sufficient may have a cost benefit as well.

Two demographic factors are independently associated with both complications and readmissions: female sex and black race. Sex-based differences are chiefly a result of higher rates of postoperative hypocalcemia in women. This may be attributed to postmenopausal women having a smaller calcium "reserve" relative to male counterparts, increasing the likelihood that parathyroid ischemia causes symptomatic hypocalcemia, which could, in turn, prompt readmission. As Wang and Yen have summarized,

up to one-third of readmissions after thyroid surgery are driven by symptomatic hypocalcemia, which can be addressed with postoperative counseling and implementation of algorithms for inpatient and outpatient management.²⁴ Race-based differences are more concerning because a biological explanation likely does not exist. Sosa et al have previously shown that, on a national level, >50% of black and Hispanic patients have thyroid surgery by the lowest-volume surgeons (1–9 cases per year), compared with 44% of white patients.²⁵ It is likely that in our study as well, race-based differences in complications and resulting readmissions were driven by inequity in access to high-quality care. Dedicated health disparities research would better address race-based differences in thyroidectomy outcomes.

The main limitations of this study stem from the nature of administrative databases, such as SEER-Medicare, which do not indicate the intent of clinical or coding decisions. For example, the multifactorial decision about extent of surgery depends on the possibility of concomitant benign thyroid disease, patient preference, size of the thyroid gland, radiation history, and preexisting dependence on thyroid supplementation, factors that SEER-Medicare does not fully capture. Furthermore, diagnosis codes may be imprecise measures of complications such as surgical hypoparathyroidism. Rates of hypocalcemia likely overestimate this value (owing to low specificity), whereas rates of hypoparathyroidism or filled calcitriol prescriptions are likely underestimates (owing to sparse documentation and limited sensitivity, respectively). Unlike other databases, however, SEER-Medicare allows us to examine all three measures. Therefore, while a precise rate of surgical hypoparathyroidism may be inaccessible, the observed consistency in trends across all three measures of this complication lends internal validity to the conclusion that surgical hypoparathyroidism is more common after total thyroidectomy.

To our knowledge, this is the first nationally representative study of the effect of extent of surgery on postoperative complications, ED visits, and unplanned readmissions in older adults (≥ 66 years) with low-risk PTC with tumors ≤ 2 cm (T1N0M0). Complications, often severe enough to warrant readmission, are independently associated with total thyroidectomy, female sex, and black race. Among older adults, more judicious use of total thyroidectomy and an increased vigilance for postoperative hypocalcemia may mitigate the morbidity of a disease with an otherwise excellent prognosis.

Funding/Support

Supported by grant TL1TR001116 from the National Center for Advancing Translational Sciences of the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest/Disclosure

Julie A. Sosa, MD, MA, is a member of the Data Monitoring Committee of the Medullary Thyroid Cancer Consortium Registry, which is supported through United BioSource Corporation by NovoNordisk, GlaxoSmithKline, Astra Zeneca, and Eli Lilly. Terry Hyslop, PhD, receives consulting funds from AbbVie to work on statistical methods related to the demonstration of interchangeability and biosimilarity for biologic products. This study used the linked SEER-Medicare database, and the interpretation and reporting of these data are the sole responsibility of the authors.

Acknowledgments

The authors acknowledge the efforts of the National Cancer Institute; the Office of Research, Development and Information, CMS; Information Management Services, Inc; and the SEER program tumor registries in the creation of the SEER-Medicare database.

References

- Lubitz CC, Sosa JA. The changing landscape of papillary thyroid cancer: Epidemiology, management, and the implications for patients. *Cancer*. 2016;122:3754–3759.
- National Cancer Institute. SEER cancer stat facts: Thyroid cancer; 2017. <http://seer.cancer.gov/statfacts/html/thyro.html>. Accessed February 11, 2017.
- Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. 2016;26:1–133.
- Cooper DS, Doherty GM, Haugen BR, et al. Management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2006;16:109–142.
- American Thyroid Association (ATA) Guidelines Taskforce on Thyroid Nodules and Differentiated Thyroid Cancer, Cooper DS, Doherty GM, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2009;19:1167–1214.
- Singer PA, Cooper DS, Daniels GH, et al. Treatment guidelines for patients with thyroid nodules and well-differentiated thyroid cancer. American Thyroid Association. *Arch Intern Med*. 1996;156:2165–2172.
- Wang TS, Goffredo P, Sosa JA, Roman SA. Papillary thyroid microcarcinoma: An over-treated malignancy? *World J Surg*. 2014;38:2297–2303.
- Adam MA, Pura J, Gu L, et al. Extent of surgery for papillary thyroid cancer is not associated with survival: An analysis of 61,775 patients. *Ann Surg*. 2014;260:601–605; discussion 605–607.
- Hauch A, Al-Qurayshi Z, Randolph G, Kandil E. Total thyroidectomy is associated with increased risk of complications for low- and high-volume surgeons. *Ann Surg Oncol*. 2014;21:3844–3852.
- Sullivan MC, Roman SA, Sosa JA. Clinical and economic outcomes of thyroid surgery in elderly patients: A systematic review. *J Thyroid Res*. 2012;2012:615846.
- Centers for Medicare & Medicaid Services. Hospital-Acquired Condition (HAC) Reduction Program; 2018. <https://go.cms.gov/2j5Hmor>. Accessed November 11, 2018.
- National Cancer Institute. SEER: Surveillance, Epidemiology, and End Results Program; 2012. <https://seer.cancer.gov/>. Accessed November 11, 2018.
- Thomusch O, Machens A, Sekulla C, Ukkat J, Brauckhoff M, Dralle H. The impact of surgical technique on postoperative hypoparathyroidism in bilateral thyroid surgery: A multivariate analysis of 5846 consecutive patients. *Surgery*. 2003;133:180–185.
- Siracuse JJ, Shah NK, Peacock MR, et al. Thirty-day and 90-day hospital readmission after outpatient upper extremity hemodialysis access creation. *J Vasc Surg*. 2017;65:1376–1382.
- Kim Y, Gani F, Lucas DJ, et al. Early versus late readmission after surgery among patients with employer-provided health insurance. *Ann Surg*. 2015;262:502–511; discussion 509–511.
- Wick EC, Shore AD, Hirose K, et al. Readmission rates and cost following colorectal surgery. *Dis Colon Rectum*. 2011;54:1475–1479.
- Stedman MR, Doria-Rose P, Warren JL, Klabunde CN, Mariotto AB. *Comorbidity technical report*. National Cancer Institute; 2018. <https://comorbidity-report.html>. Accessed November 11, 2018.
- Kandil E, Krishnan B, Noureldine SI, Yao L, Tufano RP. Hemithyroidectomy: A meta-analysis of postoperative need for hormone replacement and complications. *ORL J Otorhinolaryngol Relat Spec*. 2013;75:6–17.
- Vaiman M, Nagibin A, Olevson J. Complications in primary and completed thyroidectomy. *Surg Today*. 2010;40:114–118.
- Asari R, Passler C, Kaczirek K, Scheuba C, Niederle B. Hypoparathyroidism after total thyroidectomy: A prospective study. *Arch Surg*. 2008;143:132–137; discussion 138.
- Collier K, Sataloff J, Wirtalla C, Kuo L, Karakousis GC, Kelz RR. Understanding readmissions following operations of the thyroid and parathyroid glands. *Am J Surg*. 2017;214:501–508.
- Iannuzzi JC, Fleming FJ, Kelly KN, Ruan DT, Monson JR, Moalem J. Risk scoring can predict readmission after endocrine surgery. *Surgery*. 2014;156:1432–1438; discussion 1438–1440.
- Tuggle CT, Park LS, Roman S, Udelsman R, Sosa JA. Rehospitalization among elderly patients with thyroid cancer after thyroidectomy are prevalent and costly. *Ann Surg Oncol*. 2010;17:2816–2823.
- Wang TS, Yen TW. Readmission after thyroidectomy and parathyroidectomy: What can we learn from NSQIP? *Surgery*. 2014;156:1419–1422.
- Sosa JA, Mehta PJ, Wang TS, Yeo HL, Roman SA. Racial disparities in clinical and economic outcomes from thyroidectomy. *Ann Surg*. 2007;246:1083–1091.