



Feasibility of incorporating high-intensity interval training into physical education programs to improve body composition and cardiorespiratory capacity of overweight and obese children: A systematic review

Pedro Delgado-Floody ^{a,*}, Pedro Latorre-Román ^b, Daniel Jerez-Mayorga ^c, Felipe Caamaño-Navarrete ^d, Felipe García-Pinillos ^a

^a Department of Physical Education, Sport and Recreation, Universidad de La Frontera, Temuco, Chile

^b University of Jaen, Jaen, Spain

^c Faculty of Rehabilitation Sciences, Universidad Andres Bello, Santiago, Chile

^d Faculty of Education, Universidad Católica de Temuco, Chile

ARTICLE INFO

Article history:

Received 17 July 2018

Received in revised form

13 November 2018

Accepted 15 November 2018

Available online 15 November 2018

Keywords:

Children

High-intensity interval training

Physical fitness

ABSTRACT

Background/objective: High-intensity interval training (HIIT) can produce similar or improved results compare with traditional training, but the question as to whether HIIT can be used in the setting of physical education (PE) remains unanswered. The aim of this systematic review was to critically analyze the feasibility of incorporating HIIT programs into PE classes to improve the body compositions and cardiorespiratory fitness of overweight students.

Methods: We conducted database searches for literature dating between January 2012 and January 2017. Of the final six studies selected, three were conducted in children under 12 years old and three involved adolescents between 12 and 18 years old.

Results: The HIIT protocols consisted of 2–3 sessions per week, with intervals of 15 s and passive or active rests of 15 s, totaling up to 6 min of work with 4 min of rest. The duration of HIIT programs was 6–24 weeks. Significant changes were reported in body composition, body mass index, body fat (%), waist circumference, and sum of skinfolds; and increases in muscle mass were observed. The inclusion of HIIT programmes improved maximal oxygen uptake (VO_{2max}), performance in the intermittent Yo-Yo test and maximal aerobic speed.

Conclusions: The HIIT programmes showed improvements in the variables studied, with interventions two or three times weekly. Therefore, they can be used in schools, as a strategy to combat the childhood obesity pandemic and HIIT can be used alongside with existing PE activities within the same lesson or in specific periods during day school.

© 2018 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Childhood obesity is a complex condition that can drive many risk factors.¹ It is characterized, among other factors, by an imbalance between food input and energy expenditure which results in an increase in body mass index (BMI).² This pathology is a

worldwide health problem, increasingly prevalent in child and adolescent populations, especially over the last few decades.^{3,4} This increase has been associated with a concomitant increase in various chronic illnesses such as metabolic syndrome, type 2 diabetes, cardiovascular disease, cancer, and arthritis as well as a higher probability of obesity and low cardiorespiratory capacity in adulthood.^{5,6} Cardiorespiratory capacity is a measurement of how the body functions and develops, and should, through training, play an important role in activities related to improving physical activity. This is a fundamental component of a healthy lifestyle for children and adolescents.⁵ Furthermore, high cardiorespiratory capacity is

* Corresponding author. Universidad de La Frontera, Uruguay Street No. 1980, Temuco, Chile.

E-mail address: pedro.delgado@ufroterra.cl (P. Delgado-Floody).

associated with lower cardiometabolic risk.⁶

Effective strategies for the prevention and treatment of childhood obesity require the involvement of the school environment. Modification of physical education (PE) curricula in schools have resulted in consistent changes in the quantity of physical activity and student motivation, with positive effects on quality of life and other biomarkers.^{7,9} Adequate exposure to PE effectively contributes to the development of a healthier lifestyle among children, which continues throughout their lifespan. However, many PE programs do not comply with recommendations in terms of class times and intensity (moderate-to-vigorous physical activity) due to time lost to management-related activities.^{8,9}

To combat childhood obesity, a healthy diet and/or aerobic training are essential.¹⁰ Although continuous aerobic training is generally included in intervention programs, low volume high-intensity interval training (HIIT) can produce similar or improved responses compared with traditional continuous aerobic training, despite the short amount of time that HIIT requires.¹¹

Alvarez et al. observed that following a HIIT program for a total of 18 sessions (three times per week) improved the insulin resistance levels (assessed by homeostatic model assessment of insulin resistance [HOMA-IR]) as well as the anthropometric, cardiovascular, and physical performance of insulin-resistant school-children.¹² A systematic review on children and adolescents reported that HIIT programs are a time-efficient tool for improving the biomarkers of cardiovascular disease.¹³ However, recommendations regarding the intensity, volume, frequency of the exercise, and rest intervals remain scarce which hinders the application of HIIT in the context of school PE classes. The question of whether HIIT could be used for the treatment and prevention of obesity in school-aged children, and the feasibility of applying HIIT in school PE programs, remain to be addressed. Therefore, the aim of this systematic review was to critically analyze the feasibility of including HIIT-based programs in PE classes to improve the body compositions and cardiorespiratory fitness of overweight and obese students.

Methods

Search strategy

This systematic review was carried out following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) recommendations.¹⁴ The following databases were used: Web of Science, Scopus, MEDLINE, SPORTDiscus and PUBMED. The keywords used in the search and their possible combinations were: obesity or overweight or obese and child or adolescents or children or teen and high-intensity interval training or high-intensity interval exercise or high-intensity intermittent exercise.

Inclusion criteria

The inclusion criteria were as follows: a) children and adolescents classified as overweight or obese between 6 and 19 years old exclusively, b) studies published between 31 January 2012 and 31 January 2017, c) studies that included anthropometric and cardiorespiratory capacity evaluations in pre- and post-intervention, d) studies in English published in peer-reviewed journals, and e) studies that used HIIT alone or combined with other methods.^{15,16}

Studies where caloric restriction was applied, interventions that used pharmaceuticals, or those centered on programs aimed at adolescents with hormonal imbalances and eating disorders were excluded.

Quality assessment

Two reviewers (P.D. and D.J.) independently selected titles and abstracts of potential studies identified by the search strategy. Study quality was determined through the Evidential Physiotherapy (PEDro) database,¹⁷ with scores ranging from 0 to 11 points.

Results

The search procedure identified 158 potential articles (Web of Science, $n = 29$; SCOPUS, $n = 81$; Medline, $n = 14$; SPORTDiscus, $n = 10$; PUBMED, $n = 24$). In the initial analysis, 53 publications were excluded due to duplication. Subsequently, 70 articles were excluded after reading the title and summary. The remaining 35 articles were read in full and six were selected as eligible after quality assessment^{18–23} (Fig. 1).

High-intensity training modalities, outcome measures, and results of different interventions are presented in Table 1. Of the six studies selected, three were conducted in children under 12 years old^{18,22,23} and three focused on adolescents between 12 and 18 years old.^{19–21} The HIIT protocols consisted of 2–3 sessions per week, with working times of 15 s and passive or active rest periods of 15 s, totaling up to 6 min work with 4 min rest. The durations of the programs were 6–24 weeks. According to the studies included in this review, significant changes were reported in body composition Body mass index (BMI), BMI z-score, % body fat (%BF) and waist circumference (WC),¹⁹ in the sum of skinfolds,¹⁸ in body mass (BM), BMI, BF% and WC,²¹ and increase in muscle mass.²³ HIIT programmes improved maximal oxygen uptake ($\text{VO}_{2\text{max}}$),^{19,21–23} performance in the intermittent Yo-Yo test, maximal aerobic speed (MAS)¹⁸ and aerobic capacity.²⁰

Quality assessment

Study quality for the seven selected articles using the PEDro scale. This returned a score of seven points out of a maximum of eleven (Table 2). The lost points were mainly due to: i) assignment concealment not being entirely relevant in studies of this nature, ii) difficulty in blinding the participants, as they are previously informed of the type of activity that will be undertaken, and iii) blinding the persons who apply the interventions (specifically, teachers) was not applicable.

Discussion

The main findings of this review were: 1) HIIT programs resulted in improvements in body composition in less time than less-intense exercise, 2) HIIT programs resulted in improvements in cardiorespiratory fitness, and 3) the HIIT protocols are highly applicable to PE classes due to their short duration and adaptability to the school setting.

Body composition

This review revealed that HIIT resulted in significant changes in several parameters compared with control groups that performed other types of exercises or did not receive intervention. Racil et al.¹⁹ compared HIIT with moderate-intensity interval training (MIIT) and found that HIIT led to greater improvements on cardiometabolic variables. Lau et al.¹⁸ observed a significant decrease in the sum of skinfolds of overweight children who followed HIIT programs in comparison with low-intensity (LI) interval exercises. In another investigation, Racil et al., compared HIIT programs with HIIT-plus-plyometric programs and reported that both approaches produced significant results (BMI, BF%, WC).²¹ Hay et al.,²⁰ however,




Fig. 1. Flowchart illustrating the different phases of the search and selection of studies.

did not report any differences in adiposity measurements between groups who undertook HIIT and those who followed a moderate-intensity endurance training (ET) program. Lambrick et al.²³ through high intensity discontinuous games found that the anthropometric parameters of obese children improved significantly compared with the control group. However, McNarry,²² who performed an intervention with similar characteristics to that of Lambrick et al.,²³ reported no anthropometric variations.

In some studies, similar results were reported for body composition when comparing HIIT with ET, although the duration of HIIT sessions was 70% less than ET sessions.²⁴ In another study, HIIT proved to be a more effective approach for weight loss in obese adolescents compared with ET.²⁵ An intervention which employed different exercise methods reported that HIIT groups achieved greater increases in fat oxidation rates and greater decreases in BMI than the LI groups.²⁶ Racil et al.²⁷ observed a greater decrease in BF % in HIIT groups compared with MIIT groups. With regard to WC, the application of HIIT for obese adolescents resulted in a greater decrease than moderate exercise.²⁸ Subjects of both HIIT and ET reported similar energy expenditures in the 24 h post-session,²⁹ which could explain the similar results for body composition in some cases, despite the lower total volume of training and time commitment of HIIT.³⁰ A further benefit of high-intensity compared with LI training for obesity and cardiovascular health is the improvement of parasympathetic tone and autonomic heart

modulation in obese adolescents.³¹

Cardiorespiratory capacity

The HIIT programs included in this review improved VO_{2max} ,^{19,21–23} performance in the intermittent Yo-Yo test, MAS,¹⁸ and aerobic capacity.²⁰ When comparing the results found in this review with previous studies, we found that 4×4 min interval exercises at 90% maximal heart rate (HR) generated significant improvements in VO_{2MAX} after 3 months.³² Hottenrott et al.³³ showed that HIIT exercise caused significantly greater improvements in VO_{2MAX} than ET. According to a recent review, HIIT is a time-efficient tool for increasing cardiorespiratory fitness in adolescents.³⁴

Araujo et al.²⁴ found that the improvements in VO_{2max} for continuous training at 80% peak HR for 30–60 min are similar to those achieved by HIIT through sprints. After following an 8 week HIIT program, performing four repetitions at 90–100% max HR for 20 s with 10 s rest periods, significant increases in VO_{2max} of adolescents with low physical activity levels were observed.³⁵ Gillen et al.³⁶ reported that 12 weeks of HIIT with a total time of 10 min, resulted in similar improvements in VO_{2max} when compared with 50 min sessions of aerobic training. Similar improvements were seen in systolic and diastolic heart functions following twice weekly HIIT sessions for 13 weeks in place of standard aerobic training.³⁷ In addition, an intervention which compared HIIT with

Table 1
Results of studies examining the effects of HIIT on cardiorespiratory fitness and anthropometric parameters in obese children and adolescents (n = 6).

Studies	Subject description	Study design	Training program	Outcomes	Result
Lau et al. (2014)	n = 15 Age: 11.0 ± 0.6 years BM: 51.1 ± 6.6 kg Size: 146.7 ± 5.2 cm	Randomized clinical trial. HIIE, LIIE and CON	HIIE = 12 sets x 15-s at 120% MAS followed by a 15- second passive recovery. Duration: 6 weeks, three times a week.	BMI (kg/m ²) Skinfold thickness (mm) Sum of skinfolds (mm) YYIET (level one)	↓ Sum of skinfolds (mm) ↑ YYIET distance covered (m)
Racil et al. (2016)	n = 17 Age: 14.2 ± 1.2 years. BM: 87.3 ± 4.5 kg Size: 164.0 ± 5.0 cm	Randomized clinical trial. HIIT, MIIT and CON	HIIT = 3 sessions × 4–8 min (15 s/15 s)15-s at 100% MAS follow by 15 s recovery at 50% MAS. 3-min of inter-session passive recovery period. Duration: 12 weeks, three times a week.	BM (kg) BMI–Z-score BF (%) WC (cm) VO _{2max} (L·min ⁻¹)	↓ BM (kg) ↓ BMI Z-score ↓ WC (cm) ↓ BF (%) ↑ VO _{2max} (ml/kg/min)
Hay et al. (2015)	n = 38 Age: 15.3 ± 1.7 years BM:89.1 ± 15.7 kg Size: not reported	Randomized clinical trial. HIIT, moderate-intensity and CON	HIIT = 70–85% HRres Duration: 6- month, three times a week.	BM (kg) BMI (kg·m ²) WC (cm), BF% trunk fat % VO _{2max} (ml/kg/min)	↑ VO _{2max} (ml/kg/min)
Racil et al. (2015)	n = 23 Age: 16.6 ± 0.9 years BM: 83.9 ± 4.5 kg Size: 163 ± 5.0 cm	Randomized clinical trial HIIT, P+HIIT and CON	HIIT = 6–8 blocks per session of 30-s runs at 100%MAS, with 30-s active recovery between bouts at 50%. Duration: 12 weeks, three times a week.	BMI Z-score WC (cm) BF (%) LM (kg) VO _{2max} (ml/kg/min).	↓ BM (kg) ↓ BMI Z-score ↓ WC (cm) ↓ BF (%) ↑ VO _{2max} (ml/kg/min)
McNarry et al. (2015)	n = 15 Age: 9.3 ± 0.9 years BM: 49.1 ± 10.8 Kg Size: 143.3 ± 9.2 cm	Randomized clinical trial EX and CON	EX = 6-min high- intensity exercise followed by 2 min of recovery (6 games per session) and a 4-min circuit. Duration: 6 weeks, two times a week	BM (kg) BMI (kg·m ²) VO _{2max} (ml/kg/min).	↑ VO _{2max} (ml/kg/min)
Lambrick et al. (2015)	n = 28 Age: 9.3 ± 0.9 years BM 48.9 ± 11.0 kg Size: 143.3 ± 9.0 cm	Randomized clinical trial EX and CON	EX = 6-min high- intensity exercise followed by 2 min of recovery (6 games per session) and a 4-min circuit. Duration: 6 weeks, two times a week.	BM (kg) BMI (kg·m ²) WC(cm) BF(%) LM (kg) VO _{2max} (ml/kg/min).	↓ WC (cm) ↑ LM (kg) ↑ VO _{2max} (ml/kg/min).

BM = body mass; BMI = body mass index; WC = waist circumference; BF = body fat; LM = lean mass; FFM = free fat mass; LIIE = low intensity intermittent exercise; HIIE = high intensity intermittent exercise; HIIT = high intensity interval training; MIIT = moderate-intensity interval training; CON = control group; AT = aerobic training; P = plyometric training; P+HIIT = plyometric exercise combined with HIIT; EX = exercise; YYIET= Yo-Yo intermittent endurance test; MAS = maximal aerobic speed; HRres = reserve heart rate; VO_{2max}= maximal oxygen uptake; ↑; =Significant increase; ↓ = Significant reduction.

Table 2
Physiotherapy evidence database scale (PEDro).

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	Total Score
Lau et al. (2015)	1	1	0	1	0	0	0	1	1	1	1	7
Racilet al.(2016)	1	1	0	1	0	0	0	1	1	1	1	7
Hay et al. (2016)	1	1	0	1	0	0	0	1	1	1	1	7
Racilet al. (2015)	1	1	0	1	0	0	0	1	1	1	1	7
McNarry et al. (2015)	1	1	0	1	0	0	0	1	1	1	1	7
Lambrick et al. (2015)	1	1	0	1	0	0	0	1	1	1	1	7

Notes: 0 = item was not satisfied; 1 = item was satisfied. Item 1: eligibility criteria were specified; Item 2: subjects were randomly allocated to groups; Item 3: allocation was concealed; Item 4: the groups were similar at baseline regarding the most important prognostic indicators; Item 5: there was blinding of all subjects; Item 6: there was blinding of all therapists who administered the therapy; Item 7: there was blinding of all assessors who measured at least one key outcome; Item 8: measurements of at least one key outcome were obtained from more than 85% of the subjects initially allocated to groups; Item 9: all subjects for whom outcome measurements were available received the treatment or control condition as allocated, or where this was not the case, data for at least one key outcome were analyzed by "intention to treat"; Item 10: the results of between groups statistical comparisons are reported for at least one key outcome; Item 11: the study provides bothpoint measurements and measurements of variability for at least one key outcome.

LI aerobic training for 6 months showed significant reductions in systolic and diastolic blood pressure of participants in both groups. However, beneficial changes in HR and HR variability were only observed in the HIIT group.³¹ It is important to consider that while aerobic training is beneficial, its application requires longer session

times to achieve similar results to HIIT.

On the other hand, HIIT increase metabolic and cardiorespiratory stress. A single exercise session using a 4 × 4 min protocol at 90–95% peak power output causes increased VO₂, average HR, ventilation, and ventilatory equivalent.³⁸ These results are supported by a recent study conducted on 17 school-age children, which revealed increased VO₂ following 12 intervals of 30 s at 90% max HR.³⁹ Revised HIIT protocols can result in improved cardiorespiratory capacity, increased VO_{2max}, and significantly decreased systolic and diastolic blood pressure. Maintaining optimal levels of these parameters plays a fundamental role in limiting the risk factors in overweight adolescents.

Limitations

The limitations of this systematic review are as follows: 1) the studies considered performed HIIT with different exercises (games, running, etc.) and combined with other training methods, making it difficult to generate a specific intervention protocol; 2) studies with caloric restrictions were not accepted, mainly because this is an uncontrollable variable for the application of these methodologies in overweight and obese school-age children; and 3) sex was not considered when analyzing results.

Conclusion and practical application

In conclusion, HIIT programs can result in improvements in

body composition and cardiorespiratory capacity, as well as improvements in health markers. In addition, HIIT methods require less time than low-intensity activities, making them applicable for use in schools as a strategy to combat the childhood obesity pandemic. The incorporation of HIIT programs could be carried out alongside existing PE activities within the same lesson or in specific periods during the school day. The authors suggest that in order to guarantee HIIT effectiveness, programs should include 2–3 sessions per week, with intervals of 15–30 s and passive or active rest periods of 15–30 s. For higher volume programs, discontinuous games of up to 6 min work with 4 min rest periods for a total session time of 40 min can be used.

Despite this study highlights the importance and feasibility of HIIT methodology in a school setting, some questions need to be addressed. Future research should identify the optimal duration of HIIT sessions for improving physical, physiological and cognitive health in young children. Likewise, future studies should determine the optimal frequency within the school calendar for this type of programs.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest

The author(s) have no conflicts of interest relevant to this article.

Author contributions

PDF revised the literature and wrote the manuscript; PLR and FGP critically reviewed the study; DJM and FCN reviewed the literature and helped write the study. All authors have read and approved the final version of this paper and agree with the order of its presentation.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jesf.2018.11.003>.

References

- Karnik S, Kanekar A. Childhood obesity: a global public health crisis. *Int J Prev Med*. Jan 2012;3(1):1–7.
- Swinburn BA, Sacks G, Lo SK, et al. Estimating the changes in energy flux that characterize the rise in obesity prevalence. *Am J Clin Nutr*. Jun 2009;89(6):1723–1728.
- Lob-Corzilius T. Overweight and obesity in childhood—a special challenge for public health. *Int J Hyg Environ Health*. Oct 2007;210(5):585–589.
- Kosti RI, Panagiotakos DB. The epidemic of obesity in children and adolescents in the world. *Cent Eur J Publ Health*. Dec 2006;14(4):151–159.
- Jankowski M, Niedzielska A, Brzezinski M, Drabik J. Cardiorespiratory fitness in children: a simple screening test for population studies. *Pediatr Cardiol*. Jan 2015;36(1):27–32.
- Diez-Fernandez A, Sanchez-Lopez M, Mora-Rodriguez R, Notario-Pacheco B, Torrijos-Nino C, Martinez-Vizcaino V. Obesity as a mediator of the influence of cardiorespiratory fitness on cardiometabolic risk: a mediation analysis. *Diabetes Care*. 2014;37(3):855–862.
- Carrel AL, Clark RR, Peterson SE, Nemeth BA, Sullivan J, Allen DB. Improvement of fitness, body composition, and insulin sensitivity in overweight children in a school-based exercise program: a randomized, controlled study. *Arch Pediatr Adolesc Med*. Oct 2005;159(10):963–968.
- Hollis JL, Williams AJ, Sutherland R, et al. A systematic review and meta-analysis of moderate-to-vigorous physical activity levels in elementary school physical education lessons. *Prev Med*. May 2016;86:34–54.
- Bevans KB, Fitzpatrick LA, Sanchez BM, Riley AW, Forrest C. Physical education resources, class management, and student physical activity levels: a structure-process-outcome approach to evaluating physical education effectiveness. *J Sch Health*. Dec 2010;80(12):573–580.
- Kelishadi R, Azizi-Soleiman F. Controlling childhood obesity: a systematic review on strategies and challenges. *J Res Med Sci : Offic J Isfahan Univ Med Sci*. Oct 2014;19(10):993–1008.
- Thivel D, Masurier J, Baquet G, et al. High-intensity interval training in overweight and obese children and adolescents: systematic review and meta-analysis. *J Sports Med Phys Fit*. Mar 27 2018. <https://doi.org/10.23736/S0022-4707.18.08075-1>.
- Alvarez C, Ramirez-Campillo R, Ramirez-Velez R, Izquierdo M. Effects of 6-weeks high-intensity interval training in schoolchildren with insulin resistance: influence of biological maturation on metabolic, body composition, cardiovascular and performance non-responses. *Front Physiol*. 2017;8:444.
- Eddolls WTB, McNarry MA, Stratton G, Winn CON, Mackintosh KA. High-intensity interval training interventions in children and adolescents: a systematic review. *Sports Med*. Jun 22 2017;1–12.
- Hutton B, Catala-Lopez F, Moher D. The PRISMA statement extension for systematic reviews incorporating network meta-analysis: PRISMA-NMA. *Med Clinica*. Sep 16 2016;147(6):262–266.
- Gibala MJ, Little JP, Macdonald MJ, Hawley JA. Physiological adaptations to low-volume, high-intensity interval training in health and disease. *J Physiol*. Mar 01 2012;590(5):1077–1084.
- Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle. *Sports Med*. 2013;43(5):313–338.
- Sherrington C, Herbert RD, Maher CG, Moseley AM, PEDro. A database of randomized trials and systematic reviews in physiotherapy. *Man Ther*. Nov 2000;5(4):223–226.
- Lau PW, Wong del P, Ngo JK, Liang Y, Kim CG, Kim HS. Effects of high-intensity intermittent running exercise in overweight children. *Eur J Sport Sci*. 2015;15(2):182–190.
- Racil G, Coquart JB, Elmontassar W, et al. Greater effects of high- compared with moderate-intensity interval training on cardio-metabolic variables, blood leptin concentration and ratings of perceived exertion in obese adolescent females. *Biol Sport*. Jun 2016;33(2):145–152.
- Hay J, Wittmeier K, MacIntosh A, et al. Physical activity intensity and type 2 diabetes risk in overweight youth: a randomized trial. *Int J Obes*. Apr 2016;40(4):607–614.
- Racil G, Zouhal H, Elmontassar W, et al. Plyometric exercise combined with high-intensity interval training improves metabolic abnormalities in young obese females more so than interval training alone. *Appl Physiol Nutr Metabol*. 2015;41(1):103–109.
- McNarry MA, Lambrick D, Westrupp N, Faulkner J. The influence of a six-week, high-intensity games intervention on the pulmonary oxygen uptake kinetics in prepubertal obese and normal-weight children. *Appl Physiol Nutr Metabol*. 2015;40(10):1012–1018.
- Lambrick D, Westrupp N, Kaufmann S, Stoner L, Faulkner J. The effectiveness of a high-intensity games intervention on improving indices of health in young children. *J Sports Sci*. 2016/02/01 2016;34(3):190–198.
- Corte de Araujo AC, Roschel H, Picanco AR, et al. Similar health benefits of endurance and high-intensity interval training in obese children. *PLoS One*. 2012;7(8), e42747.
- Kargarfard M, Lam ET, Shariat A, et al. Effects of endurance and high intensity training on ICAM-1 and VCAM-1 levels and arterial pressure in obese and normal weight adolescents. *Phys Sportsmed*. Sep 2016;44(3):208–216.
- Lazzer S, Tringali G, Caccavale M, De Micheli R, Abbruzzese L, Sartorio A. Effects of high-intensity interval training on physical capacities and substrate oxidation rate in obese adolescents. *J Endocrinol Invest*. Feb 2017;40(2):217–226.
- Racil G, Ben Ounis O, Hammouda O, et al. Effects of high vs. moderate exercise intensity during interval training on lipids and adiponectin levels in obese young females. *Eur J Appl Physiol*. Oct 2013;113(10):2531–2540.
- Starkoff BE, Eneli IU, Bonny AE, Hoffman RP, Devor ST. Estimated aerobic capacity changes in adolescents with obesity following high intensity interval exercise. *Appl Physiol Nutr Metab*. 2014;2(3):1.
- Francois ME, Graham MJ, Parr EB, et al. Similar metabolic response to lower-versus upper-body interval exercise or endurance exercise. *Metabolism*. 2017;68:1–10.
- Skelly LE, Andrews PC, Gillen JB, Martin BJ, Percival ME, Gibala MJ. High-intensity interval exercise induces 24-h energy expenditure similar to traditional endurance exercise despite reduced time commitment. *Appl Physiol Nutr Metab*. Jul 2014;39(7):845–848.
- Farah B, Ritti-Dias R, Balagopal P, Hill J, Prado W. Does exercise intensity affect blood pressure and heart rate in obese adolescents? A 6-month multidisciplinary randomized intervention study. *Pediatr Obes*. 2014;9(2):111–120.
- Tjønnå AE, Stølen TO, Bye A, et al. Aerobic interval training reduces cardiovascular risk factors more than a multitreatment approach in overweight adolescents. *Clin Sci*. 2009;116(4):317–326.
- Hottenrott K, Ludyga S, Schulze S. Effects of high intensity training and continuous endurance training on aerobic capacity and body composition in recreationally active runners. *J Sports Sci Med*. 2012;11(3):483–488.
- Costigan SA, Eather N, Plotnikoff RC, Taaffe DR, Lubans DR. High-intensity interval training for improving health-related fitness in adolescents: a systematic review and meta-analysis. *Br J Sports Med*. Oct 2015;49(19):1253–1261.
- Logan GR, Harris N, Duncan S, Plank LD, Merien F, Schofield G. Low-active male adolescents: a dose response to high-intensity interval training. *Med Sci Sports Exerc*. Mar 2016;48(3):481–490.
- Gillen JB, Martin BJ, MacInnis MJ, Skelly LE, Tarnopolsky MA, Gibala MJ. Twelve weeks of sprint interval training improves indices of cardiometabolic health

- similar to traditional endurance training despite a five-fold lower exercise volume and time commitment. *PLoS One*. 2016;11(4), e0154075.
37. Ingul CB, Tjonna AE, Stolen TO, Stoylen A, Wisloff U. Impaired cardiac function among obese adolescents: effect of aerobic interval training. *Arch Pediatr Adolesc Med*. 2010;164(9):852–859.
38. Kilian Y, Engel F, Wahl P, Achtzehn S, Sperlich B, Mester J. Markers of biological stress in response to a single session of high-intensity interval training and high-volume training in young athletes. *Eur J Appl Physiol*. Dec 2016;116(11–12):2177–2186.
39. Harris NK, Dulson DK, Logan GRM, Warbrick IB, Merien FLR, Lubans DR. Acute responses to resistance and high-intensity interval training in early adolescents. *J Strength Cond Res*. May 2017;31(5):1177–1186.