



Short communication

Detection of specific immune response in sole (*Solea senegalensis*) against *Photobacterium damsela* subsp. *piscicida* antigens

A. Medina, I. García-Millán, E. Martínez-Manzanares, M.A. Moriñigo, S. Arijo*

Universidad de Málaga, Departamento de Microbiología, 29071, Málaga, Spain

ARTICLE INFO

Keywords:

Solea senegalensis

Photobacterium damsela subsp. *piscicida*

Immunization

Antibody

ELISA

2D-PAGE

Western blot

Antigen identification

ABSTRACT

The pathogenic bacteria *Photobacterium damsela* subsp. *piscicida* affects the development of *Solea senegalensis* culture. Vaccines made with inactivated cells have produced a relative protection against the sickness, however the administration of subcellular and purified antigens as vaccine could increase the effectiveness of the immune response.

Thus, the aim of this work was the determination of antigens of *P. damsela* subsp. *piscicida* involved in the specific immune response of *S. senegalensis*.

Fish were immunized by intraperitoneal injection (i.p.) with inactivated extracellular polymeric substances (ECP) and whole cells of *P. damsela* subsp. *piscicida*, and Freund's incomplete adjuvant. Two months later fish were boosted with the same antigens. Serum from fish was collected to determine by ELISA the title of antibodies against subcellular fractions of bacteria (ECP, capsule, outer membrane proteins, O antigen and formalized whole cells). Significant differences were found between control and immunized fish, but differences between first immunization and booster were only found for O antigen and capsule.

Western blots derived from 2D-PAGE of ECP and Outer Membrane Proteins (OMP), using sole immunized serum, detected two high reactive antigens from ECP. Proteins were identified, by mass spectrometry, as ATP-dependent metalloprotease and Telurite resistance proteins. In the case of OMP, three antigenic proteins were detected and identified as Nrfa Y218f, Anti-oxidant AhpC/TSA, and a protein domain DNA binding heat shock related.

Senegalense sole (*Solea senegalensis*) has a high potential for diversification of aquaculture. However, its production is limited by the emergence of pasteurellosis outbreaks caused by the pathogenic bacteria *Photobacterium damsela* subsp. *piscicida*.

Several studies have reported the protection against *P. damsela* subsp. *piscicida* by different designs of vaccines in cultured fish [1–5]. Vaccines are usually composed of inactivated whole cells (bacterin), containing a wide pool of antigens. However, not all the antigens protect against pathogens. Different studies showed that vaccines consisting of immunogenic fractions can induce higher protection than inactivated whole cell bacteria in fish [6,7]. Therefore, the detection of immunogenic bacterial antigens is vital to design an antigen-based vaccine. Extracellular substances (ECP) and outer membrane have demonstrated a high specific immune response in immunized fish [8,9] and could be good candidates to select immunogenic proteins.

The aim of this work is the identification of subcellular components of *Photobacterium damsela* subsp. *piscicida* involved in the immunogenic response of Senegalese sole (*Solea senegalensis*).

1. Fish

Solea senegalensis specimens, ranging from 50 to 100 g, were obtained from fish farms in the south of Spain (Cádiz and Motril) and maintained in aerated marine water at 17–22 °C. Feeding was to 2% body weight daily with commercial feed. All the experiments were performed using 6–8 fish as replicates.

2. Bacteria strain

Photobacterium damsela subsp. *piscicida* Lg41/01, a virulent strain isolated from a sick sole (LD50 2.8×10^4 cfu/g), was cultured in trypticase soy broth and trypticase soy agar (Oxoid) supplemented with NaCl at 2% final concentration (TSBs and TSAs respectively).

3. Fish immunization to obtain polyclonal antiserum

Fish were immunized with inactivated whole cells (bacterin) and

* Corresponding author.

E-mail address: sarijo@uma.es (S. Arijo).




Fig. 1. Specific antibody titre for Lg41/01 antigens in serum of sole immunized with *P. damselae* subsp. *piscicida* whole cells and ECP by i.p. injection using bacterine, extracellular products (ECP), outer membrane proteins (OM), somatic antigen (AgO) and capsule as antigens. The Y axis shows the values of the limit dilution of sera expressed in log₂. *Significant differences to the control, **significant differences to the control and once vaccinated fish (p < 0.05).




Fig. 2. ECP spots from 2D-PAGE (a) and Western blotting (b) using sole immunized serum against *P. damselae* subsp. *piscicida* whole cells and ECPs. A: metalloprotease ATP-dependent; B: tellurite resistance protein.

ECP of Lg41/01 strain according Arijo et al. [2]. The bacterin was obtained scraping the whole cells into phosphate-buffered saline (PBS) to obtain a suspension of 10¹⁰ cells/mL. Formalin to 1% (v/v) final concentration was added with incubation overnight at 4 °C to inactivate the cells. The suspensions were washed thrice in PBS, and then resuspended in PBS to 10⁹ cells/mL. The extracellular polymeric substances (ECP) of *P. damselae* subsp. *piscicida* were obtained using the

cellophane technique described by Liu [10]. The ECP were inactivated by the addition of formalin to a final concentration of 1% (v/v), and then heating at 100 °C for 30 min.

Suspension of inactivated whole cells was supplemented with 10% of inactivated ECP (800 µg proteins/ml). Then, sole specimens were inoculated with 0.1 mL of inactivated whole cell-ECP preparation in 0.1 mL of Freund's incomplete adjuvant (FIA; Sigma-Aldrich) by i.p.




Fig. 3. OMP spots from 2D-PAGE (a) and Western blotting (b) using sole immunized serum against *P. damsela* subsp. *piscicida* whole cells and ECPs. A: Nrfa; B: AhpC/TSA; C: protein with DNA binding heat shock domain.

injection. Two months later, the fish were re-inoculated with the same antigens and adjuvant. Two months after the first immunization and one month after the second immunization, groups of fish were anaesthetized and blood collected by venepuncture, allowed to clot to obtain sole antiserum to Lg41.01 and stored at -80°C until its use.

4. Detection of specific antibody titre of serum

Indirect ELISAs, following the method described by Arijó et al. [6], were carried out to test the antibody titre in sole serum. The following bacterial antigens were used in ELISAs: inactivated whole cells (bacterin); ECP (isolated as before); bacterial outer membrane proteins (OMP), purified according to the method of Lambert [11]; O-antigens (Ag-O), obtained following the technique described by Edwards and Ewing [12]; and capsule, isolated using the technique described by Bonet et al. [13].

Briefly, bottomed microtitre plates were coated with inactivated whole cells, ECP, OMP, Ag-O and capsule. Antigen-coated wells were sequentially incubated with serial dilutions of fish serum, polyclonal rabbit antibody anti-sole immunoglobulin (produced by the University of Malaga) (dilution 1:1000), horseradish peroxidase-conjugated antibody goat anti-rabbit immunoglobulin (Sigma) (1:5000), and developing solution (o-phenyldiamine dihydrochloride, Sigmafast OPD; Sigma-Aldrich). The reaction was stopped by $3\text{M H}_2\text{SO}_4$. Absorbance values were determined using a microplate reader (Thermo Electron Corporation) at 490 nm. The limit dilution method was used for the determination of the specific antibody titre (absorbance of the highest serum dilution \geq average of blanks absorbance + two SD) [14]. Results were compared using a Student's T-test, with significant differences at $p < 0.05$.

A significant increase of specific antibody level was detected in immunized fish for all the *P. damsela* subsp. *piscicida* subcellular compounds tested regard to unvaccinated fish (Fig. 1). Similar results for *P. damsela* subsp. *piscicida* vaccines have been reported for gilthead seabream (*Sparus aurata*) [8]. However, the booster did not significantly increase the antibody titre, comparing with one immunized

fish, except for somatic antigen and capsule fractions in which boosters increased the titre obtained respect to the first immunization (Fig. 1). This result may also be influenced by the use of Freund's incomplete adjuvant [15]. In this case, for OMP and ECP, a booster could be unnecessary.

There were no significant differences in antibody titres between the different subunits, so we can consider that any of them is optimal for vaccine development.

5. Identification of immunogenic proteins

ECP and OMP were selected for the identification of immunogenic proteins in base of that a high level of antibody titre is obtained with only one immunization. The analysis was carried out using two-dimensional PAGE [16]. Samples were added to 11 cm immobilized pH gradient strips (IPG, pH gradient 3–10, ReadyStrip IPG strips, Bio-Rad). Proteins in IPG strips were electrofocused using a Protean Isoelectrofocusing (IEF) Cell, (BioRad). The IPG strips were placed on top of 10% acrylamide gel subjected to immunoblot analysis. Second dimension separations were performed on an Omnipage platform (Cleaver Scientific). Proteins were separated according their molecular weight at 250 V. Proteins in the gels were fixed with 50% methanol/10% acetic acid. The gels were stained by silver stain (Proteo Silver Plus, Sigma) and were scanned at 700 ppi (GS-800, BioRad).

Western Blot analysis was carried out according to the method described by Arijó et al. [6]. After electrophoresis, gels were transferred into 0.45 μm pore size nitrocellulose membranes (Sigma) in a semi-dry transblotter (BioRad). Then, the membranes were incubated in Senegalese sole sera immunized against *P. damsela* subsp. *piscicida* (dilution 1:400). Sera collected from control fish were used as negative control. The membranes were sequentially incubated with rabbit anti-Senegalese sole Ig (1:1000), goat anti-rabbit Ig G horseradish peroxidase-conjugate (Sigma) diluted 1:5000, and developer 3,3',5,5'-Tetramethylbenzidine (TMB; Sigma). The 2-D image of the gel from each sample was compared with the Western Blot, selecting the spots that reacted with the sole antiserum.

Table 1
ECP and OMP spots isolated from *P. damsela* subsp. *piscicida* by 2D-PAGE and identified with confidence by MASCOT (confidence interval > 75%). MW: molecular weight; PI: isoelectric point; and Accession number according the National Center for Biotechnology (<http://www.ncbi.nlm.nih.gov/>).

Sample	Protein	Specie	Protein score	Confidence interval	Protein MW (Da)	PI	Accession number
ECP A	ATP-dependent metalloprotease protein	<i>Serratia proteamaculans</i>	78	99,768	70577,4	5,95	gi 157368730
ECP B	Tellurite resistance protein	<i>Bacillus cereus</i>	81	99,881	41656,6	5,37	gi 54298038
OMP A	ChainA, Nrfa Y218f protein (nitrite reductase)	<i>Wolinella succinogenes</i>	94	99,994	55771,8	7,56	gi 168988896
OMP B	Anti-oxidant Ahp/CTSA family protein	<i>Cobwellia psycherythraea</i> , <i>Vibrio cholerae</i> , <i>Photobacterium</i> sp...	159	100	22146,8	4,69	gi 71282444
OMP C	cold-shock DNA-binding domain-containing protein	<i>Shewanella loihica</i>	136	100	7432,74	5,09	gi 127512097

The protein identification was carried out by mass spectrometry MALDI/TOF-TOF (Matrix Assisted Laser Desorption/Time of Flight, 4700 Proteomics Analyzer, Applied Biosystems), obtaining a mass spectrum (MS) or “Footprint Protein” [17]. A database search was then performed, combining the results of the fingerprint peptide (MS) with a fragmentation (MS/MS) of peptides from each spot according to the MS spectrum, using MASCOT as the search method.

ECP showed two spots corresponding to immunogenic proteins with high reactivity close to 50 kDa (Fig. 2), while 3 OMP were highly reactive proteins from 50 kDa to 20 kDa (Fig. 3). These proteins had a Mascot Score greater than 75, representing a significant identification ($p < 0.05$) and upper confidence interval of 99.7% (Table 1). Western Blot analysis using control sera did not show any reactive protein.

The two ECP identified have both a potential role in virulence and antimicrobial resistance (Table 1): ECP-A was identified as a metallo-protease ATP-dependent. The ATP-dependent proteases break trans-membrane proteins and activate virulence factors [18,19]. Also, several metalloproteases have been identified as virulence factor secreted by *P. damsela* subsp. *piscicida* and it is abundantly [20,21]. ECP-B was a protein that confers resistance to tellurite. Despite being a rare element in the environment, the tellurite resistance gene operon (*ter*) is found in many pathogenic species. Although the *ter* operon has been associated in tellurite resistance and pathogenicity, its mode of action still remains unclear. *Ter* proteins could acts against host defenses produced by the cells of the immune system by employing mechanisms equivalent to tellurite reduction [22].

Regarding the immunogenic OMP (Table 1), OMP-A was identified as a protein Nrfa, a cytochrome c nitrite reductase. Nitrite reductase proteins diminish the impact of nitric oxide generated by the immune response [23]. The OMP-B was homologous to a protein antioxidant AhpC/TSA (peroxiredoxin). Peroxiredoxin family proteins protect the pathogen against reactive oxygen species [24]. The Ahpc genes increase their transcription when *P. damsela* subsp. *piscicida* infect *S. senegalensis* [25]. On the other hand the OMP-C was homologous to a protein domain DNA binding heat shock related. There is very little information available about the immune importance of OMP-C, a protein that protects in the low-temperature [26].

In conclusion, *P. damsela* subsp. *piscicida* subcellular components produce a specific immune response in farmed soles even with one immunization. At least two ECP proteins and three OMP are good candidates to design vaccines. However, more investigations must be developed to demonstrate their ability to protect the host against this pathogen.

Acknowledgements

This work was supported by the Ministry of Science of Spain (*Plan Nacional de Investigación*, AGL2010-17789) and the European Regional Development Fund. We also wish to thank the collaboration of Mr. Victor Fernandez, for the supply and maintenance of fish used in the experiments, and the Instituto Español de Oceanografía (IEO) of Vigo and Santander, for supplying fish serum. English language assistance provided by Mrs. Marie-Louise Smets.

References

- [1] F. Andreoni, R. Boiani, G. Serafini, G. Amagliani, S. Dominici, G. Riccioni, ... M. Magnani, Isolation of a novel gene from *Photobacterium damsela* subsp. *piscicida* and analysis of the recombinant antigen as promising vaccine candidate, *Vaccine* 31 (2013) 820–826.
- [2] S. Arijó, R. Rico, M. Chabrilion, P. Diaz-Rosales, E. Martínez-Manzanares, M.C. Balebona, B. Magariños, A.E. Toranzo, M.A. Moriñigo, Effectiveness of a divalent vaccine for sole, *Solea senegalensis* (Kaup), against *Vibrio harveyi* and *Photobacterium damsela* subsp. *piscicida*, *J. Fish. Dis.* 28 (2005) 33–38.
- [3] K. Gravingen, M. Sakai, T. Mishiba, T. Fujimoto, The efficacy and safety of an oil-based vaccine against *Photobacterium damsela* subsp. *piscicida* in yellowtail (*Seriola quinqueradiata*): a field study, *Fish Shellfish Immunol.* 24 (2008) 523–529.
- [4] L.P. Ho, C.J. Chang, H.C. Liu, H.L. Yang, J.Y. Lin, Evaluating the protective efficacy

- of antigen combinations against *Photobacterium damsela* ssp. *piscicida* infections in cobia, *Rachycentron canadum* L., J. Fish. Dis. 37 (2014) 51–62.
- [5] G. Kato, K. Yamashita, H. Kondo, I. Hirono, Protective efficacy and immune responses induced by a DNA vaccine encoding codon-optimized PPA1 against *Photobacterium damsela* subsp. *piscicida* in Japanese flounder, Vaccine 33 (2015) 1040–1045.
- [6] S. Arijo, J. Brunt, M. Chabrilón, P. Díaz-Rosales, B. Austin, Subcellular components of *Vibrio harveyi* and probiotics induce immune responses in rainbow trout, *Oncorhynchus mykiss* (Walbaum), against *V. harveyi*, J. Fish. Dis. 31 (2008) 579–590.
- [7] L.P. Ho, J. Han-You Lin, H.-C. Liu, H.-E. Chen, T.-Y. Chen, H.-L. Yang, Identification of antigens for the development of a subunit vaccine against *Photobacterium damsela* ssp. *piscicida*, Fish Shellfish Immunol. 30 (2011) 412–419.
- [8] S. Arijo, C. Balebona, E. Martínez-Manzanares, M.A. Moriñigo, Immune response of gilt-head seabream (*Sparus aurata*) to antigens from *Photobacterium damsela* subsp. *piscicida*, Fish Shellfish Immunol. 16 (2004) 65–70.
- [9] A. Medina, J.M. Mancera, E. Martínez-Manzanares, M.A. Moriñigo, S. Arijo, Identification of *Vibrio harveyi* proteins involved in the specific immune response of Senegalese sole (*Solea senegalensis*, Kaup), Fish Shellfish Immunol. 47 (2015) 377–380.
- [10] P.V. Liu, Survey of hemolysin production among species of pseudomonads, J. Bacteriol. 74 (1957) 718–727.
- [11] P.A. Lambert, Enterobacteriaceae: composition, structure and function of the cell envelope, J. Appl. Bacteriol. 65 (1988) 21S–34S.
- [12] P.R. Edwards, W.H. Ewing, Edwards and Ewing's Identification of Enterobacteriaceae, Elsevier, 1986.
- [13] R. Bonet, B. Magariños, J.L. Romalde, M.D. Simon-Pujol, A.E. Toranzo, F. Congregado, Capsular polysaccharide expressed by *Pasteurella piscicida* grown in vitro, FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Lett. 124 (1994) 285–289.
- [14] H.U. Bergmeyer, J. Bergmeyer, M. Grassl, Methods of Enzymatic Analysis: Samples, Reagents, Assessment of Results, Verlag Chemie, 1983.
- [15] D.P. Anderson, Immunostimulants, adjuvants, and vaccine carriers in fish: applications to aquaculture, Annu. Rev. Fish Dis. 2 (1992) 281–307.
- [16] A. Görg, W. Weiss, M.J. Dunn, Current two-dimensional electrophoresis technology for proteomics, Proteomics 4 (2004) 3665–3685.
- [17] P. Wilmes, P.L. Bond, The application of two-dimensional polyacrylamide gel electrophoresis and downstream analyses to a mixed community of prokaryotic microorganisms, Environ. Microbiol. 6 (2004) 911–920.
- [18] T. Defoirdt, Virulence mechanisms of bacterial aquaculture pathogens and anti-virulence therapy for aquaculture, Rev. Aquacult. 6 (2014) 100–114.
- [19] R. Zhou, C. Cusumano, D. Sui, R.M. Garavito, L. Kroos, Intramembrane proteolytic cleavage of a membrane-tethered transcription factor by a metalloprotease depends on ATP, Proc. Natl. Acad. Sci. U. S. A. 106 (2009) 16174–16179.
- [20] A. Do Vale, D. Cabanes, S. Sousa, Bacterial toxins as pathogen weapons against phagocytes, Front. Microbiol. 7 (2016) 42.
- [21] D.S. Silva, L.M. Pereira, A.R. Moreira, F. Ferreira-da-Silva, R.M. Brito, T.Q. Faria, ... P.J. Pereira, The apoptogenic toxin AIP56 is a metalloprotease AB toxin that cleaves NF-kb P65, PLoS Pathog. 9 (2) (2013) e1003128.
- [22] L. Turkovicova, R. Smidak, G. Jung, J. Turna, G. Lubec, J. Aradska, Proteomic analysis of the TerC interactome: novel links to tellurite resistance and pathogenicity, J. Proteomics 136 (2016) 167–173.
- [23] A. Vázquez-Torres, A.J. Bäuml, Nitrate, nitrite and nitric oxide reductases: from the last universal common ancestor to modern bacterial pathogens, Curr. Opin. Microbiol. 29 (2016) 1–8.
- [24] G.H. Kaihami, J.R.F. de Almeida, S.S. dos Santos, L.E.S. Netto, S.R. de Almeida, R.L. Baldini, Involvement of a 1-Cys peroxiredoxin in bacterial virulence, PLoS Pathog. 10 (2014) 1–10.
- [25] J. Núñez-Díaz, M. Fumanal, A. do Vale, C. Fernández-Díaz, M. Moriñigo, M. Balebona, Transcription of IVIAT and virulence genes in *Photobacterium damsela* subsp. *piscicida* infecting *Solea senegalensis*, Microorganisms 63 (2018) 67.
- [26] C. Zhu, B. Sun, T. Liu, H. Zheng, W. Gu, W. He, ... X. Peng, Genomic and transcriptomic analyses reveal distinct biological functions for cold shock proteins (Vpa CspA and Vpa CspD) in *Vibrio parahaemolyticus* CHN25 during low-temperature survival, BMC Genomics 18 (1) (2017) 436.