



Full length article

Catching the complexity of salmon-lice interactions

Cristian Gallardo-Escárate^{a,*}, Valentina Valenzuela-Muñoz^a, Gustavo Núñez-Acuña^a,
Crisleri Carrera^a, Ana Teresa Gonçalves^a, Diego Valenzuela-Miranda^a, Bárbara P. Benavente^a,
Steven Roberts^b

^a Interdisciplinary Center for Aquaculture Research, Laboratory of Biotechnology and Aquatic Genomics, Department of Oceanography, Universidad de Concepción, Concepción, Chile

^b School of Aquatic and Fishery Sciences (SAFS), University of Washington, Seattle, USA

ABSTRACT

The study of host-parasite relationships is an integral part of the immunology of aquatic species, where the complexity of both organisms has to be overlaid with the lifecycle stages of the parasite and immunological status of the host. A deep understanding of how the parasite survives in its host and how they display molecular mechanisms to face the immune system can be applied for novel parasite control strategies. This review highlights current knowledge about salmon and sea louse, two key aquatic animals for aquaculture research worldwide. With the aim to catch the complexity of the salmon-lice interactions, molecular information gleaned through genomic studies are presented. The host recognition system and the chemosensory receptors found in sea lice reveal complex molecular components, that in turn, can be disrupted through specific molecules such as non-coding RNAs.

1. Introduction

Host-parasite interactions are a complex relationship where one organism benefits the other and often where there is a negative impact on the host [1,2]. In the presence of a parasite, the host displays changes in its behavior, reproduction, regulation of immune response, among other processes [2]. Meanwhile, the parasite to establish itself in the host must counteract its defense mechanisms [1]. This explains why parasites have evolved to develop multiple strategies to evade the defense mechanisms of their hosts, and in parallel, the latter have had to develop strategies to evade the parasites [3]. In parasites, there are several different evasion mechanisms of the host response. For example, the plant parasite *Meloidogyne* secretes proteins such as proteases, superoxide dismutase and calreticulin [3]. In the case of ectoparasites, proteases, protease inhibitors, cathepsins, and other molecules are released. These secreted proteins can immuno-suppress the host, facilitating ectoparasite penetration in unbroken skin tissues where feeding can occur [4,5]. A large part of the life cycle of ectoparasites occurs when they are established in their hosts with different developmental stages often involving different molecular mechanism driving host-parasite interactions [6,7]. The molecular basis of host-parasite interactions has been studied in the context of different environments but there is limited information from marine systems given the emerging landscape of these interactions associated with environmental change.

Sea lice is a marine ectoparasite that mainly infests salmonid species. Given their high impact on the production of salmonid fish

worldwide the most studied sea lice species are *Lepeophtheirus salmonis* and *Caligus rogercresseyi* [8–10]. Annually the salmon industry presents estimated losses of US \$ 480 million, representing between 4 and 10% of production costs [10]. The presence of sea lice on the fish causes skin damage, immunosuppression, and facilitate the co-infection with bacterial pathogens [11–13]. Like all ectoparasites, lice spend a large part of its life cycle on a host, which makes it necessary for this ectoparasite to generate mechanisms for evading the immune response of its host.

Recently, genomic tools have allowed increasing knowledge about the molecular mechanisms associated with the parasite-host interaction process [14–17]. As lice are organisms with a significant economic impact on aquaculture, numerous studies have been aimed to understand the mechanisms of immune responses of salmonids as well as the evasion mechanisms used by lice for a successful infection. In this, several hypotheses have been proposed to explain the clear differences that exist in host with different susceptibility to lice infections. Some of these hypotheses have been associated with host skin thickness [18], mucus composition [19], rapid and effective innate immune response [20], homeostasis and iron regulation [21], adaptive immunity [22] and regulation of responses measured by non-coding RNA [23]. In this review, different mechanisms sea lice use for recognition and response to salmonid defenses are described. In addition, immune response mechanisms that have been identified in salmonid species are discussed.

* Corresponding author. Universidad de Concepción, P.O. Box 160-C, Concepción, 4030000, Chile.

E-mail address: crisgallardo@udec.cl (C. Gallardo-Escárate).

2. Recognition of host and chemosensory receptors in sea lice

Chemical or “semiochemical” signals have been crucial for the evolution of various species of animals in the marine ecosystem [24–26]. For instance, copepods are a class of marine species that communicate through various intra- or inter-specific chemical signals [27,28]. This group of aquatic crustaceans has developed various receptor structures that allow the identification of various chemical signals. Among them, the most described are the *sensilla*, which are neurons located in the main chemosensory organs where the most common are known as unimodal *chemosilla* [29,30]. However, the molecular basis for the recognition of semiochemicals from the environment remains virtually unknown in these marine invertebrates.

The life cycle of the sea louse species is complex and consist of several instars divided by molts. For instance, *L. salmonis* has eight stages, comprising five larval stages, two preadult stages and the adult stage [157]. Meanwhile, *C. rogercresseyi* comprises two larval stages (nauplius I, nauplius II and copepodite), four juvenile stages (chalimus I - IV) and one adult stage (male or female) [138]. During the copepodite stage, the process of host identification occurs, preparing the lice for infestation and settlement. From this point, the juvenile stages can initialize the infective process. The successful infestation process on the host allows the parasites the intake of nutrients for reproduction and adult development [31]. Previous studies have shown that lice species, such as *L. salmonis* that infects fish in the northern hemisphere, have developed physical mechanisms of host recognition. Among these, lice can identify the temperature of the water in which white fish species live, salinity changes, and detect small variations in water movement generated swimming fish [28,32,33]. Host identification via detection of semiochemicals has also been reported [34,35]. The presence of advanced chemoreceptors in *L. salmonis* has been described, which are capable of identifying specific molecules of different host species [29,32,35,36]. The main chemosensory receptors described in lice have been found in antennae located in the parasite's head. Neuronal activity measured through electrophysiological probes increases in the antennae exposed to water containing fish mucus [29,37]. Likewise, there is evidence that the ablation of antennae generates a reduction in the capacity to seek the host [35]. Low molecular weight compounds such as isophorone and 6-methyl-5-hepten-2-one, produced by Atlantic salmon can generate attractancy on *L. salmonis* [34]. With respect to *C. rogercresseyi*, it has been described that there is a preference for water conditioned with fish mucus, suggesting that there is a similar chemical recognition phenomenon [38]. Receptors involved in host recognition have been identified in the sea louse *C. rogercresseyi*, suggesting that the gene family of ionotropic receptors (IRs) are key molecular components for the salmon-lice interaction [39]. Notably, these genes have differential expression depending on the host species, and the use of in-feed ingredients such as odor-masking compounds [40]. The functional role related to the chemosensory system has been based on the structure highly similar to those reported for insects, with a known olfactory function [41]. Additionally, these genes have an expression profile highly linked to the infective stage of copepodid (IR25a), or to adult males (kainate receptors), which have a role in mating behavior [42]. These receptors generally act in conjunction with binding proteins, which in the case of insects are known as odorant binding proteins (OBP), which have not been identified yet in sea lice [43]. However, this does not exclude all types of binding proteins, since there are other types that could interact with chemosensory receptors, such as chemosensory proteins (CSP), which are found in various arthropod species [44]. On the other hand, in *L. salmonis* it was identified that chemosensory genes were mainly associated to the gene expression of the parasite's antennules, which are structures with chemosensory function [45]. A recent study in *L. salmonis* showed that the silencing of an ionotropic receptor (IR25a) decreases parasite activity related to host recognition, evidencing a key role related to host recognition [46]. The IR genes have significant sequence similarity for *L. salmonis* and *C.*

rogercresseyi.

3. Effect of masking compounds on sea lice infections *C. rogercresseyi*

It has been shown that there is a differential expression of genes related to the chemosensory system of *C. rogercresseyi* depending on the host species. Through a temporal scale, there is a relationship between the expression of IRs in the parasite and the dynamics of parasite load in infected salmonids [39,40]. In these studies, the Atlantic salmon (*Salmo salar*) and Coho salmon (*Oncorhynchus kisutch*) have been identified as susceptible and resistant species, respectively, for lice infestation. Salmon gene expression patterns have been correlated with in-feed additives to disrupt the parasite host-recognition, and with immunostimulants to strengthen the fish immune defense [47]. The decrease in the expression of IRs in parasites infecting Atlantic salmon fed with masked diets indicates that the infestation success is reduced by disruption of chemosensory capacity.

4. Role of host salmon antimicrobial peptides on the chemosensory system of sea lice

In salmonids, antimicrobial peptides (AMPs) have been described as important defense mechanisms that are expressed in tissues that are generally exposed to infections. These molecules are secreted by the mucus and in the case of salmon through the skin [48,49]. Fish produce various types of AMP, including defensins and cathelicidins, as well as others specific to the group such as piscidins [50]. As for salmonid fish, they produce AMPs such as those mentioned above, as well as hepcidins and histone-derived AMPs. Together these AMPs form a main basis of defense against pathogens [51]. Of these AMPs, the cathelicidin group could also be involved in long-term defense, as described in trout [52]. Cathelicidins have been widely studied in several vertebrate species, but in salmonids, they have their own characteristics at a structural level, which could confer new functions [53]. This class of AMP is highly associated with the response of Atlantic salmon to bacterial infections such as yersiniosis [54]. Cathelicidins were also found among the genes that were activated in salmon infected with the sea lice ectoparasite *L. salmonis*, while other AMPs such as defensins decreased their expression levels [55].

It is important to point out that *C. rogercresseyi* begins to feed on host fish mucus when the copepodid is attached to the skin [31], so direct contact between AMPs and lice is suggested. It has been reported in other fish species that their AMPs may come into contact with ectoparasites that infect them in target tissues [56]. The expression profiles of AMPs in *C. rogercresseyi* hosts have been shown to be expressed in early stages of infection in *O. kisutch*, which is congruent with the observation that this species is resistant given a rapid and effective activation of the defensive mechanisms [57]. One of the AMPs produced by salmonids, cathelicidin-2 (Cath-2), promotes the development of the frontal filament of *C. rogercresseyi* and activates the expression of IRs, as well as genes related to the formation of the cuticle [58]. The frontal filament is a structure highly associated with the success of infection by this parasite so that the stimulus of its development suggests the host favors the adhesion of the louse on the fish skin [59].

Regarding the antimicrobial peptide Cath-2, it has an attractant activity on lice, especially in its infective stage. Characteristics of Cath-2 including its molecular weight, three-dimensional structure, chemical nature and concentration in the water meet all criteria to be detectable by sea lice [29]. Moreover, it has been established that the movement of copepodids when stimulated by Cath2 is directed, similar to that observed in response to stimuli with light [60]. This is consistent with that reported in another crustacean, *Daphnia pulex*, which has swimming behavior addressed by the presence of stimuli that generate attraction [61]. In addition, there is a global transcriptome activation in response to Cath-2 [58,60], which is consistent with other species of arthropods

that show a broad activation at the RNA level in the presence of specific attractant molecules [62]. Previous studies in *C. rogercresseyi* has revealed that genes modulated in copepodids match with chemosensory genes and synaptic-related genes [58], which is consistent with studies in crustaceans [63,64]. With all this evidence it is inferred that the antimicrobial peptide Cath-2 acts as a kairomone that attracts sea lice, favoring the infection process.

5. Modulation of the host's immune response during the lice infestation

The development of aquaculture worldwide has made it necessary to increase efforts to understand the defense mechanisms of fish against infections. Due to the explosive advance of the sequencing technologies, the amount of genomic information of aquatic organisms, as well as the knowledge of the immune response pathways of fish has substantially increased [65]. In general, teleost organisms present an immune system similar to that of other vertebrates with elements of the innate and adaptive immune response. Studies on the immune response mechanisms of fish in the presence of copepods are scarce. In Atlantic salmon infested with *L. salmonis*, the immune response has been described and is associated with the expression of pro-inflammatory cytokines [66], secretion of proteases [5], and humoral response [22,66]. Specifically, increasing levels of IL4/13 and IL10 have been reported in families of Atlantic salmon resistant to *L. salmonis*, as well as in pink and Coho salmon that are less susceptible, suggesting a CD4⁺ T helper 2 (Th2) type response to cope with lice infestation [9,22,67] (Fig. 1). The response mechanisms of salmonids infested by *C. rogercresseyi* differ from those described for salmon infested with *L. salmonis* [20,66–69] (Table 1). Notably, Atlantic salmon has higher levels of susceptibility to *C. rogercresseyi* compared to Coho salmon, which also modulates the types of the immune response during the infestation process. Atlantic salmon responds to the presence of *C. rogercresseyi* by activating a toll-like receptor (TLR), specifically tlr22a2, which has been reported exclusively in fish [70]. This TLR has also been associated in other species of fish infested by ectoparasites, as one of the first mechanisms to activate the immune response. For example, upregulation of tlr22a2 in *L. rohita* infested by the ectoparasite *A. siamensis* has been described [71,72] (Table 1). In *L. rohita*, a direct relationship was observed between expression levels of tlr22a2 and the amount of *A. siamensis* in fish [73]. Notably, Coho salmon does not have a significant increase in the expression of tlr22a2, contrary to the increase in the expression of tlr13, which in fish has a common ancestor with TLR22 [74] (Fig. 1). Thus, overregulation of tlr13 in Coho salmon suggests that this salmon species could respond via TLR to the infestation of *C. rogercresseyi*, similarly to Atlantic salmon with tlr22a2 [75] (Table 1). Furthermore, studies conducted on salmon infected with *L. salmonis* suggest a role for MHC II in response to skin damage in fish [22,67]. This has also been reported in fish with the presence of *Paramoeba perurans* [76]. The inflammatory response is the first reaction associated with the coping against sea lice, generating variations in the transcriptional activity of genes such as interleukin 1 α , MMP13, MMP19, TNF- α , prostaglandins E2 and IL-8 [9,20,22,66,67,77,78]. Fish that have lower susceptibility to infestation with *L. salmonis* display a Th2 type immune response [22,66,67,77] (Table 1). Coho salmon, tolerant to infection with *C. rogercresseyi*, displays a low expression of genes associated with the Th2 response. However, an early response to the presence of *C. rogercresseyi* in Coho salmon shows an increase in the expression levels of genes associated with Th1 response [75] (Fig. 1). We hypothesize that the low expression of Th2 genes in Coho salmon could be a strategy to avoid the overproduction of mucus in fish, the main food source of *C. rogercresseyi*, since it has been described that the Th2 response increases secretion of mucus through the production of mucin [66,79,80] (Table 1).

6. Role of iron homeostasis mechanisms in the Salmon-lice interaction

Parasites depend on host nutrients, thus it is crucial to select those that have the greatest amount of nutritional resources for their development, survival, and reproduction [81,82]. A type of immunity called nutritional immunity has been proposed and consists of the limitation of nutrients by the host [81], generating a hostile environment for the development of parasites. Here, the availability of ions such as Fe, Mn, Zn is mainly modulated through molecular mechanisms that involve key signaling pathways and regulatory components [81]. Among the most important nutrients in this immune strategy is iron, due to its role in different physiological processes such as respiration, and enzymatic activity [81,83]. It has also been observed that iron can modulate proinflammatory cytokines and acute phase proteins [84]. Evidence of nutritional immunity has been reported in fish infested by *L. salmonis*, observing a correlation between resistance to lice infection and heme regulation [69]. In parallel, fish infested with lice display low expression levels of hepcidin, a protein responsible for the transport of iron into the cell [22].

Transcriptome studies in infected-fish with *C. rogercresseyi* have shown a large number of transcripts associated with ion binding [85] (Table 1). For instance, an increase in hepcidin has been associated with the secretion of IL-6 after the activation of pattern recognition receptors (PPRs) during bacterial infections [86]. In macrophages, it has been observed that iron limitation reduces the expression of MHCII, iNOs, and also inhibits the IFN-g signaling pathway [84]. Likewise, in fish with low susceptibility to the ectoparasite *L. salmonis*, an increase in regulatory molecules of the heme group has been observed, reflected in the increase in the expression levels of heme oxygenase [69]. From transcriptomic data of the skin and head kidney of Atlantic and Coho salmon infested *C. rogercresseyi*, over-regulation of hemoglobin subunit in both species have been reported [85]. In contrast, Coho salmon infested with *C. rogercresseyi* a decrease in expression of heme oxygenase is associated with the heme degradation, and also an increasing level of the biosynthesis pathway comprising the aminolevulinic acid synthetase and coproporphyrinogenase genes [85]. For Atlantic salmon, there has been reported an increase in the expression of BLVr, a gene associated with the degradation of the heme group at seven days post-infestation [85]. This result, similar to that reported by González et al. [87], where a reduction of hemoglobin in Atlantic salmon infested by *C. rogercresseyi* was observed, suggests a reduction of the heme group as an iron source, as a strategy to cope with the lice infestation. Iron homeostasis is not only given by the regulation of the heme group, but there are also proteins responsible for reducing circulating iron as a consequence of the degradation of the heme group [83]. Among this, hepcidin, haptoglobin, ferritin, transferrin proteins have been reported to play a role in iron regulation. Atlantic salmon infected with *C. rogercresseyi* increases of BLVr levels has been reported in conjunction with the transferrin gene. It has been also observed high levels of hepcidin, a protein that has also been associated with anemic states in fish [88,89]. The modulation of the availability to store iron through biosynthesis or degradation of the heme group, and also iron transporter genes, suggests that this mechanism is a key molecular component exploited by fish during the lice infestation [85] (Table 1).

The nutritional strategy seems to be a unique response to lice infestation in Atlantic salmon. A comparative study in skin and head-kidney in infested fish with *L. salmonis* and *C. rogercresseyi* showed significant differences in genes associated with biosynthesis/degradation of the heme group as well as iron transporter genes and oxidative stress response [21]. Among the differences observed, it can be noted that fish infested with *C. rogercresseyi* showed high expression of hepcidin, such as the one observed in salmon resistant to *L. salmonis* [22]. In addition, fish infested with *C. rogercresseyi* showed high modulation of genes associated with heme group degradation. While the genes associated with biosynthesis pathway such as ferritin and transferrin




Fig. 1. Schematic model of salmon-lice interactions. (a) Description of the molecular mechanism of sea lice for host recognition. (b) Sea lice molecular response for avoiding host response and (c) Host innate and adaptive immune mechanism and iron homeostasis modulation in response to sea lice infestation.

receptor were highly expressed in fish infested with *L. salmonis* [21]. In addition, the increase in heme biosynthesis is associated with an increase in oxidative stress levels [90], which is reflected in fish infested with *L. salmonis* through the increasing of glutathione S-transferase (GST) and superoxide dismutase (SOD) transcription expression. With this information, it is possible to confirm that there is a nutritional response mechanism in Atlantic salmon infested by sea lice. According to the results, the nutritional strategy to respond to *L. salmonis* and *C. rogercresseyi* varies in the same species of salmonids, being of greater relevance in the response to *L. salmonis*, which is a hematophagous organism [90], whereas *C. rogercresseyi* is a facultative hematophagous

organism (Fig. 1).

7. The immune system of sea lice

From the recognition process through infestation and the development of different life stages, lice must cope with the rejection mechanisms displayed by the fish. In order to understand the molecular mechanisms that lice use for its defense, transcriptomic data from different stages of development of *C. rogercresseyi* [7] have been characterized. Innate immune response pathways IMD and TLR [91], genes associated with the antioxidant system (ROS) [92] and secretory

Table 1
Transcriptome studies associated with the host's immune response during sea lice infestation.

Host species	Sea lice species	Molecular mechanism	Experimental design	Reference
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Up-regulation of MHII, COX-2, CRP and IL-8	24-48 hpi (juvenile)	Braden et al. (2015)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Down-regulation of MHII, C-reactive protein, IL-1b, IL-8, COX-2	24-48 hpi (mature)	Braden et al. (2015)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Expression of MH class II, IL-6, CRP, MMP13, IL-1 β and COX-2 genes	24-48 hpi	Braden et al. (2012)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Increase expression of TNF-alpha	24-48 hpi	Braden et al. (2015)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Pro-inflammatory response, PRRs and iron homeostasis	6 dpi	Sutherland et al. (2014)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Oxidative stress	6 dpi	Sutherland et al. (2014)
<i>O. gorbuscha</i>	<i>L. salmonis</i>	Suppressed antiviral immunity	6 dpi	Sutherland et al. (2014)
<i>O. keta</i>	<i>L. salmonis</i>	Suppressed antiviral immunity	6 dpi	Sutherland et al. (2014)
<i>O. keta</i>	<i>L. salmonis</i>	Pro-inflammatory response, PRRs and iron homeostasis	6 dpi	Sutherland et al. (2014)
<i>O. kisutch</i>	<i>L. salmonis</i>	Early pro-inflammatory TH1-type pathway and regulatory TH2-type processes	24-48 hpi	Braden et al. (2015)
<i>O. kisutch</i>	<i>C. rogercresseyi</i>	Oxidative stress	7 dpi	Vera-Bizama et al. (2015)
<i>O. kisutch</i>	<i>C. rogercresseyi</i>	TLR pathway (up-regulation of tlr13)	14 dpi	Valenzuela-Muñoz et al. (2016)
<i>O. kisutch</i>	<i>C. rogercresseyi</i>	Increase in Th1 gene related	14 dpi	Valenzuela-Muñoz et al. (2016)
<i>O. kisutch</i>	<i>C. rogercresseyi</i>	Pro-inflammatory response	7-14 dpi	Valenzuela-Muñoz et al. (2016)
<i>O. kisutch</i>	<i>C. rogercresseyi</i>	Up-regulation of Heme synthesis	7-14 dpi	Valenzuela-Muñoz et al. (2017)
<i>S. salar</i>	<i>L. salmonis</i>	Suppressive effect in pro-inflammatory mediators	24 h dpi	Braden et al. (2012)
<i>S. salar</i>	<i>L. salmonis</i>	Innate immune response reactions	1 dpi	Tadiso et al. (2011)
<i>S. salar</i>	<i>L. salmonis</i>	Decreases of values of immune-related genes	4 dpi	Holm et al. (2017)
<i>S. salar</i>	<i>L. salmonis</i>	Suppressed antiviral immunity	6 dpi	Sutherland et al. (2014)
<i>S. salar</i>	<i>L. salmonis</i>	TLR/IMD signaling Pathway	7 dpi	Valenzuela-Muñoz et al. (2016)
<i>S. salar</i>	<i>L. salmonis</i>	Up-regulation of Heme synthesis	8 dpi	Valenzuela-Muñoz et al. (2017)
<i>S. salar</i>	<i>L. salmonis</i>	Increase of Th1, Th17 and Th2 pathways	8 dpi	Holm et al. (2017)
<i>S. salar</i>	<i>L. salmonis</i>	Increase expression of IL-1b, TNF α , MHII, TGF β -like cytokine and COX-2	9 dpi	Fast et al. (2006)
<i>S. salar</i>	<i>L. salmonis</i>	Adaptative Humoral Immune Response	5-10 dpi	Tadiso et al. (2011)
<i>S. salar</i>	<i>L. salmonis</i>	Up-regulation of alternatively activated macrophages	3-33 dpi	Skugor et al. (2008)
<i>S. salar</i>	<i>L. salmonis</i>	Pro-inflammatory response (Th2-like pathway)	3-33 dpi	Skugor et al. (2008)
<i>S. salar</i>	<i>L. salmonis</i>	Increase expression of the MH class I gene	21 dpi	Fast et al. (2006)
<i>S. salar</i>	<i>L. salmonis</i>	Down-regulation of alternatively activated macrophages	22 dpi	Skugor et al. (2008)
<i>S. salar</i>	<i>L. salmonis</i>	Increase expression of MH class II and TGF β -like genes	33 dpi	Fast et al. (2006)
<i>S. salar</i>	<i>L. salmonis</i>	Increase expression of Interleukin-1b and TNF α -like genes	40 dpi	Fast et al. (2006)
<i>S. salar</i>	<i>L. salmonis</i>	Th2 immune response	3 weeks dpi	Holm et al. (2015)
<i>S. salar</i>	<i>C. rogercresseyi</i>	Early responses of TLR/IMD signaling pathway	3 dpi	Vera-Bizama et al. (2015)
<i>S. salar</i>	<i>C. rogercresseyi</i>	Oxidative stress	7 dpi	Vera-Bizama et al. (2015)
<i>S. salar</i>	<i>C. rogercresseyi</i>	TLR pathway (up-regulation of TLR22 and MHCI)	14 dpi	Valenzuela-Muñoz et al. (2016)
<i>S. salar</i>	<i>C. rogercresseyi</i>	Inflammatory and oxidative stress response (Th1 pathway)	14 dpi	Boltaña et al. (2016)
<i>S. salar</i>	<i>C. rogercresseyi</i>	Up-regulation of Heme degradation and iron transport	7-14 dpi	Valenzuela-Muñoz et al. (2017)
<i>S. salar</i>	<i>C. rogercresseyi</i>	Up-regulation of various innate immune and detoxication genes	15 dpi - in-feed additives	Núñez-Acuña et al. (2016)

products such as serpins and trypsin [93] have been described. In addition, variations in the response of *C. rogercresseyi* infesting Atlantic and Coho salmon have been observed [94]. Here, both fish species increase the transcription activity of genes associated with oxidative stress, suggesting that lice respond to a large number of reactive oxygen species (ROS) generated by infected fish. Notably, high expression of antioxidant genes in lice has been correlated with low susceptibility to *L. salmonis* infestation [69]. In addition, the modulation of ROS-related genes in fish has been linked to increased expression of cytokines [11] and metalloproteases [68] released during an infestation. In parallel, *C. rogercresseyi* infecting Atlantic salmon showed a significant increase in the expression of ferritin, suggesting a compensatory response against the ROS that in turn, can be controlled by the Fenton reaction catalyzed by ferritin [95]. It is important to note that ferritin is also highly expressed in Atlantic salmon during the infestation, so it can be suggested that both the host and the ectoparasite compete for free iron [94]. However, the iron competition will differ from the host and the parasite proposes, while the molecular mechanisms are common and highly conserved.

During the infestation process, the ectoparasites feed on the host's blood by penetrating the skin, and in parallel, release molecules such as proteases and immunosuppressive molecules. The excretory/secretory proteins (SEPs) of the pathogens change the environment of the host cell by modulation of the host's immune system, facilitating the infection and its proliferation [96]. SEPs participate in the parasite-host interaction, modulating the timing related to the life-stages development [97]. Among the main components of parasite, secretions are proteolytic enzymes including cathepsin L, cathepsin B, serine proteases (serpins) and carboxypeptidases or trypsin-like [98].

Serine proteases inhibitors (serpins) regulate the innate immunity of insects through the inhibition of cascades of serine proteinases that initiate immune responses such as melanization and the production of antimicrobial peptides [99,100]. The serpins are a superfamily of proteins widely distributed with a SERPIN domain and participate in several immune responses. From the transcriptomic database of *C. rogercresseyi* [7] several isoforms have been identified. Among them, Cr-serpin 10 is outstanding, expressed during the different stages of ectoparasite development [93] (Table 2). In one study, high levels of expression of Cr-serpin 10 and Cr-serpin 3 were observed in copepodids, suggesting a role of these serpins in the evasion process of the immune response of *C. rogercresseyi* [93]. Other proteins associated with the secretome of sea lice are the cathepsins. In *L. salmonis* the cathepsin-L has been observed within the secretory/excretory products, indicating a putative role of these proteases in immune-evasion and parasite survivor [101]. The function of these proteins appears to be essential in a variety of important parasite biological processes, such as molting, cuticle remodeling, embryogenesis, and feeding, making them an attractive target for vaccine development [102]. There are different groups of cathepsins, the cathepsins of the L group being the most abundant in arthropods [103]. In *L. salmonis*, three highly expressed cathepsins have been identified during the infective stage, including cathepsin L, suggesting a putative role for feeding and immune defense [104]. In *C. rogercresseyi*, a comprehensive study has identified 56 cathepsin-like sequences distributed in five groups of cysteine proteases (B, F, L, Z, and S) and one for the aspartic protease group (D) [105]. As in *L. salmonis*, cathepsin L is one of the most abundant cathepsins in *C. rogercresseyi* [105]. In *C. rogercresseyi*, high levels of Cr-CatL1 expression have been reported in females, associating their function with

Table 2
Studies associate to transcriptome analysis of sea lice secretome.

Host species	Sea lice species	Molecular mechanisms	Experimental design	Reference
<i>S. salar</i>	<i>C. rogerresseyi</i>	Prostaglandin Enolase Cathepsin D1, B1 PhoC Trypsin 2, 5, 11, 12, 13, 17 Serpins 3, 4, 10 PTGSE Enolase Cathepsin D1, B1 PhoC Tryp 2, 5, 11, 12, 13, 17 Serpins 3, 4, 10 Serpins 1, 2, 3, 4, 5, 6, 7, 8, 10, 11	3 and 7 dpi	Vera-Bizama et al. (2015)
<i>O. kisutch</i>	<i>C. rogerresseyi</i>		3 and dpi	Vera-Bizama et al. (2015)
<i>S. salar</i>	<i>C. rogerresseyi</i>		Ontogenic development	Maldonado-Aguayo and Gallardo-Escárate (2014)
<i>S. salar</i>	<i>C. rogerresseyi</i>	chymotrypsin 1, trypsin 1, 6, 10, 36, 41, 42	Delousing drugs treatments	Valenzuela-Miranda and Gallardo-Escárate (2016)
<i>S. salar</i>	<i>L. salmonis</i>	serine-type endopeptidases, 2 cysteine type endopeptidases, 1 phosphatase, 1 acetylcholine receptor	Collection of E/S material from adult stages of the parasite	Hamilton et al. (2018)
<i>S. salar</i>	<i>L. salmonis</i>	secretory products	E/S products evaluation from primary culture of salmon	Lewis et al. (2014)
<i>S. salar</i>	<i>L. salmonis</i>	secretory/excretory products	SHK-1 exposition	Fast et al. (2007)
<i>O. kisutch</i> , <i>S. salar</i> , <i>O. nerka</i>	<i>L. salmonis</i>	Trypsin 1, trypsin-like serine protease, collagenase, carboxypeptidase B, Chymotrypsin B1, Chymotrypsin A chain C	24 and 48 h post-infection	Braden et al. (2017)
<i>S. salar</i>	<i>C. rogerresseyi</i>	Serine proteases inhibitor, alpha 1 antiproteinase-like, antithrombin protein, heat shock protein 47	15 dpi	Núñez-Acuña et al. (2015)
<i>S. salar</i>	<i>C. rogerresseyi</i>	Cathepsin B1 Cathepsin D1, D2 Cathepsin L1, L26, L27 Cathepsin Z10	Distribution and abundance of cathepsin-like transcript identified in sea lice developmental stages.	Maldonado-Aguayo et al. (2015)
<i>Salmon</i>	<i>L. salmonis</i>	Cathepsin L1 Cathepsin L2 Cathepsin B cysteine protease Proteases and alkaline phosphatase	Library construction of various life stages (chalarimus I–IV, preadult male I–II, preadult female I–II, adult male, adult female and egg strings) Adult <i>L. salmonis</i> were collected from infected salmon found Adult and preadult sea lice	McCarthy et al. (2012)
<i>Salmon</i>	<i>O. mykiss</i> , <i>O. kisutch</i> , <i>S. salar</i> , <i>P. americanus</i>			Cunningham et al. (2010) Fast et al. (2003)

louse embryonic development. In addition, Cr-CatL1, Cr-CatL26, and Cr-CatL27, together with Cr-CatB1, are highly expressed during the larval stages, nauplius, suggesting their possible participation in the processes of molting or cuticle remodeling [105].

Among the proteins of greater abundance within the SEPs, trypsins are found [106]. Trypsin is a protease stored as zymogen, which is activated during feeding, being deposited in the saliva through autolysis or the action of proteases provided by the host [107]. In *L. salmonis*, LsTryp10 has been identified and characterized. This trypsin is distributed uniformly in the ovaries and oocytes of the ectoparasite, suggesting the regulation of the degradome of the yolk [108] (Table 2). In *C. rogercresseyi*, 44 trypsin and seven chymotrypsin were identified [109]. Notably, these proteins seem to be also modulated in adults of *C. rogercresseyi* exposed to delousing drugs [109]. Here, the authors report sex-specific modulation of Cr-Tryp1-6-10-36 and 41 transcripts differentially expressed in response to pesticides, suggesting a putative role in drug metabolism.

8. Role of non-coding transcripts during *C. rogercresseyi* infestation

The increasing availability of transcriptome data has revealed the importance of the non-coding RNAs as key regulators of the mRNA transcription. Herein, microRNAs (miRNAs) and long non-coding RNAs (lncRNAs) have been studied [110,111]. The lncRNAs are long sequences greater than 200 nucleotides, transcribed in a similar way to the coding RNAs [112,113]. The study of lncRNAs is relatively new. Genomic studies have shown that in mammals as well as other vertebrates the number of RNAs that are translated into protein is close to 20,000 and due to differences in the genomic sizes of organisms, it has been suggested that numbers of lncRNAs have been increasing during the evolutionary process, so that organisms with more complex mechanisms have acquired more lncRNAs, as a mechanism to control several biological processes [114]. lncRNAs studies report that there is a proportion that is expressed in a cell-specific manner, and even in cells of the immune system [114,115]. For instance, in zebrafish 342 lncRNAs are transcribed in more than one tissue and only 77 lncRNAs are expressed specifically in a single tissue [116]. Another example is in rainbow trout where it has been observed that of 2935 lncRNAs, about 5.2% of lncRNAs identified in this species, are expressed in a specific tissue manner [117]. In Atlantic and Coho salmon lncRNAs were identified from the skin and head-kidney tissues, observing a greater abundance of exclusive lncRNAs in the head-kidney for both fish species [23]. In addition, exclusive lncRNAs were identified in the skin, which is similar to that reported for rainbow trout [117].

lncRNAs play important roles in the modulation of biological processes, being involving in regulatory processes of the immune response associated with Toll [118] and NF- κ B [119] signaling pathways. In fish, the role of lncRNAs as regulators of the immune response has been poorly studied. However, some studies conducted in *Larimichthys polyactis* after exposure to *Vibrio anguillarum*, reveal a high number of lncRNAs upregulated in response to infection. Here, the authors suggest that the immune system can be modulated through lncRNAs closely localized to immune-related genes [120]. Another transcriptome study conducted in Atlantic salmon infected with the ISA virus identified 4967 lncRNAs differentially expressed and positively correlated with the viral burden [121]. lncRNAs have also been reported in salmon infected with the pathogenic bacteria *P. salmonis*, showing that the lncRNAs are modulated during infection. Here, lncRNAs were directly correlated with genes involved in the response to this bacterium, such as endocytosis and iron homeostasis [122]. Furthermore, a comparative study in Atlantic salmon exposed to virus, bacteria, and ectoparasites demonstrated that lncRNAs are specifically regulated in response to each type of infection [123]. With respect to the modulation of lncRNAs in Atlantic and Coho salmon exposed to *C. rogercresseyi*, it was observed that the abundance of lncRNAs expressed during the time of infestation

is similar to that observed for coding transcripts. In both salmon species, a high number of lncRNAs were expressed at 7 dpi and 14 dpi in skin and head-kidney, respectively [23]. The lncRNAs expressed in Coho salmon infested with *C. rogercresseyi* were closely located to genes associated with the response to external stimuli, tissue repair, and cell communication [23]. On the other hand, the lncRNAs identified in Atlantic salmon expressed during the infestation by *C. rogercresseyi* were associated with biological processes such as stress, immune and inflammatory response [23]. Additionally, 439 lncRNAs expressed in the head-kidney of Atlantic salmon after 15 days post-infestation with *C. rogercresseyi* were correlated with genes associated with transcriptional regulation processes [123]. From the identification and expression analysis of the lncRNAs in salmon, it is possible to establish that the lncRNAs play an important role during the lice infestation. However, it is necessary to design and conduct functional studies, uncovering the molecular mechanisms that drive the fish immune response.

Another class of ncRNAs is the microRNAs (miRNAs), which are small RNA molecules of ~22 nucleotides that bind to a messenger RNA, mainly at its 3' end UTR and block the gene translation [124,125]. Like the lncRNAs, miRNAs are expressed in a tissue-specific manner and are capable of regulating biological processes such as development, growth, metabolism, apoptosis, etc. [126–128]. In regard to the immune response, miRNAs participate in the regulation of both innate and adaptive immune responses, as well as differentiators of T cells, modulators of inflammatory response and activation of Toll-like receptors [129–132]. In Atlantic salmon, miRNAs have been identified and differentially expressed in response to lice infection [133]. Here, 1718 miRNAs from skin and head-kidney were annotated, while miRNA families named as mir-21, mir-181 and let-7 were the most enriched [133]. These families have been found to be abundant in other species of fish such as trout and flatfish [134–136]. Due to the role of miRNAs to interfere the transcriptional activity [124,125], it can be inferred that the abundance of miRNAs post-lice infestation in key tissue as head kidney and skin can modulate genes related to the immune response in Atlantic salmon. Among the highly expressed miRNAs in Atlantic salmon, mir-140 was highly expressed in the head-kidney at 7 dpi, which have as target gene the toll-like receptor *tlr22a2*, which is also negatively regulated. Another miRNA that seems to influence the response of Atlantic salmon to *C. rogercresseyi* is mir-21, which targets the C-C Motif Chemokine Receptor 3 gene (CCR3) associated with Th2 response. This receptor is highly relevant during the response of salmon to *L. salmonis* [22]. Additionally, and associated with the regulation of iron, mir-181a-2-5 was also identified with a putative regulation on ALAs. This gene has been associated with the biosynthesis of the heme group, a process which was also found to be inhibited in Atlantic salmon infested with *C. rogercresseyi*.

There are studies that show that miRNAs are not only regulators of biological processes, but also that they can participate in the parasite-host interaction processes. In insects affected by viruses, it has been observed that viruses are capable of releasing miRNAs that can regulate the expression of their host genes in order to successfully establish the infection [137]. For *C. rogercresseyi*, several miRNAs expressed during the different stages of development have been characterized [138,139]. Within the profile of miRNAs characterized in *C. rogercresseyi*, the miRNA annotated as Bantam is highly expressed in the infective stage of copepodid [138]. This suggests that Bantam has a key role in the success of the infection. Predictions of Bantam target genes have shown that this miRNA could have an important role in the parasite-host interaction process by regulating genes associated with the secretome of *C. rogercresseyi* such as trypsin and cathepsin [138]. Notably, Bantam seems to be a putative regulator of genes associated with the immune response of Atlantic salmon such as *trl22a2*, *cd83* and *IFN γ* [138].

9. Host-parasite microbiome modulation in sea lice infection

The microbiota community of an organism actively participates in vital functions with a leading role. It is associated mainly with the homeostatic, nutritional, metabolic and defensive performance [140–142]. In fish, mucosa associated microbiota have been studied thoroughly [140,141,143] and host-specific signatures have been found, especially in the skin [143]. However, several factors can disrupt mucosal microbiome stability, such as stress, water quality, and diet [144]. Since commensal microbiota constitute a resistance barrier to opportunistic pathogens colonization [145], if dysbiosis occurs, diseases can easily emerge. Sea lice impact fish health in several ways, and are a strong stressor regardless of infestation pressure [146,147], changing mucus production and composition [148,149]. However, the impact of sea lice infection in the fish epidermal mucosal microbiome has only recently been investigated [150]. Atlantic salmon's skin microbial community changed identity as infection with *L. salmonis* progressed, and the richness and beta-diversity of this community were also impacted [150]. During infection, an increase in Rhizobiales and NS10_marine_group, a member of Cryomorphaeae family, was observed together with a decrease in *Arthrobacter* genus. The biological significance of these changes is yet to be uncovered, but it indicates the modulatory effect of lice in the fish skin microbiome.

In copepods, mucosal microbiome research is not developed as in fish, but it is known that microorganism communities are more abundant in the mouth, digestive tract and egg sacs [151], and it is possible to observe a central or “core” natural microbiome that is stable, as well as host-specific associations, symbiosis, and other associations that are environment and diet-dependent [152]. In the case of *C. rogercresseyi*, diet is mainly composed by fish mucus, skin and muscle and this might influence the parasite's microbiome, especially if the host is suffering a mucosal dysbiosis. Minniti et al. [153] found that 24 h after handling, a significant effect on Atlantic salmon's microbial composition is observed, with the rising of the genus *Burkholderia* – a potentially pathogenic taxon for several animal species – whereas the abundance of *Methylobacterium* affiliated phylotypes – a poly-B-hydroxybutyrate producing beneficial bacteria - decreased. Data on the interaction on host mucus-lice microbiota is rather scarce, although Llewellyn et al. [150] found some similarities reflected mainly by the abundance of the genera *Vibrio* and NS10_marine_group. More research is necessary to properly infer interactions and host-parasite microbiome associations, and this takes special relevance in parasitic copepods. It is known that in copepods the interaction with bacteria includes permanent and transient endo and epibiotic associations that might have a significant role in copepod's health [152]. In the case of parasitic copepods such as *C. rogercresseyi*, it might have an effect on their infectious capacity and resistance and be directly related to disease outbreaks.

In addition, sea lice associated microbiota can play a role as a reservoir of secondary pathogens as seen in several studies. Sepulveda et al. [154] found that *Caligus lalandei* – an ectoparasite common in *Seriola lalandi* – cultured microbiota has a high abundance of *Vibrio alginolyticus*, a known fish pathogen, whereas Nese and Enger [155] and Novak et al. [156] reported that *L. salmonis* is a potential vector for *Aeromonas salmonicida*. The potential role of *C. rogercresseyi* as a vector for relevant pathogens such as *P. salmonis* is presently under debate, but so far it has been only reported that it is a reservoir of ISA (Infectious Salmon anaemia) virus and can act as a vector for spreading this disease [13]. Research on salmon-lice microbiome associations and interactions is underway for *C. rogercresseyi*, and information that is being generated will allow for a better understanding of possible modulations for better control of Caligidosis. Moreover, considering the impact of *C. rogercresseyi* infestations in aquaculture in Chile, the constant usage of delousing drugs and increasing observed resistance in lice the study of the effect of microbiome variations on drug disposition, action, and toxicity (pharmacomicrobiomics) will be an important field going forward.

10. Conclusion and perspectives

The understanding of how the parasite survives in its host and how they display molecular mechanisms to face with the immune system. The detection of semiochemicals has been associated to the ionotropic receptors (IRs) gene family, involving key molecular components for the salmon-lice interaction. For instance, a relevant finding is the parasite's ability to detect the antimicrobial peptide Cath-2 expressed from infected-fish, allowing the subsequent infection process. Moreover, the inflammatory response is the first reaction associated with the coping against sea lice, generating strong modulation of immune-related genes and signaling pathways (Table 1). From the recognition process through infestation and the development of different life stages, lice must cope with the rejection mechanisms displayed by the fish. Innate immune response pathways IMD and TLR, antioxidant system and secretory products are pivotal molecular components utilized by lice species to infect. The increasing availability of transcriptome data has revealed the importance of the non-coding RNAs as key regulators of the mRNA transcription. Notably, salmon-lice interactions are significantly modulated by lncRNAs and miRNAs during the infestation process. The understanding of how these molecules interact with pivotal signaling pathways involved in the immune response can be applied to develop novel control strategies.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgments

This study was funded by projects FONDAP #15110027 and FONDECYT #1180867 awarded by CONICYT-Chile.

References

- [1] R. Khan, Host-parasite interactions in some fish species, *J. Parasitol. Res.* 2012 (2012).
- [2] G. Froeschke, S. von der Heyden, A review of molecular approaches for investigating patterns of coevolution in marine host-parasite relationships, *Adv. Parasitol.* 84 (2014) 209–252.
- [3] D.G. Biron, H.D. Loxdale, Host-parasite molecular cross-talk during the manipulative process of a host by its parasite, *J. Exp. Biol.* 216 (2013) 148–160.
- [4] P. Willadsen, Vaccination against ectoparasites, *Parasitology* 133 (2006) S9–S25.
- [5] D. Lewis, D. Barker, R. McKinley, Modulation of cellular innate immunity by *Lepeophtheirus salmonis* secretory products, *Fish Shellfish Immunol.* 38 (2014) 175–183.
- [6] Y.-J. Choi, M.T. Aliota, G.F. Mayhew, S.M. Erickson, B.M. Christensen, Dual RNA-Seq of Parasite and Host Reveals Gene Expression Dynamics during Filarial Worm-Mosquito Interactions, (2014).
- [7] C. Gallardo-Escárate, V. Valenzuela-Muñoz, G. Núñez-Acuña, RNA-Seq analysis using *de novo* transcriptome assembly as a reference for the salmon louse *Caligus rogercresseyi*, *PLoS One* 9 (2014) e92239.
- [8] S.C. Johnson, M.D. Fast, G. Wiegertjes, G. Flik, Interactions between sea lice and their hosts, *Host – Parasite Interactions* (2004) 131–159.
- [9] M.D. Fast, Fish immune responses to parasitic copepod (namely sea lice) infection, *Dev. Comp. Immunol.* 43 (2014) 300–312.
- [10] M.J. Costello, Ecology of sea lice parasitic on farmed and wild fish, *Trends Parasitol.* 22 (2006) 475–483.
- [11] C. Hamilton-West, G. Arriagada, T. Yatabe, P. Valdés, L.P. Hervé-Claude, S. Urcelay, Epidemiological description of the sea lice (*Caligus rogercresseyi*) situation in southern Chile in August 2007, *Prev. Vet. Med.* 104 (2012) 341–345.
- [12] S. Bravo, The reproductive output of sea lice *Caligus rogercresseyi* under controlled conditions, *Exp. Parasitol.* 125 (2010) 51–54.
- [13] K. Oelckers, S. Vike, H. Duesund, J. Gonzalez, S. Wadsworth, A. Nylund, *Caligus rogercresseyi* as a potential vector for transmission of Infectious Salmon Anaemia (ISA) virus in Chile, *Aquaculture* 420 (2014) 126–132.
- [14] A.M. Prince, F.H. Khan, B.F. Matthews, N. Islam, S.S. Natarajan, Transcriptome and proteome alterations during the nematode-soybean interaction, *Nematropica* 44 (2015) 137–145.
- [15] S. Soto, C. López, RNA-seq as a transcriptome useful tool for the studies of plant pathogen interactions, *Fitosanidad* 16 (2012) 101–113.
- [16] H.J. Lee, A. Georgiadou, T.D. Otto, M. Levin, L.J. Coin, D.J. Conway, et al., Transcriptomic studies of malaria: a paradigm for investigation of systemic host-pathogen interactions, *Microbiol. Mol. Biol. Rev.* 82 (2018) e00071-17.
- [17] E. Videvall, C.K. Cornwallis, D. Ahrén, V. Palinauskas, G. Valkiūnas, O. Hellgren,

- The transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific gene expression, *Mol. Ecol.* 26 (2017) 2939–2958.
- [18] M. Fast, D.E. Sims, J.F. Burka, A. Mustafa, N. Ross, Skin morphology and humoral non-specific defence parameters of mucus and plasma in rainbow trout, coho and Atlantic salmon, *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 132 (2002) 645–657.
- [19] M.D. Fast, N.W. Ross, A. Mustafa, D.E. Sims, S.C. Johnson, G.A. Conboy, et al., Susceptibility of rainbow trout *Oncorhynchus mykiss*, Atlantic salmon *Salmo salar* and coho salmon *Oncorhynchus kisutch* to experimental infection with sea lice *Lepeophtheirus salmonis*, *Dis. Aquat. Org.* 52 (2002).
- [20] T.M. Tadiso, A. Krasnov, S. Skugor, S. Afanasyev, I. Hordvik, F. Nilsen, Gene expression analyses of immune responses in Atlantic salmon during early stages of infection by salmon louse (*Lepeophtheirus salmonis*) revealed bi-phasic responses coinciding with the copepod-chalimus transition, *BMC Genomics* 12 (2011) 141.
- [21] V. Valenzuela-Muñoz, C. Gallardo-Escárate, Iron metabolism modulation in Atlantic salmon infested with the sea lice *Lepeophtheirus salmonis* and *Caligus rogercresseyi*: a matter of nutritional immunity? *Fish Shellfish Immunol.* 60 (2017) 97–102.
- [22] L.M. Braden, B.F. Koop, S.R. Jones, Signatures of resistance to *Lepeophtheirus salmonis* include a T H 2-type response at the louse-salmon interface, *Dev. Comp. Immunol.* 48 (2015) 178–191.
- [23] V. Valenzuela-Muñoz, D. Valenzuela-Miranda, C. Gallardo-Escárate, Comparative analysis of long non-coding RNAs in Atlantic and Coho salmon reveals divergent transcriptome responses associated with immunity and tissue repair during sea lice infestation, *Dev. Comp. Immunol.* 87 (2018) 36–50.
- [24] M.E. Hay, Marine chemical ecology: chemical signals and cues structure marine populations, communities, and ecosystems, *Ann. Rev. Mar. Sci.* 1 (2009) 193–212.
- [25] J. Lindsey, R. Lasker, Chemical signals in the sea: marine allelochemicals and evolution, *Fish. Bull.* 72 (1974) 1.
- [26] S. Steiger, T. Schmitt, H. Scafer, The origin and dynamic evolution of chemical information transfer, *Proc. Biol. Sci.* 278 (2011) 970–979.
- [27] E.L. Preisser, D.I. Bolnick, M.F. Benard, Scared to death? The effects of intimidation and consumption in predator–prey interactions, *Ecology* 86 (2005) 501–509.
- [28] J. Heuschele, E. Selander, The chemical ecology of copepods, *J. Plankton Res.* 2014 (2014) 1–19.
- [29] D.M. Fields, M.J. Weissburg, H.I. Browman, Chemoreception in the salmon louse *Lepeophtheirus salmonis*: an electrophysiological approach, *Dis. Aquat. Org.* 78 (2007) 161–168.
- [30] Y. Barrientos, Ultrastructure of Sensory Units on the First Antennae of Calanoid Copepods, MSci Thesis University of Ottawa, Canada, 1980.
- [31] L. Gonzalez, J. Carvajal, Life cycle of *Caligus rogercresseyi*, (Copepoda: caligidae) parasite of Chilean reared salmonids, *Aquaculture* 220 (2003) 101–117.
- [32] R.L. Genna, W. Mordue, A.W. Pike, A.J. Mordue, Light intensity, salinity, and host velocity influence presettlement intensity and distribution on hosts by copepodids of sea lice, *Lepeophtheirus salmonis*, *Can. J. Fish. Aquat. Sci.* 62 (2005) 2675–2682.
- [33] P.A. Heuch, M.H. Doall, J. Yen, Water flow around a fish mimic attracts a parasitic and deters a planktonic copepod, *J. Plankton Res.* 29 (2007) i3–i16.
- [34] R.J.E. Bailey, M.A. Birkett, A. Ingvarsdóttir, A.J. Mordue, W. Mordue, B. O'Shea, et al., The role of semiochemicals in host location and non-host avoidance by salmon louse (*Lepeophtheirus salmonis*) copepodids, *Can. J. Fish. Aquat. Sci.* 63 (2006) 448–456.
- [35] M.Q. Hull, A.W. Pike, A.J. Mordue, G.H. Rae, Patterns of pair formation and mating in an ectoparasitic caligid copepod *Lepeophtheirus salmonis* (Krøyer 1837): implications for its sensory and mating biology, *Phil. Trans. Roy. Soc. Lond. B Biol. Sci.* 353 (1998) 753–764.
- [36] A.J. Mordue Luntz, M.A. Birkett, A review of host finding behaviour in the parasitic sea louse, *Lepeophtheirus salmonis* (Caligidae: copepoda), *J. Fish Dis.* 32 (2009) 3–13.
- [37] A. Ingvarsdóttir, Birkett Ma, I. Duce, R.L. Genna, W. Mordue, Ja Pickett, et al., Semiochemical strategies for sea louse control: host location cues, *Pest Manag. Sci.* 58 (2002) 537–545.
- [38] J. Pino-Marambio, A.J. Mordue, M. Birkett, J. Carvajal, G. Asencio, A. Mellado, et al., Behavioural studies of host, non-host and mate location by the sea louse, *Caligus rogercresseyi* boxshall & bravo, 2000 (copepoda: caligidae), *Aquaculture* 271 (2007) 70–6.
- [39] G. Nuñez-Acuña, V. Valenzuela-Muñoz, C. Gallardo-Escárate, High-throughput SNP discovery and transcriptome expression profiles from the salmon louse *Caligus rogercresseyi* (Copepoda: caligidae), *Comp. Biochem. Physiol. Genom. Proteonom.* 10 (2014) 9–21.
- [40] G. Nuñez-Acuña, F. Vera-Bizama, S. Boltaña, C. Hawes, J.P. Marambio, S. Wadsworth, et al., In-feed additives modulate ionotropic receptor genes from the sea louse *Caligus rogercresseyi*: a comparative analysis in two host salmonid species, *Aquaculture* 451 (2016) 99–105.
- [41] R. Benton, K.S. Vannice, C. Gomez-Diaz, L.B. Voshall, Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*, *Cell* 136 (2009) 149–162.
- [42] A. Ingvarsdóttir, M.A. Birkett, I. Duce, W. Mordue, J.A. Pickett, L.J. Wadhams, et al., Role of semiochemicals in mate location by parasitic sea louse, *Lepeophtheirus salmonis*, *J. Chem. Ecol.* 28 (2002) 2107–2117.
- [43] W.S. Leal, Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes, *Annu. Rev. Entomol.* 58 (2013) 373–391.
- [44] F.G. Vieira, J. Rozas, Comparative genomics of the odorant-binding and chemosensory protein gene families across the arthropoda: origin and evolutionary history of the chemosensory system, *Genome Biol. Evol.* 3 (2011) 476–490.
- [45] A.Z. Komisarczuk, S. Grotmol, F. Nilsen, Ionotropic receptors signal host recognition in the salmon louse (*Lepeophtheirus salmonis*, Copepoda), *PLoS One* 12 (2017) e0178812.
- [46] G. Nuñez-Acuña, C. Gallardo-Escárate, A.B. Skiftesvik, D.M. Fields, H.I. Browman, Silencing of ionotropic receptor 25a decreases chemosensory activity in the salmon louse *Lepeophtheirus salmonis* during the infective stage, *Gene* 697 (2019) 35–39.
- [47] G. Nuñez-Acuña, A.T. Gonçalves, V. Valenzuela-Muñoz, J. Pino-Marambio, S. Wadsworth, C. Gallardo-Escárate, Transcriptome immunomodulation of in-feed additives in Atlantic salmon *Salmo salar* infested with sea lice *Caligus rogercresseyi*, *Fish Shellfish Immunol.* 47 (2015) 450–460.
- [48] E.J. Noga, A.J. Ullal, J. Corrales, J.M.O. Fernandes, Application of antimicrobial polypeptide host defenses to aquaculture: exploitation of downregulation and upregulation responses, *Comp. Biochem. Physiol. Genom. Proteonom.* 6 (2011) 44–54.
- [49] V. Rajanbabu, J.Y. Chen, Applications of antimicrobial peptides from fish and perspectives for the future, *Peptides* 32 (2011) 415–420.
- [50] J.A. Masso-Silva, G. Diamond, Antimicrobial peptides from fish, *Pharmaceuticals* 7 (2014) 265–310.
- [51] Y. Valero, E. Chaves-Pozo, J. Meseguer, M.A. Esteban, A. Cuesta, Biological role of fish antimicrobial peptides, *Antimicrobial Peptides* (2013) 31–60.
- [52] E. Casadei, S. Bird, S. Wadsworth, J.L. González Vecino, C.J. Secombes, The longevity of the antimicrobial response in rainbow trout (*Oncorhynchus mykiss*) fed a peptidoglycan (PG) supplemented diet, *Fish Shellfish Immunol.* 44 (2015) 316–320.
- [53] X.-J. Zhang, X.-Y. Zhang, N. Zhang, X. Guo, K.-S. Peng, H. Wu, et al., Distinctive structural hallmarks and biological activities of the multiple cathelicidin antimicrobial peptides in a primitive teleost fish, *J. Immunol.* 194 (2015) 4974–4987.
- [54] A. Bridle, E. Nosworthy, M. Polinski, B. Nowak, Evidence of an Antimicrobial-Immunomodulatory Role of Atlantic Salmon Cathelicidins during Infection with *Yersinia ruckeri* vol. 6, (2011) e23417.
- [55] A. Krasnov, S. Skugor, M. Todorovic, K.A. Glover, F. Nilsen, Gene expression in Atlantic salmon skin in response to infection with the parasitic copepod *Lepeophtheirus salmonis*, cortisol implant, and their combination, *BMC Genomics* 13 (2012).
- [56] A. Colorni, A. Ullal, G. Heinisch, E.J. Noga, Activity of the antimicrobial polypeptide piscidin 2 against fish ectoparasites, *J. Fish Dis.* 31 (2008) 423–432.
- [57] B.J. Sutherland, K.W. Koczka, M. Yasuike, S.G. Jantzen, R. Yazawa, B.F. Koop, et al., Comparative transcriptomics of Atlantic *Salmo salar*, chum *Oncorhynchus keta* and pink salmon *O. gorbuscha* during infections with salmon lice *Lepeophtheirus salmonis*: a review of the impact of parasitic copepods on marine aquaculture, *BMC Genomics* 15 (2014) 1–17.
- [58] G. Nuñez-Acuña, J.P. Marambio, T. Valenzuela, S. Wadsworth, C. Gallardo-Escárate, Antimicrobial peptides from *Salmo salar* skin induce frontal filament development and olfactory/cuticle-related genes in the sea louse *Caligus rogercresseyi*, *Aquaculture* 464 (2016) 171–177.
- [59] A. Pike, S. Wadsworth, Sealice on salmonids: their biology and control, *Adv. Parasitol.* (1999) 233–337 Elsevier.
- [60] G. Nuñez-Acuña, C. Gallardo-Escárate, D.M. Fields, S. Shema, A.B. Skiftesvik, I. Ormazábal, et al., The Atlantic salmon (*Salmo salar*) antimicrobial peptide cathelicidin-2 is a molecular host-associated cue for the salmon louse (*Lepeophtheirus salmonis*), *Sci. Rep.* 8 (2018) 13738.
- [61] S.I. Dodson, T. Hanazato, P.R. Gorski, Behavioral responses of *Daphnia pulex* exposed to carbaryl and *Chaoborus* kairomone, *Environ. Toxicol. Chem.* 14 (1995) 43–50.
- [62] I. Eyres, L. Duvaux, K. Gharbi, R. Tucker, D. Hopkins, J.C. Simon, et al., Targeted re-sequencing confirms the importance of chemosensory genes in aphid host race differentiation, *Mol. Ecol.* 26 (2017) 43–58.
- [63] H. Miyakawa, M. Imai, N. Sugimoto, Y. Ishikawa, A. Ishikawa, H. Ishigaki, et al., Gene up-regulation in response to predator kairomones in the water flea, *Daphnia pulex*, *BMC Dev. Biol.* 10 (2010) 45.
- [64] A. Schwarzenberger, C. Courts, E. von Elert, Target gene approaches: gene expression in *Daphnia magna* exposed to predator-borne kairomones or to microcystin-producing and microcystin-free *Microcystis aeruginosa*, *BMC Genomics* 10 (2009) 527.
- [65] P. Alvarez-Pellitero, Fish immunity and parasite infections: from innate immunity to immunoprophylactic prospects, *Vet. Immunol. Immunopathol.* 126 (2008) 171–198.
- [66] S. Skugor, K.A. Glover, F. Nilsen, A. Krasnov, Local and systemic gene expression responses of Atlantic salmon (*Salmo salar* L.) to infection with the salmon louse (*Lepeophtheirus salmonis*), *BMC Genomics* 9 (2008) 498.
- [67] H. Holm, N. Santi, S. Kjøglum, N. Perisic, S. Skugor, Ø. Evensen, Difference in skin immune responses to infection with salmon louse (*Lepeophtheirus salmonis*) in Atlantic salmon (*Salmo salar* L.) of families selected for resistance and susceptibility, *Fish Shellfish Immunol.* 42 (2015) 384–394.
- [68] L.M. Braden, D.E. Barker, B.F. Koop, S.R. Jones, Comparative defense-associated responses in salmon skin elicited by the ectoparasite *Lepeophtheirus salmonis*, *Comp. Biochem. Physiol. Genom. Proteonom.* 7 (2012) 100–109.
- [69] B.J. Sutherland, K.W. Koczka, M. Yasuike, S.G. Jantzen, R. Yazawa, B.F. Koop, et al., Comparative transcriptomics of Atlantic *Salmo salar*, chum *Oncorhynchus keta* and pink salmon *O. gorbuscha* during infections with salmon lice *Lepeophtheirus salmonis*, *BMC Genomics* 15 (2014) 200.
- [70] P.R. Rauta, M. Samanta, H.R. Dash, B. Nayak, S. Das, Toll-like receptors (TLRs) in aquatic animals: signaling pathways, expressions and immune responses, *Immunol. Lett.* 158 (2014) 14–24.
- [71] R.P. Panda, V. Chakrapani, S.K. Patra, J.N. Saha, P. Jayasankar, B. Kar, et al., First evidence of comparative responses of Toll-like receptor 22 (TLR22) to relatively resistant and susceptible Indian farmed carps to *Argulus siamensis* infection, *Dev. Comp. Immunol.* 47 (2014) 25–35.

- [72] B. Kar, J. Mohanty, K. Hemaprasanth, P.K. Sahoo, The immune response in rohu, *Labeo rohita* (Actinopterygii: cyprinidae) to *Argulus siamensis* (Branchiura: argulidae) infection: kinetics of immune gene expression and innate immune response, *Aquacult. Res.* 46 (2015) 1292–1308.
- [73] S. Saurabh, B. Mohanty, P. Sahoo, Expression of immune-related genes in rohu *Labeo rohita* (Hamilton) by experimental freshwater lice *Argulus siamensis* (Wilson) infection, *Vet. Parasitol.* 175 (2011) 119–128.
- [74] S.A. Quiniou, P. Boudinot, E. Bengtén, Comprehensive survey and genomic characterization of Toll-like receptors (TLRs) in channel catfish, *Ictalurus punctatus*: identification of novel fish TLRs, *Immunogenetics* 65 (2013) 511–530.
- [75] V. Valenzuela-Muñoz, S. Boltaña, C. Gallardo-Escárate, Comparative immunity of *Salmo salar* and *Oncorhynchus kisutch* during infestation with the sea louse *Caligus rogercresseyi*: an enrichment transcriptome analysis, *Fish Shellfish Immunol.* 59 (2016) 276–287.
- [76] R. Morrison, E. Koppang, I. Hordvik, B. Nowak, MHC class II+ cells in the gills of Atlantic salmon (*Salmo salar* L.) affected by amoebic gill disease, *Vet. Immunol. Immunopathol.* 109 (2006) 297–303.
- [77] H.J. Holm, S. Skugor, A.K. Bjelland, S. Radunovic, S. Wadsworth, E.O. Koppang, et al., Contrasting expression of immune genes in scaled and scaleless skin of Atlantic salmon infected with young stages of *Lepeophtheirus salmonis*, *Dev. Comp. Immunol.* 67 (2017) 153–165.
- [78] M. Fast, D. Muise, R. Easy, N. Ross, S. Johnson, The effects of *Lepeophtheirus salmonis* infections on the stress response and immunological status of Atlantic salmon (*Salmo salar*), *Fish Shellfish Immunol.* 21 (2006) 228–241.
- [79] J.E. Allen, R.M. Maizels, Diversity and dialogue in immunity to helminths, *Nat. Rev. Immunol.* 11 (2011) 375–388.
- [80] G. Mulcahy, S. O'Neill, J. Fanning, E. McCarthy, M. Sekiya, Tissue migration by parasitic helminths—an immunoevasive strategy? *Trends Parasitol.* 21 (2005) 273–277.
- [81] M.I. Hood, E.P. Skaar, Nutritional immunity: transition metals at the pathogen–host interface, *Nat. Rev. Microbiol.* 10 (2012) 525–537.
- [82] B. Tschirren, L.L. Bischoff, V. Saladin, H. Richner, Host condition and host immunity affect parasite fitness in a bird–ectoparasite system, *Funct. Ecol.* 21 (2007) 372–378.
- [83] J.E. Cassat, E.P. Skaar, Iron in infection and immunity, *Cell Host Microbe* 13 (2013) 509–519.
- [84] G. Weiss, Modification of iron regulation by the inflammatory response, *Best Pract. Res. Clin. Haematol.* 18 (2005) 183–201.
- [85] V. Valenzuela-Muñoz, S. Boltaña, C. Gallardo-Escárate, Uncovering iron regulation with species-specific transcriptome patterns in Atlantic and coho salmon during a *Caligus rogercresseyi* infestation, *J. Fish Dis.* 40 (2017) 1169–1184.
- [86] E. Nemeth, S. Rivera, V. Gabayan, C. Keller, S. Taudorf, B.K. Pedersen, et al., IL-6 mediates hypoferremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin, *J. Clin. Invest.* 113 (2004) 1271–1276.
- [87] M.P. González, J.L. Muñoz, V. Valerio, L. Vargas-Chacoff, Effects of the ectoparasite *Caligus rogercresseyi* on *Salmo salar* blood parameters under farm conditions, *Aquaculture* 457 (2016) 29–34.
- [88] X. Hu, A.C. Camus, S. Aono, E.E. Morrison, J. Dennis, K.E. Nusbaum, et al., Channel catfish hepcidin expression in infection and anemia, *Comp. Immunol. Microbiol. Infect. Dis.* 30 (2007) 55–69.
- [89] P.G. Fraenkel, D. Traver, A. Donovan, D. Zahrieh, L.I. Zon, Ferroportin1 is required for normal iron cycling in zebrafish, *J. Clin. Invest.* 115 (2005) 1532–1541.
- [90] S. Toh, A. Glanfield, G. Gobert, M. Jones, Heme and Blood-Feeding Parasites: Friends or Foes? *Parasites and Vectors* vol. 3, (2010).
- [91] V. Valenzuela-Muñoz, C. Gallardo-Escárate, TLR and IMD signaling pathways from *Caligus rogercresseyi* (Crustacea: copepoda): *In silico* gene expression and SNPs discovery, *Fish Shellfish Immunol.* 36 (2014) 428–434.
- [92] J. Chavez-Mardones, C. Gallardo-Escárate, Deltamethrin (AlphaMax™) reveals modulation of genes related to oxidative stress in the ectoparasite *Caligus rogercresseyi*: implications on delousing drug effectiveness, *Aquaculture* 433 (2014) 421–429.
- [93] W. Maldonado-Aguayo, C. Gallardo-Escárate, Increasing transcriptome response or SERPINS during the ontogenetic stages in the salmon louse *Caligus rogercresseyi* (Copepoda: caligidae), *Marine Genomics* 15 (2014) 55–64, <https://doi.org/10.1016/j.margen.2014.04.006>.
- [94] F. Vera-Bizama, V. Valenzuela-Muñoz, A.T. Gonçalves, J.P. Marambio, C. Hawes, S. Wadsworth, et al., Transcription expression of immune-related genes from *Caligus rogercresseyi* evidences host-dependent patterns on Atlantic and coho salmon, *Fish Shellfish Immunol.* 47 (2015) 725–731.
- [95] P.M. Harrison, P. Arosio, The ferritins: molecular properties, iron storage function and cellular regulation, *Biochim. Biophys. Acta Bioenerg.* 1275 (1996) 161–203.
- [96] S. Ranganathan, G.J.G. Garg, Secretome: clues into pathogen infection and clinical applications, 1 (2009) 113.
- [97] Y. Moreno, T.G. Geary, Stage- and gender-specific proteomic analysis of *Brugia malayi* excretory-secretory products, *PLoS Neglected Trop. Dis.* 2 (2008) e326.
- [98] M.W. Robinson, R. Menon, S.M. Donnelly, J.P. Dalton, S.J.M. Ranganathan, C. Proteomics, An Integrated Transcriptomics and Proteomics Analysis of the Secretome of the Helminth Pathogen Fasciola Hepatica Proteins Associated with Invasion and Infection of the Mammalian Host vol. 8, (2009), pp. 1891–1907.
- [99] E. Clynen, L. Schoofs, M. Salzet, A review of the most important classes of serine protease inhibitors in insects and leeches, *Med. Chem. Rev. Online* 2 (2005) 197–206.
- [100] D.A. Meekins, M.R. Kanost, K. Michel, Serpins in arthropod biology, *Semin. Cell Dev. Biol.* (2017) 105–119 Elsevier.
- [101] E. McCarthy, E. Cunningham, L. Copley, D. Jackson, D. Johnston, J.P. Dalton, et al., Cathepsin L proteases of the parasitic copepod, *Lepeophtheirus salmonis*. *Aquaculture* 356 (2012) 264–271.
- [102] L. Ford, J. Zhang, J. Liu, S. Hashmi, J.A. Fuhrman, Y. Oksov, et al., Functional analysis of the cathepsin-like cysteine protease genes in adult *Brugia malayi* using RNA interference, *PLoS Neglected Trop. Dis.* 3 (2009) e377.
- [103] D. Sojka, I.M. Francischetti, E. Calvo, M. Kotsyfakis, Cysteine Proteases from Bloodfeeding Arthropod Ectoparasites. Cysteine Proteases of Pathogenic Organisms, Springer, 2011, pp. 177–191.
- [104] S. Hamilton, K. McLean, S.J. Monaghan, C. McNair, N.F. Inglis, H. McDonald, et al., Characterisation of proteins in excretory/secretory products collected from salmon lice, *Lepeophtheirus salmonis* 11 (2018) 294.
- [105] V. Maldonado-Aguayo, J. Chávez-Mardones, A.T. Gonçalves, C. Gallardo-Escárate, Cathepsin Gene Family Reveals Transcriptome Patterns Related to the Infective Stages of the Salmon Louse *Caligus rogercresseyi* vol. 10, (2015) e0123954.
- [106] C.J. Lehiy, B.S. Drolet, The salivary secretome of the biting midge, *Culicoides sonorensis*. *PeerJ.* 2 (2014) e426.
- [107] D.M. Blow, The tortuous story of asp... his... ser: structural analysis of α -chymotrypsin, *Trends Biochem. Sci.* 22 (1997) 405–408.
- [108] R. Skern-Mauritzen, P. Frost, S. Dalvin, B.O. Kvamme, I. Sommerset, F. Nilsen, A trypsin-like protease with apparent dual function in early *Lepeophtheirus salmonis* (Krøyer) development, *BMC Mol. Biol.* 10 (2009) 44.
- [109] D. Valenzuela-Miranda, C. Gallardo-Escárate, *Caligus rogercresseyi* serine proteases: transcriptomic analysis in response to delousing drugs treatments, *Aquaculture* 465 (2016) 65–77.
- [110] P. Johnsson, L. Lipovich, D. Grandér, K.V. Morris, Evolutionary conservation of long non-coding RNAs; sequence, structure, function, *Biochim. Biophys. Acta Gen. Subj.* 1840 (2014) 1063–1071.
- [111] Z. Tang, Y. Wu, Y. Yang, Y.-C.T. Yang, Z. Wang, J. Yuan, et al., Comprehensive analysis of long non-coding RNAs highlights their spatio-temporal expression patterns and evolutionary conservation in *Sus scrofa*, *Sci. Rep.* 7 (2017).
- [112] R. Cech Thomas, A. Steitz Joan, The noncoding RNA revolution—trashing old rules to forge new ones, *Cell* 157 (2014) 77–94.
- [113] C.P. Ponting, P.L. Oliver, W. Reik, Evolution and functions of long noncoding RNAs, *Cell* 136 (2009) 629–641.
- [114] T.M. Aune, C.F. Spurlock, Long non-coding RNAs in innate and adaptive immunity, *Virus Res.* 212 (2016) 146–160.
- [115] C.F. Spurlock III, J.T. Tossberg, Y. Guo, S.P. Collier, P.S. Crooke Iii, T.M. Aune, Expression and functions of long noncoding RNAs during human T helper cell differentiation, *Nat. Commun.* 6 (2015) 6932.
- [116] K. Kaushik, V.E. Leonard, K.V. Shamsudheen, M.K. Lalwani, S. Jalali, A. Patowary, et al., Dynamic expression of long non-coding RNAs (lncRNAs) in adult zebrafish, *PLoS One* 8 (2013) e83616.
- [117] R. Al-Tobasei, B. Paneru, M. Salem, Genome-wide discovery of long non-coding RNAs in rainbow trout, *PLoS One* 11 (2016) e0148940.
- [118] S. Carpenter, D. Aiello, M.K. Atianand, E.P. Ricci, P. Gandhi, L.L. Hall, et al., A long noncoding RNA mediates both activation and repression of immune response genes, *Science* 341 (2013) 789–792.
- [119] X. Zhou, X. Han, A. Wittfeldt, J. Sun, C. Liu, X. Wang, et al., Long non-coding RNA ANRIL regulates inflammatory responses as a novel component of NF- κ B pathway, *RNA Biol.* 13 (2016) 98–108.
- [120] L. Jiang, W. Liu, A. Zhu, J. Zhang, J. Zhou, C. Wu, Transcriptome analysis demonstrate widespread differential expression of long noncoding RNAs involve in *Larimichthys crocea* immune response, *Fish Shellfish Immunol.* 51 (2016) 1–8.
- [121] S. Boltaña, D. Valenzuela-Miranda, A. Aguilar, S. Mackenzie, C. Gallardo-Escárate, Long noncoding RNAs (lncRNAs) dynamics evidence immunomodulation during ISAV-Infected Atlantic salmon (*Salmo salar*), *Sci. Rep.* 6 (2016).
- [122] D. Valenzuela-Miranda, C. Gallardo-Escárate, Novel insights into the response of Atlantic salmon (*Salmo salar*) to *Piscirickettsia salmonis*: interplay of coding genes and lncRNAs during bacterial infection, *Fish Shellfish Immunol.* 59 (2016) 427–438.
- [123] E. Tarifeño-Saldivia, D. Valenzuela-Miranda, C. Gallardo-Escárate, In the shadow: the emerging role of long non-coding RNAs in the immune response of Atlantic salmon, *Dev. Comp. Immunol.* 73 (2017) 193–205.
- [124] D.P. Bartel, MicroRNAs: genomics, biogenesis, mechanism, and function, *Cell* 116 (2004) 281–297.
- [125] D.P. Bartel, MicroRNAs: target recognition and regulatory functions, *Cell* 136 (2009) 215–233.
- [126] Y. Bi, G. Liu, R. Yang, MicroRNAs: novel regulators during the immune response, *J. Cell. Physiol.* 218 (2009) 467–472.
- [127] C.C. Judice, C. Bourgard, A.C. Kayano, L. Albrecht, F.T. Costa, MicroRNAs in the host-apicomplexan parasites interactions: a review of immunopathological aspects, *Front. Cellular Infect. Microbiol.* 6 (2016), <https://doi.org/10.3389/fcimb.2016.00005>.
- [128] M.A. Lindsay, microRNAs and the immune response, *Trends Immunol.* 29 (2008) 343–351.
- [129] L.F. Lu, A. Liston, MicroRNA in the immune system, microRNA as an immune system, *Immunology* 127 (2009) 291–298.
- [130] Y.-C. Lu, J.T.-C. Chang, Y.-C. Huang, C.-C. Huang, W.-H. Chen, L.-Y. Lee, et al., Combined determination of circulating miR-196a and miR-196b levels produces high sensitivity and specificity for early detection of oral cancer, *Clin. Biochem.* 48 (2015) 115–121.
- [131] R.M. O'Connell, D.S. Rao, D. Baltimore, microRNA regulation of inflammatory responses, *Annu. Rev. Immunol.* 30 (2012) 295–312.
- [132] R.M. O'Connell, D.S. Rao, A.A. Chaudhuri, D. Baltimore, Physiological and pathological roles for microRNAs in the immune system, *Nat. Rev. Immunol.* 10 (2010) 111–122.

- [133] V. Valenzuela-Muñoz, B. Novoa, A. Figueras, C. Gallardo-Escárate, Modulation of Atlantic salmon miRNome response to sea louse infestation, *Dev. Comp. Immunol.* 76 (2017) 380–391.
- [134] R. Farlora, D. Valenzuela-Miranda, P. Alarcon-Matus, C. Gallardo-Escarate, Identification of microRNAs associated with sexual maturity in Rainbow Trout brain and testis through small RNA deep sequencing, *Mol. Reprod. Dev.* 82 (2015) 651–662.
- [135] M. Salem, C. Xiao, J. Womack, C.E. Rexroad, J. Yao, A microRNA repertoire for functional genome research in Rainbow Trout (*Oncorhynchus mykiss*), *Mar. Biotechnol.* 12 (2010) 410–429.
- [136] Z. Sha, G. Gong, S. Wang, Y. Lu, L. Wang, Q. Wang, et al., Identification and characterization of *Cynoglossus semilaevis* microRNA response to *Vibrio anguillarum* infection through high-throughput sequencing, *Dev. Comp. Immunol.* 44 (2014) 59–69.
- [137] M. Hussain, S. Asgari, MicroRNAs as mediators of insect host–pathogen interactions and immunity, *J. Insect Physiol.* 70 (2014) 151–158.
- [138] C. Gallardo-Escárate, V. Valenzuela-Muñoz, S. Boltaña, G. Nuñez-Acuña, D. Valenzuela-Miranda, A. Gonçalves, et al., The *Caligus rogercresseyi* miRNome: discovery and transcriptome profiling during the sea lice ontogeny, *Agri Gene* 4 (2017) 8–22.
- [139] D. Valenzuela-Miranda, G. Nuñez-Acuña, V. Valenzuela-Muñoz, S. Asgari, C. Gallardo-Escárate, MicroRNA biogenesis pathway from the salmon louse (*Caligus rogercresseyi*): emerging role in delousing drug response, *gene* 555 (2015) 231–241.
- [140] S. Egerton, S. Culloty, J. Whooley, C. Stanton, R.P. Ross, The gut microbiota of marine fish, *Front. Microbiol.* 9 (2018) 873.
- [141] D. Merrifield, A. Rodiles, The fish microbiome and its interactions with mucosal tissues, in: B.R. Beck, E. Peatman (Eds.), *Mucosal Health in Aquaculture*, Academic Press, 2015, pp. 273–289.
- [142] J. Romero, E. Ringø, D.L. Merrifield, *The Gut Microbiota of Fish*. Aquaculture Nutrition, John Wiley & Sons, Ltd, 2014, pp. 75–100.
- [143] A. Larsen, Z. Tao, S.A. Bullard, C.R. Arias, Diversity of the skin microbiota of fishes: evidence for host species specificity, *FEMS Microbiol. Ecol.* 85 (2013) 483–494.
- [144] J.B. Xiong, L. Nie, J. Chen, Current understanding on the roles of gut microbiota in fish disease and immunity, *Zool. Res.* 40 (2019) 70–76.
- [145] R.A. Britton, V.B. Young, Role of the intestinal microbiota in resistance to colonization by *Clostridium difficile*, *Gastroenterology* 146 (2014) 1547–1553.
- [146] M.P. Gonzalez Gomez, S.L. Marin Arribas, L. Vargas-Chacoff, Stress response of *Salmo salar* (Linnaeus 1758) facing low abundance infestation of *Caligus rogercresseyi* (Boxshall & Bravo 2000), an object in the tank, and handling, *J. Fish Dis.* 39 (2016) 853–865.
- [147] M.P. González, L. Vargas-Chacoff, S.L. Marín, Stress response of *Salmo salar* (Linnaeus 1758) when heavily infested by *Caligus rogercresseyi* (Boxshall & Bravo 2000) copepodids, *Fish Physiol. Biochem.* 42 (2016) 263–274.
- [148] R.H. Easy, N.W. Ross, Changes in Atlantic salmon (*Salmo salar*) epidermal mucus protein composition profiles following infection with sea lice (*Lepeophtheirus salmonis*), *Comp. Biochem. Physiol. Genom. Proteonom.* 4 (2009) 159–167.
- [149] F. Provan, L.B. Jensen, K.E. Uleberg, E. Larssen, T. Rajalahti, J. Mullins, et al., Proteomic analysis of epidermal mucus from sea lice-infected Atlantic salmon, *Salmo salar* L., *J. Fish Dis.* 36 (2013) 311–321.
- [150] M.S. Llewellyn, S. Leadbeater, C. Garcia, F.E. Sylvain, M. Custodio, K.P. Ang, et al., Parasitism perturbs the mucosal microbiome of Atlantic Salmon, *Sci. Rep.* 7 (2017) 43465.
- [151] K.R. Carman, F.C. Dobbs, Epibiotic microorganisms on copepods and other marine crustaceans, *Microsc. Res. Tech.* 37 (1997) 116–135.
- [152] P.H. Moisanter, A.D. Sexton, M.C. Daley, Stable Associations masked by temporal variability in the marine copepod microbiome, *PLoS One* 10 (2015) e0138967.
- [153] G. Minniti, L.H. Hagen, D. Porcellato, S.M. Jorgensen, P.B. Pope, G. Vaaje-Kolstad, The skin-mucus microbial community of farmed atlantic salmon (*Salmo salar*), *Front. Microbiol.* 8 (2017) 2043.
- [154] F.A. Sepúlveda, J.F. Torres, C.D. Infante, M.T. González, Potential role of ectoparasites (*Zeuxapta seriolae* and *Caligus lalandei*) in the transmission of pathogenic bacteria in yellowtail kingfish *Seriola lalandi*, inferred from cultivable microbiota and molecular analyses, *J. Fish Dis.* 40 (7) (2017 Jul) 979–985.
- [155] L. Nese, O. Enger, Isolation of *Aeromonas salmonicida* from salmon lice *Lepeophtheirus salmonis* and marine plankton, *Dis. Aquat. Org.* 16 (1993) 79–81.
- [156] C.W. Novak, D.L. Lewis, B. Collicutt, K. Verkaik, D.E. Barker, Investigations on the role of the salmon louse, *Lepeophtheirus salmonis* (Caligidae), as a vector in the transmission of *Aeromonas salmonicida* subsp. *salmonicida*, *J. Fish Dis.* 39 (2016) 1165–1178.
- [157] L.A. Hamre, C. Eichner, C.M.A. Caipang, S.T. Dalvin, J.E. Bron, Nilsen, R. Skern-Mauritzen, The salmon louse *Lepeophtheirus salmonis* (copepoda: caligidae) life cycle has only two chalimus stages, *PLoS One* 8 (2013) (2013) e73539.