

¹ Department of Biology, University of Waterloo, 200 University Ave W., Waterloo, ON, Canada

² Center for Genetic Engineering and Biotechnology, Havana, Cuba

³ Department of Pathobiology, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada

Abstract

Pituitary adenylate cyclase-activating polypeptide (PACAP) is a multi-functional neuropeptide that is widely distributed in mammals and is capable of performing roles as a neurotransmitter, neuromodulator and vasodilator. This polypeptide belongs to the glucagon/secretin superfamily, of which some members have been shown to act as antimicrobial peptides in both mammalian and aquatic organisms. In teleosts, PACAP has been demonstrated to have direct antimicrobial activity against several aquatic pathogens, yet this phenomenon has never been studied throughout a live bacterial infection. The present study focuses on the influence of PACAP on the rainbow trout monocyte/macrophage-like cell line, RTS11, when exposed to the coldwater bacterial pathogen *Flavobacterium psychrophilum*. PACAP was shown to have direct antimicrobial activity on *F. psychrophilum* when grown in both cytophaga broth and cell culture media (L-15). Further, the ability of teleostean PACAP to permeabilize the membrane of an aquatic pathogen, *F. psychrophilum*, was revealed for the first time. The viability of RTS11 when exposed to PACAP was also observed using a trypan blue exclusion assay to determine optimal experimental doses of the antimicrobial peptide. Interestingly, when RTS11 was pre-treated with PACAP for 24 hours before experiencing exposure to live *F. psychrophilum*, growth of the pathogen was severely inhibited in a dose-dependent manner when compared to control cells receiving no PACAP pre-treatment. Relative expression of pro-inflammatory cytokines and PACAP receptors was also observed in RTS11 following PACAP exposure alone and in conjunction with live *F. psychrophilum* challenge. These qRT-PCR findings revealed that PACAP may have a synergistic effect on RTS11 immune function. The results of this study provide evidence that PACAP has immunostimulatory activity on rainbow trout immune cells as well as direct antimicrobial activity against aquatic bacterial pathogens such as *F. psychrophilum*. As there are numerous pathogens that impact the aquaculture industry, PACAP may stimulate the teleost immune system while also providing an efficacious alternative to antibiotic use.

Keywords: Rainbow trout, antimicrobial peptide, PACAP, RTS11, *Flavobacterium psychrophilum*

Corresponding author.

E-mail address: slemple@uwaterloo.ca (S.L. Semple).

O-108.

Efficient and long-lasting protection against the pacific oyster mortality syndrome through antiviral immune priming

Maxime Lafont¹, Bruno Petton², Julien deLorgeril¹, Agnes Vergnes¹, Jeremie Vidal-Dupiol¹, Yannick Gueguen¹, Philippe Haffner¹, Guillaume Mitta¹, Benjamin Gourbal¹, Caroline Montagnani^{1, #}.

¹ IHPE, Université de Montpellier, CNRS, Ifremer, Université de Perpignan Via Domitia, France

² Ifremer, LEMAR UMR6539, Argenton-en-Landunvez, France

Abstract

The major economic and environmental consequences of recurring mortalities affecting the Pacific oyster *Crassostrea gigas* have initiated many research projects aiming at understanding these phenomena. The solutions anticipated to deal with these mortalities are mainly based on mass selection breeding programs but preventive treatments are still lacking. However, over the last decade, studies have been accumulating revealing the adaptive capabilities of innate immunity, the only component of defense mechanisms in invertebrates. Numerous findings have shown that a wide range of invertebrates can develop innate immune memory (also

called immune priming) leading to improved survival during a second encounter with a pathogen. In this context, we undertook to study the possibilities of acting against mortalities by stimulating immune capacities of oysters.

In the present study, we show that the exposure of oyster juveniles to an immunostimulant (a viral mimic called poly (I: C)) can lead to enhanced survival capacities (up to 100%) following OsHV-1 infection or during a mortality episode in the field. That protection is specific to viral protection as poly(I:C) fails to protect oyster against a pathogenic bacteria. We also show that this priming phenomenon is durable as it can last more than 4 months suggesting for the first time the existence of mechanisms of immune memory in this invertebrate species. Finally, analysis of the molecular pathways underlying that protection using dual RNAseq, revealed that priming was based on the triggering of a strong and sustained antiviral response limiting replication of the virus, thus allowing the protection of oysters on the long term. Altogether these results bring new insights into the oyster capacities to build an innate immune memory, its adaptive capacities and provide a platform to further explore novel strategies to help mitigate disease threats upon marine bivalves.

Keywords: Immunity, antiviral, OsHV-1, priming, oyster

Corresponding author.

E-mail address: cmontagn@ifremer.fr (C. Montagnani).

O-109.

The immune recognition mechanisms in molluscs

Lingling Wang^{1,2,3}, Weilin Wang^{1,3}, Chuanyan Yang^{1,3}, Jiejie Sun^{1,3}, Linsheng Song^{1,2,3, #}.

¹ Liaoning Key Laboratory of Marine Animal Immunology, Dalian Ocean University, Dalian 116023, China

² Functional Laboratory of Marine Fisheries Science and Food Production Process, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266200, China

³ Liaoning Key Laboratory of Marine Animal Immunology and Disease Control, Dalian Ocean University, Dalian 116023, China

Abstract

Immune recognition is the first step of immune response, which plays key role in immune protection in all organisms. Since the proposal of pattern recognition theory, numerous pattern recognition receptors (PRRs) have been reported, especially in molluscs where expansive PRRs have been identified by genomic prediction. Taken the model bivalve oyster as example, oysters are constantly threatened by the invasion of pathogens and they have developed a sophisticated repertoire of PRRs to recognize diversiform microorganisms. So far, seven PRRs families, such as peptidoglycan recognition proteins (PGRPs), lectins (including C-type lectins, galectins and sialic acid-binding Ig-like lectins), toll-like receptors (TLRs), C1q domain containing proteins (C1qDCs), Gram-negative binding proteins (GNBPs), scavenger receptors (SRs) and fibrinogen-related proteins (FREPs) have been identified in oysters. What's more, some novel PRRs, such as DM9 domain containing proteins (DM9CPs), Caspases, interleukin 17 (IL-17) and arginine kinase have also been characterized from oysters, which expands our knowledge of PRRs in invertebrates. These various PRRs have been partially validated to be different in recognition specificity, down-stream signal pathway and immune effects. Most of the PRRs serve as multi-functional proteins, not only in immune recognition, but also in the elimination of invading microbes. In addition, these PRRs differentially expressed in mucosal immune tissues and systemic circulatory system where immune recognition taking place. And different effects could be induced by mucosal and systemic immune recognition, such as immune memory or immune tolerance. These results uncovered in molluscs have expanded our knowledge about the classical pattern recognition theory, and also provided theoretical basis for vaccine development in mollusc breeding-species in the future.