

³ Aquaculture and Fisheries Group, Wageningen Institute of Animal Sciences, Wageningen University & Research, Wageningen, The Netherlands

⁴ Faculty of Science, University of South Bohemia in České Budějovice, České Budějovice, Czech Republic

⁵ Faculty of Engineering and Natural Sciences, Johannes Kepler University, Linz, Austria

⁶ Cell Biology and Immunology Group, Wageningen Institute of Animal Sciences, Wageningen University & Research, Wageningen, The Netherlands

Abstract

Sphaerospora molnari is a myxozoan parasite causing skin and gill sphaerosporosis in common carp (*Cyprinus carpio*) in Central Europe. Prior to spore formation, multicellular proliferative stages of *S. molnari* circulate for several weeks in the vascular system of its host despite the array of humoral and cellular immune effector mechanisms. Using our laboratory infection model, we aimed to elucidate the kinetics of presporogonic development of *S. molnari*, while simultaneously analyze the immune responses over a period of 63 days. The obtained results identified two peaks of acute parasitemia on day 28 and 42 respectively. Unexpectedly, the highest parasite load was detected in the liver, a previously unknown localization of *S. molnari*. In response to the infection, the immune system induced dynamic changes in the expression of pro- and anti-inflammatory cytokines, with a predominant role of IL-10 reaching up to 1456 fold increase compared to control fish. The haematological analysis revealed a steady increase in the number of lymphocytes from day 28 onwards, correlating with the growing number of parasites, and only marginal changes in other populations. Additionally, our data revealed a strong increase in the expression of IgM transcripts and increased number of IgM⁺ B lymphocytes, which produce specific antibodies recognizing *S. molnari* antigens in western blot. Strikingly, although the sera of infected fish exhibit potent opsonizing capacity *in vitro*, *S. molnari* isolated from the blood of infected fish are not labelled with carp IgM. These findings indicate the presence of so far unknown evasion strategy and questions the importance of *S. molnari*-specific antibodies in parasite elimination. To our knowledge, this is the first study analyzing the early myxozoan development and immune modulation mechanisms along with innate and adaptive immune responses of the fish host in a controlled laboratory system, adding important information on host-parasite interaction of early metazoans (Cnidaria) with basic vertebrate immune systems.

Keywords: *Sphaerospora molnari*, Antibodies, Cytokines, B lymphocytes, IL-10

Corresponding author.

E-mail address: tkorytar@frov.jcu.cz (T. Korytář).

O-124.

From gross morphology to gill transcriptome in farmed Atlantic salmon (*Salmo salar*): Lessons from multi-site sampling

E. Król¹, A. Douglas¹, P. Noguera², V. Valdenegro^{1,2,3,4}, K. Gajardo³, R. Bickerdike⁴, S.A.M. Martin^{1,#}

¹ School of Biological Sciences, University of Aberdeen, Aberdeen, AB24 2TZ, UK

² Aquaculture and Marine Environment, Marine Scotland Science, Aberdeen, UK

³ BioMar AS, Havnegata 9, Pirsenteret, Norway

⁴ Scottish Sea Farms, Laurel House, Laurelhill Business Park, Stirling, FK7 9JQ, UK

Abstract

The gill is a multifunctional organ involved in many physiological processes such as gas exchange, osmotic and ionic regulation, acid-base balance and excretion of nitrogenous waste. Due to its interface with the environment, the gill plays a key role as primary mucosal defence tissue

against pathogens and is equipped with the gill-associated lymphoid tissue (GIALT). In recent years, the prevalence of gill damage and gill diseases has increased significantly, leading to the substantial losses in Atlantic salmon aquaculture worldwide. Both the transition from healthy to unhealthy gill phenotypes as well the progression of various gill pathologies such as proliferative gill disease (PGD), amoebic gill disease (AGD) and complex gill disease (CGD) are commonly characterised by inflammation and epithelial cell hyperplasia. Routine monitoring for PGD relies on gross (macroscopic) evaluation of gill health, coupled with histological examination of gill sections. To explore underlying molecular events that are associated with progression of PGD, we examined Atlantic salmon from geographically diverse aquaculture sites in Scotland. Total RNA was extracted from 43 fish presenting low or medium gill PGD scores and analysed by whole transcriptome analysis using RNA-seq to determine the molecular signature of the advanced PGD. For each fish, 20M reads were generated and mapped to the Atlantic salmon genome. Importantly, we showed that the sampling site had greater effect on the gill transcriptome than the actual PGD score, providing support for a complex and multifactorial aetiology of PGD, with minimal common molecular responses between different sites. Similar pattern was found for histology, agreeing with the outcome of the RNA-seq analysis. In general, both RNA-seq and histology data clustered together based on the origin of samples, suggesting that the PGD scores may inform about the overall progression of gill damage, but not about the underlying pathology.

Keywords: Proliferative gill disease, gill histopathology, gill transcriptome, PGD scores, RNA-seq

Corresponding author.

E-mail address: sam.martin@abdn.ac.uk (S.A.M. Martin).

O-125.

A proteomic study of resistance to brown ring disease in the manila clam, *Ruditapes philippinarum*

M. Smits^{1,2,#}, S. Artigaud¹, B. Bernay³, A. Bidault¹, L. Bargelloni², C. Paillard¹

¹ Laboratoire des Sciences de l'Environnement Marin, Institut Universitaire Européen de la Mer, rue Dumont d'Urville, 29280, Plouzané, France

² Department of Comparative Biomedicine and Food Science, University of Padova, Agripolis Campus, Viale dell'Università, 16, 35020, Legnaro, PD, Italy

³ Plateforme Proteogen, SFR ICORE 4206, Université de Caen Basse-Normandie, Esplanade de la paix, 14032, Caen cedex, France

Abstract

Mollusks represent over a fifth of the global aquaculture market, accounting for USD 29.2 billion in 2016. Infectious disease is one of the main limiting factors to the development of mollusk aquaculture, and the difficulties inherent to combating pathogens through antibiotic therapies or disinfection have led to extensive research on host defense mechanisms and host-pathogen relationships. It has become increasingly clear that innate immunity and genetic variability are key factors underlying disease resistance, and that genetic selection for resistance is essential for effective disease control. The Manila clam, *Ruditapes philippinarum*, is a main cultured bivalve species of economic interest produced on a global scale. While the species originates from Japan, it was introduced to European coasts for aquaculture in the early 1970s. In 1987, mass mortalities in clam landings on the Atlantic coast of France led to the identification of Brown Ring Disease (BRD) and of its etiological agent, *Vibrio tapetis*. BRD is characterized by bacterial colonization of the host's extra-pallial compartment, provoking abnormal conchiolin deposits along the inner surface of the shell. While some clams are capable of effectively combating the pathogen, others succumb to a chronic infection characterized by thick conchiolin deposits, low condition index, and death. Although a significant body of research has allowed us to gain a better understanding the