



ELSEVIER

Contents lists available at ScienceDirect

## Fish and Shellfish Immunology

journal homepage: [www.elsevier.com/locate/fsi](http://www.elsevier.com/locate/fsi)

Full length article

# Transcriptomic analysis of clam extrapallial fluids reveals immunity and cytoskeleton alterations in the first week of Brown Ring Disease development

Alexandra Rahmani<sup>a,\*\*</sup>, Erwan Corre<sup>b</sup>, Gaëlle Richard<sup>a</sup>, Adeline Bidault<sup>a</sup>, Christophe Lambert<sup>a</sup>, Louisi Oliveira<sup>c</sup>, Cristiane Thompson<sup>c</sup>, Fabiano Thompson<sup>c</sup>, Vianney Pichereau<sup>a,\*,1</sup>, Christine Paillard<sup>a,\*\*,1</sup>

<sup>a</sup> Univ Brest, CNRS, IRD, Ifremer, UMR 6539 LEMAR, F-29280, Plouzane, France

<sup>b</sup> Sorbonne Universités, Université Pierre et Marie Curie-Paris 6, CNRS, FR2424, Station Biologique de Roscoff, Roscoff, France

<sup>c</sup> Centro de Ciências da Saúde, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

## ARTICLE INFO

## Keywords:

Brown ring disease  
*V. tapetis*  
*R. philippinarum*  
 Hemocytes  
 Actin cytoskeleton  
 $\beta$ -Thymosin  
 Coactosin  
 Resting cells

## ABSTRACT

The Brown Ring Disease is an infection caused by the bacterium *Vibrio tapetis* on the Manila clam *Ruditapes philippinarum*. The process of infection, in the extrapallial fluids (EPFs) of clams, involves alteration of immune functions, in particular on hemocytes which are the cells responsible of phagocytosis. Disorganization of the actin-cytoskeleton in infected clams is a part of what leads to this alteration. This study is the first transcriptomic approach based on collection of extrapallial fluids on living animals experimentally infected by *V. tapetis*. We performed differential gene expression analysis of EPFs in two experimental treatments (healthy-against infected-clams by *V. tapetis*), and showed the deregulation of 135 genes. In infected clams, a downregulation of transcripts implied in immune functions (lysosomal activity and complement- and lectin-dependent PRR pathways) was observed during infection. We also showed a deregulation of transcripts encoding proteins involved in the actin cytoskeleton organization such as an overexpression of  $\beta$ 12-Thymosin (which is an actin sequestration protein) or a downregulation of proteins that closely interact with capping proteins such as Coactosin, that counteract action of capping proteins, or Profilin. We validated these transcriptomic results by cellular physiological analyses that showed a decrease of the lysosome amounts and the disorganization of actin cytoskeleton in infected hemocytes.

## 1. Introduction

*Ruditapes philippinarum*, the Manila clam, has been imported in Europe from Indo-pacific coasts in the early 70's. Ten years later, a disease causing a brown organic deposit at the inner face of the shell, between the pallial line and the edge of the shell, spread mostly all over Northern coasts of Europe, causing mass mortalities to this species [1,2]. This infection has been called the Brown Ring Disease (BRD, according to its macroscopic manifestations) and is due to the bacterium *Vibrio tapetis* [3,4].

*V. tapetis* is a Gram-negative bacillus that acts, in most of cases, as an external microparasite, mainly in cold waters (optimal infection

temperature of 14 °C [5]) by spreading into the extrapallial fluids (EPF) of the Manila clam. Despite the fact that growth declines and shell deformations are frequently observed in this disease, tissue's infections are rare and only happen in case of pre-existing tissues lesions, then causing the death of animals. Nevertheless, BRD is a microparasitic infection which dynamics does not imply the death of animals [4,6].

A particular symptom of BRD is the rounding of hemocytes in infected clams, thus revealing damages to the clam's immune system [7,8]. Indeed, hemocytes are key effectors of the immune defenses of the clam that eliminate pathogens by phagocytosis [9]. In a normal phagocytosis process, hemocytes spread their pseudopods which encapsulate and internalize bacteria and, by a phagosome/lysosome

**Abbreviations:** ABP, Actin Binding Protein; BRD, Brown Ring Disease; EPF, Extrapallial Fluids; FSW, Filter Sterilized Seawater

\* Corresponding author.

\*\* Corresponding author.

\*\*\* Corresponding author.

E-mail addresses: [alexandra.rahmani@univ-brest.fr](mailto:alexandra.rahmani@univ-brest.fr) (A. Rahmani), [vianney.pichereau@univ-brest.fr](mailto:vianney.pichereau@univ-brest.fr) (V. Pichereau), [christine.paillard@univ-brest.fr](mailto:christine.paillard@univ-brest.fr) (C. Paillard).

<sup>1</sup> These authors contributed equally.

<https://doi.org/10.1016/j.fsi.2019.08.025>

Received 21 February 2019; Received in revised form 9 August 2019; Accepted 10 August 2019

Available online 13 August 2019

1050-4648/ © 2019 Elsevier Ltd. All rights reserved.

fusion, destroy the pathogen. During BRD infection, it was shown that *V. tapetis* causes a disorganization of the hemocyte actin cytoskeleton, resulting in the loss of hemocytes pseudopods (and thus, the rounding of cells) thus reducing its phagocytosis capacity [10]. *V. tapetis* is then able to multiply into hemocytes, finally resulting in cell membrane disruption. Bacteria can then spread into clams before being released by pseudo feces into the environment (sediments) where they can infect other clams [7,11–13].

This study is the first transcriptomic approach based on collection of extrapallial fluids (EPFs) on living animals experimentally infected by *V. tapetis*. This study led to the identification of a set of transcripts regulated during BRD infection, combined with physiological experiments performed in order to check the validity of some of the major interpretations raised from the differential expression analysis. This study gave new insights to understand the pathogenicity of *V. tapetis* to Manila clams.

## 2. Materials and methods

### 2.1. Clams in vivo infection, RNA extraction and sequencing

Animals and experimental design are fully described in Ref. [14]. Two years-old clams were harvested in the Marennes-Oleron Bay (Charente Maritime, France) on May 5th, 2014. A first visual health diagnostic was performed on 50 clams *in situ* and no BRD clinical signs were observed. Clams were then transferred to Ifremer facilities in Plouzané (Finistère, France) and held in a 6-day quarantine with a chloramphenicol ( $8 \text{ mg} \cdot \text{L}^{-1}$ ) treatment to avoid local contamination. After quarantine, animals were randomly divided and maintained in trays placed in eighteen 100 L-tanks equipped with air-lift systems (100 clams per tank). Water temperature was maintained at  $13^\circ\text{C}$  during the whole experiment so that infection by *V. tapetis* was favored [5]. Tanks were continuously supplied with filter sterilized seawater (FSW, filtered at  $1 \mu\text{m}$  and UV-sterilized) at a renewal rate of 50% per day and cleaned every two days. Clams were fed *ad libitum* with T-iso. Manila clams, *R. philippinarum*, were injected in the extrapallial cavity with  $100 \mu\text{L}$  of *V. tapetis* CECT4600 suspension at the concentration of  $10^7 \text{ CFU mL}^{-1}$  (Infected, 900 clams) or by Filter Sterilized Seawater (FSW, Control, 900 clams) as described in Le Bris et al. [15]. At 7 days post injection, EPFs of infected and control clams were harvested and total RNAs were extracted with TRI reagent® according to the procedure manufacturer (Sigma-Aldrich®). Presence or absence of the Brown Ring Disease have been diagnosed according to the standardized methodology established by Paillard et al. [16] based on the presence and localization of brown deposit in clam shells. We performed this diagnosis by an image analysis method using a 50-mm CANON macroscopic lens and analyzed using Visilog 6.6 image analysis software, according to image analysis methodology already performed [14,16]. This method allows to visualize the shell and to analyze the presence of a brown deposit (BRD positive diagnosis) with an image analysis software. BRD was diagnosed in all infected clams whereas no BRD was found in control animals. As ribosomal RNA (rRNA) is the most abundant component in total RNA, while not being informative in this kind of RNAseq strategy, we performed two rRNA depletion steps, the first one targeting bacterial rRNA and the second one targeting eukaryotic ones (by using the Ribo-Zero rRNA Removal Kits, bacteria and Human/Mouse/Rat, respectively, from Illumina). The RNAs from seven clams, 3 healthy and 4 infected, were subjected to cDNA libraries creation according to standard Illumina procedures (TruSeq Stranded mRNA Library Prep Kit). All the libraries were sequenced, by Illumina Miseq PE300, at the Federal University of Rio de Janeiro, Brazil. Other control and infected clams have been further analyzed for other questions [14].

### 2.2. Transcriptomic analysis

Raw data analysis and differential expression evaluation were carried out using the ABILMS Galaxy platform, Station Biologique de Roscoff, CNRS/Sorbonne Université, France (<http://galaxy.sb-roscoff.fr>, January 2019).

Reads have been trimmed using AdaptorRemoval v2 [17], Trimmomatic v0.36.3 [18] and Prinseq v0.20.4 [19] in order to remove read adaptors, low quality sequences and poly(A) tails. Persistent rRNA were removed with SortMeRNA v2.1b.4 [20]. For the differential analysis, all the reads were mapped to a dataset of 61 747 transcripts obtained from a combination of *R. philippinarum*'s stimulated hemocytes ESTs (expressed sequences tag) library [21], from *R. philippinarum*'s ESTs deposited in NCBI EST database (<https://www.ncbi.nlm.nih.gov/nucest/>) and sequences deposited in the Ruphibase database (<http://compgen.bio.unipd.it/ruphibase/>). Transcripts abundances were estimated by using the “Align and Build expression matrix” pipeline (mapping with bowtie + reads estimation with RSEM) of the Trinity suite v2.4.0.2 [22]. The statistical analysis of differentially expressed transcripts were done using the “Differential expression analysis” pipeline of Trinity suite v2.4.0 (using DESeq2) and “Extract and cluster differentially expressed transcripts” pipeline of Trinity suite v2.4.0 [22]. We have set up parameters to exclude differential expression with a LogFoldChange between  $-2$  and  $2$ . Transcripts were considered as significantly differentially expressed if the p-value adjusted by FDR (false discovery rate) is  $< 10^{-3}$ . The identified deregulated transcripts were subjected to blast2GO for re-annotation, including blastX vs the NCBI nr, refseq and swissprot (2018 version) non redundant databases, interpro, GO mapping and GO annotation, all with standard parameters. All the annotations and functions were carefully manually checked. The sequencing data have been made available at the European Nucleotide Archive (project PRJEB23385). The analysis data are given in Table S1.

### 2.3. Clams in vitro infection: bacterial exposure

Animals used in this study were Manila clams from the SATMAR shellfish aquaculture site in Marennes (Charente-Maritime, France). The clam pool was acclimatized in oxygenated seawater using a bubbler at  $14^\circ\text{C}$ . Hemolymph was harvested directly from the adductor muscle. Each hemolymph was sampled individually and the quality of the hemocytes present in this fluid was checked by observation under the microscope (presence of pseudopods, few round-shaped hemocytes). Good quality hemolymph samples were pooled and the hemocytes enumerated by using a Malassez cell. For exposure assays, hemolymph was exposed to a bacterial suspension (in FSSW) of *V. tapetis* CECT4600, at a 25/1: bacteria/hemocyte ratio. Hemocytes exposure were performed in at least 3 replicates and two independent experiments for each condition tested. Briefly,  $100 \mu\text{L}$  of hemolymph was added in 24-well plates. After a few minutes, in order to let hemocytes attach on the bottom of the plate,  $100 \mu\text{L}$  of bacterial suspension was added for the exposed samples, and  $100 \mu\text{L}$  of FSSW for the controls.

#### 2.3.1. F-actin visualization by microscopy

To visualize F-actin in control and infected hemocytes, we used the fluorescent probe Rhodamine Phalloidin (Invitrogen). Exposure assays were performed as described above. After 3 h of challenge, we collected the supernatant and added  $30 \mu\text{L}$  of Triton X-100 0.1% for 15 min. Triton permeabilizes cells, thus allowing probes to enter the cell.  $2 \mu\text{L}$  of Rhodamine-phalloidin (Stock solution dissolved in 1.5 mL methanol) was then added to each sample (hemolymph non exposed and exposed to *V. tapetis*) which were stained for 20 min, on ice, without light. Epifluorescence microscopy was then performed by using a ZEISS AXIO observer Z1 inverted microscope with a wavelength of excitation BP (band pass) 550/25 nm, Beam splitter 570 nm and of emission BP 605/70 nm. Cells were observed both in red epifluorescence and in bright field.

#### 2.3.2. LysoTracker assay

The LysoTracker® probes (Molecular Probes®) are fluorescent acidotropic probes for labeling and tracking acidic organelles in live cells, according to the manufacturer definition. We have developed and tested a method to characterize the amount of acidic organelles on hemocytes exposed to *V. tapetis* or FSSW for control. LysoTracker assay is performed during the *in vitro* bacterial exposure assay based on the

rounding of hemocytes caused by *V. tapetis* previously developed by Choquet et al. [7]. Our protocol allows to perform both tests on the same sample, and then to correlate the rounding phenotype to the cell content in acidic organelles.

Hemocytes exposure were performed as described above. The 24 wells plates were incubated 1 h at 18 °C before addition of 4 µL of 50 µM LysoTracker® Red DND-99 (Invitrogen, emission from 550 to 700 nm, final concentration 1 µM) to each well. The 24 wells plates were then let 2 h more, in dark conditions at 18 °C, to reach the 3 h exposition. The contents of the wells were then transferred in 5-mL cytometry polystyrene tubes (Falcon®, BD Biosciences, San Jose, CA, USA).

First, flow cytometry analyses were performed by using a BD FACSVerser flow cytometer using its blue laser (488 nm) as an excitation source. The mean red fluorescence level (LysoTracker fluorescence linked to acidic organelles) of the selected hemocytes was measured using the PerCP-Cy5.5 detector of the flow cytometer (700/54 nm).

Second, 2 µL of a 100X dilution in ultra pure water of a commercial solution of SYBR-Green® I nucleic acid gel stain 10 000X in DMSO (Life Technology, USA) were added in each cytometry tube for 10 min, at room temperature, in dark condition, before a second flow cytometry analysis. Hemocytes were selected according to their green fluorescence (SYBR-Green, FITC detector of the flow cytometer 527/32 nm). Addition of SYBR-Green also allowed to quantify the number of non-adherent hemocytes in our sample.

This second flow cytometry analyses of red fluorescence were performed using a red laser (640 nm) as an excitation source and fluorescence was measured using the APC detector of the flow cytometer (660/10 nm). This detector was selected instead of the PerCP-Cy5.5 detector (700/54 nm) to overcome a possible SYBR-Green fluorescence overlapping. Results are expressed in mean red fluorescence level per hemocyte in arbitrary units (UA).

### 3. Results and discussion

#### 3.1. Transcriptomic results: differential expression analysis

The total number of reads was 29.3 million, with an average length of 262.5 bp, for a total coverage of 5.62X. Differential expression analysis revealed 135 differentially expressed genes, which 90 and 45 appeared down- and up-regulated, respectively, in the EPF treatment (*V. tapetis* infected clams), as compared to the control treatment. We could obtain an annotation, and functional information, for 118 of them, which were classified according to their probable cellular role (Table S1, Table 1). In all, 85% of these transcripts belonged to four functional categories, ie. translation (34 transcripts), cytoskeleton (27), immune response (24) and metabolism (15) (Table 1). The deregulation of these transcripts in clams EPFs appears clearly related to the infection by *V. tapetis*. In the following section, we discuss the pathogen induced modulation of the immune response, which is a major point to better understand the host-pathogen interactions. Table 2 summarizes all the differentially expressed transcripts involved in cytoskeleton or immunity. It is noteworthy that the whole reads were also mapped

**Table 1**

Number and functions of the transcripts differentially expressed in clams EPFs during *V. tapetis* infection.

Functions	Total number of transcripts	Down regulated in infected clams	Up regulated in infected clams
Translation	34	23	11
Cytoskeleton	27	18	9
Immunity	24	18	6
Metabolism	15	9	6
Proteins processing	6	4	2
Unknown/ Hypothetical/Other	29	18	11

against the recently released genome of *V. tapetis* CECT4600 [23], but no significant correspondence has been found with the pathogen.

#### 3.2. Immune response

Extrapallial fluids (EPFs) play an essential role in the biomineralization of the shell of bivalves. Given the particular etiology and symptoms of BRD, this compartment, containing numerous hemocytes [24], is particularly interesting to better understand both the infection process and how the clam's hemocytes fight against it. For this reason, in this study, *V. tapetis* infection was carried out directly inside the EPFs passing through the periostracal lamina barrier to accelerate the immune response, particularly in EPFs.

A transcript encoding lysozyme was found to be upregulated in response to *V. tapetis* infection. In previous studies, lysozyme activities in cell lysates of extrapallial fluids were significantly higher 7 days after *V. tapetis* inoculation into the pallial cavity, as compared to control ones [24]. The antibacterial protein lysozyme, which is circulating in body fluids such as hemolymph and EPFs, is an important line of defense of the animal against bacterial pathogens because of their bacteriolytic properties [5,25].

Only five transcripts directly related to immunity were actually upregulated during *V. tapetis* infection. Beside the lysozyme, three genes belonging to the cyclophilin family, as well as dermatopontin, were upregulated (Table 2). Dermatopontin is a tyrosine-rich acidic matrix protein (TRAMP) that primarily acts as a structural component of the extracellular matrix. However, this protein was shown to modulate TGF-β activity as well as cell-cell aggregation during infection in invertebrates [26]. Dermatopontin was previously shown to be upregulated in the EPFs and hemolymph of clams during BRD [27]. Not less than 3 cyclophilin transcripts appeared upregulated following *V. tapetis* infection. Cyclophilin was shown to be transcriptionally upregulated during infection in the shrimp [28]. In the clam *R. philippinarum*, Chen et al. [29] previously showed that two cyclophilin genes were differentially expressed in hemolymph when exposed to pathogenic bacteria.

The cellular component of the innate immune system also includes more specific pathways that involve the recognition of specific Pathogen-Associated Molecular Patterns (PAMPs) by pattern recognition receptor (PRRs). Most of these receptors are membrane proteins (eg. Toll like receptors), but they also include soluble proteins (eg. SAP, CRP) that are activated by the C1q proteins of the complement system [30–32]. When the pathogen is recognized by either C1q proteins or Mannose Binding Proteins (MBP), these proteins activate the complement pathway through the “classical pathway” or the “lectin pathway” resulting in phagocytosis with the formation of a phagosome [30]. Pathogens are then degraded after the fusion of the phagosome and the lysosome by the action of low pH, acid hydrolases and by reactive oxygen species produced during the pathogen-induced oxidative burst [33].

In this study, 18 out of the 24 deregulated genes potentially involved in clams' immune defenses were downregulated. Nine of these transcripts encoded proteins clearly related to lysosome activity (ie., cathepsins, saposins), four were C1q complement proteins, while others encoded CD63 (a cell death-inducing P53-target protein), galectin 3, as well as two proteins annotated as Defense Proteins, that should be related to ferric iron chelation/reduction. Different transcripts of C1q proteins were also reported in the clam *R. decussatus* infected with *Perkinsus olseni* [34], and in *R. philippinarum* infected with *V. tapetis* [27].

Pathogens have developed many strategies to bypass the immune defenses of their hosts. In the particular case of BRD, it has been shown that the immune system of clams is not able to eliminate *V. tapetis* during infection, even if very high phagocytic rates of *V. tapetis* were measured when the pathogen is injected into the extrapallial space [35]. However, during phagocytosis by hemocytes, *V. tapetis* could persist and multiply within hemocytes and was also able to inhibit the phagosome-lysosome fusion, thus preventing its destruction [6]. The repression of nine genes encoding saposins (2) and cathepsins (7) could reflect the ability of the bacteria to inactivate the lysosomal

**Table 2**

Transcripts significantly regulated during infection by *V. tapetis* and implied in the clam immune system or in actin cytoskeleton organization. FDR: false discovery rate, LogFC: Log Fold-Change.

Down regulated in infected clams			Up regulated in infected clams		
Seq. Description	logFC	FDR	Seq. Description	logFC	FDR
<b>Immune system</b>			<b>Immune system</b>		
cathepsin K-like	−10.52	1.25E-05	complement C1q-like protein 4	11.02	4.29E-04
cathepsin K	−10.04	4.87E-04	cyclophilin-like protein	9.45	3.93E-04
cathepsin L	−9.93	4.16E-04	lysozyme	9.16	4.87E-04
cathepsin L1	−9.81	3.31E-04	cyclophilin A	7.60	8.56E-05
cathepsin S	−9.77	2.27E-11	dermatopontin	6.29	8.88E-04
saposin B domain-containing protein	−9.42	1.79E-11	cyclophilin-like protein	4.91	9.48E-04
saposin B domain-containing protein precursor	−8.74	2.95E-09			
cell death-inducing p53-target protein 1	−8.46	2.95E-09			
cathepsin S	−8.22	2.50E-05			
ferric-chelate reductase 1	−8.08	4.99E-04			
CD63 antigen	−7.51	1.44E-04			
defense protein 3-like	−7.30	5.66E-04			
complement C1q-like protein 4	−7.24	4.50E-05			
galectin-3 isoform 1	−7.20	6.22E-04			
c1q domain containing protein 1q13	−6.89	1.83E-06			
complement c1q TNF-related protein 3-like isoform 2	−6.34	3.82E-07			
complement c1q-like protein 2 precursor	−5.76	2.63E-04			
saposin B domain-containing protein	−5.04	1.25E-05			
<b>Actin cytoskeleton organization</b>			<b>Actin cytoskeleton organization</b>		
actin	−11.30	1.49E-04	actin	12.46	7.50E-05
actin	−11.21	8.94E-05	cytoplasmic actin	10.82	1.46E-04
actin	−10.49	5.02E-05	actin	10.82	1.49E-04
cytoplasmic actin	−10.18	4.81E-08	kinesin	10.33	4.34E-08
myophillin	−10.12	1.98E-05	actin	8.84	9.99E-04
cytoplasmic actin	−10.09	5.50E-05	thymosin beta-12	8.76	4.93E-04
coactosin-like protein	−9.73	2.07E-05	actin	8.57	6.14E-04
beta-actin	−8.86	1.27E-04	cytoplasmic actin	8.34	6.65E-04
cytoplasmic actin	−8.30	2.95E-09	myosin light chain	5.57	1.98E-05
actin 5c	−7.96	4.99E-04			
cytoplasmic actin	−7.45	4.99E-04			
cytoplasmic actin	−7.37	7.39E-04			
coactosin-like protein	−6.94	4.93E-04			
cytoplasmic actin	−6.84	2.82E-07			
cysteine and glycine-rich protein 3	−6.80	5.97E-04			
profilin	−6.63	9.08E-04			
cysteine and glycine-rich protein 3	−6.63	8.28E-04			
dynein light chain, cytoplasmic	−6.52	9.99E-04			

functioning. Indeed, Cathepsins K, L, L1 and S are cysteine proteases [36,37] that belong to lysosomal acid hydrolases, that are normally released during the phagosome-lysosome fusion. Saposins are small lysosomal proteins that participate (stimulate/activate) to the hydrolysis of sphingolipids and many other lipids [38]. Interestingly, the CD63 antigen belongs to the family of lysosomal associated proteins (LAMPs) and is also considered as an indicator of the phagosome-lysosome fusion [39]. In all, these results strongly suggest that lysosomal function plays a key role in the development of BRD.

In addition to the predominant role of lysosome inactivation, the repression of genes encoding three complement C1q/tumor necrosis factor proteins, and of the galectin 3, might suggest an important role for the complement- and lectin-dependent PRR pathways [40,41]. The downregulation of a cell death-inducing P53-target protein, which is known to regulate the TNF-alpha-mediated apoptosis, might also suggest a role for apoptotic processes during the infectious process [42].

Finally, the downregulation of a protein annotated as ‘defense protein’, that contains a ferric-chelate reductase domain, could reflect an important role for iron acquisition during the infectious process. Interestingly, the concomitant downregulation of ferritin, which is implied in iron storage [43] and reduction of reactive oxygen species (ROS) accumulation [44], could also be related to this, allowing then more availability of iron for the pathogen.

### 3.3. Ribosomal proteins

We also identified deregulated genes involved in functions other than immunity, but whose deregulation could strongly influence the host's immune response. This is especially the case for cytoskeleton reorganization, thoroughly discussed thereafter, but also for translation proteins, which is the most modified category and, in other ways, for genes encoding proteins which function was categorized in metabolism and protein processes (Table 1).

A major reorganization of the translational apparatus may have occurred in response to *V. tapetis* infection as a result of gene expression reprogramming (32 ribosomal proteins and 2 translation elongation factors). Eleven genes were induced, while 23 genes had their expression reduced (Table 1). Ribosome inactivation resemble one of the major symptoms of the so-called ‘metabolic depression’ observed in many environmental stress responses in several marine mollusks [45,46]. However, it is noteworthy that the expression of several ribosomal proteins was increased during infection (S14, S21, S30, L21, SA, P0), and others appeared both repressed and induced (L3, L8, L27). For these last proteins, we presume that the protein exists as different alternate isoforms (with different regulations).

It is noteworthy that ribosomal proteins are increasingly studied for their alternate roles, ie. independent of their ribosomal function [45]. More specifically, it should be noted that the role of ribosomal proteins




Fig. 1. Adherence test on hemocytes exposed to *Vibrio tapetis* CECT4600 (black) or non-exposed (grey) representing the mean ratio of non-adherent hemocytes in exposed/control samples. Error bars = Standard error.

in the regulation of the innate immune response has been well documented in recent years [45], especially the major role of the ribosomal protein L13A in the GAIT complex, that selectively modulates the translation of some genes involved in the interferon- $\gamma$ -mediated inflammatory response [47]. However, many other examples have been reported. For example, the ribosomal proteins S14 and L8, both found upregulated in our study, would be able to activate P53 and the major regulator NF- $\kappa$ B, respectively [48,49]. It would be particularly interesting to perform new focused investigations to better understand the role of the translation apparatus in the development of the BRD.

### 3.4. Actin cytoskeleton organization

Actin filaments (F-actin) belong to the cell cytoskeleton and are dynamically polymerized and depolymerized from monomers of actin (G-actin) into the cell to enable cellular activities such as locomotion, exocytosis or phagocytosis. The actin cytoskeleton is also responsible for the cell shape. In this study, we showed that at least 27 transcripts involved in actin cytoskeleton organization were differentially expressed during infection, among which 9 were induced while 18 had their expression reduced. Genes downregulated during infection are mainly involved in actin filament stabilization such as myophillin, a member of calponin protein family [50,51], cysteine glycine rich protein (CSR1) [52,53] and dynein light chain [54]. Furthermore, our results showed that expression regulation during infection also affected transcripts involved in cell motility such as up regulation of the Myosin light chain, which role in hemocytes phagocytosis in shrimp was already described [55].

Two transcripts downregulated during BRD might give additional information on the molecular mechanisms related to infection. The first one encodes a Coactosin like protein that counteracts the action of capping proteins that reduce or slow down actin polymerization [56]. The second one encodes Profilin, an ‘actin binding protein’ (ABP) which is known to interact with others ABPs such as  $\beta$ 4-thymosin [57]. The role of profilin in actin polymerization is quite complex. In the presence of molecules that stimulate actin assembly, profilin displaces  $\beta$ 4-thymosin, binds to G-actin and promotes actin polymerization [58]. Furthermore, profilin regulates the effects of capping proteins. Hopmann and Miller [59] demonstrated that balanced activities of capping protein and profilin are essential in the regulation of actin dynamics and organization in *Drosophila*.

Capping proteins can be associated, in resting cells, with actin filament by electrostatic interactions on the barbed (+) end. By changing the conformation of the capping protein during this interaction, the + end is no longer available for G-actin and polymerization is then inhibited [60,61].

In this study, we also highlighted an overexpression of  $\beta$ 12 thymosin transcripts during infection by *V. tapetis*.  $\beta$ 12-thymosin belongs to the ABPs family. Fifteen variants of  $\beta$ -thymosin proteins have been described in many vertebrates and invertebrates, but not on prokaryotes and yeasts. They are able to bind ATP-G-actin complex (1:1), the monomeric form of actin, and stabilized it. This actin-monomer sequestration prevents them from polymerization by changes in actin conformation [60,62]. Many factors are actually involved in the dynamics of actin in eukaryotic cells, such as e.g. the ATP concentration or the Critical Concentration ( $C_c$ ) of G-actin monomers.

### 3.5. Physiological verification of differential expression analysis

Transcriptomics analysis is a very powerful tool to give, in a single experiment, a broad overview of the multiple processes potentially involved in a biological response. In our study, we showed the deregulation of many transcripts encoding proteins involved in immunity and in the dynamics of actin cytoskeleton, thus suggesting an important role for these biological processes during infection of *R. philippinarum* by *V. tapetis*. Therefore, we performed additional experiments in order to check the validity of some of the major interpretations raised from the differential expression analysis.

First, RNA-seq data showed a downregulation of many transcripts encoding proteins involved in the lysosomal activity, thus suggesting a crucial of related biological process during infection. We thus aimed at checking the actual effect of *V. tapetis* exposure on the lysosomal activity of *R. philippinarum* hemocytes, by using the Lysotracker assay, which is commonly recognized as an efficient proxy of the amount and activity of acidic organelles within eukaryotic cells.

As previously described [7,8], we first confirmed that the exposition to *V. tapetis* CECT4600 induces a loss of hemocytes adherence as compared to the control (Fig. 1), and results in a rounded phenotype. In these experimental conditions, we observed that the loss of adherence is accompanied by a two-fold decrease of acidic organelles in hemocytes exposed to *V. tapetis* CECT4600 as compared to the control (Fig. 2). As a




Fig. 2. Mean red fluorescence of Lysotracker red used as a measure of the amount of acidic organelles in hemocytes, exposed to *V. tapetis* CECT4600 (black) or non-exposed (grey). Error bars = Standard error.

consequence, we hypothesize that the general depletion of lysosomal transcripts observed in the RNAseq experiment should result from the decreased number of lysosomes in EPFs-derived hemocyte cells.

Second, as many transcripts involved in actin-cytoskeleton dynamics were deregulated, we aimed at observing the F-actin filaments in hemolymph that had been exposed or not to *V. tapetis* (Fig. 3), by performing a phalloidin staining experiment. Results presented in Fig. 3 clearly show the loss of pseudopods and rounding of hemocytes after *V. tapetis* exposure (Fig. 3C and D). In these conditions, it is noteworthy that the F-actin filaments were clearly mainly located at the periphery of cells in healthy hemocytes (Fig. 3B), while phalloidin staining revealed a diffuse localization of F-actin in cells exposed to *V. tapetis* (Fig. 3D). These results are consistent with previous works showing F-actin network disorganization in diseased hemocytes in another clam species, *Mya arenaria* [63], as well as with the infection model developed by Paillard [6] considering that *V. tapetis* induces a loss of phagosome-lysosome formation during infection of the Manila clam *R. philippinarum*.

### 3.6. Hypothetical model of clam-*Vibrio tapetis* interaction

We took advantage of our data to propose an hypothetical model that encompasses bacterium recognition (through microbe-associated molecular pattern [MAMP]) and some relevant metabolic processes over-represented in the transcriptomic profile of clams EPFs in response to *V. tapetis*. Altogether, these results are consistent with the knowledge we have about the influence of the BRD on actin cytoskeleton [13]. Indeed, in the case of *R. philippinarum*'s infection by *V. tapetis*, the bacterium is phagocytosed into clam's hemocytes which leads to hemocytes actin-cytoskeleton disorganization. The result of this perturbation is the loss of pseudopods (ie. rounding phenotype) and then the loss of hemocytes adhesion properties. This consequence of *V. tapetis* infection on clams is commonly used to measure strains virulence by *in vitro* characterization [7].

The fact that transcripts such as coactosin or profilin, that interact with either capping proteins and/or ABPs, are downregulated during infection, let us to formulate different hypothesis. Considering our results and the knowledge on actin dynamics, we made the hypothesis





Fig. 3. Rhodamine phalloidin staining of F-actin filaments in hemocytes that had been exposed (C, D) or not (A, B) to *V. tapetis* CECT4600. A, C: bright field microscopy observations of control (A) and exposed (C) hemocytes. B, D: epifluorescence observations of control (B) and exposed (D) hemocytes. Wavelengths: excitation, 550 nm; emission, 605 nm. The scale bars (2 cm = 10 μm) correspond to the pictures A, B, C & D.

that infection might trigger a deregulation of capping protein which would lead hemocytes to be blocked in a “resting cell state”. According to our model, capping proteins would block the + end polymerization in infected hemocytes, then increasing the levels of free G-actin [60]. In this case, Mannherz and Hannappel [60] described that in case of resting cells,  $\beta$ -thymosins massively bound to free G-actin, thus decreasing its concentrations, thus stalling net polymerization [60]. On the other hand, they also report that during this process, profilin concentration is low and most of it is inactivated. This hypothesis would need additional experiments but might explain the disorganization of the F-actin network observed during infection that gives to hemocytes this ‘rounded’ phenotype after challenge with *V. tapetis*.




**Fig. 4.** The hypothetical model of clam – *V. tapetis* interaction. a) Immune response of clams to foreign bacteria in healthy, functional and active hemocytes. b) Immune response of clams to *V. tapetis* on the first steps of infection in early Infected Hemocytes. c) Immune response of clams to *V. tapetis* on the late steps of infection in infected, non-functional and, according to our model, resting hemocytes. (-) represent down regulated genes in this study.

Fig. 4A to C represent the conceptual model of the Manila clam EPF infection by *V. tapetis* in order to understand functions and interactions between the proteins that were deregulated in our experiments, and their place in the immune response to infection and actin dynamics polymerization process. Fig. 4A represents the immune response of healthy hemocytes to foreign bacteria. Foreign bacteria are opsonized and phagocytized by hemocyte. After fusion between the phagosome and the lysosome, bacteria are destroyed. Fig. 4B represents early stage of hemocytes infection by *V. tapetis* and thus downregulation of transcripts related to pathogen recognition (C1q proteins and lectin) or to decrease of lysosomal activity as demonstrated in this study. Fig. 4C represents the deregulation of the immune system in hemocytes

infected by *V. tapetis* inducing a rounding phenotype related to infection with absence of pseudopods formation.

**4. Conclusion**

This study presents the first transcriptomic approach on EPFs of experimentally infected clams. It brought novel insights on the interaction between *V. tapetis* and clam by exploring *in vivo* gene expression during the infection. This study revealed a downregulation of the immune response, especially the complement pathway and lysosomal activity, the disorganization of the actin cytoskeleton, accompanied by ribosomal genes deregulation.



**Fig. 4.** (continued)




Fig. 4. (continued)

## Fundings

This project received grants from the H2020 European project “VIVALDI” (grant agreement N°678589), CNRS, France INSU EC2CO-MicrobiEn “MICROPAL”, CAPES-COFEUCUB (MICROBIV) and PVE Brésil. This work was also supported by the ‘Université de Bretagne Occidentale’ (UBO, France), and the “investment for the future” programs LabexMER (ANR-10-LABX-19) and ISblue (ANR-17-EURE-0015).

## Author contributions

CP, FT and VP acquired funds and coordinated the study. GR, AB and CP designed and performed the experimental infection. AB performed RNA extractions and libraries preparations, with the advices of LO. LO and FT performed DNA sequencing. AR and EC performed bioinformatics analysis, with inputs from VP. AR and CL performed the flow cytometry analyses. AR, VP and CP wrote the article (the original draft was written by AR). The article was carefully reviewed by other co-authors, who all approved the final version.

## Acknowledgments

We warmly thank Fabien Fonteneau (EARL Les Claires de Bonsonge, Marennes, France) and Jean François Auvray (SATMAR Marennes, France) for providing the clams. We thank colleagues from the Federal University of Rio De Janeiro, Brazil who welcomed us (CP, VP, AB, GR) in Rio. We thank N. Le Goïc, F. Nunez, H. Hégaret, C. Fabioux, I. Paul-Pont, F. Le Grand, F. Riera, AL. Cassone, P. Le Souchu, A. Muir, E. Harney, Y. Epelboin, M. Fuhrmann, M. Provost, L. Frère, F. Boullot, B. Dubief, M. Czamanski, R. Morvezzen, V. Foulon, M. Protat, C. Quéré, and A. Huvet for their help for animals conditioning, injections and samplings.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fsi.2019.08.025>.

## References

- [1] C. Paillard, Etiologie et caractérisation de la maladie de l’anneau brun chez la palourde d’élevage *Ruditapes philippinarum*, PhD thesis Université de Bretagne Occidentale, France, 1992 194p.
- [2] C. Paillard, L. Perceley, M. Le Pennec, D. Le Picard, Origine pathogène de l’«anneau brun» chez *Tapes philippinarum* (Mollusque, bivalve), Comptes Rendus de l’Académie Des Sciences. Série 3, Sciences de La Vie 309 (1989) 235–241.
- [3] P. Maes, C. Paillard, Effect of the *Vibrio* P1, pathogene de *Ruditapes philippinarum*, sur d’autres espèces de bivalves, Les Mollusques Marins: Biologie et Aquaculture, Brest (France), 14 1992, pp. 141–148.
- [4] J.J. Borrego, D. Castro, A. Luque, C. Paillard, P. Maes, M.T. Garcia, A. Ventosa, *Vibrio tapetis* sp. nov., the causative agent of the brown ring disease affecting cultured clams, Int. J. Syst. Evol. Microbiol. 46 (1996) 480–484.
- [5] C. Paillard, B. Allam, R. Oubella, Effect of temperature on defense parameters in Manila clam *Ruditapes philippinarum* challenged with *Vibrio tapetis*, Dis. Aquat. Org. 59 (2004) 249–262.
- [6] Paillard, Rôle de l’environnement dans les interactions hôtes-pathogènes; développement d’un modèle de vibriose chez les bivalves, Habilitation à Diriger Des Recherches (HDR), Université de Bretagne Occidentale, Brest, 2004.
- [7] G. Choquet, P. Soudant, C. Lambert, J.-L. Nicolas, C. Paillard, Reduction of adhesion properties of *Ruditapes philippinarum* hemocytes exposed to *Vibrio tapetis*, Dis. Aquat. Org. 57 (2003) 109–116.
- [8] F. Jeffroy, C. Paillard, Involvement of nitric oxide in the *in vitro* interaction between Manila clam, *Ruditapes philippinarum*, hemocytes and the bacterium *Vibrio tapetis*, Fish Shellfish Immunol. 31 (2011) 1137–1141, <https://doi.org/10.1016/j.fsi.2011.10.008>.
- [9] A.F. Rowley, A. Powell, Invertebrate immune systems-specific, quasi-specific, or nonspecific? J. Immunol. 179 (2007) 7209–7214, <https://doi.org/10.4049/jimmunol.179.11.7209>.
- [10] B. Allam, S.E. Ford, Effects of the pathogenic *Vibrio tapetis* on defence factors of susceptible and non-susceptible bivalve species: I. Haemocyte changes following *in vitro* challenge, Fish Shellfish Immunol. 20 (2006) 374–383, <https://doi.org/10.1016/j.fsi.2005.05.012>.
- [11] C. Paillard, An Ecological Approach to Understanding Host-Pathogen-Environment Interactions: the Case of Brown Ring Disease in Clams, in: J.L. Romalde (Ed.), Oysters and clams: cultivation, habitat threats and ecological impact, 7 Microbiology Research Advances, Nova Publishers, 2016228p.
- [12] E. Lane, T.H. Birkbeck, Species specificity of some bacterial pathogens of bivalve molluscs is correlated with their interaction with bivalve haemocytes, J. Fish Dis. 23 (2000) 275–279.
- [13] F. Brulle, F. Jeffroy, S. Madec, J.-L. Nicolas, C. Paillard, Transcriptomic analysis of *Ruditapes philippinarum* hemocytes reveals cytoskeleton disruption after *in vitro* *Vibrio tapetis* challenge, Dev. Comp. Immunol. 38 (2012) 368–376, <https://doi.org/10.1016/j.dci.2012.03.003>.
- [14] G. Richard, Approche mécanistique de la réponse de la palourde japonaise, *Ruditapes philippinarum*, exposée à la bactérie *Vibrio tapetis*: influence de la température et du régime algal, PhD thesis Université de Bretagne Occidentale, France, 2015 322p.
- [15] C. Le Bris, G. Richard, C. Paillard, C. Lambert, C. Segueineau, O. Gauthier, F. Pernet, F. Guérard, Immune responses of phenoloxidase and superoxide dismutase in the manila clam *Venerupis philippinarum* challenged with *Vibrio tapetis* – Part I: spatio-temporal evolution of enzymes’ activities post-infection, Fish Shellfish Immunol. 42 (2015) 16–24, <https://doi.org/10.1016/j.fsi.2014.10.021>.
- [16] C. Paillard, P. Maes, Brown ring disease in the Manila clam *Ruditapes philippinarum*: establishment of a classification system, Dis. Aquat. Org. 19 (1994) 137–146.
- [17] S. Lindgreen, AdapterRemoval: easy cleaning of next-generation sequencing reads, BMC Res. Notes 5 (2012) 337.
- [18] A.M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data, Bioinformatics 30 (2014) 2114–2120, <https://doi.org/10.1093/bioinformatics/btu170>.
- [19] R. Schmieder, R. Edwards, Quality control and preprocessing of metagenomic datasets, Bioinformatics 27 (2011) 863–864, <https://doi.org/10.1093/bioinformatics/btr026>.
- [20] E. Kopylova, L. Noé, H. Touzet, SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data, Bioinformatics 28 (2012) 3211–3217, <https://doi.org/10.1093/bioinformatics/bts611>.
- [21] R. Moreira, P. Balseiro, J.V. Planas, B. Fuste, S. Beltran, B. Novoa, A. Figueras, Transcriptomics of *in vitro* immune-stimulated hemocytes from the Manila clam *Ruditapes philippinarum* using high-throughput sequencing, PLoS One 7 (2012) e35009.
- [22] M.G. Grabherr, B.J. Haas, M. Yassour, J.Z. Levin, D.A. Thompson, I. Amit, X. Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, et al., Full-length transcriptome assembly from RNA-Seq data without a reference genome, Nat. Biotechnol. 29 (2011) 644–652, <https://doi.org/10.1038/nbt.1883>.
- [23] G.M. Dias, A. Bidault, P. Le Chevalier, G. Choquet, C. Der Sarkissian, L. Orlando, C. Medigue, V. Barbe, S. Manganot, C.C. Thompson, *Vibrio tapetis* displays an original type IV secretion system in strains pathogenic for bivalve molluscs, Front. Microbiol. 9 (2018) 227.
- [24] B. Allam, C. Paillard, M. Auffret, Alterations in hemolymph and extrapallial fluid parameters in the Manila clam, *Ruditapes philippinarum*, challenged with the pathogen *Vibrio tapetis*, J. Invertebr. Pathol. 76 (2000) 63–69, <https://doi.org/10.1006/jipa.2000.494>.
- [25] C. López, M.J. Carballeda, C. Azevedo, A. Villalba, Enzyme characterisation of the circulating haemocytes of the carpet shell clam, *Ruditapes decussatus* (Mollusca:bivalvia), Fish Shellfish Immunol. 7 (1997) 595–608, <https://doi.org/10.1006/fsim.1997.0112>.
- [26] G. Wang, Z. Zhang, S. Lin, L. Zhang, B. Wang, S. Wang, Y. Wang, A homologue of dermatopontin from *Haliotis diversicolor* and its response to pathogenic infection, Aquacult. Res. 46 (2015) 1537–1549, <https://doi.org/10.1111/are.12305>.
- [27] B. Allam, E. Pales Espinosa, A. Tanguy, F. Jeffroy, C. Le Bris, C. Paillard, Transcriptional changes in manila clam (*Ruditapes philippinarum*) in response to Brown ring disease, Fish Shellfish Immunol. 41 (2014) 2–11, <https://doi.org/10.1016/j.fsi.2014.05.022>.
- [28] L. Qiu, S. Jiang, J. Huang, W. Wang, C. Zhu, T. Su, Molecular cloning and mRNA expression of cyclophilin A gene in black tiger shrimp (*Penaeus monodon*), Fish Shellfish Immunol. 26 (2009) 115–121, <https://doi.org/10.1016/j.fsi.2008.03.022>.
- [29] L. Chen, C. Mu, J. Zhao, C. Wang, Molecular cloning and characterization of two

- isoforms of cyclophilin A gene from *Venerupis philippinarum*, Fish Shellfish Immunol. 31 (2011) 1218–1223, <https://doi.org/10.1016/j.fsi.2011.07.001>.
- [30] S.S. Bohlsón, S.D. O'Conner, H.J. Hulsebus, M.-M. Ho, D.A. Fraser, Complement, C1q, and C1q-related molecules regulate macrophage polarization, Front. Immunol. (2014) 5, <https://doi.org/10.3389/fimmu.2014.00402>.
- [31] T.W. Du Clos, C. Mold, Pentraxins (CRP, SAP) in the process of complement activation and clearance of apoptotic bodies through Fc $\gamma$  receptors, Curr. Opin. Organ Transplant. 16 (2011) 15–20, <https://doi.org/10.1097/MOT.0b013e32834253c7>.
- [32] W.-C. Song, Crosstalk between complement and toll-like receptors, Toxicol. Pathol. 40 (2012) 174–182, <https://doi.org/10.1177/0192623311428478>.
- [33] P. Parham, C. Janeway, The Immune System, fourth ed., Garland Science, Taylor & Francis Group, New York, NY, 2015.
- [34] R.B. Leite, M. Milan, A. Coppe, S. Bortoluzzi, A. dos Anjos, R. Reinhardt, C. Saavedra, T. Patarnello, M. Cancela, L. Bargelloni, mRNA-Seq and microarray development for the Grooved carpet shell clam, *Ruditapes decussatus*: a functional approach to unravel host-parasite interaction, BMC Genomics 14 (2013) 741, <https://doi.org/10.1186/1471-2164-14-741>.
- [35] B. Allam, C. Paillard, S.E. Ford, Pathogenicity of *Vibrio tapetis*, the etiological agent of brown ring disease in clams, Dis. Aquat. Org. 48 (2002) 221–231.
- [36] J. Liu, G.-P. Shi, W.-Q. Zhang, G.-R. Zhang, W.-H. Xu, Cathepsin L function in insect moulting: molecular cloning and functional analysis in cotton bollworm, *Helicoverpa armigera*, Insect Mol. Biol. 15 (2006) 823–834, <https://doi.org/10.1111/j.1365-2583.2006.00686.x>.
- [37] D. Niu, K. Jin, L. Wang, B. Feng, J. Li, Molecular characterization and expression analysis of four cathepsin L genes in the razor clam, *Sinonovacula constricta*, Fish Shellfish Immunol. 35 (2013) 581–588, <https://doi.org/10.1016/j.fsi.2013.06.001>.
- [38] A. Darmoise, P. Maschmeyer, F. Winau, The immunological functions of saposins, Advances in Immunology, Elsevier, 2010, pp. 25–62, [https://doi.org/10.1016/S0065-2776\(10\)05002-9](https://doi.org/10.1016/S0065-2776(10)05002-9).
- [39] A. Welin, J. Raffetseder, D. Eklund, O. Stendahl, M. Lerm, Importance of phagosomal functionality for growth restriction of *Mycobacterium tuberculosis* in primary human macrophages, Journal of Innate Immunity 3 (2011) 508–518, <https://doi.org/10.1159/000325297>.
- [40] U. Kishore, C. Gaboriaud, P. Waters, A.K. Shrive, T.J. Greenhough, K.B.M. Reid, R.B. Sim, G.J. Arlaud, C1q and tumor necrosis factor superfamily: modularity and versatility, Trends Immunol. 25 (2004) 551–561, <https://doi.org/10.1016/j.it.2004.08.006>.
- [41] G.R. Vasta, M. Quesenberry, H. Ahmed, N. O'Leary, C-type lectins and galectins mediate innate and adaptive immune functions: their roles in the complement activation pathway, Dev. Comp. Immunol. 23 (1999) 401–420, [https://doi.org/10.1016/S0145-305X\(99\)00020-8](https://doi.org/10.1016/S0145-305X(99)00020-8).
- [42] N.D. Marchenko, A. Zaika, U.M. Moll, Death Signal-induced Localization of p53 Protein to Mitochondria a potential role in apoptotic signaling, J. Biol. Chem. 275 (2000) 16202–16212, <https://doi.org/10.1074/jbc.275.21.16202>.
- [43] P. Ponka, C. Beaumont, D.R. Richardson, Function and regulation of transferrin and ferritin, Semin. Hematol. 35 (1998) 35–54.
- [44] K. Orino, L. Lehman, Y. Tsuji, H. Ayaki, S.V. Torti, F.M. Torti, Ferritin and the response to oxidative stress, Biochem. J. 357 (2001) 241–247, <https://doi.org/10.1042/bj3570241>.
- [45] X. Zhou, W.-J. Liao, J.-M. Liao, P. Liao, H. Lu, Ribosomal proteins: functions beyond the ribosome, J. Mol. Cell Biol. 7 (2015) 92–104, <https://doi.org/10.1093/jmcb/mjv014>.
- [46] M. Achard-Joris, P. Gonzalez, V. Marie, M. Baudrimont, J.-P. Bourdineaud, cDNA cloning and gene expression of ribosomal S9 protein gene in the mollusk *Corbicula fluminea*: a new potential biomarker of metal contamination up-regulated by cadmium and repressed by zinc, Environ. Toxicol. Chem. 25 (2006) 527, <https://doi.org/10.1897/05-211R.1>.
- [47] B. Mazumder, D. Poddar, A. Basu, R. Kour, V. Verbovetskaya, S. Barik, Extraribosomal L13a is a specific innate immune factor for antiviral defense, J. Virol. 88 (2014) 9100–9110, <https://doi.org/10.1128/JVI.01129-14>.
- [48] L. Wang, J. Luo, Q. Nian, Q. Xiao, Z. Yang, L. Liu, Ribosomal protein S14 silencing inhibits growth of acute myeloid leukemia transformed from myelodysplastic syndromes via activating p53, Hematology 19 (2014) 225–231, <https://doi.org/10.1179/1607845413Y.0000000127>.
- [49] H.-X. Zhang, Z.-X. Liu, Y.-P. Sun, J. Zhu, S.-Y. Lu, X.-S. Liu, Q.-H. Huang, Y.-Y. Xie, H.-B. Zhu, S.-Y. Dang, H.-F. Chen, G.-Y. Zheng, Y.-X. Li, Y. Kuang, J. Fei, S.-J. Chen, Z. Chen, Z.-G. Wang, Rig-I regulates NF- $\kappa$ B activity through binding to Nf-b1 3'-UTR mRNA, Proc. Natl. Acad. Sci. 110 (2013) 6459–6464, <https://doi.org/10.1073/pnas.1304432110>.
- [50] R.M. Martin, N.B. Chilton, M.W. Lightowlers, R.B. Gasser, *Echinococcus granulosus* myophilin—relationship with protein homologues containing “calponin-motifs, Int. J. Parasitol. 27 (1997) 1561–1567.
- [51] A. Goodman, B.L. Goode, P. Matsudaira, G.R. Fink, The *Saccharomyces cerevisiae* calponin/transgelin homolog Scp1 functions with fimbrin to regulate stability and organization of the actin cytoskeleton, Mol. Biol. Cell 14 (2003) 2617–2629, <https://doi.org/10.1091/mbc.e03-01-0028>.
- [52] T.C. Tran, C. Singleton, T.S. Fraley, J.A. Greenwood, Cysteine-rich protein 1 (CRP1) regulates actin filament bundling, BMC Cell Biol. 6 (2005) 45, <https://doi.org/10.1186/1471-2121-6-45>.
- [53] H.A. Louis, J.D. Pino, K.L. Schmeichel, P. Pomiès, M.C. Beckerle, Comparison of three members of the cysteine-rich protein family reveals functional conservation and divergent patterns of gene expression, J. Biol. Chem. 272 (1997) 27484–27491, <https://doi.org/10.1074/jbc.272.43.27484>.
- [54] A. Ghosh-Roy, B.S. Desai, K. Ray, Dynein light chain 1 regulates dynamin-mediated F-actin assembly during sperm individualization in *Drosophila*, Mol. Biol. Cell 16 (2005) 3107–3116, <https://doi.org/10.1091/mbc.e05-02-0103>.
- [55] F. Han, Z. Wang, X. Wang, Characterization of myosin light chain in shrimp hemocytic phagocytosis, Fish Shellfish Immunol. 29 (2010) 875–883, <https://doi.org/10.1016/j.fsi.2010.07.030>.
- [56] U. Röhrig, G. Gerisch, L. Morozova, M. Schleicher, A. Wegner, Coactosin interferes with the capping of actin filaments, FEBS (Fed. Eur. Biochem. Soc.) Lett. 374 (1995) 284–286, [https://doi.org/10.1016/0014-5793\(95\)01130-7](https://doi.org/10.1016/0014-5793(95)01130-7).
- [57] E. Korenbaum, P. Nordberg, C. Björkregren-Sjögren, C.E. Schutt, U. Lindberg, R. Karlsson, The role of profilin in actin polymerization and Nucleotide exchange  $\dagger$ , Biochemistry 37 (1998) 9274–9283, <https://doi.org/10.1021/bi9803675>.
- [58] H. Lodish (Ed.), Molecular Cell Biology, 4, Freeman, New York, NY, 2002[Nachdr.].
- [59] R. Hopmann, K.G. Miller, A balance of capping protein and profilin functions is required to regulate actin polymerization in *Drosophila* bristle, Mol. Biol. Cell 14 (2003) 118–128, <https://doi.org/10.1091/mbc.E02-05-0300>.
- [60] H.G. Mannherz, E. Hannappel, The  $\beta$ -thymosins: intracellular and extracellular activities of a versatile actin binding protein family, Cell Motil. Cytoskelet. 66 (2009) 839–851, <https://doi.org/10.1002/cm.20371>.
- [61] A. Weber, Actin binding proteins that change extent and rate of actin monomer-polymer distribution by different mechanisms, Mol. Cell. Biochem. 190 (1999) 67–74.
- [62] M. Hertzog, E.G. Yarmola, D. Didry, M.R. Bubb, M.-F. Carlier, Control of actin dynamics by proteins made of  $\beta$ -thymosin repeats - the actobindin family, J. Biol. Chem. 277 (2002) 14786–14792, <https://doi.org/10.1074/jbc.M112064200>.
- [63] C.A. Moore, N. Beckmann, M. Patricia Morse, Cytoskeletal structure of diseased and normal hemocytes of *Mya arenaria*, J. Invertebr. Pathol. 60 (1992) 141–147, [https://doi.org/10.1016/0022-2011\(92\)90087-K](https://doi.org/10.1016/0022-2011(92)90087-K).