



Full length article

Gene expression analysis of the innate immune system during early rearing and weaning of meagre (*Argyrosomus regius*)Cindy Campoverde^a, Douglas J. Milne^b, Christopher J. Secombes^b, Alicia Estévez^a, Enric Gisbert^a, Karl B. Andree^{a,*}^a IRTA, San Carlos de La Rápita, 43540, Tarragona, Spain^b Scottish Fish Immunology Research Centre, University of Aberdeen, Zoology Building, Tillydrone Avenue, Aberdeen AB24 2TZ, Scotland, UK

ARTICLE INFO

Keywords:

Innate immunity
Argyrosomus regius
 Meagre
 PAMP
 Ontogeny

ABSTRACT

The present study is the first report of some representative innate immune genes in meagre (*Argyrosomus regius*) larvae. This study has specifically focused on the growth period from hatching to the juvenile stage, a critical time in marine fish development when reliance on innate immune mechanisms are required for survival. We report molecular cloning of partial open reading frames and expression patterns for some innate immune genes (*c3*, *cox2*, *met*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3*). In addition, phylogenetic analyses of some of the sequences obtained was performed where confusion among closely allied isoforms may have existed. These results show the *met* isoform from meagre is *met II*, an isoform more similar to a homolog described in *Larimichthys crocea*; lysozyme (*lyzc*) corresponds to the c-type and NOD isoforms (*nod2*, *nod3*) separate into different clades confirming their distinctness within a common evolutionary history. Gene expression profiles of innate genes were investigated, for nine developmental stages, from 8 days post-hatching (dph) to 120 dph. Present results demonstrated that *c3*, *cox2*, *met II*, *lyzc*, *mxp*, *myd88*, *nod2*, and *nod3* were expressed in all stages of larval development and displayed distinct expression profiles in separate tissues (kidney, spleen gut and gill). Moreover, expression patterns suggested these innate immune genes may be influenced by feeding practices, *i.e.* switching from live prey (rotifer and *Artemia*) and weaning onto an inert commercial diet. In addition to evaluating changes in gene expression during early development, this study evaluated the modulation of gene expression by means of *in vivo* trials in juveniles that were stimulated with PAMPs (LPS, poly I:C, β-glucan). These results revealed significant changes in mRNA levels of target genes in the kidney, spleen, gut and gills. However, expression profiles differed in magnitude depending on the stimulant and/or tissue. These results are discussed in terms of their relevance and potential application in aquaculture practices.

1. Introduction

In recent decades, the development of the aquaculture industry has been achieved by important advances in production techniques, as well as the introduction of new species [1]. To achieve greater sustainability for the industry, more enhanced production is needed to boost efficiency; thus, proper health management is a key issue in actual fish farming operations. One research area that could improve health management is vaccine development, among other preventive measures (*i.e.* functional feeds and/or therapeutic agents). However, in addition to the significant research and financial investments required for vaccine development for each particular infectious agent, there is a requirement for proper understanding of the immune functions and timing of their development in order to synchronize rearing practices

(*i.e.* vaccination) to the stage of fish development.

Meagre (*Argyrosomus regius*) is an emerging species in aquaculture and is currently receiving a lot of attention within European region. Meagre can be found throughout the Mediterranean and Black Seas, as well as in the East Atlantic coasts [2]. As is common in production of a new species in aquaculture, the artificial rearing of meagre larvae needs to be optimized to improve survival rate in an effort to reduce larval rearing costs [3]. The successful culture of larvae from marine species not only requires optimized rearing and feeding protocols based on the species' nutritional requirements and digestive capabilities, but it also relies on an understanding of the function and timing of maturation of the immune system to aid management of the occurrence of infectious disease, which cause high losses in aquaculture, especially during early development [4].

* Corresponding author.

E-mail address: karl.andree@irta.es (K.B. Andree).<https://doi.org/10.1016/j.fsi.2019.10.009>

Received 26 August 2019; Received in revised form 3 October 2019; Accepted 5 October 2019

Available online 06 October 2019

1050-4648/ © 2019 Published by Elsevier Ltd.

Fish larvae are not capable of eliciting a specific immune response during early development since immune memory and a fully competent adaptive response develops later during the juvenile stage. Therefore, larvae are dependent on innate defenses against pathogens or opportunistic agents during their early life stages. While the epithelial layers of gills skin and digestive tract provide an initial physical barrier, if breached by a pathogen the innate immune system, composed of various effector molecules, provides the first line of defense. The innate immune system includes membrane-bound pathogen recognition receptors (PRRs), such as NOD-like receptors, and cytoplasmic proteins that bind pathogen-associated molecular patterns (PAMPs) expressed on the surfaces of invading microbes [5]. Innate immunity exerts its effect by employing a variety of cells and effector proteins able to destroy or inhibit the growth of infectious micro-organisms [6]. The complement system is one of the first lines of immune defense and is a regulator of acquired immunity. The complement system is activated through three different, but partially overlapping routes: the classical, alternative and lectin binding pathways [7]. The classical pathway is triggered by antibody binding to the cell surface that in turn binds a cascade of complement factors; the alternative pathway is independent of antibody binding and is activated directly by foreign microorganisms; and the lectin pathway, which is activated by the binding of a protein complex consisting of mannose/mannan-binding lectin to oligosaccharide ligands on bacterial cells [8]. Lysozyme is expressed in a wide variety of tissues of the innate immune system of vertebrates and it is an important defense molecule that possesses lytic activity against microbial cell walls preventing invasion [9]. The role of Mx protein (mxp) in resistance to negative-strand RNA viruses has been well established, despite the precise mechanism of viral inactivation by Mxp not being fully understood [10]. Mx proteins are highly conserved in vertebrates, and are able to inhibit the virus life cycle in different phases, whether they are localized in the cytoplasm or in the nucleus of cells [11]. Cyclooxygenase (cox2) is responsible for synthesis of prostaglandins to maintain homeostatic functions and it also plays an important role in the inflammatory response, among other relevant functions [12]. Metallothionein is an important protein contributing to resistance to heavy metal contamination in aquatic organisms, but also has important immune functions [13]. Metallothionein plays a relevant role in the detoxification of heavy metal ions and plays a role in buffering action against heavy metals by binding essential metals such as Cu and Zn, which has an inhibitory effect on systemic pathogens and provides protection against oxidative stress [14]. For these effector molecules to carry-out their function in a coordinated manner there is a need for intra- and inter-cellular signaling. For this, the innate immune response relies on signaling by members of the toll-like receptor (TLR) family that are highly conserved in vertebrates, and associated adaptor molecules, such as MyD88 (myeloid differentiation primary response gene). MyD88 has been implicated in the downstream signaling of TLRs with the possible exception of TLR3 [15], and thus, it is implicated in defense against a variety of pathogens. MyD88 contains a Toll/Interleukin receptor (TIR) domain and is required for activation of mitogen-activated protein (MAP) kinase family members as well as nuclear factor κ B (NF- κ B) translocation, which in turn activates transcription of pro-inflammatory cytokines such as interleukin 1 beta (IL1B) [16]. Innate immunity additionally has the potential to respond to endogenous molecules [17] that are released by host cells as a result of necrosis, pathogen infection, and certain pathological conditions, which are directly or indirectly recognized by NOD-like receptors. The NOD-like receptors (NLRs) function as cytoplasmic sensors of pathogen presence. NOD2 detects muramyl dipeptide (MDP) found in the peptidoglycan of the cell wall of both gram-positive and negative bacteria [17]. NOD2 in mammals is highly expressed in epithelial cells or macrophages associated with the intestine. Studies in zebrafish have detected NOD3 (*nod3*) which has been shown to be an orthologue of mammalian NOD3 (NLR3) and found to have similar NACHT domains and an equal number of Leucine-Rich-Repeat (LRR) domains, likely required for

binding to pathogen associated molecular patterns [18], but its specific ligand remains to be identified.

The aim of this study is to improve understanding of the innate immune response of meagre during larval rearing when the adaptive immune response remains incomplete. To this end this study will examine i) the ontogenic changes in innate immune-related gene expression of *c3*, *cox2*, *metl1*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3* during larval and juvenile development in meagre, and ii) how their expression is modulated by different PAMPs *in vivo* using adult fish; and iii) examine the modulation using PAMPs in isolated cells from specific organs *in vitro*. This information will be of value for better understanding early ontogeny of the immune system in this fish species, as well as serve as a basis for proper health maintenance in hatchery and nursery management practices.

2. Materials and methods

2.1. Larval rearing and sample collection

Larvae used in this study were obtained from a meagre broodstock held at IRTA-SCR (San Carlos de la Rapita, Spain). The broodstock was maintained under controlled simulated natural water temperature and photoperiod using a recirculation system (IRTamar®). During the natural reproductive period (April–June), mature fish were selected based on oocyte size (< 550 μ m) and spermiating condition. Pairs of mature fish (21.2 ± 3.7 kg females and 16.1 ± 2.6 kg males) were hormonally induced (15μ g/kg of des-Gly10, [D-Ala6]-gonadotropin-releasing hormone ethylamide (Sigma, Spain)) to spawn in 10,000 L tanks. The resulting fertilized eggs were collected using a passive egg collector placed in the outflow of the tank. Batches of 50,000 eggs were incubated at 18 – 19 °C in 35 L mesh-bottomed (300 μ m mesh) incubators with aeration and gentle air-lift water exchange that were placed within 2,000 L tanks. Hatching rate was determined by estimating the number of larvae (three 100 ml sub-samples) obtained from the stocked eggs.

Larvae were then transferred to two 1.5 m³ tanks and reared using a mesocosm system. Water temperature was maintained at 20 °C. From two days post-hatching (dph), larvae were fed enriched rotifers (*Brachionus sp.*) until 11 dph. Freshly enriched *Artemia* metanauplii were introduced from 9 dph until 31 dph. The experimental tanks were fed with a dry commercial feed for fish larvae containing 60% protein, 15% lipids, 9% ash, 0.5% fibre, 0.5% phosphorus, (Gemma Wean 0.2, Trouw France S.A, France). The diet was incorporated from 21 dph until the end of the experiment. Random samples of larvae and juvenile meagre were taken at 8, 15, 29, 40, 43, 60, 85, 96, and 120 dph. Each sample consisted of ten animals that were placed in an Eppendorf tube on ice, containing RNAlater™ (Ambion, Austin, Texas), then preserved at -80 °C until RNA extraction. Time points for analysis were chosen to coincide with the specific periods of change in rearing practices; primarily changes in diet, but also rapid changes in organogenesis, all of which increase stress and enhance susceptibility to diseases. Before immersion in RNAlater™ (Ambion, Austin, Texas) the fish were euthanized using a high concentration of MS222 (1 g/l) (Aldrich, E10521). Larger larvae (post-29 dph) had an excess of tissue, mainly muscular tissue that was trimmed to reduce signal dilution from non-target tissues. The anterior section from the gills forward and the posterior part from the anus to the tail were removed (Fig. 1), whereas younger larvae were processed entire. At days 85, 96, and 120 individual tissues (gill, kidney, spleen, and intestine) were excised aseptically.

2.2. Total RNA extraction and reverse transcription

Total RNA was extracted using Trizol reagent (Invitrogen) according to manufacturer's instruction. RNA concentration and purity was determined by spectrophotometry (NanoDrop 2000, Thermo Fisher Scientific, Madrid, Spain) measuring the absorbance at 260 and 280 nm. The quality of extracted RNA was verified with visualization of




Fig. 1. Prior to the extraction of RNA from the samples, the rostrum and caudal portions were removed by cutting with scalpel as illustrated with dashed lines. Only the remaining ventral part was used for further analysis.

the 28S and 18S ribosomal RNA bands by agarose gel electrophoresis. Prior to reverse transcription total RNA was treated with the DNase 1 AMPD1 kit, (Sigma–Aldrich), according to manufacturer's instructions to remove possible contaminating genomic DNA. Total RNA then underwent reverse transcription by adding 2 µg total RNA using the ThermoScript™ Reverse Transcriptase (Invitrogen) with oligo-dT (0.5 µg/µl) and random hexamer primers (50 ng/µl) 10X RT buffer [200 mM Tris-HCl (pH8.4), 500 mM KCl] 1.5 mM MgCl₂, 800 mM dNTP mix, RNase inhibitor, SuperScript™ II RT, followed by RNase H (Invitrogen) treatment. Once reverse transcription reactions were prepared they were placed in a Mastercycle® nexus GSX1 (Eppendorf AG, Hamburg, Germany) to complete first-strand cDNA synthesis. All the samples were diluted 1:20 in molecular biology grade water (Sigma–Aldrich) and stored at –20 °C. Negative controls (no RT enzyme) were included to confirm absence of genomic DNA contamination.

2.3. Gene isolation

Nucleotide sequences of target genes from the Genebank (www.ncbi.gov) were chosen from different marine teleost species, such as large yellow croaker (*Larimichthys crocea*), mandarin fish (*Synchiropus splendidus*), Humphead snapper (*Lutjanus sanguineus*), European seabass (*Dicentrarchus labrax*), orange spotted grouper (*Epinephelus coioides*), turbot (*Scophthalmus maximus*), Asian sea bass (*Lates calcarifer*) and gilthead sea bream (*Sparus aurata*). The sequences were aligned using CLUSTAL W (BioEdit package; <http://www.mbio.ncsu.edu/BioEdit/bioedit.html>) for designing degenerate/consensus primers for amplification from cDNA produced from meagre tissues as described previously (Campoverde et al., 2017). The fragments amplified were separated by gel electrophoresis and resulting bands of the expected length were excised, purified using the QIAquick PCR purification kit (Qiagen) and sequenced by Sistemas Genomicos (Valencia, Spain). The transcript sequences obtained were then used as templates for designing primers for quantitative PCR assays.

2.4. Phylogenetic analysis

Some of the genes chosen for analysis are known to be members of larger more diverse and previously studied families of genes. As such, there exists in genetic databases other homologs for comparison to establish the member of such a gene family that has been identified for study. To clarify this, evolutionary analyses were conducted in MEGA5 [19] after alignment of all selected homologs using CLUSTAL W. Nucleotide substitution models were chosen for each analysis using the utility included in MEGA 5. The evolutionary reconstruction of the NOD isoforms was performed using the Kimura 2-parameter nucleotide substitution model [20] and Neighbor-Joining method [21]. The evolutionary history for metallothionein and lysozyme were inferred by using the Maximum Likelihood method based on the Jukes-Cantor model [22]. For these analyses, the initial trees for the heuristic search were obtained automatically as follows: when the number of common sites was < 100 or less than one fourth of the total number of sites, the maximum parsimony method was used; otherwise Neighbor-Joining method with MCL distance matrix was used. A discrete gamma

distribution was used to model evolutionary rate differences among sites [5 categories (+G, parameter = 200,000)]. Statistical robustness of data for all analyses was analyzed using the bootstrap method [23] with bootstrap confidence values from 1,000 replicates shown at branch nodes (values < 40 not shown). During analyses all ambiguous positions were removed for each sequence pair. The analysis of NOD isoforms (*nod2* and *nod3*) involved 129 positions from 25 sequences. For metallothionein (*metII*), a total of 22 sequences were included in the final dataset each with 71 positions. The analysis of lysozyme (*lyzc*) included 25 sequences with 148 positions in each.

2.5. PAMP stimulation

To investigate the effect of PAMP stimulation on the expression of the selected genes, 38 healthy juvenile individuals (30–40 g each) were held in a recirculating water tank (100 l) at 20 °C. Fish were fed with a commercial diet, 58% protein, 17% lipids, 10% ash, 0.6% fibre, 1.3% phosphorous (Gemma Wean 0.3, Trouw France S.A, France) twice a day (09:00 and 16: 00) for two weeks to acclimate them to the indoor culture environment. For primary cell culture and collection of samples, fish were euthanized with an overdose of anesthesia (50 mg/l MS-222).

2.5.1. In vivo stimulation

Fish were injected ($n = 8$) intraperitoneally with 100 µl PBS containing either 100 µg poly I: C (3.3 mg/kg) (Sigma, UK. P1530), 400 µg LPS (13.3 mg/kg) (Sigma, UK. L3129) or 100 µg β-glucan (3.3 mg/kg) (Sigma, UK. 89862). The control animals were injected with PBS only. After 24 h, the individuals were dissected and tissues (head kidney, spleen, gut and gill) sampled. Total RNA was isolated and cDNA was prepared as described above. The mRNA expression level of *metII*, *c3*, *cox2*, *lyzc*, *mxd*, *myd88*, *nod2*, *nod3* were determined by qPCR. Relative expression was normalized to GAPDH expression and calculated as arbitrary units and converted to a proportion relative to the PBS control samples.

2.5.2. In vitro stimulation

Tissues (kidney, spleen, gills, and gut) from apparently healthy fish ($n = 6$) were collected from euthanized fish for primary cell culture. Tissue from gills and gut were pre-treated with collagenase in HBSS with FBS (0.37mg/ul – Sigma Aldrich, Ref# C0130-100 MG) for 30 min at 37 °C to facilitate dissociation of cells from intercellular connective tissue. Afterwards these tissues, together with the spleen and kidney samples, were passed through a 100 µm nylon mesh cell strainer (SefarNytal PA-13xxx/100, Spain) in Leibovitz L15 medium (Gibco) containing penicillin/streptomycin (Gibco, #15140-122) at 1:1000 and 2% foetal calf serum (Gibco, #10270-098). The resulting cell suspension was collected and centrifuged at 400 × g for 10 min. The supernatant was removed and replaced with 10 ml of previously described L15 media. The cell suspension was again centrifuged and supernatants removed and replaced with 30 ml of media. Cells were distributed to 12 well microtiter plates in 5 ml aliquots. Wells were stimulated using LPS (Sigma, #L3129-100 MG) at 50 µg/ml, poly (I:C) (Sigma, #P1530-25 MG) at 100 µg/ml, and B-glucan (Sigma, #89862-1G-F) at 50 µg/ml in triplicate. Control samples included 250 µl of PBS. Four, 12 and 24 h after stimulation the cells were harvested and centrifuged at 400 × g for 10 min, the supernatant discarded, and the pellet suspended in RNAlater. Total RNA was isolated and cDNA was prepared as described above.

2.6. Real-time qPCR

The qPCR reactions for innate immune gene expression were carried out using a LightCycler® 480 Real-Time (Applied Biosystems, Roche). A master mix was prepared containing: 5 µl SYBR Green Supermix (Life Technologies), 1 µl of molecular biology grade water (Sigma) containing 10 µM of forward and reverse gene-specific primers and 4 µl of

Table 1

Primers used for gene expression analysis by qPCR, including the amplicon size and primer sequences. Abbreviations: *gpdh*, glyceraldehyde 3-phosphate dehydrogenase; *β-act*, beta-actin; *hprt*, hypoxanthine-guanine phosphoribosyltransferase; *metII*, methallothionein; *c3*, complement; *cox2*, cyclooxygenase; *lyzc*, lysozymes; *mxp*, mx protein; *myd88*, myeloid differentiation primary response gene 88; *nod2*, nucleotide-binding oligomerization domain-containing protein 2; *nod3*, nucleotide-binding oligomerization domain-containing protein 3.

	Genes	E (%)	Forward primer (5' → 3')	Reverse primer (5' → 3')	Size (bp)
Reference genes	<i>gpdh</i>	100.0	CCAGTACGTGGTGGAGTCCACTG	AGCGTCAGCGGTGGGTGCAGAG	109
	<i>β-act</i>	100.0	TGGGGGAGCAATGATCTTGATCTTCA	AGCCCTCTTTCTCGGTATGGAGTC	212
	<i>hprt</i>	100.9	CATGGACTCATCTTGACAGGACAGA	GCCTTGATGTAGTCCAGCAGGTC	137
Immune genes	<i>metII</i>	102.2	GATCCTGCAATTGCAAAGACTGTTC	CCGGATGGGCGCATGGGCAG	70
	<i>c3</i>	100.1	AACCCATACGCTGTGCCATGACG	CACGTCCTTTAGGTACTGGCCAG	120
	<i>cox2</i>	100.5	GGAAGTTGGTGTGACATGCACTAC	CAATCAGGATGAGCCGTGTGGTC	211
	<i>lyzc</i>	100.0	GATGGATCCACTGACTACGGCATC	AAGCTGGCTGCCTGGATGTGGC	100
	<i>mxp</i>	96.2	AGTCAGTGGTTGACATTGTTTCATAATG	AACAGTGGCATGACCGTCATTGTAG	187
	<i>myd88</i>	102.2	GCTACTGCCAGAGTACTTCGAGT	TCCATACACACGAACCCGGGAGG	120
	<i>nod2</i>	100.1	CTCAATACTGTCTGATGCCATGG	CAAGTGTAACCTTTGGAGTAAGGTAG	145
	<i>nod3</i>	100.8	CAGCTTGGTGAACCTGTTCATCAC	TAAATCAGTCAGATCTCAGTGTTC	130

sample cDNA in a final volume of 10 µl. The real-time qPCR cycling was carried out as follows: 10 min at 95 °C, 40 cycles of 95 °C for 25 s, followed by an annealing step of approximately 59 °C for 25 s (annealing temperatures were adjusted for each specific primer pair), followed by 72 °C for 15 s, with a final dissociation stage of 0.5 °C increments from 75 °C to 95 °C. Each sample on the qPCR plate had two methodological replicates. Primer sequences used and amplicon lengths for each assay are shown in Table 1. The specificity of the primers was checked by confirming that only one melt peak was produced, and also by running a subsample on an agarose gel (2%) to confirm the presence of a single band of the expected size. The efficiency of amplification (E, %) of each primer pair was assessed from five serial ten-fold dilutions of cDNA from individual tissues, then calculated following the equation: $E (\%) = 10^{(-1/\text{slope})} - 1$, where the slope is that calculated from the regression line of the standard curve. Efficiencies of the gene expression assays ranged from 96.2% to 100.9% (100.27 ± 1.58 , mean \pm SD) (Table 1). The absence of primer-dimer formation in the NTC (non-template control) was also confirmed. Relative expression of genes from larvae, and each tissue from juveniles (ontogeny study), was normalized using the three endogenous controls glyceraldehyde phosphate dehydrogenase (*gpdh*), beta-actin (*β-act*), and hypoxanthine-guanine phosphoribosyltransferase (*hprt*) as determined using geNORM (<http://www.primersdesign.co.uk/products/9461-genormplus-kits/>), while the relative quantification at 120 dph, showing the lowest expression, was used as the calibrator.

For analyzing the relative expression *in vivo* and *in vitro* after PAMP stimulation, data was normalized using just GAPDH expression values, as this endogenous control gene showed less variability under the conditions tested. After normalization to this endogenous control as arbitrary units, results were converted to a proportion relative to the control group (PBS injected fish).

2.7. Statistical analysis

All data sets were checked for homogeneity of variances by subjecting them to a Levene's test using univariate analysis in a general linear model, based on a Tukey HSD post-hoc test, with a sample size of $n = 10$ (larva) and $n = 8$ (juvenile) to determine differences between time points ($P \leq 0.05$). The Kruskal-Wallis test was used for analyzing the expression for the *in vivo* ($n = 8$) samples after PAMP stimulation, while the ANOVA test was used for analyzing expression for the *in vitro* ($n = 6$) based on a Tukey HSD post-hoc test, using the statistical software package SPSS 20.0 (SPSS Inc., US).

3. Results

3.1. Gene isolation

Specific gene products were obtained using degenerate or consensus-primed PCR assays. All amplified fragments were sequenced bidirectionally and analyzed using the BLAST utility via the National Center for Biotechnology Information (NCBI) database (<http://www.ncbi.nlm.nih.gov/>) to confirm their identity as proper orthologues of the intended target gene. Sequences for transcripts from *met*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3*, are reported for the first time in this species. These sequences have been uploaded to GenBank under the following accession numbers: *met* = MF281966, *c3* = MF281960, *cox2* = MF281967, *lyzc* = MF281968, *mxp* = MF281965, *myd88* = MF281964, *nod2* = MF281970, *nod3* = MF281969. Sequences were also obtained for GAPDH, *β-actin* and *HPRT* as endogenous control genes for gene expression assays and have been entered into the GenBank database under the following accession numbers MF281962, MF186587 and MF186588, respectively.

3.2. Phylogenetic analysis

Some of the genes of interest for this study are known to have additional isoforms. For this reason, phylogenetic analysis was performed to properly identify such genes. Homologs of metallothionein, *nod2*, *nod3*, and lysozyme from meagre were further characterized as to class or isoform to give clarity to resulting gene expression analyses. To achieve this, extant sequences from GeneBank were aligned using CUSTAL W and the alignment edited manually. For all analyses, the percentage of trees in which the associated taxa clustered together (bootstrap values) is shown next to the branch nodes. Phylogenetic analysis of representative homologs of *nod2* and *nod3* sequences from GenBank were compared to that obtained from meagre resulting in an optimal tree with a sum branch length = 1.89832313. The two different Nod isoforms, *nod2* and *nod3*, clearly separated into distinct and differing clades. Analysis of metallothionein II confirmed its segregation from metallothionein I and generated a tree with the highest log likelihood of -157.7290 . The tree resulting from the analysis of lysozyme had highest log likelihood value of -2526.8685 and demonstrated its membership in the clade with *lyzc*.

3.3. Gene expression analyses

3.3.1. Gene expression during ontogeny

In order to better understand the biological role of several significant innate immune genes in meagre, mRNA levels for *met II*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3* were analyzed by quantitative real-




Fig. 2. The evolutionary history for (a) *met II*, (b) *lyz C* (c) *nod2*, *nod3* from meagre (shown in bold) was inferred using the maximum likelihood (a), (b) and neighbor-joining (c) method conducted in MEGA5. All coding positions were included and values for the bootstrap confidence values (1,000 replicates) are shown at branch nodes.

time PCR from 8 until 120 dph (Fig. 2). The expression of the above-mentioned genes were variable throughout development and growth of larvae and fry. However, among genes studied three general patterns of

expression were evident. In the first group, including *cox2*, there was a general decrease over time between 8 and 60 dph. A second group, composed of *mxp* and *myd88*, showed an expression pattern

c)




Fig. 2. (continued)

characterized by an initial low expression that increased over time with an important peak in expression at 60 dph. A third group of genes, consisting of *lyzc*, *nod2*, *nod3*, was also discernible, which showed low levels initially with peaks at 29 and 60 dph. The pattern of *met II* and *c3* expression was clearly at a high level at day 8, and then it decreased to a moderate level at day 29 and followed by a substantial increase at day 60. However, a different expression pattern was found for *mxp* and *myd88* transcripts. They were detected at day 8 and subsequently the expression level increased gradually until day 60. In contrast, the expression of *cox2* gradually decreased during the study period, possibly due to the expression signal being diluted by accumulation of non-expressing cells as the fish larva grew. A comparison of *nod2* and *nod3* show their expression profiles were similar. In fact, the levels of expression of both transcripts were higher at day 29 and then, decreased to moderate levels (40–43 dph), followed by an increase at day 60. The pattern of expression seen with *lyzc* expression was similar, but with greater differences in expression between 29 and 60 dph. Interestingly, changes in the expression of these genes seemed to correlate to changes in the diet during on-growing (Fig. 3).

As larvae increased in size during the latter part of the ontogeny study (85, 96, or 120 dph) the transcript expression was monitored in individual tissues (kidney, spleen, gill and gut). As shown in Fig. 4, the eight genes exhibited different patterns of tissue specific expression. The expression of *cox2* showed a significant increase in gill, gut and kidney at day 120 compared to earlier time points (85–96 dph). Expression of *myd88* transcripts showed a significant increase in the kidney at 96 dph that was maintained high and stable until 120 dph. However, *met II* and *mxp* transcripts showed significant increases in the gut during 85–96 dph. Finally, *nod3* transcript levels increased significantly ($P \leq 0.05$) in the spleen (85 dph) and gut (120 dph) in comparison with other preceding time points (Fig. 4).

3.3.2. Response of meagre innate immune genes to *in vivo* PAMP stimulation

To better understand how different pathogenic stimuli influence changes in the expression of these genes, an *in vivo* challenge with various PAMPs was performed, with samples collected for analysis after 24 h. Post stimulation, *c3* mRNA levels were most highly up-regulated

in the kidney, spleen, gut and gills by β -glucan. However, LPS seemed to be a more potent stimulator of expression in the gut and gills. Lysozyme was significantly up-regulated in the kidney, spleen, gut and gill by LPS, but, curiously, there was significant down-regulation in mucosal tissues (gut and gill) when poly I:C was used, while there was a significant up-regulation in systemic tissues (kidney and spleen) by β -glucan stimulation. The expression of *met II* was potentially induced in the spleen, gut, and gills by LPS stimulation and by poly I:C and β -glucan stimulation in the spleen and gut. The expression of *myd88* was significantly up-regulated in the gut by all the PAMP stimulants. The abundance of *mxp* transcripts were increased in all the tissues tested when stimulated by poly I:C and in the gut there was also a significant up-regulation ($P \leq 0.05$) when stimulated by LPS and β -glucan. The expression level of *cox2* was significantly up-regulated in kidney and spleen by LPS. The expression of *nod2* transcripts showed the highest up-regulation in spleen and gut when stimulated by LPS and β -glucan. Different expression profiles were observed for *nod3*; there was significant up-regulation ($P \leq 0.05$) in the kidney and spleen by β -glucan, while the other PAMP stimulants had little significant effect on mRNA levels, as seen in Fig. 5.

3.3.3. Response of meagre innate immune genes to *in vitro* PAMP stimulation

Only the lymphoid organs of kidney and spleen were analyzed for the *in vitro* part of this study. The kidney is the first major lymphoid organ to develop during early ontogeny and the spleen plays an important role in hematopoietic and immune functions. Cells were isolated from these lymphoid organs (kidney and spleen) and challenged with 3 PAMPs for 4, 12 and 24 h. The expression of *met II* was significantly upregulated (Fig. 6) in splenocytes for all stimulants after 24 h. The expression of *mxp* was up-regulated significantly in the spleen and kidney cells after 4 and 12 h stimulation with poly I:C, but in the kidney cells the only significant increase by all of the stimulants was at 24 h. Similarly, with kidney cells a differential expression profile was seen with increased expression of *c3* in response to all three PAMPs at 24 h, but splenocytes showed an up-regulation after 24 h stimulation with LPS and β -glucan. No change in expression of *myd88* was found in the stimulated kidney and spleen cells at 4 h, but at 12 h there was up-




Fig. 3. The temporal change in gene expression of larval and juvenile ($n = 10$) meagre for *met II*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3*. Transcripts were normalized using an arithmetic mean of three housekeeping genes: GAPDH, HPRT, and β -actin. Different letters above the bars indicate significant differences among different time points (Tukey's test, $P \leq 0.05$) and the larval feeding schedule also are shown. Results are expressed as the mean \pm SEM.

regulation using poly I:C. However, at 24 h up-regulation occurred in spleen cells with LPS. The mRNA expression of *cox2* in kidney cells was up-regulated at 4 h by all 3 stimulants. In splenocytes, significant expression levels were only detected at 12 h after the stimulation by poly I; C and β -glucan. A significant up-regulation of *lyzc* expression was seen in kidney cells at 4 h after stimulation by LPS and β -glucan and at 24 h after stimulation using poly I:C. In splenocytes, up-regulation was seen at 12 h after stimulation by poly I:C and β -glucan at 24 h after stimulation by LPS and β -glucan. For *nod2*, transcription levels in kidney cells was significantly up-regulated at 24 h post-stimulation with LPS and β -glucan. This contrasted with *nod3* expression, which was up-regulated in kidney cells after 4 h stimulation with LPS and β -glucan and had returned to control level by the other later time points. For splenocytes, *nod3* expression was significantly up-regulated at 24 h post-stimulation with LPS and β -glucan.

4. Discussion

4.1. Ontogeny of innate gene expression

The present study is the first report of some representative innate immune genes in meagre and the characterization of their expression patterns in larvae and early juveniles, which is a critical time in fish development, when reliance on innate immune mechanisms is required for combatting infectious disease. Although previous studies have focused on identifying immune genes and/or characterizing their response to certain pathogens, there are few comparative studies reporting their temporal appearance during development. Further we have characterized the expression of these genes under specific immune stimulation using four different PAMPs. This immune stimulation study was performed in whole animals with all the consequent communication among various tissues, and in primary cell isolations from those same organs to see how organ-specific responses might differ from the response obtained from the intact animal.

Phylogenetic analysis was used to confirm the specific identity of




Fig. 4. The temporal change in gene expression of *met II*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3* in the kidney, spleen, gill, and gut ($n = 8$) from 85 to 120 dph in meagre. Transcripts were normalized using an arithmetic mean of three housekeeping genes: *gpdh*, *hppt*, and β -act. Different letters above the bars indicate significant differences among different time points (Tukey's test, $P \leq 0.05$). Results are expressed as the mean \pm SEM. K = kidney, Sp = spleen, Gu = gut, Gi = gill.

genes where closely allied isoforms are known that could confound gene expression analyses. The phylogenetic analysis of *met* revealed that the isoform identified from meagre was *met II*, and it is an isoform more similar to a homolog described in *Larimichthys crocea* as expected, since both species are quite close, from a taxonomical point of view. Metallothionein has been shown to be a potential biomarker for metal contamination, but it also has an important role in Zn regulation during development [24]. Zinc is one of the major trace elements indispensable for functioning of, among other things, zinc-finger transcription factors. In this study, *met II* showed a higher expression level from 8 dph (4.49 ± 0.39 mm SL), but slowly decreased during development in

meagre. In specific tissues during the early juvenile period, the abundance of *met II* transcripts increased, particularly in the kidney and gut. In these tissues, metallothionein is especially important in development due to cellular signals that are required during cell growth and differentiation, which require nuclear localization of this intracellular zinc regulator [25] and maintaining Zn homeostasis at the cytoplasmic level for regulation of several crucial processes implicated in the innate immune [26]. For example, Zn ions are important for the production of pro-inflammatory cytokines such as interleukins IL-1 β , IL-6 and tumor necrosis factor α (TNF- α), and Zn deficiency leads to reduced chemotaxis by polymorphonuclear cells [26], and impaired formation,




Fig. 5. The *in vivo* expression of *met II*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2*, *nod3* in meagre kidney, spleen, gill and gut following PAMP stimulation. Fish ($n = 8$) were injected intraperitoneally with PBS, poly I:C, LPS and β -glucan and sampled at 24 h post-injection. The gene for GAPDH, was employed as an internal reference. Asterisks (*) mark significant between stimulated and control groups (Kruskal Wallis test, $P \leq 0.05$). Data are means \pm SEM. Abbreviations: K = kidney, Sp = spleen, Gu = gut, Gi = gill.

activation, and maturation of lymphocytes, disturbed intercellular communication via cytokines, and weakened innate host defense via phagocytosis and oxidative burst [27].

The complement system is known as an integral aspect of innate immunity, of which *c3* is a central component. This system also links with the adaptive immune response and has been shown to be important for C3-trophic driven morphogenesis of developing embryos/larvae [28]. In this study, the expression of *c3* was detected from the initial time point of the study (8 dph), but transcript levels of *c3* reached their lowest levels at 29 dph (14.41 ± 1.62 mm SL), coinciding with the end of the weaning period (the transition from artemia to an artificial diet) where as a result, larvae may be immunosuppressed in comparison to other life development periods and possibly more sensitive to potential pathogenic organism. As larvae matured, the

expression level of *c3* in tissues increased in the spleen and gut tissues, probably acting to facilitate hematopoietic development [29], stem-cell differentiation [30] and defense against allochthonous bacteria ingested with feed, some of which may be potential pathogens [28]. These previous findings support the idea that complement and its recognized role in immune defense, may also play a role in the formation and generation of different organs during development [31].

Cyclooxygenase, is the enzyme responsible for the initial rate-limiting conversion of arachidonic acid to prostaglandins (PGE2 and PGH₂). Mammals contain two *cox* genes, *cox1* and *cox2*. Homologs of these genes have been found in many fish species, including zebrafish (*Danio rerio*) [32]. Although, it is known that both isoforms are similar, *cox1* is generally assumed to be constitutively expressed performing homeostatic and maintenance functions, while *cox2* is induced by




Fig. 6. The expression patterns *in vitro* of a) *met II*, b) *mxp*, c) *c3*, d) *myd88*, e) *cox2*, f) *lyzc*, g) *nod2*, h) *nod3* mRNA after PAMP stimulation of cell suspensions isolated from meagre (*A. regius*) kidney and spleen tissues, collected from healthy fish ($n = 6$). The cells were stimulated with poly I:C, LPS and β -glucan and sampled at different time points post-stimulation (4, 12 and 24 h), for RNA extraction and qPCR. All samples were compared to PBS treated controls. Asterisks (*) mark significant between stimulated and PBS control groups (Tukey's test, $P \leq 0.05$). Data are means \pm SEM. Abbreviations: K = kidney, Sp = spleen.

inflammation, cytokines, endotoxin and other pathophysiological processes [33]. In the present study, *cox2* displayed a decrease in expression between 8 dph and 60 dph (during which period whole larvae were analyzed), that may be due to the expression signal being diluted as the larvae grew, since the abundance of *cox2*-positive cells likely did not increase in the same proportion as all other somatic cells. Although Ishikawa et al. [34] observed *cox2* expression in zebrafish, it exhibited a more restricted pattern of expression and no evidence of an association to morphogenic development. In Atlantic salmon (*Salmo salar*) [33] and mummichug (*Fundulus heteroclitus*) [35], *cox2* was observed to have the highest level of expression in the gills, and it was suggested that there was a possible higher requirement for osmoregulation processes and/or

stress responses. In the present study, high *cox2* expression levels were seen in the kidney, gills and gut in early juveniles aged 120 dph, although the relation of this expression to PGE production in these tissues was not evaluated. High *cox2* expression with a potential increase in production of PGs [12,36] in some tissues such as gills possibly aid the process of osmoregulation [12] and ion transport in euryhaline species [35], and in the gut to maintain the integrity of the mucosal barrier [37]. Significantly, expression of *cox2* is associated with several cell types, e.g. monocytes and macrophages [38] that likely aid in surveillance of mucosal barriers.

In fish, there exist two variants of lysozyme: c and g-types. The lysozyme isoform identified in meagre was shown to be the c-type, which

has been reported previously in many teleost species [39–41]. Lysozyme possesses lytic activity against both gram-positive and gram-negative bacteria. For this reason, it is an important defense molecule of the immune system, widely present at host/environment interfaces including gills, and gastrointestinal tract, but it is also present in lymphoid tissues, serum, mucus, and other body fluids [29]. Aside from its antibacterial function, it promotes phagocytosis by directly activating polymorphic nuclear leucocytes and macrophages, or indirectly, by an opsonizing foreign bodies. At 29 dph, transcript levels of *lyzc* were found to be significantly up-regulated around the time of weaning (transfer to *Artemia* and dry feed). Possibly, this increased expression during the transition of diet may reflect influence from certain nutrients or exposure to new cohorts of bacterial species in different feed sources that are stimulating expression of this immune effector. It may also be that specific components supplemented in the feed modulate lysozyme expression [42]. Lysozyme gene expression in teleost has been assessed in tissues from healthy fish, but the levels and pattern of expression varied for individual species, the isoform under study, and tissues examined [40,43]. In the current study, analysis of *lyzc* in tissues showed an increase in the kidney, spleen and mucosal tissues (gut and gill) in juveniles aged 60–85 dph, whereas its expression was even higher at later time points (120 dph). This observation could be correlated with the structure of the organs and increase in mucus/goblets cells in mucosal tissues during this period in this species [44]. Further, some reports have found the c type lysozyme in rainbow trout, (*Oncorhynchus mykiss*) [45] was specifically expressed in the liver and kidney [46], whereas in Japanese flounder, *Paralichthys olivaceus*, the c-type lysozyme gene was expressed in head kidney, spleen, brain and ovary, while in brill (*Scophthalmus rhombus*) expression was observed in liver and stomach [39]. Therefore, differences in the *lyzc*-positive tissue distribution and levels of expression of *lyzc* among fish species might suggest some variation in function, and potentially multiple roles.

The induction of the gene *mxp* in mammals is specifically induced by type I-IFN [47]. The antiviral effect of type I IFN is exerted through its binding to the IFN- α/β receptor, which triggers a signal transduction through the Jak-Stat pathway resulting in expression of *mxp* and other antiviral proteins. In the present study, during early development the peak in expression of this gene was observed at 60 dph and it was also expressed in all tissues examined during later stages of development, with a prominent increase in the gut, coincident with an abundance of granulocytes in this tissue at later developmental periods [44]. The gut epithelium is a continuation of the interface between the external environment and internal host environment, therefore it requires defense mechanisms to be operating continuously to act rapidly for limiting infection and maintaining gut homeostasis. *Mxp* is among those proteins that responds quickly to a viral attack, but as the immune system matures might be maintained at a constitutively higher level in the gut relative to other tissues (Fig. 4).

Myeloid differentiation factor 88 (Myd88) is an important adaptor protein in the Toll-like receptor (TLR) signaling pathway and is used by all TLRs except for TLR3. Orthologues of *myd88* have been found in multiple fish species [15,48,49], demonstrating among fish the wide evolutionarily distribution of this intracellular immune mediator due to its crucial role in host immunologic surveillance where it functions as an accessory protein to ligand receptors. The gene for Myd88 was found expressed continuously in all stages of larval development with a sharp increase at 60 dph. In more developed juveniles where individual tissues were examined, *myd88* transcripts were detected in all tissues, while high levels of transcripts were observed in the gill, spleen and kidney at 85, 96 and 120 dph.

The phylogenetic analyses of NOD isoforms from meagre demonstrated the two isoforms identified reside hierarchically in different clades, consistent with their presumed separate functions. The genes *nod2* and *nod3* belong to the subfamily of NOD-like receptors (NLRs) characterized by CARD-containing effector-binding domains. *Nod2* is a member of the cytoplasmic pattern recognition receptors (PRRs) family

that recognizes muramyl dipeptides derived from peptidoglycan, present both in gram-positive and negative bacteria. Upon ligand recognition, *Nod2* induces the activation of the NF- κ B and MAPK pathways. Activation of NF κ B and Mapk induce transcription and production of inflammatory cytokines, chemokines and antimicrobial peptides which mediate the antimicrobial response. *Nod3* (NLRC3) belonging to the NLR-C subfamily, plays a role in the innate immune response against bacteria and virus. Studies have also suggested that it has a role in modulating T cells and inhibiting inflammatory mechanisms, although studies in species such as catfish have shown *nod3* is present in many tissues [50]. Little is known about the precise mechanism, activation and signaling cascades of members of the NLRC subfamily. During the first 60 days of this study, *nod2* and *nod3* genes showed similar expression patterns during larval development, with the transcripts exhibiting abrupt up-regulation at 29 dph and 60 dph, which could suggest a coordinated activity between the two peptides these genes encode. In contrast, differences were observed in the individual tissues examined. Expression of *nod2* increased in the kidney and spleen while *nod3* transcripts were highly expressed in gill and gut at 120 dph. Similar results for *nod2* expression in the kidney were obtained from grass carp (*Ctenopharyngodon idella*) [51], and grouper (*Epinephelus coioides*) [52], while in rainbow trout (*Oncorhynchus mykiss*), *nod2* isoforms (*nod2a* and *nod2b*) were detected highly expressed in muscle [53].

4.2. *In vivo* stimulation

While studies of immune gene expression are becoming more common with fish, the totality of data is somewhat limited and comparative studies on molecular effectors of the innate immune system often rely on the greater abundance of literature from mammalian studies as a starting point. Herein is presented a study of the expression of key innate immune genes as modulated by PAMPs. An *in vivo* study was performed with three different PAMPs injected intraperitoneally and measured the immune gene expression responses to each PAMP 24 h after their administration.

In our study, we found an increase in expression of *met II* in the spleen, after stimulation by the three different PAMPs, while in gut the expression of *met II* was only up-regulated by LPS and β -glucan. In contrast, it was only slightly stimulated in the gills by LPS. These results suggest that the spleen is somewhat more responsive with regard to *met II* activation where recruitment of lymphocyte lineages are an important response for preventing systemic infections. The response in the gut was much more pronounced using LPS and β -glucan as stimulants, demonstrating that this tissue is functioning as a first line of defense with a much faster, stronger response against invading pathogens. Interestingly, there was a slight, but statistically significant, decrease in expression in the kidney under stimulation by LPS. The induction of *met II* is mediated by immune cells and inflammatory signals. Additional gene products may be working to suppress the expression in kidney tissue where filtration of excess ions and their removal from the blood occurs.

In this study, *c3* transcripts were up-regulated in all the tissues exposed to β -glucan. Interestingly, Campoverde et al. [54] detected a similar response in piscidin (*pisc*), after β -glucan stimulation, which could suggest a common up-stream response mechanism when meagre are exposed to β -glucan, such as a shared regulation by pro-inflammatory cytokines or toll-like receptors.

With regard to *cox2* expression, significant differences were observed in the kidney, spleen and gut after injection with LPS and β -glucan, but there were no changes in gill tissue. This makes sense physiologically, as there is a need to maintain respiratory functions and avoid unnecessary damage to respiratory epithelia from an excessive inflammatory response. Therefore, in gills more stimulation might be needed to elicit significant up-regulation of *cox2* and the concomitant increase in prostaglandin synthesis. These results demonstrate that *cox2*

can be induced by PAMPs binding to PRRs in the kidney, spleen, and gut; all tissues that need a thorough humoral and cellular response to clear pending systemic infections.

Multiple genes encoding the c-type lysozyme have been identified in different teleost species, and in some species different isoforms have been shown to perform distinct functions, or have tissue-specific expression patterns [40]. In the case of meagre, *lyzc* transcripts were found to increase significantly in the kidney and spleen after injection with LPS and β -glucan, while *lyzc* expression in the gut and gills was up-regulated only by LPS stimulation. However, poly I:C stimulation down-regulated *lyzc* expression in gills and gut. The higher levels of expression detected in the gill after injection with LPS and β -glucan are coherent with the idea of epithelia cells forming the primary barrier against a multitude of potential pathogens requiring a quick and rapid response to obviate any ingress of invaders [9]. These data from meagre are consistent with other studies from teleosts, but contrasting results have been obtained; studies of c-type lysozyme in brill showed significant changes in mRNA levels in stomach and liver after a bacterial challenge [40] possibly indicating this isoform might provide some direct protection in the digestive system in brill, whereas for g-type lysozyme similar tissue expression profiles were observed.

A relatively high *mxp* expression was observed in all tissues stimulated by poly I:C, but *mxp* expression was also stimulated to a lesser degree in the gut by LPS and β -glucan. High expression of *mxp* transcripts in mucosal tissues (gill and gut) was expected as these are the main entrance route for viral infections in aquatic organisms, while systemic tissues like the kidney and spleen have a high abundance of lymphocytes that respond to viral attack and produce Mx protein.

The expression of *myd88* was found to be up-regulated in the gut by all PAMPs tested. The expression of *myd88* is ubiquitous as it functions as a mediator for intracellular signaling in immune cells. Increased expression in the gut may facilitate and improve the integrity of the intestinal barrier following PAMP recognition [55]. Toll-like receptors (TLRs) recruit adaptor molecules, like Myd88, for signal transduction to activate nuclear factor-kappa B (NF- κ B).

In the case of *nod2*, transcripts were up-regulated in the spleen following injection of LPS and β -glucan, whereas in the gut an up-regulation was observed following stimulation with each PAMP. In gills, there was a weak response when fish were injected with LPS, which might again be explained as a protective measure to avoid an unregulated pro-inflammatory response leading to tissue trauma. The aquatic habitat is a rich source of bacterial LPS and the immune responses activated by the presence of LPS would need to be under tight control to differentiate presence of LPS liberated from natural bacterial cell death from an active invasion.

Gene expression levels for *nod3* were generally weak in all tissues; although there was some significant up-regulation, in the kidney by LPS and β -glucan, in spleen by β -glucan, in gut by poly I:C and LPS, and in gill tissue by LPS.

This *in vivo* trial using poly I:C, LPS, and beta-glucan has demonstrated that gene expression was effectively modulated in response to these PAMPs for all the studied genes, while the induction of gene expression by each stimulant demonstrated tissue-specific differences.

4.3. *In vitro* stimulation

Due to some difficulties in obtaining cell suspensions of sufficient quality from gills and intestine, we analyzed *in vitro* results only for the spleen and kidney. Exposure of kidney and spleen cells to different PAMPs elicited differential responses as compared to control cell suspensions treated with only PBS, demonstrating that these cells have the ability to recognize molecular patterns mimicking diverse pathogen types.

Early modulation of expression was seen for *mxp*, *cox2*, *lyzc*, *nod3* in kidney cells. An important component of the fish antiviral response is the IFN-inducible protein *mxp*. In Atlantic salmon and Japanese

flounder *mxp* was found to significantly protect fish cells against viral infection and induce the gene expression levels of type 1 IFN and *mxp* [56,57]. In the present study kidney and spleen cells responded strongly to poly I:C stimulation with an increase in mRNA transcripts of *mxp* as early as 4 h, and up to 12 h following exposure. At the same time expression *myd88* reached maximum levels following exposure to poly I:C in both systemic tissues, then recovered its basal levels at 24 h. These results suggest stimulation of kidney and spleen cells with poly I:C elicits a significantly increased expression of *mxp* and *myd88* where they may function in synergy against viral infections [58].

The antimicrobial activity of c-type lysozyme is due to its hydrolytic activities, inhibition of viral genomic RNA or RNA transcripts, and degradation of viral polysaccharides. In orange-spotted grouper, *E. coioides* and Japanese flounder, *Paralichthys olivaceus* the expression of c-type lysozyme transcripts was most abundant in head kidney [59,60] and in grouper cells when they were exposed to Singapore grouper iridovirus (SGIV). In the present study, expression of lysozyme was up-regulated in kidney cells at 4 h post exposure to LPS and β -glucan. However, this level increased significantly in spleen at 12 h post exposure to poly I:C and β -glucan demonstrating lysozyme is more active as an immediate response element when bacteria (LPS) is the source of stimulation.

The complement system, present in the blood plasma, plays a central role in recognizing foreign antigens and subsequent microbial killing, and phagocytosis. In the present study, when kidney cells were exposed to PAMPs, *c3* gene expression increased at 24 h exposure, but was more highly expressed in spleen cells stimulated with LPS and β -glucan. The difference in expression between these tissues may be due in part to the types of cells present in each organ since kidney tissue has significant osmoregulatory roles as well as functioning in hematopoiesis.

Prostaglandin synthesis via *cox2* (a.k.a. prostaglandin synthase) activity, among other things, functions as a mediator of inflammation. It has been reported that the expression of *cox2* is not constitutive in rainbow trout and sea bass, but can be induced by factors such as LPS [61,62]. In this study, there was an immediate and early up-regulation of expression of *cox2* in stimulated kidney cells using all three PAMPs, whereas there was a later, but more prolonged increase of expression, in spleen cells stimulated with LPS and beta-glucan. By 24 h post exposure, the levels of expression in kidney cells were not significantly different from the control PBS-treated cells. The significance of the response in kidney cells may be related to the need for production of O₂ radicals generated as by-products of prostaglandin synthesis that would negatively impact bacterial agents potentially found in the blood during an episode of bacteremia.

The *met II* gene, has been used as a biomarker of heavy metals exposure in aquatic animals [63] and can also be implicated in protection from cellular and oxidative stress. Met II has been detected mainly in liver and kidney in mandarin fish and catfish [64,65] after exposure to cadmium and selenium, respectively, however this study found that spleen cells from meagre responded late to all 3 PAMPs at 24 h post-exposure; in contrast to kidney cells which did not show any significant stimulation by the three PAMPs used. Chelation of metals such as iron and zinc may provide a significant edge in combatting pathogens that require such trace metals for regulation of gene expression (eg. - zinc-finger transcription factor proteins) or metabolic processes requiring biologically available forms of iron. The response from the spleen cells seen with this perspective has sense. That there is no response seen from kidney cells is an area that could be profitable for future study.

NLRs are important intracellular cytosolic sensors for the initiation of innate immune responses against infectious agents [17]. Recently, genes grouped in the NLR family have been characterized in Japanese flounder [66] and seabass [67]. High levels of *nod2* mRNA have been found in the spleen and different immune cell populations and in particular neutrophils [68]. *Nod3* plays a role in modulating T cell responsiveness and inhibits inflammatory mechanisms [67]. The current

work with meagre has shown that only the expression of *nod3* in kidney cells increased at 4 h following an exposure to LPS and β -glucan. However both kidney and spleen cells respond strongly at 24 h to LPS and β -glucan with an increase of mRNA transcripts of *nod2* and *nod3*. The difference in the timing of expression of *nod2* and *nod3* suggests different regulatory pathways may be controlling each in the immediate early response. In no instance was poly I:C seen to function as a stimulant for either NLR. This is in agreement with previous findings, considering that muramyl dipeptides are previously described as a ligand for *nod2* [17].

The findings from both the *in vivo* and *in vitro* experimental data confirm the genes studied are responsive to pathogen associated molecular patterns and the host immune response is tissue specific. The timing of expression during on-growing of larval and juvenile meagre, can depend on the state of maturity of the organs and their requisite cellular composition. In conclusion, the current study is the first to identify the genes for *met II*, *c3*, *cox2*, *lyzc*, *mxp*, *myd88*, *nod2* and *nod3*, in meagre and examine how these immune genes are expressed during larval and juvenile development, and the effects of PAMP stimulation on the modulation of expression *in vivo*. Interestingly, the data from this study suggests that the transition from live feed to a commercial dry feed might be a time when the innate immune response is in flux, and that it may be the change in diet and the nutritional and/or compositional changes of the feed that are inducing the significant alterations in expression seen at these times. This study, by using poly I:C, LPS and β -glucan as proxies for actual viral, bacterial and fungal pathogens pathogens, also provides valuable insight into how the meagre innate immune response may respond in comparison to other species. However, further studies focused on larval nutrition, specifically identifying changes in the components in the diet that can beneficially modulate gene expression of the immune system, will enhance larval rearing protocols by reducing mortalities.

Declaration of competing interest

None.

Acknowledgements

This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration (KBBE-2013-07 single stage, GA 603121, DIVERSIFY). CC received funding from a Ph.D. grant awarded by the Secretaría Nacional de Educación Superior, Ciencia, Tecnología e Innovación (SENESCYT), Ecuador. We would like to thank G. Macià and M. Monllaó, S. Molas for their help in live prey production and meagre larval and juvenile rearing.

References

- [1] F. Asche, K.H. Roll, T. Trolsvik, New aquaculture species—the whitefish market. *Aquacult. Econ. Manag.* 13 (2009) 76–93, <https://doi.org/10.1080/13657300902881641>.
- [2] L. Quémer, *Le maigre commun (Argyrosomus regius): biologie, pêche, marché et potentiel aquacole*, Editions Quae (2002) 1–32.
- [3] J. Roo, C.M. Hernández-Cruz, C. Borrero, D. Schuchardt, H. Fernandez-Palacios, Effect of larval density and feeding sequence on meagre (*Argyrosomus regius*; Asso, 1801) larval rearing. *Aquaculture* 302 (2010) 82–88, <https://doi.org/10.1016/j.aquaculture.2010.02.015>.
- [4] M. Seppola, H. Johnsen, S. Mennen, B. Myrnes, H. Tveiten, Maternal transfer and transcriptional onset of immune genes during ontogenesis in Atlantic cod, *Dev. Comp. Immunol.* 33 (2009) 1205–1211, <https://doi.org/10.1016/j.dci.2009.06.013>.
- [5] L. Zhu, L. Nie, G. Zhu, L. Xiang, J. Shao, Advances in research of fish immune-relevant genes: a comparative overview of innate and adaptive immunity in teleosts, *Spec. Issue Comp. Immunol.* China 39 (2013) 39–62, <https://doi.org/10.1016/j.dci.2012.04.001>.
- [6] B. Beutler, Innate immunity: an overview, *Mol. Immunol.* 40 (2004) 845–859, <https://doi.org/10.1016/j.molimm.2003.10.005>.
- [7] H. Boshra, J. Li, J.O. Sunyer, Recent advances on the complement system of teleost fish, *Rev. Fish Immunol.* Fish Immunol 20 (2006) 239–262, <https://doi.org/10.1016/j.fsi.2005.04.004>.
- [8] M.C.H. Holland, J.D. Lambris, The complement system in teleosts, *Fish Shellfish Immunol.* 12 (2002) 399–420, <https://doi.org/10.1006/fsim.2001.0408>.
- [9] S. Saurabh, P.K. Sahoo, Lysozyme: an important defence molecule of fish innate immune system, *Aquacult. Res.* 39 (2008) 223–239, <https://doi.org/10.1111/j.1365-2109.2007.01883.x>.
- [10] A. Garcia-Sastre, Inhibition of interferon-mediated antiviral responses by influenza A viruses and other negative-strand RNA viruses, *Virology* 279 (2001) 375–384, <https://doi.org/10.1006/VIRO.2000.0756>.
- [11] P. Novel, M.A. Fernández-Trujillo, J.B. Gallardo-Gálvez, I. Cano, M. Manchado, F. Buonocore, E. Randelli, G. Scapigliati, M.C. Álvarez, J. Béjar, Two Mx genes identified in European sea bass (*Dicentrarchus labrax*) respond differently to VNNV infection, *Vet. Immunol. Immunopathol.* 153 (2013) 240–248, <https://doi.org/10.1016/j.vetimm.2013.03.003>.
- [12] F. Norambuena, S. Mackenzie, J.G. Bell, A. Callol, A. Estévez, N. Duncan, Prostaglandin (F and E, 2- and 3-series) production and cyclooxygenase (COX-2) gene expression of wild and cultured broodstock of senegalese sole (*Solea senegalensis*), *Gen. Comp. Endocrinol.* 177 (2012) 256–262, <https://doi.org/10.1016/j.ygcen.2012.04.009>.
- [13] H. Mao, D.-H. Wang, W.-X. Yang, The involvement of metallothionein in the development of aquatic invertebrate, *Aquat. Toxicology* 110 (2012) 208–213, <https://doi.org/10.1016/j.aquatox.2012.01.018>.
- [14] G.K. Andrews, Regulation of metallothionein gene expression by oxidative stress and metal ions, *Biochem. Pharmacol.* 59 (2000) 95–104, [https://doi.org/10.1016/S0006-2952\(99\)00301-9](https://doi.org/10.1016/S0006-2952(99)00301-9).
- [15] O.W. S, A.M. van der Sar, C. van der Laan, H.P. Spaink, W. Bitter, A.H. Meijer, MyD88 innate immune function in a zebrafish embryo infection model, *Infect. Immun.* 74 (2006) 2436–2441, <https://doi.org/10.1128/IAI.74.4.2436-2441.2006>.
- [16] S. Janssens, R. Beyaert, A universal role for MyD88 in TLR/IL-1R-mediated signaling, *Trends Biochem. Sci.* 27 (2002) 474–482, [https://doi.org/10.1016/S0968-0004\(02\)02145-X](https://doi.org/10.1016/S0968-0004(02)02145-X).
- [17] B. Swain, M. Basu, B.R. Sahoo, N.K. Maiti, P. Routray, A.E. Eknath, M. Samanta, Molecular characterization of nucleotide binding and oligomerization domain (NOD)-2, analysis of its inductive expression and down-stream signaling following ligands exposure and bacterial infection in rohu (*Labeo rohita*), *Dev. Comp. Immunol.* 36 (2012) 93–103, <https://doi.org/10.1016/j.dci.2011.06.018>.
- [18] K.J. Laing, M.K. Purcell, J.R. Winton, J.D. Hansen, A genomic view of the NOD-like receptor family in teleost fish: identification of a novel NLR subfamily in zebrafish, *BMC Evol. Biol.* BMC Evol. Biol. 8 (2008) 42.
- [19] K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, S. Kumar, MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods, *Mol. Biol. Evol.* 28 (2011) 2731–2739, <https://doi.org/10.1093/molbev/msr121>.
- [20] M. Kimura, A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences, *J. Mol. Evol.* 16 (1980) 111–120.
- [21] N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.* 4 (1987) 406–425.
- [22] T.H. Jukes, C.R. Cantor, Evolution of protein molecules, *Mamm. Protein Metab* 3 (1969) 132.
- [23] J. Felsenstein, Confidence limits on phylogenies: an approach using the bootstrap, *Evolution (N. Y.)* 39 (1985) 783–791.
- [24] W.-Y. Chen, J.A.C. John, C.-H. Lin, H.-F. Lin, S.-C. Wu, C.-H. Lin, C.-Y. Chang, Expression of metallothionein gene during embryonic and early larval development in zebrafish, *Aquat. Toxicol.* 69 (2004) 215–227, <https://doi.org/10.1016/j.aquatox.2004.05.004>.
- [25] M.G. Cherian, M.D. Apostolova, Nuclear localization of metallothionein during cell proliferation and differentiation, *Cell. Mol. Biol.* 46 (2000) 347–356.
- [26] P. Bonaventura, G. Benedetti, F. Albarède, P. Miossec, Zinc and its role in immunity and inflammation, *Autoimmun. Rev.* 14 (2015) 277–285, <https://doi.org/10.1016/j.autrev.2014.11.008>.
- [27] A.S. Prasad, Zinc: role in immunity, oxidative stress and chronic inflammation, *Curr. Opin. Clin. Nutr. Metab. Care* 12 (2009) 646–652.
- [28] S. Lange, S. Bambir, A.W. Dodds, B. Magnadóttir, The ontogeny of complement component C3 in Atlantic cod (*Gadus morhua* L.)—an immunohistochemical study, *Fish Shellfish Immunol.* 16 (2004) 359–367, <https://doi.org/10.1016/j.fsi.2003.06.001>.
- [29] B. Magnadóttir, S. Lange, S. Gudmundsdóttir, J. Bøgwald, R.A. Dalmo, Ontogeny of humoral immune parameters in fish, *Fish Larval Immunity* Fish Larval Immunol 19 (2005) 429–439, <https://doi.org/10.1016/j.fsi.2005.03.010>.
- [30] D. Mastellos, J.C. Papadimitriou, S. Franchini, P.A. Tsonis, J.D. Lambris, A novel role of the complement: mice deficient in the fifth component of complement (C5) exhibit impaired liver regeneration, *J. Immunol.* 166 (2001) 2479–2486.
- [31] K.-C. Peng, C.-Y. Pan, H.-N. Chou, J.-Y. Chen, Using an improved Tol2 transposon system to produce transgenic zebrafish with epinecidin-1 which enhanced resistance to bacterial infection, *Fish Shellfish Immunol.* 28 (2010) 905–917, <https://doi.org/10.1016/j.fsi.2010.02.003>.
- [32] T. Ishikawa, K.J.P. Griffin, U. Banerjee, H.R. Herschman, The zebrafish genome contains two inducible, functional cyclooxygenase-2 genes, *Biochem. Biophys. Res. Commun.* 352 (2007) 181–187, <https://doi.org/10.1016/j.bbrc.2006.11.007>.
- [33] R. Olsen, A. Svardal, T. Eide, A. Wargelius, Stress and expression of cyclooxygenases (cox1, cox2a, cox2b) and intestinal eicosanoids, in Atlantic salmon, *Salmo salar* L., *Fish Physiol. Biochemistry* 38 (2012) 951, <https://doi.org/10.1007/s10695-011-9581-1>.

- [34] T. Grosser, S. Yusuff, E. Cheskis, M.A. Pack, G.A. FitzGerald, Developmental expression of functional cyclooxygenases in zebrafish, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 8418–8423, <https://doi.org/10.1073/pnas.112217799>.
- [35] K.P. Choe, J. Havird, R. Rose, K. Hyndman, P. Piermarini, D.H. Evans, COX2 in a euryhaline teleost, *Fundulus heteroclitus*: primary sequence, distribution, localization, and potential function in gills during salinity acclimation, *J. Exp. Biol.* 209 (2006) 1696–1708, <https://doi.org/10.1242/jeb.02198>.
- [36] D. Bagga, L. Wang, R. Farias-Eisner, J.A. Gaspy, S.T. Reddy, Differential effects of prostaglandin derived from ω -6 and ω -3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion, *Proc. Natl. Acad. Sci.* 100 (2003) 1751–1756.
- [37] M. Agard, S. Asakrah, L.A. Morici, PGE2 suppression of innate immunity during mucosal bacterial infection, *Front. Cell. Infect. Microbiol.* 3 (2013).
- [38] J.K. Chettri, M.K. Raida, L. Holten-Andersen, P.W. Kania, K. Buchmann, PAMP induced expression of immune relevant genes in head kidney leukocytes of rainbow trout (*Oncorhynchus mykiss*), *Dev. Comp. Immunol.* 35 (2011) 476–482.
- [39] J. Hikima, I. Hirono, T. Aoki, Characterization and expression of c-type lysozyme cDNA from Japanese flounder (*Paralichthys olivaceus*), *Mol. Mar. Biol. Biotechnol.* 6 (1997) 339–344.
- [40] R.M. Jiménez-Cantizano, C. Infante, B. Martín-Antonio, M. Ponce, I. Hachero, J.I. Navas, M. Machado, Molecular characterization, phylogeny, and expression of c-type and g-type lysozymes in brill (*Scophthalmus rhombus*), *Fish Shellfish Immunol.* 25 (2008) 57–65, <https://doi.org/10.1016/j.fsi.2007.12.009>.
- [41] S.P. Nayak, B.R. Mohanty, J. Mishra, P.R. Rauta, A. Das, A.E. Eknath, P.K. Sahoo, Ontogeny and tissue-specific expression of innate immune related genes in rohu, *Labeo rohita* (Hamilton) (2011), <https://doi.org/10.1016/j.fsi.2011.02.014>.
- [42] H. Niu, Y. Jia, P. Hu, Z. Meng, J. Lei, Effect of dietary vitamin E on the growth performance and nonspecific immunity in sub-adult turbot (*Scophthalmus maximus*), *Fish Shellfish Immunol.* 41 (2014) 501–506.
- [43] B. Grinde, Ø. Lie, T. Poppe, R. Salte, Species and individual variation in lysozyme activity in fish of interest in aquaculture, *Aquaculture* 68 (1988) 299–304, [https://doi.org/10.1016/0044-8486\(88\)90243-8](https://doi.org/10.1016/0044-8486(88)90243-8).
- [44] C. Campoverde, K.B. Andree, D.J. Milne, A. Estévez, E. Gisbert, F. Carella, Ontogeny of lymphoid organs and mucosal associated lymphoid tissues in meagre (*Argyrosomus regius*), *Fish Shellfish Immunol.* 84 (2018) 509–520.
- [45] A. Dautigny, E.M. Prager, D. Pham-Dinh, J. Jollès, F. Pakdel, B. Grinde, P. Jollès, cDNA and amino acid sequences of rainbow trout (*Oncorhynchus mykiss*) lysozymes and their implications for the evolution of lysozyme and lactalbumin, *J. Mol. Evol.* 32 (1991) 187–198.
- [46] S. Cecchini, M. Paciolla, E. Biffali, M. Borra, M. V Ursini, M.B. Lioi, Ontogenetic profile of innate immune related genes and their tissue-specific expression in brown trout, *Salmo trutta* (Linnaeus, 1758), *Fish Shellfish Immunol.* 35 (2013) 988–992, <https://doi.org/10.1016/j.fsi.2013.05.026>.
- [47] R. Nygaard, S. Husgard, A.-I. Sommer, J.-A.C. Leong, B. Robertsen, Induction of Mx protein by interferon and double-stranded RNA in salmonid cells, *Fish Shellfish Immunol.* 10 (2000) 435–450, <https://doi.org/10.1006/fsim.1999.0249>.
- [48] Z.-H. Qi, Y.-F. Liu, W.-N. Wang, X. Wu, Y. Xin, Y.-F. Lu, A.-L. Wang, Molecular Characterization and Functional Analysis of a Complement C3 Molecule in the Orange-Spotted Grouper (*Epinephelus coioides*), (2011), <https://doi.org/10.1016/j.fsi.2011.09.018>.
- [49] C.-L. Yao, P. Kong, Z.-Y. Wang, P.-F. Ji, X.-D. Liu, M.-Y. Cai, X.-Z. Han, Molecular cloning and expression of MyD88 in large yellow croaker, *Pseudosciaena crocea*, *Fish Shellfish Immunol.* 26 (2009) 249–255, <https://doi.org/10.1016/j.fsi.2008.10.014>.
- [50] K. V Rajendran, J. Zhang, S. Liu, H. Kucuktas, X. Wang, H. Liu, Z. Sha, J. Terhune, E. Peatman, Z. Liu, Pathogen recognition receptors in channel catfish: I. Identification, phylogeny and expression of NOD-like receptors, *Dev. Comp. Immunol.* 37 (2012) 77–86, <https://doi.org/10.1016/j.dci.2011.12.005>.
- [51] W.Q. Chen, Q.Q. Xu, M.X. Chang, J. Zou, C.J. Secombes, K.M. Peng, P. Nie, Molecular characterization and expression analysis of the IFN-gamma related gene (IFN- γ rel) in grass carp *Ctenopharyngodon idella*, *Vet. Immunol. Immunopathol.* 134 (2010) 199–207, <https://doi.org/10.1016/j.vetimm.2009.09.007>.
- [52] Q.-H. Hou, S.-B. Yi, X. Ding, H.-X. Zhang, Y. Sun, Y. Zhang, X.-C. Liu, D.-Q. Lu, H.-R. Lin, Differential expression analysis of nuclear oligomerization domain proteins NOD1 and NOD2 in orange-spotted grouper (*Epinephelus coioides*), *Fish Shellfish Immunol.* 33 (2012) 1102–1111, <https://doi.org/10.1016/j.fsi.2012.08.015>.
- [53] M. Chang, T. Wang, P. Nie, J. Zou, C.J. Secombes, Cloning of two rainbow trout nucleotide-binding oligomerization domain containing 2 (NOD2) splice variants and functional characterization of the NOD2 effector domains, *Fish Shellfish Immunol.* 30 (2011) 118–127, <https://doi.org/10.1016/j.fsi.2010.09.014>.
- [54] C. Campoverde, D.J. Milne, A. Estévez, N. Duncan, C.J. Secombes, K.B. Andree, Ontogeny and modulation after PAMPs stimulation of β -defensin, hepcidin, and piscidin antimicrobial peptides in meagre (*Argyrosomus regius*), *Fish Shellfish Immunol.* 69 (2017) 200–210, <https://doi.org/10.1016/j.fsi.2017.08.026>.
- [55] J. Galindo-Villegas, D. Garcí\`a-Moreno, S. de Oliveira, J. Meseguer, V. Mulero, Regulation of immunity and disease resistance by commensal microbes and chromatin modifications during zebrafish development, *Proc. Natl. Acad. Sci.* 109 (2012) E2605–E2614.
- [56] M.J.T. Kibenge, K. Munir, F.S.B. Kibenge, Constitutive expression of Atlantic salmon Mx1 protein in CHSE-214 cells confers resistance to Infectious Salmon Anaemia virus, *Virology* 42 (2005) 75, <https://doi.org/10.1186/1743-422X-2-75>.
- [57] C.M.A. Caipang, I. Hirono, T. Aoki, In vitro inhibition of fish rhabdoviruses by Japanese flounder, *Paralichthys olivaceus* Mx, *Virology* 317 (2003) 373–382.
- [58] J. Zou, S. Bird, C. Secombes, Antiviral sensing in teleost fish, *Curr. Pharmaceut. Des.* 16 (2010) 4185–4193.
- [59] S. Wei, Y. Huang, J. Cai, X. Huang, J. Fu, Q. Qin, Molecular cloning and characterization of c-type lysozyme gene in orange-spotted grouper, *Epinephelus coioides*, *Fish Shellfish Immunol.* 33 (2012) 186–196.
- [60] S. Minagawa, J. Hikima, I. Hirono, T. Aoki, H. Mori, Expression of Japanese flounder c-type lysozyme cDNA in insect cells, *Dev. Comp. Immunol.* 25 (2001) 439–445.
- [61] S.A. MacKenzie, N. Roher, S. Boltaña, F.W. Goetz, Peptidoglycan, not endotoxin, is the key mediator of cytokine gene expression induced in rainbow trout macrophages by crude LPS, *Mol. Immunol.* 47 (2010) 1450–1457.
- [62] F. Buonocore, E. Randelli, D. Casani, M. Mazzini, I. Cappuccio, C.J. Secombes, G. Scapigliati, cDNA cloning and expression analysis of a cyclooxygenase-2 from sea bass (*Dicentrarchus labrax* L.) after vaccination, *Aquaculture* 245 (2005) 301–310, <https://doi.org/10.1016/j.aquaculture.2004.11.015>.
- [63] M. Tom, N. Chen, M. Segev, B. Herut, B. Rinkevich, Quantifying fish metallothionein transcript by real time PCR for its utilization as an environmental biomarker, *Mar. Pollut. Bull.* 48 (2004) 705–710.
- [64] J.-H. Kim, J.-S. Rhee, H.-U. Dahms, Y.-M. Lee, K.-N. Han, J.-S. Lee, The yellow catfish, *Pelteobagrus fulvidraco* (Siluriformes) metallothionein cDNA: molecular cloning and transcript expression level in response to exposure to the heavy metals Cd, Cu, and Zn, *Fish Physiol. Biochemistry* 38 (2012) 1331–1342.
- [65] D. Gao, G.T. Wang, X.T. Chen, P. Nie, Metallothionein-2 gene from the Mandarin fish *Siniperca chuatsi*: cDNA cloning, tissue expression, and immunohistochemical localization, *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* 149 (2009) 18–25, <https://doi.org/10.1016/j.cbpc.2008.05.014>.
- [66] S. Unajak, M.D. Santos, J. Hikima, T.-S. Jung, H. Kondo, I. Hirono, T. Aoki, Molecular characterization, expression and functional analysis of a nuclear oligomerization domain proteins subfamily C (NLR) in Japanese flounder (*Paralichthys olivaceus*), *Fish Shellfish Immunol.* 31 (2011) 202–211, <https://doi.org/10.1016/j.fsi.2011.05.007>.
- [67] A. Paria, A. Deepika, K. Sreedharan, M. Makesh, A. Chaudhari, C.S. Purushothaman, A.R. Thirunavukkarasu, K.V. Rajendran, Identification of Nod like receptor C3 (NLR3) in Asian seabass, *Lates calcarifer*: characterisation, ontogeny and expression analysis after experimental infection and ligand stimulation, *Fish Shellfish Immunol.* 55 (2016) 602–612.
- [68] J. Xie, M. Belosevic, Functional characterization of receptor-interacting serine/threonine kinase 2 (RIP2) of the goldfish (*Carassius auratus* L.), *Dev. Comp. Immunol.* 48 (2015) 76–85.