

FMISO-PET-based lymph node hypoxia adds to the prognostic value of tumor only hypoxia in HNSCC patients

Anna Bandurska-Luque^{a,b,*}, Steffen Löck^{a,b,c}, Robert Haase^{a,1}, Christian Richter^{a,b,c,d}, Klaus Zöphel^{a,e,f,j}, Nasreddin Abolmaali^{a,2}, Annetkatrin Seidlitz^{a,b}, Steffen Appold^{a,b}, Mechthild Krause^{a,b,c,d,f,g}, Jörg Steinbach^{i,h}, Jörg Kotzerke^e, Daniel Zips^{j,k}, Michael Baumann^{a,b,c,f,1,3}, Esther G.C. Troost^{a,b,c,d,f,3}

^a OncoRay – National Center for Radiation Research in Oncology, Faculty of Medicine and University Hospital Carl Gustav Carus, Technische Universität Dresden, Helmholtz-Zentrum Dresden – Rossendorf; ^b Department of Radiotherapy and Radiation Oncology, Faculty of Medicine and University Hospital Carl Gustav Carus, Technische Universität Dresden; ^c German Cancer Consortium (DKTK), Partner Site Dresden and German Cancer Research Center (DKFZ), Heidelberg; ^d Helmholtz-Zentrum Dresden – Rossendorf, Institute of Radiooncology – OncoRay, Dresden; ^e Department of Nuclear Medicine, Faculty of Medicine and University Hospital Carl Gustav Carus, Technische Universität Dresden; ^f National Center for Tumor Diseases, Partner Site Dresden, Germany, German Cancer Research Center (DKFZ), Heidelberg, Germany, Faculty of Medicine and University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany, Helmholtz Association/Helmholtz-Zentrum Dresden – Rossendorf (HZDR), Dresden, Germany; ^g Deutsches Krebsforschungszentrum (DKFZ), Heidelberg; ^h Helmholtz-Zentrum Dresden, Rossendorf, Institute of Radiopharmaceutical Cancer Research, Rossendorf; ⁱ Faculty of Chemistry and Food Chemistry, Technische Universität Dresden; ^j Department of Radiation Oncology, Eberhard Karls Universität Tübingen; and ^k German Cancer Consortium (DKTK), Partner Site Tübingen, Germany

ARTICLE INFO

Article history:

Received 11 April 2018

Received in revised form 10 September 2018

Accepted 12 September 2018

Available online 4 October 2018

Keywords:

Hypoxia

FMISO-PET

Lymph node

Locally advanced HNSCC

Prognostic biomarker

ABSTRACT

Purpose: This secondary analysis of the prospective study on repeat [¹⁸F]fluoromisonidazole (FMISO)-PET in patients with locally advanced head and neck squamous cell carcinomas (HNSCC) assessed the prognostic value of synchronous hypoxia in primary tumor (Tu) and lymph node metastases (LN), and evaluated whether the combined reading was of higher prognostic value than that of primary tumor hypoxia only.

Methods: This analysis included forty-five LN-positive HNSCC patients. FMISO-PET/CTs were performed at baseline, weeks 1, 2 and 5 of radiochemotherapy. Based on a binary scale, Tu and LN were categorized as hypoxic or normoxic, and two prognostic parameters were defined: Tu-hypoxia (independent of the LN oxygenation status) and synchronous Tu-and-LN-hypoxia. In fifteen patients with large LN (N = 21), additional quantitative analyses of FMISO-PET/CTs were performed. Imaging parameters at different time-points were correlated to the endpoints, i.e., locoregional control (LRC), local control (LC), regional control (RC) and time to progression (TTP). Survival curves were estimated using the cumulative incidence function. Univariable and multivariable Cox regression was used to evaluate the prognostic impact of hypoxia on the endpoints.

Results: Synchronous Tu-and-LN-hypoxia was a strong adverse prognostic factor for LC, LRC and TTP at any of the four time-points ($p \leq 0.004$), whereas Tu-hypoxia only was significantly associated with poor LC and LRC in weeks 2 and 5 ($p \leq 0.047$), and with TTP in week 1 ($p = 0.046$). The multivariable analysis confirmed the prognostic value of synchronous Tu-and-LN-hypoxia regarding LRC (HR = 14.8, $p = 0.017$). The quantitative FMISO-PET/CT parameters correlated with qualitative hypoxia scale and RC ($p < 0.001$, $p \leq 0.033$ at week 2, respectively).

Conclusions: This secondary analysis suggests that combined reading of primary tumor and LN hypoxia adds to the prognostic information of FMISO-PET in comparison to primary tumor assessment alone in particular prior and early during radiochemotherapy. Confirmation in ongoing trials is needed before using this marker for personalized radiation oncology.

© 2018 Elsevier B.V. All rights reserved. Radiotherapy and Oncology 130 (2019) 97–103

* Corresponding author at: University Hospital Carl Gustav Carus Dresden, Fetscherstrasse 74, 01307 Dresden, Germany.

E-mail address: anna.bandurska@uniklinikum-dresden.de (A. Bandurska-Luque).

¹ Current address: Myers Lab, Max Planck Institute CBG, Dresden, Germany.

² Current address: Department of Radiology, Städtisches Klinikum Dresden-Friedrichstadt, Dresden, Germany.

³ Shared last authorship.

Patients with locally advanced head and neck squamous cell carcinoma (HNSCC) have a prognosis with a 3-year overall survival rate of approx. 40% after primary radiochemotherapy (RCT) with loco-regional relapse being the prime cause of treatment failure [1,2]. In order to improve treatment outcome for high risk HNSCC patients, local radiation dose escalation and/or intensified (concurrent) systemic treatment strategies have been developed [3–8]. In

the era of personalized medicine, patient selection based on prognostic biomarkers is potentially avoiding both under- and overtreatment. Tumor cell hypoxia is one of the most promising prognostic markers in HNSCC, as hypoxia correlates with poor outcome following R(C)T and surgery, and hypoxia-modification was shown to improve treatment results [5,9–16].

The adverse impact of hypoxia on treatment outcome was initially shown with invasive Eppendorf electrode measurements in accessible primary tumors (Tu) or lymph node metastases (LN). This invasive technique is limited regarding spatial information and repeatability [17–21]. Positron emission tomography (PET) using hypoxia-related markers, e.g., [^{18}F]fluoromisonidazole (FMISO) and [^{18}F]fluoroazomycin-araboside (FAZA), enables non-invasive imaging of the entire tumor metabolism at repeat time-points [10,22–29]. (Pre)clinical data have shown that hypoxia-PET readings vary depending on the level of oxygenation, that radiation dose escalation is feasible, and that FMISO-PET findings are of prognostic value [3,27,30–37].

Recently, we have successfully validated the prognostic value of FMISO-PET/computed tomography (CT) imaging of the primary tumor for loco-regional tumor control (LRC) in advanced stage HNSCC patients [38]. Until now, the majority of the analyses has focused on the prognostic value of hypoxia-PET measured in the primary tumor or index lesions [10,21–27,38,39]. Therefore, in our study we qualitatively and quantitatively assessed FMISO-PET-based hypoxia in Tu and LN at baseline and during the course of RCT. The aim was to evaluate whether combined assessment of FMISO-PET in Tu and LN increased the prognostic value of FMISO-PET in comparison with the Tu assessment only, as well as to investigate the correlation between quantitative LN parameters and regional control (RC).

Patients and methods

Patients

Between July 2006 and August 2013, 50 advanced stage HNSCC patients were evaluated in the prospective Dresden FMISO-PET imaging trial (NCT00180180). Of these, 45 patients diagnosed with metastatic regional lymph nodes ($\geq \text{N1}$) were included in this secondary analysis. All patients had histologically-proven, (functionally) irresectable HNSCC and provided written informed consent. Further inclusion criteria and approval by authorities and the local Ethics Committee have previously been described [22,38].

Work-up, radiochemotherapy and follow-up

The protocol for staging, treatment, imaging and follow-up has previously been described in detail [22,38]. Briefly, the total radiation dose of 72 Gy to the Tu and affected LN was combined with concurrent chemotherapy consisting of intravenous 5-fluorouracil with cisplatin, or with mitomycin C [1,22].

Image acquisition and analysis

Patients received pre-therapeutic (baseline) [^{18}F]fluorodeoxyglucose (FDG-) and FMISO-PET/CT as well as FMISO-PET/CT after 8–10 Gy (week 1), 18–20 Gy (week 2) and 50–60 Gy (week 5). The pre-treatment FDG-PET/CT scans analyzed in this study were acquired 60 min post injection (p.i.), and the FMISO-PET scans 4 h p.i. (except for one data set with a 2 h p.i. scan only). Details on imaging protocol, registration and image analysis were described in [22,38,40–42]. The gross tumor volume of the Tu (GTV_{TU}) and affected LN (GTV_{LN}), as well as their sum ($\text{GTV}_{\text{total}}$), were delineated on the pre-treatment CT taking into account clinical findings as well as the FDG-positive volume automatically

segmented using an adaptive thresholding algorithm [43,44]. To improve the analyses of the FMISO-PET/CT scans prone to therapy-induced longitudinal changes, an ellipsoidal volume of interest (VOI) was placed around each GTV_{TU} and GTV_{LN} in each scan. The background activity for subsequent qualitative and quantitative analyses was assessed within an ellipsoidal VOI ($\text{Background}_{\text{VOI}}$) in the deep neck muscles (Fig. 1 A–D) [22].

Qualitative hypoxia analysis

One investigator, blinded for treatment outcome, performed the qualitative analysis of hypoxia for each Tu and LN VOI using a visual binary scale: “hypoxic” defined as FMISO uptake higher than $\text{Background}_{\text{VOI}}$ and “normoxic” defined as equal to/lower than $\text{Background}_{\text{VOI}}$ (Fig. 1 A–D) [30,36,45,46]. To ensure reproducibility of the scoring system, all FMISO-scans were evaluated twice with an interval of approximately 10 weeks, reaching an agreement rate of 94%. In case of unequivocal findings, visual binary scoring was discussed with a second observer. The results of binary hypoxia were validated against quantitative FMISO-PET/CT readings performed in a subset of patients (see next paragraph).

Two parameters were defined based on the qualitative hypoxia scale: Tu-hypoxia (patients with hypoxic Tu) and synchronous Tu- and LN-hypoxia (patients with a hypoxic primary tumor and at least one hypoxic lymph node). Based on these parameters two analyses were performed: (a) patients with Tu-hypoxia versus patients with Tu normoxia, irrespective of the LN oxygenation status; (b) patients with Tu- and LN-hypoxia versus Tu and/or LN normoxia (both tumor and LN normoxia, Tu normoxia and LN hypoxia, or Tu hypoxia and LN normoxia).

Quantitative hypoxia analysis for validation of the qualitative parameters

In order to avoid underestimation of quantitative FMISO-PET parameters due to the partial volume effect in small lesions, the quantitative analysis was only performed in large lymph node metastases defined as FDG-PET/CT positive volume >5 ml (volume of sphere structure with diameter >2 cm using automatic FDG-PET segmentation). In patients fulfilling this criterion, quantitative FMISO-PET parameters, i.e., peak standardized uptake value [SUV_{peak} ; the mean SUV within $5 \times 5 \times 5$ voxels (1.26 ml) of highest FMISO uptake], peak tumor-to-background-ratio (TBR_{peak} ; ratio of the SUV_{peak} in Tu or LN and the mean SUV in $\text{Background}_{\text{VOI}}$), as well as hypoxic volumes (HV) encompassing those voxels with SUVs above different thresholds of multiples of the mean SUV in $\text{Background}_{\text{VOI}}$, 1.4 or 1.6 ($\text{HV}_{1.4}$, $\text{HV}_{1.6}$, respectively) were extracted from LN VOIs, and SUV_{peak} and TBR_{peak} were extracted from Tu VOI (for validation of binary hypoxia scale only, see below). Validation of the binary hypoxia scale was performed using Mann–Whitney–U tests comparing differences of SUV_{peak} and TBR_{peak} values extracted from Tu and LN VOIs between groups defined by the binary hypoxia score.

Endpoints and statistics

The primary endpoint of the prospective study was LRC and secondary endpoints were freedom from distant metastases (DM), overall survival (OS), local control (LC, defined as absence of progression or recurrence in Tu), regional control (RC, defined as absence of progression or recurrence in LN) and time-to-progression (TTP; defined as time to any progression). The endpoints were calculated from the first day of radiotherapy to the date of event or censoring [22,38]. Corresponding survival curves were estimated by the Kaplan–Meier method for OS and by the cumulative incidence function accounting for the competing risk




Fig. 1. FDG-PET/CT (A,C) and FMISO-PET (B,D) in two patients with metabolically active FDG-avid primary tumor and lymph node metastasis. (A) FDG-avid primary tumor and lymph node metastasis, (B) hypoxia in the primary tumor only. (C) + (D) Synchronous FDG- and FMISO-uptake in both primary tumor and lymph node [red – primary tumor volume of interest (VOI), green – lymph node VOI, blue – gross tumor volume of primary tumor (GTV_{Tu}), purple – GTV of lymph node (GTV_{LN})]. (E) Box-and-whiskers plots showing the discriminative power of the qualitative binary hypoxia scale (normoxia vs. hypoxia) when correlated to the quantitative FMISO-PET/CT parameter TBR_{peak} in primary tumors and LNs in all patients with lymph node metastases >5 ml on FDG-PET/CT. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

of death for the other endpoints. Log rank tests (OS) and Gray's test (other endpoints) were performed for patient groups stratified by qualitative hypoxia parameters. Univariable Cox regression was used to assess the impact of prognostic factors on the endpoints. Clinical factors found to be of significance on univariable analysis and qualitative hypoxia parameters were included in multivariable analyses. Mann–Whitney–U tests were used to compare continuous parameters between two groups. The statistical analyses were performed with SPSS Statistics 22 (IBM Corporation, Armonk, NY). Two-sided tests were performed and *p*-values <0.05 were considered statistically significant.

Results

The patient and treatment characteristics of the 45 patients included in this secondary analysis are summarized in Table 1. The median follow-up of these patients was 21 months (range 1–65 months). At the time of analysis, 13 (29%) patients were alive with 38 months median follow-up of (range 24–66 months). The 2-year and 3-year OS rates were 49% and 33%, respectively, and the cumulative incidence of loco-regional recurrences was 33% at both time points. Disease progression was observed in 19 (42%) patients, of whom 18 patients presented with synchronous Tu- and-LN-hypoxia in the baseline FMISO-PET/CT scan (Table 2).

Validation of qualitative hypoxia scale

Assessment of the two parameters of the qualitative hypoxia scale is shown in Fig. 1. Highly significant differences in the quantitative FMISO-PET image parameters SUV_{peak} and TBR_{peak} were found between groups defined by the qualitative binary hypoxia

Table 1

Patient, tumor and treatment characteristics (*n* = 45).

Characteristic	Value
Age [mean, (range)]	55 (42–74)
Male/female	38/7
Primary site	
Oral cavity/oropharynx/hypopharynx/larynx	8/18/15/4
cT-stage	
cT2/cT3/cT4a/T4b	1/15/24/5
cN-stage	
cN1/cN2a/cN2b/cN2c/cN3	6/2/11/24/2
Histological grade	
G1/G2/G3	2/24/19
UICC-stage	
III/IVA/IVb	5/35/5
HPV: p16-status (positive/negative/not available)	2/37/6
History of tobacco use (yes/no)	42/3
Mean radiation dose (range)	72 Gy (69–72 Gy)
Mean overall treatment time (range)	42 days (40–65 days)
Chemotherapy cisplatin plus 5-FU/mitomycin C plus 5-FU	39/6 [†]

Abbreviations: cT, clinical tumor stage; cN, clinical nodal stage; G, grade; UICC, Union International Contre le Cancer; HPV, human papilloma virus.

* One patient prematurely terminated treatment due to health-related issues.

[†] One patient received both cisplatin (2 cycles) and mitomycin C (1 cycle).

scale (*p* < 0.001; Fig. 1E). For normoxic vs. hypoxic Tu and LN combined, the median SUV_{peak} and TBR_{peak} were 1.48 vs. 2.32, and 1.32 vs. 1.97, respectively.

Prognostic value of qualitative FMISO readings

In the cohort, 24 patients presented with Tu- and LN-hypoxia, whereas 21 patients were allocated to the Tu and/or LN normoxia group (17 with hypoxic primary tumor and normoxic LN, 3

Table 2
Tumor and LN hypoxia status as detected by qualitative FMISO-PET reading at baseline, in weeks 1, 2 and week 5 in all patients with progression during follow up; 1 – hypoxia, 0–normoxia.

Patient	Baseline		Week 1		Week 2		Week 5		Progression
	Tu	LN	Tu	LN	Tu	LN	Tu	LN	
1	1	1	1	1	NA	NA	0	1	LRR
2	1	1	1	1	1	1	0	0	LR, DM
3	1	1	1	1	1	0	1	0	LR
4	1	1	NA	NA	1	1	1	1	LR
5	1	1	1	1	1	1	0	1	LR
6	1	1	1	1	1	1	1	1	LR
7	1	1	1	1	1	1	1	1	LR
8	1	1	1	1	1	0	1	1	LRR
9	1	1	1	1	1	1	1	1	LR
10	1	1	NA	NA	1	1	0	1	LRR
11	1	0	1	0	1	0	0	0	LR
12	1	1	1	1	1	1	0	0	LR, RR*, DM
13	1	1	1	1	0	0	0	0	LR
14	1	1	1	1	1	1	0	0	LR
15	1	1	1	1	1	1	1	1	RR, DM
16	1	1	1	1	0	1	0	0	DM
17	1	1	NA	NA	0	1	0	0	DM
18	1	1	1	1	0	0	0	0	DM
19	1	1	1	1	0	0	0	0	DM

Abbreviations: LR, local recurrence; LRR, loco-regional recurrence; DM, distant metastases; RR, regional recurrence; *, regional recurrence outside of FMISO-PET/CT field of view; NA, not available for FMISO-PET/CT scan is missing.

patients with normoxic tumor and normoxic LN, and one patient with normoxic tumor and hypoxic LN). The various patterns of reoxygenation during radiochemotherapy can be found in [Figs. S1–2](#).

Patients with synchronous Tu-and-LN-hypoxia at any of the four FMISO-PET time-points were found to have worse LC, LRC and TTP compared to patients with Tu and/or LN normoxia ($p \leq 0.004$; [Fig. 2](#) and [Table 3](#)). Conversely Tu-hypoxia, the parameter based on hypoxia assessment of the primary tumor only, was of prognostic relevance for LRC and LC in weeks 2 and 5, and for TTP at all of the per-treatment time points ($p \leq 0.047$, [Fig. 2](#) and [Table 3](#)). Synchronous Tu-and-LN-hypoxia based on pre-treatment FMISO-PET correlated with poor LC, LRC, TTP and DM while Tu-hypoxia only did not ($p < 0.001$, $p < 0.001$, $p < 0.001$ and $p = 0.009$, respectively, vs. $p = 0.18$, $p = 0.16$, $p = 0.091$ and $p = 0.37$, respectively; [Table 3](#)). Finally, Tu-hypoxia was found to be associated with poor overall survival at all time points, whereas Tu-and-LN-hypoxia was at baseline, weeks 2 and 5 ([Table 3](#)).

On univariable Cox analysis, the GTV_{total} and the number of hypoxic LN were significantly associated with LRC at baseline ($p = 0.02$, HR = 1.01; and $p = 0.01$, HR = 1.5, respectively), whereas the other parameters assessed showed no significant correlations ([Table S1](#)). On multivariable Cox analysis, including GTV_{total} and the number of hypoxic LN, synchronous Tu-and-LN-hypoxia was confirmed as prognostic factor ($p = 0.017$, HR = 14.8, [Table S1](#)).

Prognostic value of quantitative FMISO-analysis

The quantitative analysis of FMISO-PET parameters was possible in 15 patients with 21 lymph nodes exceeding the minimum required volume. The quantitative FMISO-PET parameters measured in these large LN during RCT statistically significantly correlated with RC at numerous FMISO-PET time-points ([Table 4](#)). The parameters HV_{1,4} and HV_{1,6} at baseline, at weeks 1 and 5 were significantly larger in LN developing a regional recurrence (LN non-responder) than in those with a complete regional remission during follow-up (responder; [Table 4](#) and [Fig. S3](#)). In accordance with the findings for the primary tumors by Zips et al. [22] and Löck et al. [38], LN responders and non-responders revealed different reoxygenation patterns and the risk of a regional recurrence was

higher in large LN with higher level of or with persisting hypoxia ([Table 4](#) and [Fig. S3](#)).

Discussion

This is the largest analysis focusing on the prognostic value of synchronous hypoxia in the primary tumor and lymph node metastases based on repeat FMISO-PET imaging in advanced stage HNSC patients. The results of this secondary analysis of a prospective imaging trial showed that the combined assessment of tumor and lymph nodes hypoxia improves the prognostic value of hypoxia PET/CT imaging at several time-points prior to and during RCT [22,38]. Since the nature of this investigation was explorative, these findings need to be validated in an independent cohort.

Past research has mainly focused on assessing the prognostic value of hypoxia-PET imaging or invasive Eppendorf measurements of one index lesion, most often the primary tumor [10,17,20–26,38]. In this secondary analysis we showed that synchronous primary tumor and lymph node hypoxia at baseline and during RCT is highly prognostic for worse LRC and LC and that it is so at an earlier time-point compared to tumor hypoxia only, which is found in the second week of RCT. The per-treatment prognostic factor is in accordance with preclinical data suggesting that the hypoxic volume correlates with local tumor control after 20 Gy but not at baseline [47]. Previously, few authors reported on differences in the oxygenation status of primary tumors and lymph node metastases in individual patients [10,28,29,48,49] but correlations with the clinical outcome are rare [46]. The incidence of synchronous Tu-and-LN-hypoxia prior to initiation of RCT of 44–58% corresponds well to the values in our cohort (53%). Servagi-Vernat et al. [28] acquired FAZA-PET imaging in twelve patients prior to and in the second and fourth weeks of RCT, but did not correlate the PET-readings with outcome. In the cohort ($N = 20$) reported by Lee et al. [46], almost exclusively consisting of oropharyngeal cancer patients, the 3-year LC rate was 100% and consequently FMISO-imaging performed 4 weeks after start of RCT did not correlate with patient outcome. The forty-five patients included in the prospective clinical study on radiochemotherapy in combination with the hypoxic cytotoxin tirapazamine by




Fig. 2. Cumulative incidence function for loco-regional recurrences after radiochemotherapy for cohorts of HNSCC discriminated by hypoxia qualitatively determined in the primary tumor (Tu; top row) versus primary tumor and lymph nodes (LN; bottom row) at four time-points before and during treatment. Tu-hypoxia was of significant prognostic value for LRC in weeks 2 and 5, whereas synchronous Tu-and-LN-hypoxia revealed prognostic significance at all time points.

Table 3

Time-points of FMISO-PET/CT acquisition, number of qualitatively assessed primary tumors and LNs, and prognostic value of the hypoxia parameters at the respective time points (cumulative incidence corrected for competing risks; p -values < 0.05 presented in bold).

	Baseline	Week 1	Week 2	Week 5
Number of assessed patients = primary tumors	$n = 45$	$n = 40$	$n = 44$	$n = 45$
Number of assessed LN on FMISO PET/CT	103	95	101	103
<i>Local control</i>				
Tu-hypoxia	0.18	0.098	0.006	0.047
Tu-and-LN-hypoxia	<0.001	0.004	<0.001	0.003
<i>Loco-regional control</i>				
Tu-hypoxia	0.16	0.082	0.003	0.013
Tu-and-LN-hypoxia	<0.001	0.004	<0.001	<0.001
<i>Time-to-progression</i>				
Tu-hypoxia	0.091	0.046	0.14	0.100
Tu-and-LN-hypoxia	<0.001	0.001	<0.001	0.001
<i>Freedom from distant metastases</i>				
Tu-hypoxia	0.37	0.28	0.33	0.47
Tu-and-LN-hypoxia	0.009	0.022	0.40	0.93
<i>Overall survival</i>				
Tu-hypoxia	0.014	0.050	<0.001	0.003
Tu-and-LN-hypoxia	0.036	0.12	0.011	0.008

Abbreviations: Tu-hypoxia, primary tumor hypoxia (independent of LN oxygenation); Tu-and-LN-hypoxia, synchronous primary tumor and lymph node hypoxia assessed by the qualitative hypoxia scale.

Rischin et al. [30] had detectable hypoxia in either the tumor or lymph nodes or both in 71% of the patients. The authors revealed that FMISO-based primary tumor hypoxia in the non-tirapazamine receiving patients was associated with a high risk of locoregional failure but they did not investigate prognostic value of synchronous tumor and LN hypoxia.

We observed a reoxygenation kinetics in lymph nodes similar to that published by other groups [28,49]. Noteworthy, the quantitative FMISO-PET, measurements of sufficiently large LN performed on the PET/CT-scan obtained in the second week of radiochemotherapy correlated with regional control. In agreement with previously published results based on Eppendorf electrode

Table 4

Results of the quantitative analysis: FMISO-PET/CT parameters measured in large LNs ($N = 21$) for regional control: univariable Cox test (top) and Mann–Whitney U test (bottom), both at baseline and during radiochemotherapy (p -values <0.05 are given in bold).

FMISO-PET/CT parameter in LN-metastases ($N = 21$)		Baseline		Week 1		Week 2		Week 5	
		p	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)
SUV _{peak}		0.142	1.40 (0.89–2.19)	0.64	2.61 (0.94–7.19)	0.016	3.43 (1.26–9.31)	0.006	12.8 (2.09–79.8)
TBR _{peak}		0.13	1.61 (0.88–2.95)	0.037	2.24 (1.05–4.77)	0.020	2.83 (1.18–6.8)	0.012	8.75 (1.61–47.6)
HV _{1.4}		0.021	1.02 (1.00–1.01)	0.017	1.04 (1.01–1.06)	0.033	1.03 (1.00–1.06)	0.058	1.03 (1.00–1.07)
HV _{1.6}		0.034	1.03 (1.00–1.06)	0.035	1.04 (1.00–1.07)	0.032	1.03 (1.00–1.07)	0.073	1.04 (1.00–1.09)
		p	Average (SE)	p	Average (SE)	p	Average (SE)	p	Average (SE)
SUV _{peak}	Responder	0.015	2.4 (0.2)	0.079	2.1 (0.2)	0.049	1.8 (0.1)	0.001	1.5 (<0.1)
	Non-Responder		3.5 (0.4)		3.2 (0.7)		3.5 (0.7)		2.4 (0.2)
TBR _{peak}	Responder	0.040	2.1 (0.5)	0.13	1.9 (0.2)	0.080	1.7 (0.1)	0.003	1.4 (0.1)
	Non-Responder		3.1 (0.5)		3.3 (0.9)		3.1 (0.8)		2.0 (0.2)
HV _{1.4}	Responder	0.040	15.1 (6.2)	0.035	10.3 (4.7)	0.22	6.8 (2.7)	0.008	1.4 (0.6)
	Non-Responder		53.5 (21.6)		52.3 (21.0)		37.4 (20.7)		17.6 (12.0)
HV _{1.6}	Responder	0.032	10.1 (4.5)	0.046	4.8 (2.0)	0.12	2.9 (1.3)	0.003	0.2 (0.1)
	Non-Responder		35.4 (14.9)		29.8 (16.8)		28.3 (17.7)		11.4 (9.1)

Abbreviations: LN, lymph node; HR, Hazard Ratio; CI, confidence interval; SE, standard error; SUV, standardized uptake value; TBR, tumor-to-background-ratio; HV, hypoxic volume; Responder, LN with regional control; Non-Responder, LN with regional failure.

measurements, we observed that patients with at least one hypoxic LN at pretreatment FMISO-PET, have a significantly worse LRC and a higher rate of DM, and that LRC worsened with increasing number of hypoxic LNs [18,19,50]. Importantly, in our cohort no regional recurrence occurred in any of the lymph nodes ($N = 82$) not taken into account for the quantitative analysis based on the volumetric cut-off criteria (data not shown).

In agreement with the recently published data of the primary results of the present study by Löck et al. and with Mortensen et al. [10], hypoxia determined in the primary tumor before start of treatment was not of significant prognostic value for LC. This is in contrast to results by other groups that found a correlation between baseline FMISO-PET parameters measured in the FDG-positive tumor volume and LC [23,26,27]. Welz et al. [31] have recently published the planned interim analysis of their dose-escalation study (NCT02352792) in 25 advanced stage HNSCC patients based on pre-treatment FMISO-PET imaging. Toxicity was favorable and LRC increased from 44% in ten patients with hypoxic primary tumors receiving standard RCT to 70% in ten patients with hypoxic primary tumors receiving dose-escalated RCT. Based on the findings of our own prospective explorative and validation phase II study [38], we have designed a randomized phase II multicenter trial. Patients with residual tumor hypoxia, as depicted by repeat FMISO-PET imaging prior to and in the second week of RCT, will receive either a standard dose or an escalated dose to the entire tumor volume using photons or particles. This cohort will be the ideal population for validation of the findings on synchronous primary and lymph node hypoxia before and during RCT presented here.

In summary, the role of measuring PET-hypoxia and its optimal timing as a prognostic and predictive marker for radiotherapy is still subject of controversy and ought to be further examined. Taken together the results presented here on the prognostic value of synchronous primary tumor and lymph node hypoxia are promising, and they will, together with other biomarkers, serve as tools for personalized radiation oncology [51–53].

In conclusion, FMISO-based synchronous primary tumor and lymph node hypoxia in advanced stage HNSCC patients undergoing radiochemotherapy was found to be of high prognostic relevance prior to and during radiochemotherapy. This finding ought to be validated in ongoing prospective clinical trials prior to using it for personalized radiation oncology.

Conflict of interest statement

There are no conflicts of interest to declare.

Acknowledgements

The authors thank Dr. Rosalind Perrin for her thoughtful comments and suggestions during this study. Prof. Howard Thames is gratefully acknowledged for the statistical support. The authors thank Dr. Kristin Gurtner for her comments on the patients' lymph node status. This study was supported by the German Federal Ministry of Education and Research (BMBF contract 03ZIK42/OncoRay).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.radonc.2018.09.008>.

References

- [1] Budach V, Stuschke M, Budach W, et al. Hyperfractionated accelerated chemoradiation with concurrent fluorouracil-mitomycin is more effective than dose-escalated hyperfractionated accelerated radiation therapy alone in locally advanced head and neck cancer: final results of the radiotherapy cooperative clinical trials group of the German Cancer Society 95–06 Prospective Randomized Trial. *J Clin Oncol* 2005;23:1125–35.
- [2] Bourhis J, Sire C, Graff P, et al. Concomitant chemoradiotherapy versus acceleration of radiotherapy with or without concomitant chemotherapy in locally advanced head and neck carcinoma (GORTEC 99–02): an open-label phase 3 randomised trial. *Lancet Oncol* 2012;13:145–53.
- [3] Schütze C, Bergmann R, Bruchner K, et al. Effect of [¹⁸F]FMISO stratified dose-escalation on local control in FaDu hSCC in nude mice. *Radiother Oncol* 2014;111:81–7.
- [4] Rischin D, Peters LJ, O'Sullivan B, Tirapazamine, cisplatin, and, et al. radiation versus cisplatin and radiation for advanced squamous cell carcinoma of the head and neck (TROG 02.02, HeadSTART): a phase III trial of the Trans-Tasman Radiation Oncology Group. *J Clin Oncol* 2010;28:2989–95.
- [5] Overgaard J. Hypoxic modification of radiotherapy in squamous cell carcinoma of the head and neck—a systematic review and meta-analysis. *Radiother Oncol* 2011;100:22–32.
- [6] Madani I, Duthoy W, Derie C, et al. Positron emission tomography-guided, focal-dose escalation using intensity-modulated radiotherapy for head and neck cancer. *Int J Radiat Oncol Biol Phys* 2007;68:126–35.
- [7] Rasmussen JH, Hakansson K, Vogelius IR, et al. Phase I trial of 18F-Fludeoxyglucose based radiation dose painting with concomitant cisplatin in head and neck cancer. *Radiother Oncol* 2016;120:76–80.
- [8] Jentsch C, Bergmann R, Bruchner K, et al. Impact of pre- and early per-treatment FDG-PET based dose-escalation on local tumour control in fractionated irradiated FaDu xenograft tumours. *Radiother Oncol* 2016;121:447–52.
- [9] Nordmark M, Bentzen SM, Rudat V, et al. Prognostic value of tumor oxygenation in 397 head and neck tumors after primary radiation therapy. An international multi-center study. *Radiother Oncol* 2005;77:18–24.
- [10] Mortensen LS, Johansen J, Kallehauge J, et al. FAZA PET/CT hypoxia imaging in patients with squamous cell carcinoma of the head and neck treated with radiotherapy: results from the DAHANCA 24 trial. *Radiother Oncol* 2012;105:14–20.
- [11] Horsman MR, Mortensen LS, Petersen JB, Busk M, Overgaard J. Imaging hypoxia to improve radiotherapy outcome. *Nat Rev Clin Oncol* 2012;9:674–87.

- [12] Horsman MR, Overgaard J. The impact of hypoxia and its modification of the outcome of radiotherapy. *J Radiat Res* 2016;57:i90–8.
- [13] Linge A, Lock S, Gudziol V, Low, et al. Cancer stem cell marker expression and low hypoxia identify good prognosis subgroups in HPV(-) HNSCC after postoperative radiochemotherapy. A multicenter study of the DTK-ROG. *Clin Cancer Res* 2016;22:2639–49.
- [14] Janssens GO, Rademakers SE, Terhaard CH, et al. Accelerated radiotherapy with carbogen and nicotinamide for laryngeal cancer. Results of a phase III randomized trial. *J Clin Oncol* 2012;30:1777–83.
- [15] Zeman EM, Brown JM, Lemmon MJ, Hirst VK, Lee WW. SR-4233: a new bioreductive agent with high selective toxicity for hypoxic mammalian cells. *Int J Radiat Oncol Biol Phys* 1986;12:1239–42.
- [16] Jentsch C, Beuthien-Baumann B, Troost EGC, Shakinin G. Validation of functional imaging as a biomarker for radiation treatment response. *Br J Radiol* 2015;88:20150014.
- [17] Rudat V, Vanselow B, Wollensack P, et al. Repeatability and prognostic impact of the pretreatment pO(2) histography in patients with advanced head and neck cancer. *Radiother Oncol* 2000;57:31–7.
- [18] Nordsmark M, Overgaard M, Overgaard J. Pretreatment oxygenation predicts radiation response in advanced squamous cell carcinoma of the head and neck. *Radiother Oncol* 1996;41:31–9.
- [19] Nordsmark M, Overgaard J. A confirmatory prognostic study on oxygenation status and loco-regional control in advanced head and neck squamous cell carcinoma treated by radiation therapy. *Radiother Oncol* 2000;57:39–43.
- [20] Brizel DM, Sibley GS, Prosnitz LR, Scher RL, Dewhirst MW. Tumor hypoxia adversely affects the prognosis of carcinoma of the head and neck. *Int J Radiat Oncol Biol Phys* 1997;38:285–9.
- [21] Stadler P, Becker A, Feldmann HJ, et al. Influence of the hypoxic subvolume on the survival of patients with head and neck cancer. *Int J Radiat Oncol Biol Phys* 1999;44:749–54.
- [22] Zips D, Zophel K, Abolmaali N, et al. Exploratory prospective trial of hypoxia-specific PET imaging during radiochemotherapy in patients with locally advanced head-and-neck cancer. *Radiother Oncol* 2012;105:21–8.
- [23] Thorwarth D, Eschmann S, Scheiderbauer J, Paulsen F, Alber M. Kinetic analysis of dynamic 18F-fluoromisonidazole PET correlates with radiation treatment outcome in head-and-neck cancer. *BMC Cancer* 2005;5:152.
- [24] Rajendran JG, Schwartz DL, O'Sullivan J, et al. Tumor hypoxia imaging with F-18 fluoromisonidazole positron emission tomography in head and neck cancer. *Clin Cancer Res* 2006;12:5435–41.
- [25] Kikuchi M, Yamane T, Shinohara S, et al. 18F-fluoromisonidazole positron emission tomography before treatment is a predictor of radiotherapy outcome and survival prognosis in patients with head and neck squamous cell carcinoma. *Ann Nucl Med* 2011;25:625–33.
- [26] Dirix P, Vandecaveye V, de Keyser F, Stroobants S, Hermans R, Nuyts S. Dose painting in radiotherapy for head and neck squamous cell carcinoma: value of repeated functional imaging with (18)F-FDG PET, (18)F-fluoromisonidazole PET, diffusion-weighted MRI, and dynamic contrast-enhanced MRI. *J Nucl Med* 2009;50:1020–7.
- [27] Eschmann SM, Paulsen F, Bedesheim C, et al. Hypoxia-imaging with (18)F-Misonidazole and PET: changes of kinetics during radiotherapy of head-and-neck cancer. *Radiother Oncol* 2007;83:406–10.
- [28] Servagi-Vernat S, Differding S, Hanin F, et al. A prospective clinical study of (1) (8)F-FAZA PET-CT hypoxia imaging in head and neck squamous cell carcinoma before and during radiation therapy. *Eur J Nucl Med Mol Imaging* 2014;41:1544–52.
- [29] Wiedenmann NE, Bucher S, Hentschel M, et al. Serial 18F-fluoromisonidazole PET during radiochemotherapy for locally advanced head and neck cancer and its correlation with outcome. *Radiother Oncol* 2015;117:113–7.
- [30] Rischin D, Hicks RJ, Fisher R, et al. Prognostic significance of 18F-misonidazole positron emission tomography-detected tumor hypoxia in patients with advanced head and neck cancer randomly assigned to chemoradiation with or without tirapazamine: a substudy of Trans-Tasman Radiation Oncology Group Study 98.02. *J Clin Oncol* 2006;24:2098–104.
- [31] Welz S, Monnich D, Pfannenberger C, et al. Prognostic value of dynamic hypoxia PET in head and neck cancer: Results from a planned interim analysis of a randomized phase II hypoxia-image guided dose escalation trial. *Radiother Oncol* 2017.
- [32] Gagel B, Reinartz P, Dimartino E, et al. pO(2) Polarography versus positron emission tomography ((18)F fluoromisonidazole, (18)F-2-fluoro-2'-deoxyglucose). An appraisal of radiotherapeutically relevant hypoxia. *Strahlenther Onkol* 2004;180:616–22.
- [33] Zimny M, Gagel B, Dimartino E, et al. FDG—a marker of tumour hypoxia? A comparison with 18F-fluoromisonidazole and pO2-polarography in metastatic head and neck cancer. *Eur J Nucl Med Mol Imaging* 2006;33:1426–31.
- [34] Troost Esther GC, Laverman P, Philippens Marielle EP, et al. Correlation of [18F] FMISO autoradiography and pimonidazole corrected immunohistochemistry in human head and neck carcinoma xenografts. *Eur J Nucl Med Mol Imaging* 2008;35:1803–11.
- [35] Troost Esther GC, Laverman P, Kaanders Johannes HAM, et al. Imaging hypoxia after oxygenation-modification: comparing [18F]FMISO autoradiography with pimonidazole immunohistochemistry in human xenograft tumors. *Radiother Oncol* 2006;80:157–64.
- [36] Hicks RJ, Rischin D, Fisher R, Binns D, Scott AM, Peters LJ. Utility of FMISO PET in advanced head and neck cancer treated with chemoradiation incorporating a hypoxia-targeting chemotherapy agent. *Eur J Nucl Med Mol Imaging* 2005;32:1384–91.
- [37] Jakobi A, Lühr A, Stutzer K, et al. Increase in tumor control and normal tissue complication probabilities in advanced head-and-neck cancer for dose-escalated intensity-modulated photon and proton therapy. *Front Oncol* 2015;5:256.
- [38] Löck S, Perrin R, Seidlitz A, et al. Residual tumour hypoxia in head-and-neck cancer patients undergoing primary radiochemotherapy, final results of a prospective trial on repeat FMISO-PET imaging. *Radiother Oncol* 2017;124:533–40.
- [39] Thorwarth D, Eschmann S, Holzner F, Paulsen F, Alber M. Combined uptake of [18F]FDG and [18F]FMISO correlates with radiation therapy outcome in head-and-neck cancer patients. *Radiother Oncol* 2006;80:151–6.
- [40] Haase R. A modular software system for image analysis in positron emission tomography. Workshop on Tomography, data processing and image reconstruction for medicine and engineering. Dresden; 2012.
- [41] Haase R. Optimization and validation of a swarm intelligence based segmentation algorithm for low contrast positron emission tomography. Dissertation, Dresden; 2014.
- [42] Tang G, Wang M, Tang X, Gan M, Luo L. Fully automated one-pot synthesis of [18F]fluoromisonidazole. *Nucl Med Biol* 2005;32:553–8.
- [43] Jakobi A, Bandurska-Luque A, Stützer K, et al. Identification of patient benefit from proton therapy for advanced head and neck cancer patients based on individual and subgroup normal tissue complication probability analysis. *Int J Radiat Oncol Biol Phys* 2015;92:1165–74.
- [44] Hofheinz F, Pöttsch C, Oehme L, et al. Automatic volume delineation in oncological PET. Evaluation of a dedicated software tool and comparison with manual delineation in clinical data sets. *Nuklearmedizin* 2012;51:9–16.
- [45] Bucher SA. F-18-Fluoromisonidazol PET zur Evaluation der Hypoxie in Kopf-Hals-Tumoren im Verlauf einer primären Radiochemotherapie. Dissertation, Freiburg; 2012.
- [46] Lee N, Nehmeh S, Schöder H, et al. Prospective trial incorporating pre-/mid-treatment [18F]-misonidazole positron emission tomography for head-and-neck cancer patients undergoing concurrent chemoradiotherapy. *Int J Radiat Oncol Biol Phys* 2009;75:101–8.
- [47] Yaromina A, Kroeber T, Meinzer A, et al. Exploratory study of the prognostic value of microenvironmental parameters during fractionated irradiation in human squamous cell carcinoma xenografts. *Int J Radiat Oncol Biol Phys* 2011;80:1205–13.
- [48] Rischin D, Fisher R, Peters L, Corry J, Hicks R. Hypoxia in head and neck cancer: studies with hypoxic positron emission tomography imaging and hypoxic cytotoxins. *Int J Radiat Oncol Biol Phys* 2007;69:S61–3.
- [49] Zegers Catharina ML, Hoebbers Frank JP, van Elmpt W, et al. Evaluation of tumour hypoxia during radiotherapy using (18)FHX4 PET imaging and blood biomarkers in patients with head and neck cancer. *Eur J Nucl Med Mol Imaging* 2016;43:2139–46.
- [50] Bandurska-Luque A. [18F]FMISO-PET/CT-based hypoxia in lymph node metastases in advanced head and neck cancers: baseline oxygenation status, reoxygenation and prognostic value. Dissertation, Dresden; 2016.
- [51] Zips D, Baumann M. Place of proton radiotherapy in future radiotherapy practice. *Semin Radiat Oncol* 2013;23:149–53.
- [52] Baumann M, Krause M, Overgaard J, et al. Radiation oncology in the era of precision medicine. *Nat Rev Cancer* 2016;16:234–49.
- [53] Linge A, Löck S, Gudziol V, et al. Low cancer stem cell marker expression and low hypoxia identify good prognosis subgroups in HPV(-) HNSCC after postoperative radiochemotherapy: a multicenter study of the DTK-ROG. *Clin Cancer Res* 2016;22:2639–49.