



Original Research

Functional movement screen and Y balance tests in adolescent footballers with hip/groin symptoms

Pawel Linek^{a, b, *}, Nadine Booyens^{b, d}, Damian Sikora^c, Maria Stokes^{b, d}^a Institute of Physiotherapy and Health Sciences, Musculoskeletal Elastography and Ultrasonography Laboratory, The Jerzy Kukuczka Academy of Physical Education, Katowice, Poland^b School of Health Sciences, University of Southampton, Southampton, UK^c Department of Kinesitherapy and Special Methods in Physiotherapy, The Jerzy Kukuczka Academy of Physical Education, Katowice, Poland^d Centre for Sport, Exercise and Osteoarthritis Research Versus Arthritis, UK

ARTICLE INFO

Article history:

Received 1 May 2019

Received in revised form

29 June 2019

Accepted 2 July 2019

Keywords:

Football

Soccer

Hip and groin pain

Children

Adolescent

Movement control

ABSTRACT

Objectives: To assess the correlation between the functional movement screen (FMS) and Y balance test (Y-BT) performance, and the self-reported hip/groin problems, and to compare healthy with hip/groin pain participants.

Design: a cross-sectional study.

Setting: Sports hall in a football club.

Participants: 43 elite adolescent football (soccer) players.

Main outcome: The Copenhagen Hip and Groin Outcome Score (HAGOS), Anterior, posterolateral and posteromedial distance of Y-BT, FMS score.

Measures: Y balance and FMS test kit, HAGOS questionnaire.

Results: The posterolateral, posteromedial distance and composite reach score of the Y-BT were lower in participants with hip/groin problems ($p < 0.05$). FMS performance was similar in healthy and hip/groin groups. There was a weak correlation ($r = 0.32$, $p = 0.03$) between the HAGOS Activities of Daily Living subscale and FMS_{stabil}. In the Y-BT the posterolateral reach was weakly correlated with Symptoms ($r = 0.35$ – 0.44 , $p < 0.02$) and moderately correlated with Pain ($r = 0.44$, $p < 0.01$) subscales. Posteromedial direction reach with the right leg was weakly correlated with Symptoms ($r = 0.32$, $p = 0.04$) and Pain ($r = 0.39$, $p = 0.01$). The Y-BT composite score was moderately correlated with Symptoms and Pain ($r = 0.42$ – 0.44 , $p < 0.01$).

Conclusions: The Y-BT and the FMS subtests were weakly or moderately correlated with self-reported hip/groin problems. Thus, these tests should be investigated further in adolescent footballers because they may have potential to predict hip and groin problems.

© 2019 Elsevier Ltd. All rights reserved.

1. Introduction

Hip/groin pain is common in footballers, as football is characterised by quick accelerations and decelerations as well as sprinting, jumping, turning and kicking, which create high loads and torsional forces on the hip (Saw & Villar, 2004). A recent epidemiological study by Kerbel, Smith, Prodromo, Nzeogu, and Mulcahey (2018) found that the overall hip/groin injury rate

amongst athletes was above 53 per 100,000 athlete-exposures and that football had the highest rates of hip/groin injuries. The prevalence of hip/groin symptoms was high amongst elite, sub-elite and amateur players, with 59% of male and 45% of female football players at difference levels of player reporting at least one episode of hip/groin symptoms in a study by Harøy et al. (2017). Two studies of adolescent footballers found that the incidence of hip/groin pain was about 14–22% during 9–14 weeks of observation (Crow et al., 2010; Lovell, Galloway, Hopkins, & Harvey, 2006). Additionally, the risk of experiencing hip/groin symptoms is similar between elite, sub-elite and amateur footballers (Harøy et al., 2017).

Studies have attempted to identify factors that differentiate between athletes with and without hip/groin pain by assessing,

* Corresponding author. Institute of Physiotherapy and Health Sciences, Musculoskeletal Elastography and Ultrasonography Laboratory, The Jerzy Kukuczka Academy of Physical Education, 40-065, Mikołowska 72B, Poland.

E-mail address: linek.fizjoterapia@vp.pl (P. Linek).

e.g.: a) performance on tests such as the hip adductor squeeze test, bent knee fall out test and active straight-leg raise test; b) hip adductor and abductor strength; c) range of motion of the hips; d) trunk muscle function; e) plain radiography (Besjakov, von Scheele, Ekberg, Gentz, & Westlin, 2003; Jansen et al., 2010; Malliaras, Hogan, Nawrocki, Crossley, & Schache, 2009; Mens, Inklaar, Koes, & Stam, 2006; Mohammad, Abdelraouf, Elhafez, Abdel-Aziem, & Nassif, 2014). A systematic review by Mosler, Agricola, Weir, Hölmich, and Crossley (2015) concluded that of these variables tested in different studies, reduced strength on the adductor squeeze test, reduced internal hip rotation and bent knee fall out were the measures that best differentiated between athletes with and without hip/groin pain.

The functional movement screen (FMS) (Cook, Burton, & Hoogenboom, 2006a; 2006b) and Y balance test (Y-BT) (Gribble, Hertel, & Plisky, 2012) are considered to assess athletes' risk of becoming injured and identify deficiencies in functional movement, neuromuscular control, balance and trunk stability. Some studies have shown that college athletes and American professional football players with a low composite score (≤ 14) and movement asymmetry score on the FMS are at high risk of injury (Garrison, Westrick, Johnson, & Benenson, 2015; Kiesel, Butler, & Plisky, 2014). Some systematic reviews presented controversy about the potential ability of the FMS to be used as an injury prediction tool (Bonazza, Smuin, Onks, Silvis, & Dhawan, 2017; Moran, Schneiders, Mason, & Sullivan, 2017; Whittaker et al., 2017). Two reviews (Moran et al., 2017; Whittaker et al., 2017) concluded the FMS was a poor predictor of injury-risk because of small and heterogeneous samples, inconsistent injury definitions, and lack of control for confounding variables (e.g. previous history of injury). However, the third review (Bonazza et al., 2017) concluded the FMS composite score demonstrates the injury predictive value. Additionally, a recent study found that adolescent athletes demonstrating asymmetrical movement during pre-season FMS testing were more likely to sustain an injury during the regular season than players without asymmetry (Chalmers et al., 2017). Asymmetry of more than 4 cm in anterior reach distance and a normalized composite reach score $\leq 89.6\%$ on the Y-BT were found to be associated with increased risk of noncontact injury in athletes (Butler, Lehr, Fink, Kiesel, & Plisky, 2013; Smith, Chimera, & Warren, 2015). Recently, Stiffler et al. (2017) showed that the assessment of side-to-side reach asymmetry in the anterior direction, which is part of the star excursion balance test (SEBT; the Y-BT is a modified version of the SEBT), may assist in identifying athletes who are at risk of sustaining noncontact injuries to the knee or ankle. Additionally, adolescent athletes with a history of lateral ankle sprain exhibit functional performance deficits on the SEBT (Ko, Rosen, & Brown, 2018). Furthermore another study demonstrated that performance of 15 × 30 m sprints with a period of 1 min rest between repetitions, was related to football players' SEBT results (Khan et al., 2016).

To the best of our knowledge there have not been any studies using the FMS or Y-BT in adolescent football players complaining of hip/groin symptoms. Given that both these functional tests are used routinely in athletes' management (injury risk screen, corrective and rehabilitation exercise programs, pre-participation examination), the authors considered that it would be useful to investigate how well they discriminated between athletes with and without hip/groin symptoms and pain. This knowledge may: a) improve understanding of factors contributing to the development of hip/groin pain; and b) help with the development rehabilitation strategies. The existing literature indicates that athletes with hip/groin pain have limited range of motion (internal rotation), weaker hip adductor (Mosler et al., 2015) and abductor (Kloskowska, Morrissey, Small, Malliaras, & Barton, 2016) muscles than peers

without such pain, so we hypothesized that athletes with more problematic hip/groin (demonstrated by diminished self-reported questionnaire score) would show reduced performance on the FMS and Y-BT. Such a relationship may be expected as both these tests are used to assess balance, postural control, and identify postural stability imbalances in mobility and stability during functional tasks. The aims of this preliminary study were, therefore, a) to assess the correlation between FMS and Y-BT performance, and the self-reported hip/groin problems; b) to compare adolescent football players with and without hip/groin pain.

2. Material and methods

2.1. Setting and study design

This was a cross-sectional study conducted at a professional football club in the Silesian region of Poland. All outcomes (FMS, Y-BT and HAGOS) were collected by experienced, qualified physiotherapists who were not informed of the purpose of the research, so as not to bias the results. The study was designed in accordance with the Declaration of Helsinki and was approved by the local medical ethics committee (Ethics Approval number: 4/2017, 18 May 2017). All participants and their parents and/or legal guardians received oral and written information about all procedures and gave written, informed consent to participation.

2.2. Sample

The sample consisted of 45 elite adolescent athletes (mean age: 15.6 years, min: 15, max: 17; mean body mass: 65.4 ± 8.22 kg; mean body height 176.6 ± 6.52 cm; BMI: 20.9 ± 1.86 kg/m²). Prior to the study all athletes had been playing football regularly for at least 3 years (mean: 7.74 years, min: 3, max: 11). This was a sample of convenience. The inclusion criterion were: a) had not experienced an injury that prevented participation in training or competition for longer than one week during the four months prior to the examination; b) able to perform the FMS and Y-BT. Athletes were excluded if they had any history of surgery or were unwilling to participate.

Included athletes were also divided into subgroups based on their Copenhagen Hip and Groin Outcome Score (HAGOS). The healthy athlete (HA) group consisted of participants who scored ≤ 1 (no more than 1 positive answer out of all questions contained in Symptoms, Activities of daily living - ADL, Sport and recreational activities - Sport, Participation in physical activity -PA, Quality of Life - QOL subscales) in the HAGOS and no positive answers out of all questions contained in Pain subscale of HAGOS. The hip and groin (H/G) group consisted of those who had at least 5 scores among all questions in the HAGOS questionnaire and at least one positive answer contained in the Pain subscale. Athletes who had 2, 3 or 4 scores among all questions in the HAGOS were not included in the between-group comparisons. Thus, those with a HAGOS score of 2–4 were arbitrarily excluded from between-group analysis. This categorization was to achieve sufficient separation between the two groups examined, and to enable a clear comparison.

2.3. The functional movement screen

The FMS consists of seven motor tasks: a deep squat, a hurdle step, an in-line lunge, a shoulder mobility test, an active straight-leg raise, a rotary stability test and a trunk stability push-up. Performance on all tasks was assessed on a 0 to 3 scale, where 3 indicates appropriate execution of the locomotor pattern, 2 indicates execution of the locomotor pattern with some compensatory adjustments, 1 indicates inability to perform the motor pattern and

0 indicates pain during movement. Each task was performed three times, and the best result was used for further analysis (Cook et al., 2006a; 2006b). In the case of tasks completed on left and right side, the lower score was used in the calculation of the total FMS score. The procedure used in similar studies of adolescent footballers was followed (Campa, Semprini, Júdece, Messina, & Toselli, 2019a; Portas, Parkin, Roberts, & Batterham, 2016). Three separate FMS scores were calculated for stability (FMSstab: the sum of scores on the 2 stability tests, trunk stability push-up and rotatory stability), flexibility (FMSflex: the sum of scores on the 2 mobility tests, shoulder mobility and active straight-leg raise) and movement (FMSmove: the sum of the 3 movement tests, the overhead squat, hurdle step and inline lung). In the case of tests completed on both the left and right sides the asymmetry in performance was also considered in further analysis.

2.4. Y balance test

The Y-BT was performed using the Y-balance test kit, which consists of a single central plastic plate to which three tubes are attached in anterior, posteromedial and posterolateral positions (Plisky et al., 2009). All Y-BT attempts were performed in the same order: anterior direction, posterolateral direction and finally posteromedial direction. Participants always started moving the pointer with their dominant leg. The measurement procedure consisted of nine attempts. The first six attempts were performed to familiarize the athlete with the test procedure; only the final three measurements were used for statistical analysis. The Y-BT protocol has been described in detail elsewhere (Linek, Sikora, Wolny, & Saulicz, 2017). The Y-BT was conducted by a qualified, experienced physiotherapist who was not informed of the purpose of the research.

To assess the normalized Y-BT results we measured the athlete's lower limb length - the distance from the anterior superior iliac spine to the medial malleolus. The measurement was performed with a tape measure, with the athlete in a supine position on a therapeutic table. This allowed the percentage value of the distance obtained in Y-BT to be calculated (normalization by the length of the lower limb). The normalization was performed by using following formula: (the average Y-BT distance from the last three trials/relative length of the limb) x 100. Additionally, the sum of the reach distance for each of the three directions was divided by three times the limb length and multiplied by 100 = [(anterior + posterolateral + posteromedial)/(limb length x3)] x 100, to calculate composite reach distance for each leg (Plisky, Rauh, Kaminski, & Underwood, 2006).

2.5. The Copenhagen Hip and Groin Outcome Score (HAGOS)

Before the FMS and Y-BT were performed, all players completed paper form of the English version of HAGOS questionnaire (Thorborg, Hölmich, Christensen, Petersen, & Roos, 2011). There are six HAGOS subscales: Pain, Symptoms, ADL, Sport, PA and QOL. Responses were given using a five-point Likert scale ranging from 0 to 4 (0 = no hip/groin problem; 4 = extreme hip/groin symptoms). Raw scores were then transformed into percentages, so all subscales were scored on a 0–100 scale, where lower scores indicate more hip/groin symptoms. The questions were answered considering hip and groin function during the past week.

One week prior to the study, all participants were familiarized with the HAGOS. They were informed that one week later they would be asked to fill out the HAGOS. They were encouraged to focus on how their hip and groin were functioning during the forthcoming week. During the meeting all items and questions were also detailed and explained by a polish speaking physiotherapist fluent in English. Questionnaires were filled out in a separate

room, with only one athlete and the physiotherapist present. The physiotherapist was available to answer any questions whilst participants were completing the questionnaire.

2.6. Statistical analysis

Levene's test was used to assess the homogeneity of variances. Demographic, FMS and Y-BT differences were assessed with the independent *t*-test or, in the case of variables for which the variances were unequal, Welch's test. The results are presented as mean differences with 95% confidence intervals (CIs). Mean values, standard deviations and number of participants were used to assess effect sizes. Hedges's *g* (H_g) was used as an index of effect size for between-subject effects and common language (CL) effect sizes (Cumming, 2012; McGraw & Wong, 1992) are also reported. The criteria for H_g effect size were as follows: trivial (<0.20), small (0.20–0.49), medium (0.50–0.79) and large (≥ 0.80).

Data from all participants was used to examine correlations between HAGOS subscales and Y balance and FMS data were analysed using Spearman's *r* and interpreted as negligible (0.00–0.10), weak (0.10–0.39), moderate (0.40–0.69), strong (0.70–0.89), very strong (0.90–1.00) (Schober, Boer, & Schwarte, 2018). All statistical analyses were performed with the Statistica 12 PL software and *p*-values < 0.05 were considered significant.

3. Results

3.1. Hip/groin symptoms

Two of the original 45 athletes were excluded from the sample because they did not answer all the HAGOS questions. Only 10 of the 43 athletes had scores of 100% on all the subscales; the remaining participants (77.8%) reported some hip/groin symptoms. Mean scores on the subscales, with minimum scores in parentheses, were as follows, Symptoms 91.9 (67.8); Pain 97.1 (72.5); ADL 97.8 (85); Sport 94.9 (43.7); PA 97.1 (75); QOL 92.4 (35). Nineteen athletes met criteria to be included in the healthy athlete (HA) group, whereas 16 athletes met criteria to be included in hip and groin (H/G) group. Detailed profiles of the HA and H/G groups are given in Table 1. Thus, between-group comparisons were performed on data from 35 athletes, whereas correlation analyses were performed with the full sample of 43 athletes.

Among the HAGOS questions, the most common positive answers were for question no. 1 (Do you feel discomfort in your hip and/or groin?) and 3 (Do you have difficulties stretching your legs far out to the side?) in the Symptoms subscale.

3.2. Functional movement screen score

There was no difference between the FMS performance of the HA and H/G groups. Detailed FMS data with 95% CIs are presented in Table 2. Analysis of asymmetry in performance on the bilateral tests also revealed no group differences. Additionally, there were no participants reporting pain during the FMS.

3.3. Y balance test

The posterolateral distance gained was lower in the H/G group than the HA group (left leg: $H_g = 0.77$, CL = 71%). Posteromedial distance was also lower in the H/G group, but for the right lower leg ($H_g = 0.72$, CL = 70%). Composite-reach score (total Y score) was lower in the H/G group than the HA group ($H_g = 0.72$, CL = 70%). There were no group differences in side-to-side asymmetry with respect to any of the distances investigated. The anterior distance was similar in both groups. Detailed Y balance data with 95% CIs

Table 1
Basic data of participants.

Characteristic	Athletes ¹		p value
	HA (n = 19)	H/G (n = 16)	
Age (yr)	15.8 (0.53)	15.5 (0.51)	0.11 ^b
Weight (kg)	66.6 (7.43)	63.7 (10.4)	0.33 ^b
Height (cm)	177.2 (4.80)	175.2 (9.27)	0.46 ^c
BMI (kg/m ²)	21.2 (2.00)	20.6 (1.82)	0.35 ^b
Sports practice (yr)	7.76 (1.58)	7.3 (1.95)	0.75 ^b
Right dominant leg ^a	77.8%	92.8%	0.35 ^d
HAGOS (%)			
Symptoms	98.5	82.6	<0.001 ^{e*}
Pain	100	92.2	<0.001 ^{e*}
ADL	99.7	94.7	0.02 ^{e*}
Sport	100	86.3	<0.001 ^{e*}
PA	100	93.8	0.06 ^c
QOL	99.7	82.8	<0.01 ^{e*}

¹without those with 2, 3 or 4 scores among all HAGOS questions; HA – Healthy Athlete Group (no more than 1 positive answer out of all HAGOS questions); H/G – Hip and Groin Group (at least 5 positive answers among all HAGOS questions) HAGOS – Hip and Groin Outcome Score; ADL – Activities of Daily Living; Sport – Sport and Recreational Activities; PA – Participation in Physical Activity; QOL – Quality of Living.

^a Which leg do you prefer when playing football?

^b test t for independent samples.

^c Welch's test.

^d from Fischer's exact test.

^e from Mann–Whitney test; *significant differences.

Table 2
The mean (SD) of groups and the mean (95% CI) of differences between groups in FMS test.

FMS	Groups		Difference between groups	
	HA	H/G	HA minus H/G	p value
Total score	15.8 (2.45)	14.6 (2.33)	1.22 (–0.54 – 2.98)	0.17 ^b
Move score	6.52 (1.47)	6.18 (1.11)	0.33 (–0.57 – 1.25)	0.45 ^b
Flex score	4.53 (1.26)	4.00 (1.21)	0.52 (–0.32 – 1.38)	0.22 ^b
Stabil score	4.79 (0.63)	4.44 (0.89)	0.35 (–0.17 – 0.88)	0.18 ^b
Asymmetry incidence – number (%):				
HS	11 (61%)	6 (43%)	–	0.48 ^d
INLL	9 (50%)	5 (36%)	–	0.49 ^d
SM	7 (39%)	6 (43%)	–	1.00 ^d
ASLR	8 (44%)	9 (64%)	–	0.31 ^d
RS	5 (28%)	3 (21%)	–	1.00 ^d

HA – Healthy Athlete Group; H/G – Hip and Groin Group; HS – hurdle step; INLL – in-line lunge; SM – shoulder mobility; ASLR – active straight leg raise; RS – rotation stability.

^b test t for independent samples.

^d from Fischer's exact test.

Table 3
The mean (SD) of groups and the mean (95% CI) of differences between groups in Y balance test.

Reach direction (%)	Groups		Difference between groups	
	HA	H/G	HA minus H/G	p value
Anterior – R	66.9 (6.98)	66.7 (5.12)	0.18 (–3.99 – 4.35)	0.93 ^b
Posterolateral – R	110.2 (10.4)	104.9 (6.17)	5.29 (–0.52 – 11.1)	0.07 ^c
Posteromedial – R	108.5 (9.06)	103.1 (5.77)	5.42 (0.25 – 10.6)	0.04 ^{b*}
Anterior – L	67.8 (7.21)	66.8 (5.92)	0.95 (–3.57 – 5.47)	0.67 ^b
Posterolateral – L	113.3 (11.7)	105.7 (6.31)	7.59 (1.21 – 13.9)	0.02 ^{c*}
Posteromedial – L	108.6 (10.5)	105.1 (7.01)	3.50 (–2.59 – 9.59)	0.26 ^b
Anterior – A	2.45 (2.26)	2.41 (1.38)	0.04 (–1.23 – 1.31)	0.95 ^b
Posterolateral – A	5.65 (8.81)	3.65 (2.70)	1.99 (–2.42 – 6.42)	0.36 ^c
Posteromedial – A	3.11 (2.92)	4.14 (2.54)	–1.03 (–2.91 – 0.84)	0.27 ^b
Composite score – T	96.7 (7.4)	92.2 (4.57)	4.50 (0.33 – 8.68)	0.04 ^{b*}

HA – Healthy Athlete Group; H/G – Hip and Groin Group; R – reach distance by right leg; L – reach distance by left leg; A – relative side-to-side asymmetry; T – sum of anterior, posteromedial and posterolateral distance divided by 3; *significant differences.

^b Test t for independent samples.

^c Welch's test.

and p values are presented in Table 3.

3.4. Correlations between HAGOS and other variables

The only correlation found between HAGOS and the FMS was a positive weak correlation between FMS_{stabil} and the ADL subscale of HAGOS ($r = 0.32$, $p = 0.03$). There were no correlations between FMS_{total}, FMS_{move}, FMS_{flex} and the HAGOS subscales (all $p > 0.05$).

The Y-BT results were only correlated with the Symptoms and Pain subscales of HAGOS. Posterolateral reach was correlated (from weak to moderate) with Symptoms (right leg: $r = 0.35$, $p = 0.02$; left leg: $r = 0.44$, $p = 0.01$) and Pain (right: $r = 0.42$, $p < 0.01$; left: $r = 0.43$, $p < 0.01$) (Fig. 1). Posteromedial direction reach with the right leg was weak correlated with Symptoms ($r = 0.32$, $p = 0.04$) and Pain ($r = 0.38$, $p = 0.01$) (Fig. 2). Composite score was moderate correlated with Symptoms ($r = 0.44$, $p < 0.01$) and Pain ($r = 0.42$, $p < 0.01$) (Fig. 3). In all cases athletes with lower HAGOS subscale values received lower scores in Y-BT. Posteromedial reach with the left leg and anterior reach (both legs) were not correlated with the HAGOS (all $p > 0.05$).

4. Discussion

The present study has shown that 10 out of 43 of the adolescent footballers examined did not have any hip/groin symptoms as measured by the self-reported HAGOS (100% on all subscales). A further 9 athletes reported one positive answer (among all HAGOS subscales one was below 100%). Minor problems were detected among 8 participants (2–4 scores among all questions) and 16 footballers reported more serious hip/groin symptoms (at least 5 scores among all questions in HAGOS questionnaire) confirming that hip/groin problems are common in adolescent athletes (Karlsson, Dahan, Magnusson, Nyquist, & Rosengren, 2014). It also confirms the need for further studies to identify factors that differentiate healthy athletes from those complaining of hip/groin symptoms. The aims of the study were a) to assess the correlation between FMS and Y-BT performance, and the HAGOS subscales in adolescent football players; and b) to compare adolescent football players with and without hip/groin pain. To the best of our knowledge, there have not been any other studies of FMS and Y-BT performance in this population. We found that adolescent football players with hip/groin symptoms and pain had reduced posterolateral reach (left leg) and posteromedial reach (right leg) on the Y-BT compared with healthy athletes and this was reflected in lower composite scores in the H/G group. Additionally, posteromedial, posterolateral and composite Y-BT results were correlated with hip/




Fig. 1. Posterolateral reach distance and Symptoms and Pain distribution. Each participant was labelled according to the group they belonged to for between-group comparisons: 'square'- hip and groin pain; 'short horizontal line' - healthy athletes; letter 'R' - removed from between-group comparison due to not meeting the criteria.




Fig. 2. Posteromedial reach distance and Symptoms and Pain distribution. Each participant was labelled according to the group they belonged to for between-group comparisons: 'square'- hip and groin pain; 'short horizontal line' - healthy athletes; letter 'R' - removed from between-group comparison due to not meeting the criteria.

groin Symptoms and Pain subscales of HAGOS, whereas the trunk stability push-up and rotatory stability tests of the FMS were correlated with hip/groin Symptoms subscale of HAGOS.

The effect sizes for group differences in reaching in all directions and in composite score were medium, and there was a 70% probability in randomly selected pairs of healthy and symptomatic adolescent footballers that the symptomatic footballer would have a shorter posterolateral reach distance (left leg) and shorter posteromedial reach distance (right leg). The mean between-group differences varied from 5% to 8%. A reliability study of the Y-BT in adolescent football players (Linek et al., 2017) showed fair and good intra-rater reliability for Y-BT measurements with the minimal

detectable change being 7–14%. The mean group differences in Y-BT performance in the present sample were below the minimum practically significant difference. Although the between-group differences did not reach the benchmark of 7–14% for Y-BT measurements, the changes found in the current study may potentially be higher (and clinically relevant) among participants with more serious hip/groin problems and pain. Our correlation analyses suggest this was the case, because the Symptoms and Pain subscales of HAGOS were positively correlated with Y-BT results, which means that reduced Y-BT performance may be partly due to higher hip/groin symptoms and pain. This is plausible, because athletes with hip/groin symptoms have reduced internal hip rotation and




Fig. 3. Composite score of Y balance test and Symptoms and Pain distribution. Each participant was labelled according to the group they belonged to for between-group comparisons: 'square' - hip and groin pain; 'short horizontal line' - healthy athletes; letter 'R' - removed from between-group comparison due to not meeting the criteria.

reduced hip adductor (Mosler et al., 2015) and abductor (Kloskowska et al., 2016) strength. Recently, López-Valenciano, Ayala, De Ste Croix, Barbado, and Vera-García (2018) revealed that along with core stability and hip abduction isometric peak torque, passive hip flexion is one of the main variables associated with Y-BT performance in football players. Studies of other populations have reported that Y-BT performance is correlated with hip abductor strength and extension and external rotation of the hips (Wilson et al., 2018) as well as the combination of hip flexion and lateral trunk bending (Kang et al., 2015).

The Y-BT can also be considered as a postural control test. In our study most athletes in the H/G group showed rather minor hip/groin problems. These minor symptoms in such a young population maybe be the start of motor control adaptations, which then in turn may lead to more movement alterations, pain and injury. Additionally, side-to side asymmetry with respect to any of the distances investigated in the Y-BT was similar in the healthy and hip/groin athletes. This suggests that the presence of symptoms in the hip/groin group did not significantly affect the distance reached on only one side of the body (symptomatic or asymptomatic).

Hodges (2011) claims that pain lead to changes in motor control, involving redistribution of muscle activity, changes to the mechanical behaviour causing altered movements and stiffness, leading to protection from further pain or injury and multiple level motor system changes. This may explain the correlation found between the HAGOS Pain subscale and Y-BT results in the present study. Thus, the Y-BT is a complex movement test and many aspects of the tasks depend directly on the hip joint, so it may be useful in further studies of adolescent footballers complaining of hip/groin symptoms.

In the present study the FMS did not differentiate between healthy athletes and those complaining of hip/groin symptoms. There was, however, a weak correlation between FMS_{stabil} and the ADL subscale of HAGOS. The FMS_{stabil} score is based on the trunk stability push-up and rotatory stability tests. The trunk stability push-up requires a combination of trunk and upper extremity stability, whilst the rotatory stability test requires multi-plane trunk stability in conjunction with simultaneous motion of the upper and lower extremities (Cook et al., 2006a; 2006b). Improper performance of these tests may be due to inadequate stability of the trunk. According to Agresta, Slobodinsky, and Tucker (2014), athletes with lower rotatory stability FMS scores have impaired control of the trunk, pelvis and hip muscles, which contributes to abnormalities of lower extremity mechanics (Powers, 2010). Some studies have shown that athletes with hip/groin symptoms have

changes to the deepest trunk stability muscle – the transversus abdominis muscle (Cowan et al., 2004; Jansen et al., 2010). Thus the tests that contribute to the FMS_{stabil} score may have the greatest potential of all the FMS tests to discriminate between footballers with and without hip/groin problems. Recently, Campa, Spiga, and Toselli (2019b) have shown that a 20-week exercise program improved all FMS components (including FMS_{stabil}) in adolescent footballers. Similarly, exercise protocols used by Dinc, Kilinc, Bulat, Erten, and Bayraktar (2017) improved trunk stability push-up (test included in FMS_{stabil}) in adolescent footballers. It would therefore be worth examining whether exercise programs could change FMS and functional scores in footballers with hip and groin symptoms.

A limitation of the present findings is that they may only apply to the group examined, i.e. professional male adolescent football players. They may not generalise to athletes involved in other sports or to female footballers and any extrapolations should be made with caution. Another limitation of the study is the relatively small sample size and the minor nature of the hip/groin symptoms reported. However, other studies using the FMS and/or functional tests used similar sample sizes (Agresta et al., 2014; Butler et al., 2013; Kang et al., 2015; Ko et al., 2018; McCann et al., 2017). We would expect footballers with more prominent hip/groin symptoms to show poorer performance on the Y-BT.

5. Conclusions

Adolescent football players with hip/groin problems (including pain) had reduced posterolateral reach, posteromedial reach and composite scores on the Y-BT compared with healthy peers. These Y-BT results were also correlated (from weak to moderate) with hip/groin symptoms and pain. The FMS performance was not different between the healthy and hip/groin pain groups. However, the trunk stability push-up and rotatory stability tests of the FMS were weakly correlated with hip/groin symptoms. These tests are quick and easy to use in the field or clinical practice, having the potential to screen and/or predict players with hip/groin symptoms, as suggested by the present results. This could assist clinicians in determining which players may require further evaluation, and when to initiate early management and preventative interventions. Also, pain may lead to motor control changes, which could be a modifiable risk factor and, if recognised early, could be managed by timely implementation of preventative movement control exercises. This could consequently have an impact on preventing the development of hip and groin pain in this population.

Thus, performance of movement control tests and interventions should be investigated further in future studies of adolescent footballers.

Conflicts of interest

None declared.

Ethical approval

Attached in separate file.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Agresta, C., Slobodinsky, M., & Tucker, C. (2014). Functional movement ScreenTM–normative values in healthy distance runners. *Int J Sports Med*, 35, 1203–1207. <https://doi.org/10.1055/s-0034-1382055>.
- Besjakov, J., von Scheele, C., Ekberg, O., Gentz, C. F., & Westlin, N. E. (2003). Grading scale of radiographic findings in the pubic bone and symphysis in athletes. *Acta Radiologica*, 44, 79–83.
- Bonazza, N. A., Smuin, D., Onks, C. A., Silvis, M. L., & Dhawan, A. (2017). Reliability, validity, and injury predictive value of the functional movement screen: A systematic review and meta-analysis. *The American Journal of Sports Medicine*, 45, 725–732. <https://doi.org/10.1177/0363546516641937>.
- Butler, R. J., Lehr, M. E., Fink, M. L., Kiesel, K. B., & Plisky, P. J. (2013). Dynamic balance performance and noncontact lower extremity injury in college football players: An initial study. *Sport Health*, 5, 417–422. <https://doi.org/10.1177/1941738113498703>.
- Campa, F., Semprini, G., Júdece, P., Messina, G., & Toselli, S. (2019a). Anthropometry, physical and movement features, and repeated-sprint ability in soccer players. *Int J Sports Med*, 40, 100–109. <https://doi.org/10.1055/a-0781-2473>.
- Campa, F., Spiga, F., & Toselli, S. (2019b). The effect of a 20-week corrective exercise program on functional movement patterns in youth elite male soccer players. *Journal of Sport Rehabilitation*, 1–6. <https://doi.org/10.1123/jsr.2018-0039>.
- Chalmers, S., Fuller, J. T., DeBenedictis, T. A., Townsley, S., Lynagh, M., Gleeson, C., et al. (2017). Asymmetry during preseason Functional Movement Screen testing is associated with injury during a junior Australian football season. *Journal of Science and Medicine in Sport*, 20, 653–657. <https://doi.org/10.1016/j.jsams.2016.12.076>.
- Cook, G., Burton, L., & Hoogenboom, B. (2006a). Pre-participation screening: The use of fundamental movements as an assessment of function - part 1. *N Am J Sports Phys Ther*, 1, 62–72.
- Cook, G., Burton, L., & Hoogenboom, B. (2006b). Pre-participation screening: The use of fundamental movements as an assessment of function - part 2. *N Am J Sports Phys Ther*, 1, 132–139.
- Cowan, S. M., Schache, A. G., Brukner, P., Bennell, K. L., Hodges, P. W., Coburn, P., et al. (2004). Delayed onset of transversus abdominus in long-standing groin pain. *Medicine & Science in Sports & Exercise*, 36, 2040–5.
- Crow, J. F., Pearce, A. J., Veale, J. P., VanderWesthuizen, D., Coburn, P. T., & Pizzari, T. (2010). Hip adductor muscle strength is reduced preceding and during the onset of groin pain in elite junior Australian football players. *Journal of Science and Medicine in Sport*, 13, 202–204. <https://doi.org/10.1016/j.jsams.2009.03.007>.
- Cumming, G. (2012). *Understanding the new statistics: Effect sizes, confidence intervals, and meta-analysis*. New York, NY: Routledge.
- Dinc, E., Kilinc, B. E., Bulat, M., Erten, Y. T., & Bayraktar, B. (2017). Effects of special exercise programs on functional movement screen scores and injury prevention in preprofessional young football players. *Journal of Exerc Rehabilitation*, 13, 535–540. <https://doi.org/10.12965/jer.1735068.534>.
- Garrison, M., Westrick, R., Johnson, M. R., & Benenson, J. (2015). Association between the functional movement screen and injury development in college athletes. *Int J Sports Phys Ther*, 10, 21–28.
- Gribble, P. A., Hertel, J., & Plisky, P. (2012). Using the star excursion balance test to assess dynamic postural-control deficits and outcomes in lower extremity injury: A literature and systematic review. *Journal of Athletic Training*, 47, 339–357. <https://doi.org/10.4085/1062-6050-47.3.08>.
- Harøy, J., Clarsen, B., Thorborg, K., Hölmich, P., Bahr, R., & Andersen, T. E. (2017). Groin problems in male soccer players are more common than previously reported. *The American Journal of Sports Medicine*. , 036354651668753. <https://doi.org/10.1177/0363546516687539>.
- Hodges, P. W. (2011). Pain and motor control: From the laboratory to rehabilitation. *Journal of Electromyography and Kinesiology*, 21, 220–228. <https://doi.org/10.1016/j.jelekin.2011.01.002>.
- Jansen, J., Weir, A., Denis, R., Mens, J., Backx, F., & Stam, H. (2010). Resting thickness of transversus abdominis is decreased in athletes with longstanding adduction-related groin pain. *Man Ther*, 15, 200–205. <https://doi.org/10.1016/j.math.2009.11.001>.
- Kang, M.-H., Kim, G.-M., Kwon, O.-Y., Weon, J.-H., Oh, J.-S., & An, D.-H. (2015). Relationship between the kinematics of the trunk and lower extremity and performance on the Y-balance test. *PM&R*, 7, 1152–1158. <https://doi.org/10.1016/j.pmrj.2015.05.004>.
- Karlsson, M. K., Dahan, R., Magnusson, H., Nyquist, F., & Rosengren, B. E. (2014). Groin pain and soccer players: Male versus female occurrence. *J Sports Med Phys Fitness*, 54, 487–493.
- Kerbel, Y. E., Smith, C. M., Prodrromo, J. P., Nzeogu, M. I., & Mulcahey, M. K. (2018). Epidemiology of hip and groin injuries in collegiate athletes in the United States. *Orthop J Sport Med*, 6. <https://doi.org/10.1177/2325967118771676>, 2325967118771676.
- Khan, M. A., Moiz, J. A., Raza, S., Verma, S., Shareef, M. Y., Anwer, S., et al. (2016). Physical and balance performance following exercise induced muscle damage in male soccer players. *Journal of Physical Therapy Science*, 28, 2942–2949. <https://doi.org/10.1589/jpts.28.2942>.
- Kiesel, K. B., Butler, R. J., & Plisky, P. J. (2014). Prediction of injury by limited and asymmetrical fundamental movement patterns in American football players. *Journal of Sport Rehabilitation*, 23, 88–94. <https://doi.org/10.1123/jsr.2012-0130>.
- Kloskowska, P., Morrissey, D., Small, C., Malliaras, P., & Barton, C. (2016). Movement patterns and muscular function before and after onset of sports-related groin pain: A systematic review with meta-analysis. *Sports Medicine*, 46, 1847–1867. <https://doi.org/10.1007/s40279-016-0523-z>.
- Ko, J., Rosen, A. B., & Brown, C. N. (2018). Functional performance deficits in adolescent athletes with a history of lateral ankle sprain(s). *Physical Therapy in Sport*, 33, 125–132. <https://doi.org/10.1016/j.ptsp.2018.07.010>.
- Linek, P., Sikora, D., Wolny, T., & Saulicz, E. (2017). Reliability and number of trials of Y Balance Test in adolescent athletes. *Musculoskeletal Sci Pract*, 31. <https://doi.org/10.1016/j.msksp.2017.03.011>.
- López-Valenciano, A., Ayala, F., De Ste Croix, M., Barbado, D., & Vera-García, F. J. (2018). Different neuromuscular parameters influence dynamic balance in male and female football players. *Knee Surgery, Sport Traumatol Arthrosc*. <https://doi.org/10.1007/s00167-018-5088-y>.
- Lovell, G., Galloway, H., Hopkins, W., & Harvey, A. (2006). Osteitis pubis and assessment of bone marrow edema at the pubic symphysis with MRI in an elite junior male soccer squad. *Clinical Journal of Sports Medicine*, 16, 117–122.
- Malliaras, P., Hogan, A., Nawrocki, A., Crossley, K., & Schache, A. (2009). Hip flexibility and strength measures: Reliability and association with athletic groin pain. *British Journal of Sports Medicine*, 43, 739–744. <https://doi.org/10.1136/bjism.2008.055749>.
- McCann, R. S., Kosik, K. B., Terada, M., Beard, M. Q., Buskirk, G. E., & Gribble, P. A. (2017). Associations between functional and isolated performance measures in college women's soccer players. *Journal of Sport Rehabilitation*, 26, 376–385. <https://doi.org/10.1123/jsr.2016-0016>.
- McGraw, K. O., & Wong, S. P. (1992). A common language effect size statistic. *Psychological Bulletin*, 111, 361–365. <https://doi.org/10.1037/0033-2909.111.2.361>.
- Mens, J., Inklaar, H., Koes, B. W., & Stam, H. J. (2006). A new view on adduction-related groin pain. *Clinical Journal of Sports Medicine*, 16, 15–19.
- Mohammad, W. S., Abdelraouf, O. R., Elhafez, S. M., Abdel-Aziem, A. A., & Nassif, N. S. (2014). Isokinetic imbalance of hip muscles in soccer players with osteitis pubis. *Journal of Sports Science*, 32, 934–939. <https://doi.org/10.1080/02640414.2013.868918>.
- Moran, R. W., Schneiders, A. G., Mason, J., & Sullivan, S. J. (2017). Do functional movement screen (FMS) composite scores predict subsequent injury? A systematic review with meta-analysis. *British Journal of Sports Medicine*, 51, 1661–1669. <https://doi.org/10.1136/bjsports-2016-096938>.
- Mosler, A. B., Agricola, R., Weir, A., Hölmich, P., & Crossley, K. M. (2015). Which factors differentiate athletes with hip/groin pain from those without? A systematic review with meta-analysis. *British Journal of Sports Medicine*, 49, 810. <https://doi.org/10.1136/bjsports-2015-094602>.
- Plisky, P. J., Rauh, M. J., Kaminski, T. W., & Underwood, F. B. (2006). Star excursion balance test as a predictor of lower extremity injury in high school basketball players. *Journal of Orthopaedic & Sports Physical Therapy*, 36, 911–919. <https://doi.org/10.2519/jospt.2006.2244>.
- Plisky, P. J., Gorman, P. P., Butler, R. J., Kiesel, K. B., Underwood, F. B., & Elkin, B. (2009 May). The reliability of an instrumented device for measuring components of the star excursion balance test. *North American Journal of Sports Physical Therapy*, 4(2), 92–99. <https://doi.org/10.2519/jospt.2006.2244>.
- Portas, M. D., Parkin, G., Roberts, J., & Batterham, A. M. (2016). Maturational effect on Functional Movement ScreenTM score in adolescent soccer players. *Journal of Science and Medicine in Sport*, 19, 854–858. <https://doi.org/10.1016/j.jsams.2015.12.001>.
- Powers, C. M. (2010). The influence of abnormal hip mechanics on knee injury: A biomechanical perspective. *Journal of Orthopaedic & Sports Physical Therapy*, 40, 42–51. <https://doi.org/10.2519/jospt.2010.3337>.
- Saw, T., & Villar, R. (2004). Footballer's hip a report of six cases. *J Bone Joint Surg Br*, 86, 655–658.
- Schober, P., Boer, C., & Schwarte, L. A. (2018). Correlation coefficients. *Anesthesia & Analgesia*, 126, 1763–1768. <https://doi.org/10.1213/ANE.0000000000002864>.
- Smith, C. A., Chimeria, N. J., & Warren, M. (2015). Association of y balance test reach asymmetry and injury in division I athletes. *Medicine & Science in Sports & Exercise*, 47, 136–141. <https://doi.org/10.1249/MSS.0000000000000380>.
- Stiffler, M. R., Bell, D. R., Sanfilippo, J. L., Hettzel, S. J., Pickett, K. A., & Heiderscheit, B. C. (2017). Star excursion balance test anterior asymmetry is

- associated with injury status in division I collegiate athletes. *Journal of Orthopaedic & Sports Physical Therapy*, 47, 339–346. <https://doi.org/10.2519/jospt.2017.6974>.
- Thorborg, K., Hölmich, P., Christensen, R., Petersen, J., & Roos, E. M. (2011). The copenhagen hip and groin outcome score (HAGOS): Development and validation according to the COSMIN checklist. *British Journal of Sports Medicine*, 45, 478–491. <https://doi.org/10.1136/bjism.2010.080937>.
- Whittaker, J. L., Booyesen, N., de la Motte, S., Dennett, L., Lewis, C. L., Wilson, D., et al. (2017). Predicting sport and occupational lower extremity injury risk through movement quality screening: A systematic review. *British Journal of Sports Medicine*, 51, 580–585. <https://doi.org/10.1136/bjsports-2016-096760>.
- Wilson, B. R., Robertson, K. E., Burnham, J. M., Yonz, M. C., Ireland, M. L., & Noehren, B. (2018). The relationship between hip strength and the Y balance test. *Journal of Sport Rehabilitation*, 27, 445–450. <https://doi.org/10.1123/jsr.2016-0187>.