




Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en



SFORL Guidelines

Guidelines (short version) of the French Society of Otorhinolaryngology (SFORL) on pediatric cochlear implant indications



F. Simon^{a,*}, S. Roman^b, E. Truy^c, P. Barone^d, J. Belmin^e, C. Blanchet^f, S. Borel^g, A. Charpiot^h, A. Coezⁱ, O. Deguine^j, A. Farinetti^b, B. Godey^k, D. Lazard^{l,m}, M. Marxⁿ, I. Mosnier^o, Y. Nguyen^p, N. Teissier^q, B. Virole^r, E. Lescanne^s, N. Loundon^a

^a Service d'ORL pédiatrique, hôpital Necker – Enfants-Malades, université Paris Descartes, 75015 Paris, France

^b Service d'ORL pédiatrique, hôpital La Timone, Aix-Marseille université, 13005 Marseille, France

^c Service ORL et chirurgie cervico-faciale, hôpital Édouard-Herriot, université Claude-Bernard Lyon 1, 69003 Lyon, France

^d Université de Toulouse, CerCo UMR 5549 CNRS, faculté de médecine de Purpan, 31000 Toulouse, France

^e Hôpital universitaire Charles-Foix, université Pierre-et-Marie-Curie, 75013 Paris, France

^f Service d'ORL pédiatrique, hôpital Arnaud-de-Villeneuve, 34090 Montpellier, France

^g Université de Tours, 37000 Tours, France

^h Service d'ORL, CHU de Strasbourg, 67200 Strasbourg, France

ⁱ CEA-Inserm U1000 Neuroimaging and Psychiatry, service hospitalier Frédéric-Joliot, 91400 Orsay, France

^j Service d'ORL, CHU de Toulouse, 31300 Toulouse, France

^k Service d'oto-rhino-laryngologie et de chirurgie maxillo-faciale, CHU de Rennes, 35000 Rennes, France

^l Institut Arthur-Vernes, 75006 Paris, France

^m Nottingham University, Nottingham, UK

ⁿ Service d'ORL, hôpital Purpan, CHU de Toulouse, université Paul-Sabatier, 31000 Toulouse, France

^o AP-HP, groupe hospitalier Pitié-Salpêtrière, service d'ORL, otologie, implants auditifs et chirurgie de la base du crâne, Sorbonne universités, université Pierre-et-Marie-Curie, 75013 Paris, France

^p Groupe hospitalier Pitié-Salpêtrière, unité otologie, implants auditifs et chirurgie de la base du crâne, UMR-S 1159 Inserm, université Paris 6 Pierre-et-Marie-Curie, 75013 Paris, France

^q Service d'ORL pédiatrique, hôpital Robert-Debré, université Paris-Diderot, Paris, France

^r Service d'ORL et audiophonologie infantile, hôpital Robert-Debré, 75019 Paris, France

^s Service d'ORL, CHU de Tours, 37000 Tours, France

ARTICLE INFO

Keywords:

Child
 Pediatric
 Cochlear implant
 Indications
 Guidelines
 France

ABSTRACT

Objectives: The authors present the guidelines of the French Society of Otorhinolaryngology – Head and Neck Surgery (Société française d'oto-rhino-laryngologie et de chirurgie de la face et du cou – SFORL) on the indications for cochlear implantation in children.

Methods: A multidisciplinary work group was entrusted with a review of the scientific literature on the above topic. Guidelines were drawn up, based on the articles retrieved and the group members' individual experience. They were then read over by an editorial group independent of the work group. The guidelines were graded as A, B, C or expert opinion, by decreasing level of evidence.

Results: The SFORL recommends that children with bilateral severe/profound hearing loss be offered bilateral cochlear implantation, with surgery before 12 months of age. In sequential bilateral cochlear implantation in children with severe/profound hearing loss, it is recommended to reduce the interval between the two implants, preferably to less than 18 months. The SFORL recommends encouraging children with unilateral cochlear implants to wear contralateral hearing aids when residual hearing is present, and recommends assessing perception with hearing-in-noise tests. It is recommended that the surgical technique should try to preserve the residual functional structures of the inner ear as much as possible.

© 2019 Elsevier Masson SAS. All rights reserved.

* Corresponding author. Service d'ORL pédiatrique, hôpital Necker – Enfants-Malades, 149, rue de Sèvres, 75015 Paris, France.
 E-mail address: f.simon@aphp.fr (F. Simon).

SFORL work group

Co-chairs

Pr. Emmanuel Lescanne, ORL, Tours
Pr. Natalie Loundon, ORL, Paris
Pr. Éric Truy, ORL, Lyon
Pr. Stéphane Roman, ORL, Marseille

Work group members

Pascal Barone, researcher, CNRS, Toulouse
Dr. Joël Belmin, geriatrician, Ivry-sur-Seine
Dr. Catherine Blanchet, ENT, Montpellier
Stéphanie Borel, speech therapist, Tours
Pr. Anne Charpiot, ENT, Strasbourg
Arnaud Coez, hearing care specialist, Paris
Pr. Olivier Deguine, ENT, Toulouse
Dr. Anne Farinetti, ENT, Marseille
Pr. Benoît Godey, ENT, Rennes
Dr. Diane S. Lazard, ENT, Paris
Pr. Mathieu Marx, ENT, Toulouse
Dr. Isabelle Mosnier, ENT, Paris
Dr. Yann Nguyen, ENT, Paris
Pr. Natacha Teissier, ENT, Paris
Dr. Benoît Virole, psychologist, Paris

Editorial group members

Solveig Chapuis, speech therapist, Lyon
Pr. Christophe Vincent, ENT, Lille
Pr. Thierry Mom, ENT, Clermont-Ferrand
Nadine Cochard, speech therapist, Toulouse
Dr. Ruben Hermann, ENT, Lyon
Dr. Françoise Sterkers-Artieres, ENT, Palavas-les-Flots
Caroline Rebichon, psychologist, Paris
Dr. Christine Poncet-Wallet, ENT, Paris
Dr. Arnaud Deveze, ENT, Marseille
Pr. Françoise Denoyelle, ENT, Paris
Pr. Thierry Van den Abbeele, ENT, Paris
Pr. Naima Deggouj, ENT, Bruxelles, Belgium
Aude de Lamaze, speech therapist, Paris
Pr. Cécile Parietti-Winkler, ENT, Nancy
Isabelle Prang, speech therapist, Paris
Dr. Xavier Dubernard, ENT, Reims
Martial Franzoni, speech therapist, Paris
Pr. Karine Aubry, ENT, Limoges
Dr. Pascale Henrion, ENT, Paris
Pr. Valérie Franco, ENT, Bordeaux
Pr. Philippe Bordure, ENT, Nantes
Dr. Didier Bouccara, ENT, Paris
Dr. Christine Pol, ENT, Le Kremlin-Bicêtre
Dr. Yannick Lerozey, ENT, Rouen
Pr. Michel Mondain, ENT, Montpellier

1. Introduction

A first set of guidelines for cochlear implantation (CI) was drawn up by the French Health Authority (HAS) in 2012 [1]. Briefly, it was recommended that implantation be as early as possible in prelingual hearing-impaired children, to optimize language comprehension and production. In non-progressive congenital profound/total hearing loss, implantation was not recommended in

over-5 year-olds unless oral communication was desired. Conversely, children acquiring oral communication benefit from CI at whatever age. Audiometric indications in children were mainly defined as profound hearing loss in which hearing aids do not enable language development, or severe hearing loss with $\leq 50\%$ discrimination on speech audiometry. Bilateral CI was recommended in bilateral hearing loss secondary to bacterial meningitis or temporal bone fracture to prevent bilateral cochlear ossification. CI efficacy in improving hearing, communication and language was reported to be major in the months following implantation. Surgical complications rates were low ($< 5\%$), mainly consisting in bacterial meningitis.

Updating the guidelines for CI indications in children was confined to a French national multidisciplinary work group, using the French Health Authority's formalized expert consensus methodology for good practice guidelines (<http://www.has-sante.fr/>).

A pilot group organized the consensus conference logistics, choice of editorial group members and literature analysis by PubMed search. Articles were graded A, B, C or expert opinion according to decreasing level of evidence, following the literature analysis and guideline grading guide of the French National Agency for Evaluation and Accreditation in Health (ANAES). A rationale was drawn up as a basis for a first set of guidelines, assessed by the editorial group and modified according to feed-back.

2. Results

Bilateral cochlear implantation is recommended for children with bilateral severe to profound hearing loss (grade B).

The aim of early bilateral CI in children is to restore binaural hearing, thereby improving hearing in silence and noise and achieving a certain degree of acoustic location. Children with bilateral CI were shown to have better speech perception in noise, auditory memory and verbal intelligence than with unilateral CI; they have better performance in all fields (level of evidence, 2) [2].

In congenital bilateral profound hearing loss, CI should be proposed before 12 months of age (grade B).

Early CI optimizes results for auditory perception and oral language, especially when the first CI is performed before 12 months of age. A study (level of evidence, 2) found surgical risk to be comparable between under-12 month-old infants and over-12 month-old children [3]. A prospective study (level of evidence, 2) of 160 implanted children, 99 of whom had first CI before 30 months of age and 62 between 30 months and 5 years, found better language comprehension and expression in the former group. Another study (level of evidence, 2) found that children implanted before 12 months of age had better speech and language comprehension and production development, in a series of 28 hearing-impaired children all implanted before 24 months of age [4]. Finally, 2 studies by the same team (levels of evidence, 2) found that children implanted before 12 months of age had significantly better language comprehension and production than those implanted later, with no difference in scores compared to normal-hearing children [5,6]. Regarding CI before 6 months of age, there have been too few studies for any recommendations to be made.

In sequential bilateral cochlear implantation for severe to profound hearing loss, the interval between implantations should be short (grade B), preferably less than 18 months (expert opinion).

Bilateral CI is more effective in restoring binaural hearing when simultaneous or with a short interval. A 2013 review of the literature (level of evidence, 1) demonstrated an early critical period for dedicated brain area development [7]. A study (level of evidence, 1) comparing cortical potentials between 16 children with sequential bilateral CI, 10 with simultaneous bilateral CI, 8 with unilateral CI and 7 normal-hearing control subjects found that auditory cortex curves were not lateralized as in normal-hearing subjects if bilateral stimulation was introduced later than 18 months of age [8]. Demasking in noise and acoustic location vary with hearing history, inter-CI interval and age at second CI (level of evidence, 2) [9,10].

In sequential implantation, it is recommended to keep the interval as short as possible if a contralateral hearing aid fails to provide benefit (grade B).

In a 2017 study (level of evidence, 2) of 160 children with sequential CI, the longer the inter-CI interval the greater the risk of poor perceptual outcome and non-use of the second CI in the medium-term, unless a contralateral hearing aid provided benefit [11].

In case of residual hearing, tests in noise should be included in the pre-CI work-up as soon as feasible (grade C). In case of CI with contralateral hearing aid, perception should be assessed as early as possible, with speech-in-noise and acoustic location tests, and hearing aid efficacy should be regularly reassessed (grade B). In very young children with unilateral CI and useful residual hearing, caution is recommended in indicating a second CI, as binaural function is difficult to assess (expert opinion).

Perception of speech in noise indicates the efficacy of bimodal rehabilitation (CI + hearing aid) and guides indications for a second CI. A study (level of evidence, 2) showed that, while speech-in-silence perception in children with unilateral CI was comparable with or without contralateral hearing aid, the second group showed significantly poorer perception in noise [12]. A study (level of evidence, 1) showed that good post-implantation results did not correlate with residual hearing [13]. A 2015 study (level of evidence, 2) found that 1-syllable word perception in noise was better with bilateral CI than with bimodal rehabilitation [14]. Finally, another team reported that speech perception thresholds were comparable between bilateral CI and bimodal rehabilitation, while speech discrimination in noise and acoustic location were better in bilateral CI (level of evidence, 2) [15]. Speech discrimination with hearing aid and speech-therapy language-level work-up are the key examinations identifying candidates for CI. The objective contribution of a

contralateral hearing aid, however, is more difficult to determine in children than in adults: in case of residual hearing or partial hearing loss, the audiometric threshold at which CI is preferable to a hearing aid is hard to define; several authors recommend systematic assessment of hearing in noise to this end (level of evidence, 3) [16] or else test of sentence recognition in noise (levels of evidence, 2 and 3) [17,18].

Vestibular assessment is recommended ahead of each CI (grade C).

Up to 50% of children with profound bilateral hearing loss show vestibular impairment (level of evidence, 4) [19,20]. CI surgery affects vestibular function in 0–71% of cases. A 2009 study (level of evidence, 3) of 71 CI children found a 71% rate of post-CI vestibular function modification, with 10% areflexia [21]. The risk of bilateral vestibular damage in bilateral CI needs to be taken into account, and surgery should be as atraumatic as possible, especially in children below walking age. Bilateral vestibular impairment should be avoided to prevent developmental disorder, notably cognitive. Sequential bilateral CI with a 3–6 month interval allows vestibular recovery to be checked, as impairment is often transient, ahead of second CI.

Pedopsychiatric assessment and brain MRI are recommended for children who are candidates for CI, or following CI in case of signs suggestive of autism spectrum disorder (grade B).

In a retrospective study of 785 children followed for hearing impairment, 17 were identified as having autism spectrum disorder (ASD): i.e., 2.2% in hearing-impaired children, twice the rate in the general population (level of evidence, 1) [22]. Incidence was especially high in hearing loss related to pregnancy pathology (13.8%) such as congenital cytomegalovirus infection. The children also showed higher incidence of associated pathologies such as mental (15.5%) or psychomotor retardation or visual deficit. ASD is hard to diagnose in hearing-impaired children, as specific tools are lacking; classical instruments such as the Autism Diagnostic Observation Schedule (ADOS) cannot be used in the standard manner (level of evidence, 2) [23]. In CI, results and progression are to be compared within the child rather than to children without behavioral disorder. Two studies (levels of evidence, 3) found improved perception and expression capacity (60% and 67%, respectively), better environmental awareness, and no increase in autistic hyperactivity following CI in ASD children [24,25]. Brain MRI is part of the work-up in suspected ASD (level of evidence, 3) [25].

It is recommended to assess the resources, difficulties and needs of the family environment of hearing-impaired children and to encourage the child to be involved in the management of hearing loss and rehabilitation and to accompany him or her in this (grade B).

Poor family investment is a factor for poor perceptual and linguistic outcome in CI (level of evidence, 2) [26,27]. Accompaniment should consist in dialogue, support, guidance and supervision from diagnosis through rehabilitation (level of evidence, 4) [28].

Parents of children with pathologies associated with hearing loss (multi-disability) should be informed that CI results may be limited; they should consider their expectations in dialogue with a multidisciplinary team (grade B).

Hearing loss is sometimes part of a complex clinical presentation including other psychological, psychiatric and physical aspects. A study (level of evidence, 2) found poorer speech intelligibility ratings (SIR) 5 years post-CI in case of associated abnormality compared to isolated hearing loss (70% vs. 96%); SIR was inversely proportional to the number of associated abnormalities [29]. Low IQ does not in itself contraindicate CI, but should be included in the multidisciplinary assessment. Audiophonologic rehabilitation does not significantly improve IQ, but improved hearing can facilitate schooling and learning (level of evidence, 2) [30].

It is recommended that CI surgery should so far as possible spare external, middle and inner ear structures that remain functional (with or without preoperatively preserved low-frequency air-conduction thresholds) (grade C).

Electroacoustic stimulation is now applicable in certain pediatric CI candidates with low-frequency residual hearing, using cautious surgery and last-generation implants. It is important to avoid intraoperative perilymph aspiration and blood or debris penetration in the cochlea (level of evidence, 4) [31]. Patients with postoperative residual hearing show better speech performance on pure electrical stimulation (level of evidence, 2) [32].

It is recommended that the parents be informed of the risk of immediate postoperative and long-term loss of residual hearing following CI (grade C).

A lateral electrode through the cochlear window optimally preserves residual hearing (level of evidence, 3) [33,34], although no reported surgical techniques guarantee complete short- or long-term preservation.

Rehabilitation should combine acoustic and electrical stimulation in case of low-frequency residual hearing (grade B).

Combined electrical and acoustic stimulation enables better speech discrimination in noise and tune recognition than electrical stimulation alone (level of evidence, 2) [35,36].

Perioperative corticosteroids are recommended if residual hearing is intended to be preserved (expert opinion).

Low-level studies (level of evidence, 4) showed moderate benefit of perioperative corticotherapy [37]. The literature does not support any particular mode of administration.

Children with unilateral CI should be encouraged to use a contralateral hearing aid in case of residual hearing (grade B).

CI + hearing aid compared to unilateral CI alone improves recognition of 2-syllable words, sentences in silence and noise and everyday communication (level of evidence, 3) [38] and acoustic location (Level of evidence, 1) [39].

It is recommended that studies should assess the benefit of CI in children with acquired or congenital severe to profound unilateral sensorineural hearing loss (expert opinion).

In France, CI has no formal or consensual indications in unilateral childhood hearing loss. The first small-scale studies reported a certain efficacy for speech perception in noise, but results were variable and sometimes disappointing (level of evidence, 4) [40,41]. The medium-term risk in patients with generally effective hearing is that the CI will be abandoned (by up to 3 in 5 patients) (level of evidence, 4) [42].

It is recommended to assess pre- and post-CI quality of life alongside audiometry (grade A).

Quality of life assessment gives the patient and family an active role in the care pathway, and is especially relevant to patients with chronic disease and/or disability without hope of cure. Taking real-life needs and feelings into account via quality of life questionnaires places the patient at the center of care. Correlations between perceptual and quality of life results are limited, which only reinforces the interest of exploring this subjective dimension in complement to audiometric assessment (level of evidence, 1) [43].

Several versions of a given quality of life instrument should be available, adapted to age (expert opinion).

Generic questionnaires are based on those used in the general population, enabling comparison between populations or disease groups. They have the advantage of having validation and normal values for interpretation, and are more meaningful for public health decision-makers; however, they fail to give detailed account of the specificities of each pathology. Specific questionnaires tend to be

Table 1
Generic quality of life instruments used in pediatric cochlear implantation.

Instruments	Original language	Validation in French	Target age (years)	Respondents	Versions	Domains explored	Studies
EuroQoL-5D (EQ-5D)	English	Yes	Not stated	C and P	C: EQ-5D-Youth P: EQ-5D-5L/3L	QoL	The EuroQOL Group, 1990
Health Utility Index 3 (HUI 3)	English	Yes	> 5	P	P	Utility scores QALY	Sach, 2007 Cheng, 2000 Horsman, 2003 Lovett, 2010 Spareboom, 2012
Child Health and Illness Profile (CHIP)	English	Yes (version AC and PRF)	6–17	C and P	C: CRF (6–11) and AE (11–17) P: PRF (6–11)	QoL	Starfield, 1993 Clark, 2010
Child Health Questionnaire (CHQ)	English	Yes	5 mo–18	C and P	C: PF-28 (5–18) P: PF-50 (5 mo–15 yrs)	QoL	Riley, 2004 Landgraf, 1998 Ravens-Sieberer, 2001 Wake, 2004
Pediatric Quality of Life 4 (PedsQL 4.0)	English	Yes	2–18	C and P	C: (5–18) P: (2–18)	QoL	Borton, 2010 Lovett, 2010 Sparreboom, 2012 [50] Varni, 2011, 2007 Beijen, 2007 Rajendran, 2010
Kinder Lebensqualitäts-fragebogen (KINDL)	German	Yes	4–16	C and P	Kiddy (4–7) Kid (8–11) Kiddo (12–16)	QoL	Huber, 2005 Warner-Czyz, 2009 Bullinger, 2008 [44] Warner-Czyz, 2011
Youth Quality of Life Instrument (YQOL)	English	No	11–18	C	C	QoL	Loy, 2010 Edwards, 2002 Patrick, 2002 Kushalnagar, 2011
KIDSCREEN-27	English	Yes	8–18	C and P	27 items	QoL	Schick, 2013 Ravens-Sieberer, 2005 Robitail, 2007 Razafimahefa-Raelina, 2016
Glasgow Children's Benefit Inventory (GCBi)	English	No	1–15	C and P	C P	QoL after H&N surgery	Kubba, 2004 Lovett, 2010

C: child; P: parent; QALY: quality-adjusted life years; QoL: quality of life.

Table 2
Specific quality of life instruments used in pediatric cochlear implantation.

Instruments	Original language	Validation in French	Target age (years)	Respondents	Versions	Domains explored	Studies
Paediatric Audiology Quality of Life Questionnaire (PAQL)	English	No	3–18	P	P	QoL with associated comorbidity	Edwards, 2012 [45]
Parent's Views and Experiences with pediatric Cochlear Implant Questionnaire (PVECIQ)	English	No	5–16	P	P	Parental perception of CI-associated benefit/limitations and child's QoL	Archbold, 2002 O'Neill, 2004 Nunes, 2005 [48] Incesulu, 2003 Nicholas, 2003 [47]
Kinder Lebensqualitäts-fragebogen – Cochlear Implant module (KINDL-CI module)	German	Yes	4–16	C and P	Version 4–7 Version 8–16	QoL	Bullinger, 2008 [44] Warner-Czyz, 2011
Youth Quality of Life – Deaf and Hard of Hearing (YQOL-DHH)	English	No	11–18	C and P	C: YQOL-DHH P: YQOL-DHH-PROBE	QoL	Kushalnagar, 2011 Schick, 2013 Patrick, 2011 [49] Hintermair, 2011 [46]

C: child; P: parent; QoL: quality of life.

more sensitive to clinical variations, notably related to treatment, and thus to functional change. Generic and specific questionnaires used in the literature for assessment of pediatric CI are presented in Tables 1 and 2 [43–50].

3. Conclusions

Management of hearing-impaired children should be multidisciplinary, and parents should be accompanied to optimize outcome. It should be adapted to age, severity and comorbidity. In case of CI, implementation should be early (before 12 months of age) to optimize outcome, with a short interval (less than 18 months) between sequential bilateral implantations. Hearing-in-noise tests are important to assess residual hearing or efficacy of bimodal rehabilitation. Surgery should be sparing, to preserve residual cochlear and vestibular function.

Disclosure of interest

The authors declare that they have no competing interest.

Acknowledgments

Solveig Chapuis, speech therapist, Lyon
Pr. Christophe Vincent, ENT, Lille
Pr. Thierry Mom, ENT, Clermont-Ferrand
Nadine Cochard, speech therapist, Toulouse
Dr. Ruben Hermann, ENT, Lyon
Dr. Françoise Sterkers-Artieres, ENT, Palavas-les-Flots
Caroline Rebichon, psychologist, Paris
Dr. Christine Poncet-Wallet, ENT, Paris
Dr. Arnaud Deveze, ENT, Marseille
Pr. Françoise Denoyelle, ENT, Paris
Pr. Thierry Van den Abbeele, ENT, Paris
Pr. Naima Deggouj, ENT, Brussels, Belgium
Aude de Lamaze, speech therapist, Paris
Pr. Cécile Parietti-Winkler, ENT, Nancy
Isabelle Prang, speech therapist, Paris
Dr. Xavier Dubernard, ENT, Reims
Martial Franzoni, speech therapist, Paris
Pr. Karine Aubry, ENT, Limoges
Dr. Pascale Henrion, ENT, Paris
Pr. Valérie Franco, ENT, Bordeaux
Pr. Philippe Bordure, ENT, Nantes
Dr. Didier Bouccara, ENT, Paris
Dr. Christine Pol, ENT, Le Kremlin-Bicêtre
Dr. Yannick Lerosey, ENT, Rouen
Pr. Michel Mondain, ENT, Montpellier

References

- [1] Le traitement de la surdité par implants cochléaires ou du tronc cérébral. HAS; 2012 [https://www.has-sante.fr/portail/upload/docs/application/pdf/fiche_bon_usage_implants_cochleaires.pdf].
- [2] Jacobs E, Langereis MC, Frijns JH, Free RH, Goedegeure A, Smits C, et al. Benefits of simultaneous bilateral cochlear implantation on verbal reasoning skills in prelingually deaf children. *Res Dev Disabil* 2016;58:104–13.
- [3] Kim Y, Patel VA, Isildak H, Carr MM. An analysis of safety and adverse events following cochlear implantation in children under 12 months of age. *Otol Neurotol* 2017;38(10):1426–32.
- [4] May-Mederake B. Early intervention and assessment of speech and language development in young children with cochlear implants. *Int J Pediatr Otorhinolaryngol* 2012;76(7):939–46.
- [5] Colletti L, Mandala M, Colletti V. Cochlear implants in children younger than 6 months. *Otolaryngol Head Neck Surg* 2012;147(1):139–46.
- [6] Colletti L, Mandala M, Zocante L, Shannon RV, Colletti V. Infants versus older children fitted with cochlear implants: performance over 10 years. *Int J Pediatr Otorhinolaryngol* 2011;75(4):504–9.
- [7] Kral A. Auditory critical periods: a review from system's perspective. *Neuroscience* 2013;247:117–33.
- [8] Gordon KA, Wong DD, Papsin BC. Bilateral input protects the cortex from unilaterally-driven reorganization in children who are deaf. *Brain* 2013;136(Pt 5):1609–25.
- [9] Grieco-Calub TM, Litovsky RY. Sound localization skills in children who use bilateral cochlear implants and in children with normal acoustic hearing. *Ear Hear* 2010;31(5):645–56.
- [10] Litovsky RY, Misurelli SM. Does bilateral experience lead to improved spatial unmasking of speech in children who use bilateral cochlear implants? *Otol Neurotol* 2016;37(2):e35–42.
- [11] Myhrum M, Strom-Roum H, Heldahl MG, Rodvik AK, Eksveen B, Landsvik B, et al. Sequential bilateral cochlear implantation in children: outcome of the second implant and long-term use. *Ear Hear* 2017;38(3):301–13.
- [12] Choi JE, Moon JJ, Kim EY, Park HS, Kim BK, Chung WH, et al. Sound localization and speech perception in noise of pediatric cochlear implant recipients: bimodal fitting versus bilateral cochlear implants. *Ear Hear* 2017;38(4):426–40.
- [13] Chioffi JSC, Hyppolito MA. Effects of residual hearing on cochlear implant outcomes in children: a systematic review. *Int J Pediatr Otorhinolaryngol* 2017;100:119–27.
- [14] Jeong SW, Kang MY, Kim LS. Criteria for selecting an optimal device for the contralateral ear of children with a unilateral cochlear implant. *Audiol Neurotol* 2015;20(5):314–21.
- [15] Litovsky RY, Johnstone PM, Godar SP. Benefits of bilateral cochlear implants and/or hearing aids in children. *Int J Audiol* 2006;45(Suppl. 1):S78–91.
- [16] Skarzynski H, Lorens A. Electric acoustic stimulation in children. *Adv Otorhinolaryngol* 2010;67:135–43.
- [17] Gifford RH, Dorman MF, Skarzynski H, Lorens A, Polak M, Driscoll CL, et al. Cochlear implantation with hearing preservation yields significant benefit for speech recognition in complex listening environments. *Ear Hear* 2013;34(4):413–25.
- [18] Gratacap M, Thierry B, Rouillon I, Marlin S, Garabedian N, Loundon N. Pediatric cochlear implantation in residual hearing candidates. *Ann Otol Rhinol Laryngol* 2015;124(6):443–51.
- [19] Thierry B, Blanchard M, Leblouanger N, Parodi M, Wiener-Vacher SR, Garabedian EN, et al. Cochlear implantation and vestibular function in children. *Int J Pediatr Otorhinolaryngol* 2015;79(2):101–4.
- [20] Verbecque E, Marijnissen T, De Belder N, Van Rompaey V, Boudewyns A, Van de Heyning P, et al. Vestibular (dys)function in children with sensorineural hearing loss: a systematic review. *Int J Audiol* 2017;56(6):361–81.
- [21] Jacot E, Van Den Abbeele T, Debre HR, Wiener-Vacher SR. Vestibular impairments pre- and post-cochlear implant in children. *Int J Pediatr Otorhinolaryngol* 2009;73(2):209–17.
- [22] Fitzpatrick EM, Lambert L, Whittingham J, Leblanc E. Examination of characteristics and management of children with hearing loss and autism spectrum disorders. *Int J Audiol* 2014;53(9):577–86.
- [23] Szymanski CA, Brice PJ, Lam KH, Hottot SA. Deaf children with autism spectrum disorders. *J Autism Dev Disord* 2012;42(10):2027–37.
- [24] Eshraghi AA, Nazarian R, Telischi FF, Martinez D, Hodges A, Velandia S, et al. Cochlear implantation in children with autism spectrum disorder. *Otol Neurotol* 2015;36(8):e121–8.
- [25] Lachowska M, Pastuszka A, Lukaszewicz-Moszynska Z, Mikolajewska L, Niemczyk K. Cochlear implantation in autistic children with profound sensorineural hearing loss. *Braz J Otorhinolaryngol* 2018;84(1):15–9 [pii: S1808-8694(16)30231-2].
- [26] Black J, Hickson L, Black B, Khan A. Paediatric cochlear implantation: adverse prognostic factors and trends from a review of 174 cases. *Cochlear Implants Int* 2014;15(2):62–77.
- [27] Percy-Smith L, Busch GW, Sandahl M, Nissen L, Jøssvassen JL, Bille M, et al. Significant regional differences in Denmark in outcome after cochlear implants in children. *Dan Med J* 2012;59(5):A4435.
- [28] Cicieliello E, Bolzonello P, Marchi R, Falzone C, Muzzi E, Orzan E. Empowering the family during the first months after identification of permanent hearing impairment in children. *Acta Otorhinolaryngol Ital* 2016;36(1):64–70.
- [29] Nikolopoulos TP, Archbold SM, Wever CC, Lloyd H. Speech production in deaf implanted children with additional disabilities and comparison with age-equivalent implanted children without such disorders. *Int J Pediatr Otorhinolaryngol* 2008;72(12):1823–8.
- [30] Meinzen-Derr J, Wiley S, Phillips J, Altaye M, Choo DI. The utility of early developmental assessments on understanding later nonverbal IQ in children who are deaf or hard of hearing. *Int J Pediatr Otorhinolaryngol* 2017;92:136–42.
- [31] Khater A, El-Anwar MW. Methods of hearing preservation during cochlear implantation. *Int Arch Otorhinolaryngol* 2017;21(3):297–301.
- [32] Carlson ML, Driscoll CL, Gifford RH, Service GJ, Tombers NM, Hughes-Borst BJ, et al. Implications of minimizing trauma during conventional cochlear implantation. *Otol Neurotol* 2011;32(6):962–8.
- [33] Frayssy B, Macias AR, Sterkers O, Burdo S, Ramsden R, Deguine O, et al. Residual hearing conservation and electroacoustic stimulation with the nucleus 24 contour advance cochlear implant. *Otol Neurotol* 2006;27(5):624–33.
- [34] Wanna GB, O'Connell BP, Francis DO, Gifford RH, Hunter JB, Holder JT, et al. Predictive factors for short- and long-term hearing preservation in cochlear implantation with conventional-length electrodes. *Laryngoscope* 2018;128(2):482–9.
- [35] Buchner A, Illg A, Majdani O, Lenarz T. Investigation of the effect of cochlear implant electrode length on speech comprehension in quiet and noise compared with the results with users of electroacoustic-stimulation, a retrospective analysis. *PLoS One* 2017;12(5):e0174900.

- [36] Gifford RH, Davis TJ, Sunderhaus LW, Menapace C, Buck B, Crosson J, et al. Combined electric and acoustic stimulation with hearing preservation: effect of cochlear implant low-frequency cutoff on speech understanding and perceived listening difficulty. *Ear Hear* 2017;38(5):539–53.
- [37] Sweeney AD, Carlson ML, Zuniga MG, Bennett ML, Wanna GB, Haynes DS, et al. Impact of perioperative oral steroid use on low-frequency hearing preservation after cochlear implantation. *Otol Neurotol* 2015;36(9):1480–5.
- [38] Ching TY, Psarros C, Hill M, Dillon H, Incerti P. Should children who use cochlear implants wear hearing aids in the opposite ear? *Ear Hear* 2001;22(5):365–80.
- [39] Ching TY, Incerti P, Hill M, van Wanrooy E. An overview of binaural advantages for children and adults who use binaural/bimodal hearing devices. *Audiol Neurotol* 2006;11(Suppl. 1):6–11.
- [40] Greaver L, Eskridge H, Teagle HFB. Considerations for pediatric cochlear implant recipients with unilateral or asymmetric hearing loss: assessment, device fitting, and habilitation. *Am J Audiol* 2017;26(2):91–8.
- [41] Polonenko MJ, Papsin BC, Gordon KA. Children with single-sided deafness use their cochlear implant. *Ear Hear* 2017;38(6):681–9.
- [42] Thomas JP, Neumann K, Dazert S, Voelter C. Cochlear implantation in children with congenital single-sided deafness. *Otol Neurotol* 2017;38(4):496–503.
- [43] McRackan TR, Bauschard M, Hatch JL, Franko-Tobin E, Droghini HR, Nguyen SA, et al. Meta-analysis of quality of life improvement after cochlear implantation and associations with speech recognition abilities. *Laryngoscope* 2018;128(4):982–90.
- [44] Bullinger M, Brutt AL, Erhart M, Ravens-Sieberger U. Psychometric properties of the KINDL-R questionnaire: results of the BELLA study. *Eur Child Adolesc Psychiatry* 2008;17(Suppl. 1):125–32.
- [45] Edwards L, Hill T, Mahon M. Quality of life in children and adolescents with cochlear implants and additional needs. *Int J Pediatr Otorhinolaryngol* 2012;76(6):851–7.
- [46] Hintermair M. Health-related quality of life and classroom participation of deaf and hard of hearing students in general schools. *J Deaf Stud Deaf Educ* 2011;16(2):254–71.
- [47] Nicholas JG, Geers AE. Personal, social, and family adjustment in school-aged children with a cochlear implant. *Ear Hear* 2003;24(1 Suppl.):69s–81s.
- [48] Nunes T, Pretzlik U, Ilıcak S. Validation of a parent outcome questionnaire from pediatric cochlear implantation. *J Deaf Stud Deaf Educ* 2005;10(4):330–56.
- [49] Patrick DL, Edwards TC, Skalicky AM, Schick B, Topolski TD, Kushalnagar P, et al. Validation of a quality of life measure for deaf or hard of hearing youth. *Otolaryngol Head Neck Surg* 2011;145(1):137–45.
- [50] Sparreboom M, Snik AF, Mylanus EA. Sequential bilateral cochlear implantation in children: quality of life. *Arch Otolaryngol Head Neck Surg* 2012;138(2):134–41.