



Hepatoprotective and antioxidant activity of watercress extract on acetaminophen-induced hepatotoxicity in rats



Nahid Azarmehr^a, Parisa Afshar^a, Mona Moradi^b, Heibatollah Sadeghi^c, Hossein Sadeghi^c, Behnam Alipour^c, Bahman Khalvati^c, Zahra Barmoudeh^a, Kazem Abbaszadeh-Goudarzi^d, Amir Hossein Doustimotlagh^{c,e,*}

^a Student Research Committee, Yasuj University of Medical Sciences, Yasuj, Iran

^b Department of Clinical Biochemistry and Genetics, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Mazandaran, Iran

^c Medicinal Plants Research Center, Yasuj University of Medical Sciences, Yasuj, Iran

^d Cellular and Molecular Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran

^e Department of Clinical Biochemistry, Faculty of Medicine, Yasuj University of Medical Sciences, Yasuj, Iran

ARTICLE INFO

Keywords:

Cell biology
Pharmaceutical science
Biochemistry
Molecular biology
Oxidative stress
Antioxidant
Acetaminophen
Watercress
Hepatotoxicity

ABSTRACT

Introduction: Acetaminophen (APAP) as an analgesic and antipyretic drug can result to liver damages while using more than 4 g/day. Therefore, APAP toxicity causes the liver to dysfunction. This study aims to investigate the hepatoprotective and antioxidant activity of hydroalcoholic extract of watercress (WC) in APAP-induced hepatotoxicity in rats.

Materials and methods: Randomly, twenty-four Wistar rats were divided into four groups of six each. Groups named as control, APAP, APAP + WC and APAP + S for group 1, 2, 3, and 4, respectively. Group 1 received distilled water 1 ml/kg for 7 days. In group 2, 3, and 4, rats pretreated by receiving distilled water (1 ml/kg), WC extract (500 mg/kg), silymarin extract (mg/kg) for 7 days, respectively. Of note, to induce acute hepatotoxicity in groups 2, 3, and 4, rats posttreated by orally intoxicated with single dose of APAP (2 g/kg) on the sixth day. The animals were sacrificed on the seventh day. Alanine amino transferase (ALT), aspartate amino transferase (AST), ferric reducing ability of plasma (FRAP), protein carbonyl (PCO), total thiol (T-SH), glutathione peroxidase (GPx), superoxide dismutase (SOD) and catalase (CAT) activities were measured in plasma. It should be noted that the chemical composition of WC extract was identified by GC-MS analysis.

Results: The results have shown that there was a significant increase in AST, ALT, FRAP and PCO content in APAP group in comparison to control. Also, there was a significant reduction in T-SH levels and GPx activity in APAP group compared to control. However, administration of WC extract and silymarin not only causes a significant decrease in AST activity, but they markedly increased T-SH content and GPx activity compared to APAP group. GC-MS analysis showed the major compositions were found to be benzenepropanenitrile (48.30 %), Phytol (10.10 %), α -cadinene (9.50%) and linolenic acid (8.0).

Conclusions: It is concluded that the WC extract reduces APAP-induced toxicity through its hepatoprotective and antioxidant activity in rats.

1. Introduction

Liver has been known as a central place to detoxify and metabolize different drugs and xenobiotics [1]. The excessive or inaccurate use of drugs leads to liver dysfunction [2]. Acetaminophen (N-acetyl-p-aminophenol, APAP) as an analgesic and antipyretic drug can result to liver damages while using more than 4 g/day [3, 4]. In liver, more than 90% of

APAP is metabolized to non-toxic sulfate and glucuronide, and the rest, converted to a toxic metabolite called N-acetyl-p-benzoquinone imine (NAPQI) in which the reaction catalyzed by cytochrome P450 under normal conditions [5]. NAPQI, a hepatotoxic reactive molecule, is detoxified to various non-reactive molecules through reaction with glutathione peroxidase. At time of excessive production of NAPQI while using more acetaminophen, it causes a reduction in glutathione

* Corresponding author.

E-mail address: amirhossein.dousti@yums.ac.ir (A.H. Doustimotlagh).

<https://doi.org/10.1016/j.heliyon.2019.e02072>

Received 13 March 2019; Received in revised form 5 May 2019; Accepted 8 July 2019

2405-8440/© 2019 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

concentration which leads to hepatotoxicity [2, 5]. However, in time of inadequate glutathione, the production of reactive oxygen species (ROSs) like superoxide, hydrogen peroxide and hydroxyl radicals has been increased by NAPQI. Therefore, oxidative stress was induced by imbalance in the formation and removal of free radicals [6]. Of note, normal hepatocytes against free radicals and other reactants were protected by non-enzymatic and enzymatic defense systems, including glutathione peroxidase (GPx), superoxide dismutase (SOD) and catalase (CAT) [7]. Although mild oxidative stress in cells is easily handled, in a condition with severe oxidative stress, oxidant compounds react with cell components such as lipid, protein, DNA and cell membrane, and lead to pathological complications [8].

Over the past few years, many researchers have focused the use of antioxidant extracts from herbal products as hepatoprotective substances in order to reduce the effects of APAP-induced toxicity [4, 9, 10]. Watercress (WC, or *Nasturtium officinale*), a green leafy vegetable from the family of Brassicaceae, has been used in herbal medicine as remedy for several complications [9]. Study results showed that WC used in the treatment of diabetes, bronchitis, diuresis, influenza, asthma [11]. It was reported that WC had anticancer activity by reducing oxidative stress [12]. Isothiocyanates and carotenoids, the main components of WC, are responsible to regulate antioxidant by decreasing free radicals and therefore prevent the injury [10, 12, 13]. In a current study, we are going to clarify the hepatoprotective effects of WC extract in intoxicated rats with APAP.

2. Materials and methods

2.1. Drugs and chemicals

Some materials were purchased from Sigma Chemical Co (St Louis, MO, USA), including acetaminophen, (5, 5'-dithiols-(2-nitrobenzoic acid)) (DTNB) and ethylenediaminetetraacetic acid (EDTA). And also remaining components tested were standard laboratory materials, provided primarily from Merck (Germany), including trichloroacetic acid (TCA), 2, 4-dinitrophenylhydrazine (DNPH), guanidine hydrochloride and 2,4,6-tris (2-pyridyl) -s-triazine (TPTZ). It should be noted that all chemicals and reagents used in the experiment, had analytical grade.

2.2. Preparation of WC extract

In December 2017, stems and leaves of WC were collected from the Sisakht region located in Yasuj, Iran. The plant was recognized by the botanist, where a voucher sample (Herbarium No. HYU30230) was deposited. Plant was cleaned, dust and soil removed by using deionized water. And then mentioned parts were cut off using scissors. To prepare WC extract, 70% ethanol was used at room temperature ($25 \pm 2^\circ\text{C}$) for 48 h. Subsequently, hydroalcoholic extract was filtered, and plant deposit was totally vaporized under reduced pressure by using a rotary at 60°C . Finally, to do the experiment, the concentrated extract was dried and stored at -20°C .

2.3. Animals

Male Wistar rats (8 weeks old, weight 200–250 g) were obtained from the animal house of Yasuj University of Medical Sciences. The Ethics Committee approved this study (Code number: IR.YUMS.REC.1397.114). The rats were kept under constant temperature and humidity ($23 \pm 2^\circ\text{C}$, $50 \pm 5\%$) on a controlled light–dark cycle with free access to food and water. The animals were kept in accordance with the 'Guide for the Care and Use of Laboratory Animals' (NIH US publication no. 85–23 revised 1985).

2.4. Study design

Randomly, twenty-four Wistar rats were divided into four groups of

six each. Group 1 received distilled water 1 ml/kg for 7 days as a control group. In group 2, 3, and 4, rats pretreated by receiving distilled water (1 ml/kg), WC extract (500 mg/kg) [14] silymarin (100 mg/kg) [15] for 7 days, respectively. Of note, to induce acute hepatotoxicity, in group 2, 3, and 4, rats posttreated by orally intoxicated with single dose of APAP (2 g/kg) on the sixth day [16].

Blood samples were taken from the animals by cardiac puncture, collected in heparin-containing tubes, centrifuged at $3000 \times g$ for 10 minutes. For biochemical analysis, the plasma was separated and stored at -20°C . In order to evaluate antioxidant enzyme markers in liver, the rats were sacrificed on day 7. Liver tissue sectioned to small pieces and homogenized (10% w/v) by IKA Werke Ultra-Turrax T25 basic homogenizer in PBS (pH 7.4 and 50 mM), centrifuged at $10000 \times g$ for 10 min at 4°C , and finally stored in -20°C .

2.5. Biochemical analysis

To evaluate biochemical markers in liver injury, alanine aminotransferase (ALT) (and aspartate aminotransferase (AST) were measured in plasma samples using enzymatic activity assay kits (Pars Azmoon Co., Tehran, Iran).

2.6. Determination of oxidative stress markers

2.6.1. Determination of ferric reducing ability of plasma (FRAP)

In this experiment, the scale of antioxidant plasma capacity in the reduction of ferric (Fe^{+3}) ions was measured according to Benzie and Strain [17]. The blue-colored Fe-TPTZ (Tripyridyl-s-triazine) complex is formed by reducing Fe^{+3} into ferrous (Fe^{+2}) ions in acidic pH. Color intensity was measured by spectrophotometric at 593 nm. In comparison to standard solution of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0–1500 $\mu\text{mol/L}$), the plasma level of FRAP was reported.

2.6.2. Determination of protein carbonyl (PCO)

The total protein levels of plasma and liver tissue homogenate were measured by the biuret reaction [18]. The method proposed by Kholari *et al.*, was applied to measure an index of protein oxidation [19]. To treat liver tissue homogenates and plasma samples, DNPH (10 mmol/L) in HCl (2 mol/L) was used in darkness for 1 hour at room temperature. The precipitate was treated with 50% TCA, rinsed with a solution of ethanol and ethyl acetate (1:1 v/v) and then dissolved in guanidine hydrochloride (6 mol/L) for 15 minutes at 37°C . A molar absorption coefficient of $2.2 \times 10^4 \text{ M}^{-1}\text{cm}^{-1}$ was used to measure PCO content in $\mu\text{mol/g}$ and $\mu\text{mol/mg}$ level in tissue and plasma, respectively.

2.6.3. Determination of total thiol (T-SH)

According to previously described method, the level of T-SH group was evaluated with slight modifications [20]. To do the experiment, the sample (25 μL) was added to a microtube containing Tris-EDTA buffer (150 μL), absolute methanol (790 μL) and 10 mM DTNB (10 μL). It was gently mixed by pipette and then incubated for 15 minutes at the room temperature. Finally, the optical density was read at 412 nm versus DTNB blank and blank tube. A molar absorption coefficient of $13,600 \text{ M}^{-1}\text{cm}^{-1}$ was considered to express the T-SH.

2.7. Determination of antioxidant enzymes activity

According to the instructions, GPx, SOD and CAT activity were measured in the liver tissue homogenate using commercial colorimetric assay kits (Zell Bio GmbH, Ulm, Germany).

2.8. Chromatographic conditions

GC-MS analysis was carried out on a Hewlett-Packard 5973 connected with a mass detector HP6890 using a DB-1 column (55 m $\times$ 0.25 mm, film thickness 0.25 μm). The oven temperature was adjusted from

40 °C (1 min) to 250 °C (30 min) at 3 °C min⁻¹. The transferor gas was helium with the flow rate of 1.0 ml min⁻¹. The mass spectrometer (Hewlett-Packard 5973, USA) was activating in EI mode at 70 eV.

2.9. Statistical analysis

Statistical analysis was performed using SPSS 17 (SPSS Inc., Chicago, IL, USA). Data was presented as mean $\pm$ standard error of the mean (SEM). For post hoc analysis, multiple comparisons were analyzed using one-way ANOVA and Tukey test. A P value less than 0.05 was considered as statistically significant.

3. Results

3.1. Biochemical parameters

To evaluate Liver injury, the level of liver enzymes was determined in plasma. As shown in Table 1, the plasma levels of AST and ALT were significantly higher in the APAP groups compared to control rats ($p < 0.05$). Administration of WC extract and silymarin at dose of 500 mg/kg and 100 mg/kg, respectively, caused a significant decrease in AST level compared to APAP group ($P \leq 0.05$). However, they had no effect on the ALT activity compared to APAP group.

3.2. Oxidative stress markers

The results showed not only markedly increased in FRAP levels, but slightly increased in protein carbonyl levels in APAP group compared to control. However, WC extract had no effect on the FRAP content compared to APAP group (Table 2). Plasma PCO was decreased by 31.6% in WC + APAP in comparison to APAP group. Our findings showed a significant reduction in T-SH content in APAP group compared to control. Of note, a significant increase in T-SH level in WC + APAP and S + APAP groups was seen, compared to APAP group ($P \leq 0.05$).

3.3. Antioxidant enzymes activity

Antioxidant enzymes activity was measured in groups treated compared to control. The GPx activity had a significant decrease in the APAP group compared to control ($P < 0.05$). However, its activity indicated a significant increase in the APAP group receiving WC extract (500 mg/kg) (WC + APAP) compared to the APAP group ($P < 0.05$) (Fig. 1 A). More importantly, CAT and SOD activities were not significantly changed in the rats treated with WC + APAP and S + APAP groups compared to APAP group (Fig. 1 B-C).

3.4. Separation and identification of the hydroalcoholic extract of WC.

The WC extract was analyzed by HPLC in the previous study [21]. The chemical composition obtained from WC extract, presented in Table 3. However, eleven components have been identified with their percentage composition, represented 99.0 % of total WC. Of which the major ones,

Table 1

Effect of WC extract on plasma biochemical parameters in APAP intoxicated rats.

Groups	AST (U/L)	ALT (U/L)
Control	110.83 $\pm$ 12.76	29.85 $\pm$ 3.78
APAP	226.62 $\pm$ 13.92 ^a	74.31 $\pm$ 7.90 ^a
WC + APAP	156.04 $\pm$ 21.18 ^b	65.33 $\pm$ 11.84 ^a
S + APAP	128.13 $\pm$ 18.23 ^b	63.35 $\pm$ 8.78 ^a

ALT: alanine aminotransferase; AST: aspartate aminotransferase. Each value represents the mean $\pm$ SEM. APAP: acetaminophen; WC: 500 mg/kg of hydroalcoholic extract of watercress; S: 100 mg/kg of silymarin.

^astatistically significant compared to control group, P-value ≤ 0.05 .

^bstatistically significant compared to APAP group, P-value ≤ 0.05 .

Table 2

Effects of WC extract on oxidative stress markers in APAP intoxicated rats.

Groups	FRAP (μ mol/L)	PCO (μ mol/mg protein)	T-SH (μ mol/mg protein)	PCO (μ mol/g tissue)
Control	1039.41 $\pm$ 57.04	1.39 $\pm$ 0.07	3.29 $\pm$ 0.10	3.04 $\pm$ 0.08
APAP	2284.83 $\pm$ 123.46 ^a	1.74 $\pm$ 0.22	2.27 $\pm$ 0.27 ^a	3.15 $\pm$ 0.04
WC + APAP	2418.16 $\pm$ 184.23 ^a	1.19 $\pm$ 0.10	3.10 $\pm$ 0.17 ^b	2.98 $\pm$ 0.02
S + APAP	1639.00 $\pm$ 142.00 ^{a,b}	1.37 $\pm$ 0.17	3.27 $\pm$ 0.19 ^b	3.04 $\pm$ 0.09

FRAP: ferric reducing antioxidant power; PCO: protein carbonyl; T-SH: total thiol. PCO content in μ mol/g and μ mol/mg level in tissue and plasma, respectively. Each value represents the mean $\pm$ SEM. APAP: acetaminophen; WC: 500 mg/kg WC extract; S: 100 mg/kg silymarin.

^astatistically significant compared to control group, P-value ≤ 0.05 .

^bstatistically significant compared to APAP group, P-value ≤ 0.05 .

were found to be benzenepropanenitrile (48.30 %), Phytol (10.10 %), α -cadinene (9.50%) and linolenic acid (8.0).

4. Discussion

Oxidative stress is the most prevalent cause of drug-induced hepatotoxicity [5, 22]. The present study investigated the hepatoprotective effect of WC extract as an antioxidant on liver injury induced by APAP. In various studies, APAP-induced hepatotoxicity is a reliable model to evaluate hepatoprotective agents, administered orally or intraperitoneally [5, 23, 24].

In current study, the aminotransferase enzymes were investigated to evaluate the biochemical properties of the liver function after exposure to antioxidant component. ALT and AST as hepatic leakage enzymes after hepatocellular injury, released into peripheral blood [25]. Results showed significant elevation in the levels of ALT and AST in the APAP group compared to control. This increase may be due to APAP-induced liver damage which leads to a reduction in cell integrity and leakage as a consequence of this treatment, causes enzymes release into the bloodstream [15]. Karami *et al.* reported hepatoprotective effect of WC against gamma-radiation-induced hepatotoxicity in mice [26].

In previous studies, hepatoprotective activity of Clerodendron inerme, Adansonia digitata and Piper retrofractum extract was confirmed against APAP-induced hepatic injury in rats [15, 27, 28]. For the first time, the protective effects of WC extract were evaluated in rats with APAP-induced hepatotoxicity. Of note, the level of AST enzyme was significantly reduced in WC + APAP group in which WC extract orally administered, and also this group represented hepatocytes regeneration and repair. However, it had no significant effect on ALT level.

Excessive generation of NAPQI in APAP-induced hepatotoxicity, can covalently binds to lipids, proteins and DNA molecules. Additionally, NAPQI led to a lower glutathione concentration in liver and caused depletion of this antioxidant compound [24]. Proteins are critical targets of oxidative stress products, and more vulnerable due to enzymatic activities in the cell. Among oxidative changes in proteins, PCO formation is probably the first protein adduct generated due to free radical reaction with proteins [29]. Furthermore, it has been demonstrated that PCO level in homogenized liver and renal tissue significantly elevated in the rats orally treated with 3 g/kg APAP compared to control group [24]. It is also reported that oral administration of 3 g/kg APAP significantly increased PCO formation in homogenized liver tissue [4]. Results showed that PCO level elevated in the intoxicated rats with 2 g/kg of APAP compared to the control group. Nevertheless, this increase was not statistically significant. This was mainly due to using lower doses of APAP. Although WC extract (500 mg/kg) reduced PCO level in homogenized liver tissue and plasma, the difference was not statistically significant. However, this reduction suggests that WC extract may play as an antioxidant




Fig. 1. Effect of WC extract on GPx (A), SOD (B) and CAT (C) activity in the liver tissue of APAP intoxicated rats. APAP: acetaminophen; WC: 500 mg/kg of hydroalcoholic extract of watercress; S: 100 mg/kg of silymarin. ^astatistically significant compared to control group, P-value ≤ 0.05 . ^bstatistically significant compared to APAP group, P-value ≤ 0.05 .

Table 3

GC-MS components of the hydroalcoholic extract of WC.

NO	Compound	Molecular Formula	Percentage (%)
1	Benzenepropanenitrile	C ₉ H ₉ N	48.3
2	Phytol	C ₂₀ H ₄₀ O	10.1
3	α -Cadinene	C ₁₅ H ₂₄	9.5
4	Linolenic acid	C ₁₈ H ₃₀ O ₂	8.0
5	Palmitic acid	C ₁₆ H ₃₂ O ₂	7.1
6	3-Methyl-4-isopropylphenol	C ₁₀ H ₁₄ O	6.0
7	Durenol	C ₁₀ H ₁₄ O	2.5
8	Methyl ester group	-	2.4
9	Methyl 11,14,17-icosatrienoate	C ₂₁ H ₃₆ O ₂	2.0
10	Myristicin	C ₁₁ H ₁₂ O ₃	1.6
11	Thymol	C ₁₀ H ₁₄ O	1.5
Total			99.0

component by inhibiting the oxidation process in proteins.

All thiol groups are related to proteins in the plasma, play not only as sensitive indicators of oxidative stress but defense mechanism against ROS [26]. T-SH consists of two main parts: protein thiol groups (P-SH) and glutathione (GSH). Due to the negligible amount of GSH, there is a slight difference between T-SH and P-SH levels [30]. Current results are consistent with report that showed that T-SH level of hepatic tissue significantly decreased in APAP intoxicated rats compared to control group [31]. More importantly, NAQPI production as an active metabolite of APAP causes a reduction in thiol-associated proteins like GSH. Therefore, it leads to a decrease in the number of thiol groups in our experiment. Furthermore, T-SH content was increased in WC + APAP group in compared to APAP group. Altogether, administration of WC extract (500 mg/kg) increased T-SH content which was related to its high antioxidant properties as a free radical scavenger.

Antioxidant enzymes including SOD, GPx and CAT play as free radical scavengers [32]. In a research by Aycan *et al.* in the rats with APAP-induced hepatotoxicity, a decrease in GPx activity was seen [5]. Interestingly, it was reported that APAP administration significantly reduced SOD, GPx and CAT activities [33]. In this study, APAP administration showed a reduction in GPx activity compared to control group. Of note, treatment with WC increased enzyme activity in WC + APAP group. However, there was no significant difference in SOD and CAT activities in liver tissue among groups. In accordance with our results, it has been revealed that WC supplementation had no effect on these enzymes in healthy adults [12]. Free radicals, including superoxide anion, radical hydroxyl and H₂O₂ [34], are the major substrates for SOD and GPx antioxidant enzymes [5, 35]. After APAP overdose, GPx activity was reduced probably because of increasing in the formation of free radicals. The mechanism by which WC extract effects on antioxidant enzymes activity have not been fully understood. It is probably due to an increase in GPx activity by reducing hydroxyl radical formation.

5. Conclusion

In summary, it was for the first time that the effects of WC extract were evaluated on APAP-induced hepatotoxicity. The results showed that WC extract reduced oxidative stress by increasing T-SH content as well as enhancing GPx activity. It should be noted that the hepatoprotective effects of WC extract is probably because of its antioxidant activity as a free radical scavenger. More investigations are needed to suggest WC extract as an efficient supplementation to treat APAP-induced hepatotoxicity.

Declarations

Author contribution statement

Nahid Azarmehr: Performed the experiments; Contributed reagents, materials, analysis tools or data.

Parisa Afshar: Performed the experiments.

Mona Moradi, Kazem Abbaszadeh-Goudarzi: Analyzed and interpreted the data; Wrote the paper.

Heibatollah Sadeghi: Conceived and designed the experiments.

Hossein Sadeghi: Conceived and designed the experiments; Contributed reagents, materials, analysis tools or data.

Behnam Alipoor: Conceived and designed the experiments; Analyzed and interpreted the data.

Bahman Khalvati: Contributed reagents, materials, analysis tools or data.

Zahra Barmoudeh: Performed the experiments; Contributed reagents, materials, analysis tools or data.

Amir Hossein Doustimotlagh: Conceived and designed the experiments; Analyzed and interpreted the data; Wrote the paper.

Funding statement

This work is supported by grant number 960283 from by council of research of Yasuj University of Medical Sciences, Iran.

Competing interest statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

- G.S.B. Aseervatham, T. Sivasudha, J. Sasikumar, P.H. Christabel, R. Jeyadevi, D.A. Ananth, Antioxidant and hepatoprotective potential of *Pouteria campechiana* on acetaminophen-induced hepatic toxicity in rats, *J. Physiol. Biochem.* 70 (1) (2014) 1–14.
- A. Omid, N. Riahinia, M.B.M. Torbati, M.-A. Behdani, Hepatoprotective effect of *Crocus sativus* (saffron) petals extract against acetaminophen toxicity in male Wistar rats, *Avicenna J. Phytomed.* 4 (5) (2014) 330.
- M. Blieden, L.C. Paramore, D. Shah, R. Ben-Joseph, A perspective on the epidemiology of acetaminophen exposure and toxicity in the United States, *Expert Rev. Clin. Pharmacol.* 7 (3) (2014) 341–348.
- M. Cristani, A. Speciale, F. Mancari, T. Arcoraci, D. Ferrari, D. Fratanantonio, et al., Protective activity of an anthocyanin-rich extract from bilberries and blackcurrants on acute acetaminophen-induced hepatotoxicity in rats, *Nat. Prod. Res.* 30 (24) (2016) 2845–2849.
- İ.Ö. Aycan, A. Tüfek, O. Tokgöz, O. Evliyaoglu, U. Firat, G.Ö. Kavak, et al., Thymoquinone treatment against acetaminophen-induced hepatotoxicity in rats, *Int. J. Surg.* 12 (3) (2014) 213–218.
- A.H. Tezcan, O. Ozturk, S. Ustebay, Y. Adali, H. Yagmurdu, The beneficial effects of ozone therapy in acetaminophen-induced hepatotoxicity in mice, *Pharmacol. Rep.* 70 (2) (2018) 340–345.
- J.Y. Lee, S.H. Lee, H.J. Kim, J.-M. Ha, S.-H. Lee, J.-H. Lee, et al., The preventive inhibition of chondroitin sulfate against the CCl₄-Induced oxidative stress of subcellular level, *Arch. Pharm. Res.* 27 (3) (2004) 340.
- A. Geetha, L. Priya, S.A. Jeyachristy, R. Surendran, Level of oxidative stress in the red blood cells of patients with liver cirrhosis, *Indian J. Med. Res.* 126 (3) (2007) 204–211.
- S. Bahramikia, R. Yazdanparast, Antioxidant efficacy of *Nasturtium officinale* extracts using various in vitro assay systems, *J. Acupunct. Meridian Stud.* 3 (4) (2010) 283–290.
- N.A. Casanova, M.F. Simoniello, M.M. López Nigro, M.A. Carballo, Modulator effect of watercress against cyclophosphamide-induced oxidative stress in mice, *Medicina (Buenos Aires)* 77 (2017) 201–206.
- T. Ozen, Investigation of antioxidant properties of *Nasturtium officinale* (watercress) leaf extracts, *Acta Pol. Pharm.* 66 (2) (2009) 187–193.
- C.I. Gill, S. Haldar, L.A. Boyd, R. Bennett, J. Whiteford, M. Butler, et al., Watercress supplementation in diet reduces lymphocyte DNA damage and alters blood antioxidant status in healthy adults, *Am. J. Clin. Nutr.* 85 (2) (2007) 504–510.
- J.M. Pullar, S.J. Thomson, M.J. King, C.I. Turnbull, R.G. Midwinter, M.B. Hampton, The chemopreventive agent phenethyl isothiocyanate sensitizes cells to Fas-mediated apoptosis, *Carcinogenesis* 25 (5) (2004) 765–772.
- R. Yazdanparast, S. Bahramikia, A. Ardestani, *Nasturtium officinale* reduces oxidative stress and enhances antioxidant capacity in hypercholesterolaemic rats, *Chem. Biol. Interact.* 172 (3) (2008) 176–184.
- A. Hanafy, H.M. Aldawsari, J.M. Badr, A.K. Ibrahim, S.E.-S. Abdel-Hady, Evaluation of hepatoprotective activity of *Adansonia digitata* extract on acetaminophen-induced hepatotoxicity in rats, *Evid. Based Complement Altern. Med.* 2016 (2016).
- M.A. Dkhil, A.E. Abdel Moneim, *Myristica fragrans* kernels prevent paracetamol-induced hepatotoxicity by inducing anti-apoptotic genes and Nrf2/HO-1 pathway, *Int. J. Mol. Sci.* 20 (4) (2019) 1–15.
- I.F. Benzie, J. Strain, The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay, *Anal. Biochem.* 239 (1) (1996) 70–76.
- G. Drochizoiu, C.I. Ciobanu, S. Bancela, L. Ion, B.A. Petre, C. Andries, et al., Ultrasound-based protein determination in maize seeds, *Ultrason. Sonochem.* 29 (2016) 93–103.
- F.S. Kholari, A.R. Dehpour, M. Nourbakhsh, A.H. Doustimotlagh, M. Bagherieh, A. Golestani, Erythrocytes membrane alterations reflecting liver damage in CCl₄-Induced cirrhotic rats: the ameliorative effect of naltrexone, *Acta Med. Iran.* 54 (10) (2016) 631–639.
- A.H. Doustimotlagh, A.R. Dehpour, M. Nourbakhsh, A. Golestani, Alteration in membrane protein, antioxidant status and hexokinase activity in erythrocytes of CCl₄ sub 4-induced cirrhotic rats, *Acta Med. Iran.* 52 (11) (2014) 795.
- A. Asfaram, M. Arabi, A. Ostovan, H. Sadeghi, M. Ghaedi, Simple and selective detection of quercetin in extracts of plants and food samples by dispersive-micro-solid phase extraction based on core-shell magnetic molecularly imprinted polymers, *New J. Chem.* 42 (19) (2018) 16144–16153.
- A.-Y. Wang, L.-H. Lian, Y.-Z. Jiang, Y.-L. Wu, J.-X. Nan, *Gentiana manshurica* Kitagawa prevents acetaminophen-induced acute hepatic injury in mice via inhibiting JNK/ERK MAPK pathway, *World J. Gastroenterol.* 16 (3) (2010) 384.
- R. Lucarini, W.A. Bernardes, M.G. Tozatti, A.A. da Silva Filho, M.L.A. e Silva, C. Momo, et al., Hepatoprotective effect of *Rosmarinus officinalis* and rosmarinic acid on acetaminophen-induced liver damage, *Emir. J. Food Agric.* 26 (10) (2014) 878–885.
- H.-T. Yao, Y.-C. Yang, C.-H. Chang, H.-T. Yang, M.-C. Yin, Protective effects of (-)-epigallocatechin-3-gallate against acetaminophen-induced liver injury in rats, *Biomedicine* 5 (3) (2015).
- S.K. Ramaiah, A toxicologist guide to the diagnostic interpretation of hepatic biochemical parameters, *Food Chem. Toxicol.* 45 (9) (2007) 1551–1557.
- M. Karami, A. Nosrati, M. Naderi, M. Makhloogh, S. Shahani, Protective effects of *Nasturtium officinale* against gamma-irradiation-induced hepatotoxicity in C57 mice, *Res. J. Pharmacogn.* 2 (2) (2015) 19–25.
- H. Rabiul, M. Subhasish, S. Sinha, M.G. Roy, D. Sinha, S. Gupta, Hepatoprotective activity of *Clerodendron inerme* against paracetamol induced hepatic injury in rats for pharmaceutical product, *Int. J. Drug Dev. Res.* 3 (1) (2011) 118–126.
- K. Mahaldar, A. Hossain, F. Islam, S. Islam, M.A. Islam, M. Shahriar, et al., Antioxidant and hepatoprotective activity of *Piper retrofractum* against Paracetamol-induced hepatotoxicity in Sprague-Dawley rat, *Nat. Prod. Res.* (2019) 1–7.
- H. Sadeghi, F. Jahanbazi, H. Sadeghi, N. Omidifar, B. Alipoor, E.P. Kokhdan, et al., Metformin attenuates oxidative stress and liver damage after bile duct ligation in rats, *Res. Pharmaceut. Sci.* 14 (2) (2019) 122–129.
- M.-L. Hu, Measurement of protein thiol groups and glutathione in plasma, *Methods Enzymol.* (1994) 380–385.
- R. Kabirifar, F. Safari, A. Karimollah, A. Moradi, E. Eskandari-nasab, Quercetin protects liver injury induced by bile duct ligation via attenuation of Rac1 and NADPH oxidase1 expression in rats, *Hepatobiliary Pancreat. Dis. Int.* 16 (1) (2017) 88–95.
- O. Bayrak, N. Bavbek, O.F. Karatas, R. Bayrak, F. Catal, E. Cimentep, et al., *Nigella sativa* protects against ischaemia/reperfusion injury in rat kidneys, *Nephrol. Dial. Transplant.* 23 (7) (2008) 2206–2212.
- M. Aydogan, A. Polat, N. Vardi, M. Erdogan, A. Yucel, A. Yildiz, et al., Protective effects of melatonin and and #914;-D-Glucan against acetaminophen toxicity in rats, *Med. Sci. Int. Med. J.* 5 (2) (2016) 539.
- R. Fukuhara, T. Kageyama, Structure, gene expression, and evolution of primate glutathione peroxidases, *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 141 (4) (2005) 428–436.
- E.K. Vieira, S. Bona, F.C. Di Naso, M. Porawski, J. Tieppo, N.P. Marroni, Quercetin treatment ameliorates systemic oxidative stress in cirrhotic rats, *ISRN Gastroenterol.* 2011 (2011) 604071.